FIRST AID FOR THE®

USMLE® STEP 1

2020

30th ANNIVERSARY EDITION

SMTEBOOKS
Ketabton.com

Time-proven blueprint for Step 1 success

1,300+ must-know concepts with many new high-yield facts

1,200+ color photos and illustrations, expanded and revised

Student-proven exam strategies designed to boost your score

Mc Graw Hill

TAO LE ■ VIKAS BHUSHAN ■ MATTHEW SOCHAT ■ VAISHNAVI VAIDYANATHAN

(c) ketabton.com: The Digital Library

FIRST AID FOR THE®

USMLE STEP 1 2020

TAO LE, MD, MHS
Founder, ScholarRx
Associate Clinical Professor, Department of Medicine
University of Louisville School of Medicine

VIKAS BHUSHAN, MD
Boracay

MATTHEW SOCHAT, MD
Fellow, Department of Hematology/Oncology
St. Louis University School of Medicine

VAISHNAVI VAIDYANATHAN, MD
Resident, Department of Pediatric Neurology
Barrow Neurological Institute at Phoenix Children's Hospital

SARAH SCHIMANSKY, MB BCh BAO
Resident, Department of Ophthalmology
Royal United Hospitals Bath

JORDAN ABRAMS
St. George's University School of Medicine
Class of 2020

KIMBERLY KALLIANOS, MD
Assistant Professor, Department of Radiology and Biomedical Imaging
University of California, San Francisco School of Medicine

SMTEBOOKS

Mc Graw Hill

New York / Chicago / San Francisco / Athens / London / Madrid / Mexico City
Milan / New Delhi / Singapore / Sydney / Toronto

(c) ketabton.com: The Digital Library

Copyright © 2020 by Tao Le and Vikas Bhushan. All rights reserved. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written permission of the publisher.

ISBN: 978-1-26-046205-0
MHID: 1-26-046205-6

The material in this eBook also appears in the print version of this title: ISBN: 978-1-26-046204-3,
MHID: 1-26-046204-8.

eBook conversion by codeMantra
Version 1.0

All trademarks are trademarks of their respective owners. Rather than put a trademark symbol after every occurrence of a trademarked name, we use names in an editorial fashion only, and to the benefit of the trademark owner, with no intention of infringement of the trademark. Where such designations appear in this book, they have been printed with initial caps.

McGraw-Hill Education eBooks are available at special quantity discounts to use as premiums and sales promotions or for use in corporate training programs. To contact a representative, please visit the Contact Us page at www.mhprofessional.com.

Notice

Medicine is an ever-changing science. As new research and clinical experience broaden our knowledge, changes in treatment and drug therapy are required. The authors and the publisher of this work have checked with sources believed to be reliable in their efforts to provide information that is complete and generally in accord with the standards accepted at the time of publication. However, in view of the possibility of human error or changes in medical sciences, neither the authors nor the publisher nor any other party who has been involved in the preparation or publication of this work warrants that the information contained herein is in every respect accurate or complete, and they disclaim all responsibility for any errors or omissions or for the results obtained from use of the information contained in this work. Readers are encouraged to confirm the information contained herein with other sources. For example and in particular, readers are advised to check the product information sheet included in the package of each drug they plan to administer to be certain that the information contained in this work is accurate and that changes have not been made in the recommended dose or in the contraindications for administration. This recommendation is of particular importance in connection with new or infrequently used drugs.

TERMS OF USE

This is a copyrighted work and McGraw-Hill Education and its licensors reserve all rights in and to the work. Use of this work is subject to these terms. Except as permitted under the Copyright Act of 1976 and the right to store and retrieve one copy of the work, you may not decompile, disassemble, reverse engineer, reproduce, modify, create derivative works based upon, transmit, distribute, disseminate, sell, publish or sublicense the work or any part of it without McGraw-Hill Education's prior consent. You may use the work for your own noncommercial and personal use; any other use of the work is strictly prohibited. Your right to use the work may be terminated if you fail to comply with these terms.

THE WORK IS PROVIDED "AS IS." McGRAW-HILL EDUCATION AND ITS LICENSORS MAKE NO GUARANTEES OR WARRANTIES AS TO THE ACCURACY, ADEQUACY OR COMPLETENESS OF OR RESULTS TO BE OBTAINED FROM USING THE WORK, INCLUDING ANY INFORMATION THAT CAN BE ACCESSED THROUGH THE WORK VIA HYPERLINK OR OTHERWISE, AND EXPRESSLY DISCLAIM ANY WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. McGraw-Hill Education and its licensors do not warrant or guarantee that the functions contained in the work will meet your requirements or that its operation will be uninterrupted or error free. Neither McGraw-Hill Education nor its licensors shall be liable to you or anyone else for any inaccuracy, error or omission, regardless of cause, in the work or for any damages resulting therefrom. McGraw-Hill Education has no responsibility for the content of any information accessed through the work. Under no circumstances shall McGraw-Hill Education and/or its licensors be liable for any indirect, incidental, special, punitive, consequential or similar damages that result from the use of or inability to use the work, even if any of them has been advised of the possibility of such damages. This limitation of liability shall apply to any claim or cause whatsoever whether such claim or cause arises in contract, tort or otherwise.

(c) ketabton.com: The Digital Library

Dedication

To the contributors to this and past editions, who took time to share their knowledge, insight, and humor for the benefit of students and physicians everywhere.

(c) ketabton.com: The Digital Library

This page intentionally left blank

(c) ketabton.com: The Digital Library

Contents

Contributing Authors	vii	General Acknowledgments	xv
Associate Authors	viii	How to Contribute	xvii
Faculty Advisors	ix	How to Use This Book	xix
Thirtieth Anniversary Foreword	xi	Selected USMLE Laboratory Values	xx
Preface	xiii	First Aid Checklist for the USMLE Step 1	xxii
Special Acknowledgments	xiv		

SECTION I GUIDE TO EFFICIENT EXAM PREPARATION 1

Introduction	2	Test-Taking Strategies	22
USMLE Step 1—The Basics	2	Clinical Vignette Strategies	23
Defining Your Goal	12	If You Think You Failed	24
Learning Strategies	13	Testing Agencies	24
Timeline for Study	16	References	25
Study Materials	20		

SECTION I SUPPLEMENT SPECIAL SITUATIONS 27

SECTION II HIGH-YIELD GENERAL PRINCIPLES 29

How to Use the Database	30	Pathology	205
Biochemistry	33	Pharmacology	229
Immunology	95	Public Health Sciences	255
Microbiology	123		

FAS1_2019_00_Frontmatter.indd 5 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

SECTION III HIGH-YIELD ORGAN SYSTEMS 275

Approaching the Organ Systems 276
Cardiovascular 279
Endocrine 325
Gastrointestinal 357
Hematology and Oncology 403
Musculoskeletal, Skin, and Connective Tissue 445
Neurology and Special Senses 489
Psychiatry 553
Renal 577
Reproductive 611
Respiratory 659
Rapid Review 689

SECTION IV TOP-RATED REVIEW RESOURCES 711

How to Use the Database 712
Question Banks and Books 714
Web and Mobile Apps 714
Comprehensive 715
Anatomy, Embryology, and Neuroscience 715
Behavioral Science 716
Biochemistry 716
Cell Biology and Histology 716
Microbiology and Immunology 717
Pathology 717
Pharmacology 718
Physiology 718

Abbreviations and Symbols 719
Image Acknowledgments 727
Index 749
About the Editors 808

FAS1_2019_00_Frontmatter.indd 6 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Contributing Authors

MAJED H. ALGHAMDI, MBBS
Resident, Joint Program of Preventive Medicine
Jeddah, Saudi Arabia

LILIT ASLANYAN
New York Institute of Technology College of Osteopathic Medicine
Class of 2020

HUMOOD BOQAMBAR, MB BCh BAO
Assistant Registrar, Department of Orthopaedic Surgery
Farwaniya Hospital

WEELIC CHONG
Sidney Kimmel Medical College at Thomas Jefferson University
MD/PhD Candidate

KRISTINA DAMISCH
University of Iowa Carver College of Medicine
Class of 2020

YUMI KOVIC, MD
Resident, Department of Psychiatry
University of Massachusetts Medical School

KAITLYN MELNICK, MD
Resident, Department of Neurological Surgery
University of Florida College of Medicine, Gainesville

MARY KATHERINE MONTES de OCA, MD
Resident, Department of Obstetrics and Gynecology
Duke University Hospital

SCOTT MOORE, DO
Assistant Professor of Medical Laboratory Sciences
Weber State University

VASILY OVECHKO, MD
Resident, Department of Surgery
Russian Medical Academy of Continuous Professional Education

VIVEK PODDER
MBBS Student
Tairunnessa Memorial Medical College and Hospital, Bangladesh

CONNIE QIU
Lewis Katz School of Medicine at Temple University
MD/PhD Candidate

Image and Illustration Team

CAROLINE COLEMAN
Emory University School of Medicine
Class of 2020

MATTHEW HO ZHI GUANG
University College Dublin (MD), DFCI (PhD)
MD/PhD Candidate

VICTOR JOSE MARTINEZ LEON, MD
Central University of Venezuela

ALIREZA ZANDIFAR, MD
Research Fellow
Isfahan University of Medical Sciences, Iran

FAS1_2019_00_Frontmatter.indd 7 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Associate Authors

HUZAIFA AHMAD, MD
Resident, Department of Medicine
Georgetown University Hospital/MedStar Washington Hospital Center

ALEXANDER R. ASLESEN
Kirksville College of Osteopathic Medicine
Class of 2020

ANUP K. BHATTACHARYA, MD
Resident, Mallinckrodt Institute of Radiology
Washington University School of Medicine

ANUP CHALISE, MBBS
Resident, Department of General Surgery
Nepal Medical College and Teaching Hospital

ASHTEN R. DUNCAN, MPH
University of Oklahoma-Tulsa School of Community Medicine
Class of 2021

SARINA KOILPILLAI
St. George's University School of Medicine
Class of 2020

LAUREN N. LESSOR, MPH, MD
Resident, Department of Pediatrics
Mercy Health – St. Vincent Medical Center

ROHAN BIR SINGH, MD
Fellow, Department of Ophthalmology
Massachusetts Eye and Ear
Harvard Medical School

Image and Illustration Team

YAMNA JADOON, MD
Research Associate
Aga Khan University

DANA M. JORGENSON
Chicago College of Osteopathic Medicine
Class of 2020

MITCHELL A. KATONA
University of Texas Health Science Center, Long School of Medicine
Class of 2020

TAYLOR MANEY, MD
Resident, Department of Anesthesiology
Brigham and Women's Hospital

FAS1_2019_00_Frontmatter.indd 8 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Faculty Advisors

DIANA ALBA, MD
Clinical Instructor
University of California, San Francisco School of Medicine

MARK A.W. ANDREWS, PhD
Professor of Physiology
Lake Erie College of Osteopathic Medicine at Seton Hill

MARIA ANTONELLI, MD
Assistant Professor, Division of Rheumatology
MetroHealth Medical Center, Case Western Reserve University

HERMAN SINGH BAGGA, MD
Urologist, Allegheny Health Network
University of Pittsburgh Medical Center Passavant

SHIN C. BEH, MD
Assistant Professor, Department of Neurology & Neurotherapeutics
UT Southwestern Medical Center at Dallas

JOHN R. BUTTERLY, MD
Professor of Medicine
Dartmouth Geisel School of Medicine

SHELDON CAMPBELL, MD, PhD
Professor of Laboratory Medicine
Yale School of Medicine

BROOKS D. CASH, MD
Professor of Medicine, Division of Gastroenterology
University of South Alabama School of Medicine

SHIVANI VERMA CHMURA, MD
Adjunct Clinical Faculty, Department of Psychiatry
Stanford University School of Medicine

BRADLEY COLE, MD
Assistant Professor of Basic Sciences
Loma Linda University School of Medicine

LINDA S. COSTANZO, PhD
Professor, Physiology & Biophysics
Virginia Commonwealth University School of Medicine

MANAS DAS, MD, MS
Director, Clinical Anatomy, Embryology, and Histology
University of Massachusetts Medical School

ANTHONY L. DeFRANCO, PhD
Professor, Department of Microbiology and Immunology
University of California, San Francisco School of Medicine

CHARLES S. DELA CRUZ, MD, PhD
Associate Professor, Department of Pulmonary and Critical Care Medicine
Yale School of Medicine

SAKINA FARHAT, MD
Consulting Gastroenterologist
State University of New York Downstate Medical Center

CONRAD FISCHER, MD
Associate Professor, Medicine, Physiology, and Pharmacology
Touro College of Medicine

RAYUDU GOPALAKRISHNA, PhD
Associate Professor, Department of Integrative Anatomical Sciences
Keck School of Medicine of University of Southern California

RYAN C.W. HALL, MD
Assistant Professor, Department of Psychiatry
University of South Florida School of Medicine

LOUISE HAWLEY, PhD
Immediate Past Professor and Chair, Department of Microbiology
Ross University School of Medicine

JEFFREY W. HOFMANN, MD, PhD
Resident, Department of Pathology
University of California, San Francisco School of Medicine

CLARK KEBODEAUX, PharmD
Clinical Assistant Professor, Pharmacy Practice and Science
University of Kentucky College of Pharmacy

KRISTINE KRAFTS, MD
Assistant Professor, Department of Basic Sciences
University of Minnesota School of Medicine

MATTHEW KRAYBILL, PhD
Clinical Neuropsychologist
Cottage Health, Santa Barbara, California

GERALD LEE, MD
Assistant Professor, Departments of Pediatrics and Medicine
Emory University School of Medicine

FAS1_2019_00_Frontmatter.indd 9 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

KACHIU C. LEE, MD, MPH
Assistant Clinical Professor, Department of Dermatology
The Warren Alpert Medical School of Brown University

WARREN LEVINSON, MD, PhD
Professor, Department of Microbiology and Immunology
University of California, San Francisco School of Medicine

JAMES LYONS, MD
Professor of Pathology and Family Medicine
Alabama College of Osteopathic Medicine

PETER MARKS, MD, PhD
Center for Biologics Evaluation and Research
US Food and Drug Administration

DOUGLAS A. MATA, MD, MPH
Brigham Education Institute and Brigham and Women's Hospital
Harvard Medical School

VICKI M. PARK, PhD, MS
Assistant Dean
University of Tennessee College of Medicine

SOROUSH RAIS-BAHRAMI, MD
Assistant Professor, Departments of Urology and Radiology
University of Alabama at Birmingham School of Medicine

SASAN SAKIANI, MD
Fellow, Transplant Hepatology
Cleveland Clinic

MELANIE SCHORR, MD
Assistant in Medicine
Massachusetts General Hospital

SHIREEN MADANI SIMS, MD
Chief, Division of Gynecology, Gynecologic Surgery, and Obstetrics
University of Florida School of Medicine

NATHAN W. SKELLEY, MD
Assistant Professor, Department of Orthopaedic Surgery
University of Missouri, The Missouri Orthopaedic Institute

HOWARD M. STEINMAN, PhD
Assistant Dean, Biomedical Science Education
Albert Einstein College of Medicine

SUPORN SUKPRAPRUT-BRAATEN, PhD
Director of Research, Graduate Medical Education
Unity Health, Searcy, Arkansas

RICHARD P. USATINE, MD
Professor, Dermatology and Cutaneous Surgery
University of Texas Health Science Center San Antonio

J. MATTHEW VELKEY, PhD
Assistant Dean, Basic Science Education
Duke University School of Medicine

TISHA WANG, MD
Associate Clinical Professor, Department of Medicine
David Geffen School of Medicine at UCLA

SYLVIA WASSERTHEIL-SMOLLER, PhD
Professor Emerita, Department of Epidemiology and Population Health
Albert Einstein College of Medicine

ADAM WEINSTEIN, MD
Assistant Professor, Pediatric Nephrology and Medical Education
Geisel School of Medicine at Dartmouth

ABHISHEK YADAV, MBBS, MSc
Associate Professor of Anatomy
Geisinger Commonwealth School of Medicine

KRISTAL YOUNG, MD
Clinical Instructor, Department of Cardiology
Huntington Hospital, Pasadena, California

FAS1_2019_00_Frontmatter.indd 10 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Thirtieth Anniversary Foreword

Our exam experiences remain vivid in our minds to this day as we reflect on 30 years of *First Aid*. In 1989, our big idea was to cobble together a "quick and dirty" study guide so that we would never again have to deal with the USMLE Step 1. We passed, but in a Faustian twist, we now relive the exam yearly while preparing each new edition.

Like all students before us, we noticed that certain topics tended to appear frequently on examinations. So we compulsively bought and rated review books and pored through a mind-numbing number of "recall" questions, distilling each into short facts. We had a love-hate relationship with mnemonics. They went against our purist desires for conceptual knowledge, but remained the best way to absorb the vocabulary and near-random associations that unlocked questions and eponyms.

To pull it all together, we used a then "state-of-the-art" computer database (Paradox/MS DOS 4) that fortuitously limited our entries to 256 characters. That length constraint (which predated Twitter by nearly two decades) imposed extreme brevity. The three-column layout created structure—and this was the blueprint upon which *First Aid* was founded.

The printed, three-column database was first distributed in 1989 at the University of California, San Francisco. The next year, the official first edition was self-published under the title *High-Yield Basic Science Boards Review: A Student-to-Student Guide*. The following year, our new publisher dismissed the *High-Yield* title as too confusing and came up with *First Aid for the Boards*. We thought the name was a bit cheesy, but it proved memorable. Interestingly, our "High-Yield" name resurfaced years later as the title of a competing board review series.

We lived in San Francisco and Los Angeles during medical school and residency. It was before the Web, and before med students could afford cell phones and laptops, so we relied on AOL e-mail and bulky desktops. One of us would drive down to the other person's place for multiple weekends of frenetic revisions fueled by triple-Swiss white chocolate lattes from the Coffee Bean & Tea Leaf, with R.E.M. and the Nusrat Fateh Ali Khan playing in the background. Everything was marked up on 11- by 17-inch "tearsheets," and at the end of the marathon weekend we would converge at the local 24-hour Kinko's followed by the FedEx box near LAX (10 years before these two great institutions merged). These days we work with our online collaborative platform A.nnotate, GoToMeeting, and ubiquitous broadband Internet, and sadly, we rarely get to see each other.

What hasn't changed, however, is the collaborative nature of the book. Thousands of authors, editors, and contributors have enriched our lives and made this book possible. Most helped for a year or two and moved on, but a few, like Ted Hon, Chirag Amin, and Andi Fellows, made lasting contributions. Like the very first edition, the team is always led by student authors who live and breathe (and fear) the exam, not professors years away from that reality.

We're proud of the precedent that *First Aid* set for the many excellent student-to-student publications that followed. More importantly, *First Aid* itself owes its success to the global community of medical students and international medical graduates (IMGs) who each year contribute ideas, suggestions, and new content. In the early days, we used book coupons and tear-out business reply mail forms. These days, we get many thousands of comments and suggestions each year via our blog FirstAidTeam.com and A.nnotate.

FAS1_2019_00_Frontmatter.indd 11 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

At the end of the day, we don't take any of this for granted. Students are expected to synthesize an ever increasing amount of information, and we have a bigger challenge ahead of us to try to keep *First Aid* indispensable to students and IMGs. We want and need your participation in the *First Aid* community. (See How to Contribute, p. xvii.) With your help, we hope editing *First Aid* will continue to be just as fun and rewarding as the past 30 years have been.

Louisville Tao Le
Boracay Vikas Bhushan

First Aid for the USMLE Step 1 Through the Years















FAS1_2019_00_Frontmatter.indd 12 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Preface

With the 30th edition of *First Aid for the USMLE Step 1*, we continue our commitment to providing students with the most useful and up-to-date preparation guide for the USMLE Step 1. This edition represents an outstanding revision in many ways, including:

- 50 entirely new or heavily revised high-yield topics reflecting evolving trends in the USMLE Step 1.
- Reorganization of high-yield topics in Pharmacology, Endocrine, and Reproductive chapters for improved study.
- Extensive text revisions, new mnemonics, clarifications, and corrections curated by a team of more than 30 medical student and resident physician authors who excelled on their Step 1 examinations and verified by a team of expert faculty advisors and nationally recognized USMLE instructors.
- Updated with 178 new and revised diagrams and illustrations as part of our ongoing collaboration with USMLE-Rx and ScholarRx (MedIQ Learning, LLC).
- Updated with 75 new and revised photos to help visualize various disorders, descriptive findings, and basic science concepts. Additionally, revised imaging photos have been labeled and optimized to show both normal anatomy and pathologic findings.
- Updated study tips on the opening page of each chapter.
- Improved integration of clinical images and illustrations to better reinforce and learn key anatomic concepts.
- Improved organization and integration of text, illustrations, clinical images, and tables throughout for focused review of high-yield topics.
- Revised and expanded ratings of current, high-yield review resources, with clear explanations of their relevance to USMLE review.
- Real-time Step 1 updates and corrections can be found exclusively on our blog, www.firstaidteam.com.

We invite students and faculty to share their thoughts and ideas to help us continually improve *First Aid for the USMLE Step 1* through our blog and collaborative editorial platform. (See How to Contribute, p. xvii.)

Louisville	Tao Le
Boracay	Vikas Bhushan
St. Louis	Matthew Sochat
Phoenix	Vaishnavi Vaidyanathan
Bristol	Sarah Schimansky
New York City	Jordan Abrams
San Francisco	Kimberly Kallianos

FAS1_2019_00_Frontmatter.indd 13 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Special Acknowledgments

This has been a collaborative project from the start. We gratefully acknowledge the thousands of thoughtful comments, corrections, and advice of the many medical students, international medical graduates, and faculty who have supported the authors in our continuing development of *First Aid for the USMLE Step 1*.

For support and encouragement throughout the process, we are grateful to Thao Pham, Jinky Flang, and Jonathan Kirsch, Esq. Thanks to Louise Petersen for organizing and supporting the project. Thanks to our publisher, McGraw-Hill, for the valuable assistance of its staff, including Bob Boehringer, Jeffrey Herzich, and Christina Thomas.

We are also very grateful to Dr. Fred Howell and Dr. Robert Cannon of Textensor Ltd for providing us extensive customization and support for their powerful Annotate.co collaborative editing platform (www.annotate.co), which allows us to efficiently manage thousands of contributions. Thanks to Dr. Richard Usatine and Dr. Kristine Krafts for their outstanding image contributions. Thanks also to Jean-Christophe Fournet (www.humpath.com), Dr. Ed Uthman, and Dr. Frank Gaillard (www.radiopaedia.org) for generously allowing us to access some of their striking photographs.

For exceptional editorial leadership, enormous thanks to Kathleen Naylor, Christine Diedrich and Emma Underdown. Thank you to our USMLE-Rx/ScholarRx team of editors, Jessie Schanzle, Ruth Kaufman, Janene Matragrano, Susan Mazik, Isabel Nogueira, Sharon Prevost, Jen Shimony, and Hannah Warnshuis. Special thanks to our indexer Dr. Anne Fifer. We are also grateful to our medical illustrator, Hans Neuhart, for his creative work on the new and updated illustrations. Lastly, tremendous thanks to Graphic World, especially Anne Banning, Sandy Brown, Gary Clark, and Cindy Geiss.

Louisville	Tao Le
Boracay	Vikas Bhushan
St. Louis	Matthew Sochat
Phoenix	Vaishnavi Vaidyanathan
Bristol	Sarah Schimansky
New York City	Jordan Abrams
San Francisco	Kimberly Kallianos

FAS1_2019_00_Frontmatter.indd 14 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

General Acknowledgments

Each year we are fortunate to receive the input of thousands of medical students and graduates who provide new material, clarifications, and potential corrections through our website and our collaborative editing platform. This has been a tremendous help in clarifying difficult concepts, correcting errata from the previous edition, and minimizing new errata during the revision of the current edition. This reflects our long-standing vision of a true, student-to-student publication. We have done our best to thank each person individually below, but we recognize that errors and omissions are likely. Therefore, we will post an updated list of acknowledgments at our website, www.firstaidteam.com/bonus/. We will gladly make corrections if they are brought to our attention.

For submitting contributions and corrections, many thanks to Raed Ababneh, Antara Afrin, Rasim Agaev, Vanya Aggarwal, Ataa Ahmed, Hasan Alarouri, Basim Ali, Muhammad Faizan Ali, Moatasem Al-Janabi, Mohamed Almahmodi, Chima Amadi, Arman Amin, Jacqueline Aredo, Ranya Baddourah, Daniel Badin, Nida Bajwa, Dileni Bandarage, Jerrin Bawa, Esra Bayram, Craig Beavers, Jacqueline Bekhit, Matthias Bergmann, Stephanie Biecker, Aaron Birnbaum, Prateek Bommu, Nathaniel Borochov, Susan Brands, Olivia W. Brooks, Meghan Brown, Stanley Budzinski, Kevin Budziszewski, Pavel Burski, Elisa M. Cairns, Sergio Camba, Katie Carsky, Esteban Casasola, Marielys Castro, Jesse Chait, Bliss Chang, Santosh Cherian, Heewon Choi, Charilaos Chourpiliadis, Maruf Chowdhury, Matthew J. Christensen, Matthew Yat Hon Chung, Alexander Ciaramella, Dillon Clancy, Sofija Conic, M. Marwan Dabbagh, Parag Das, Ketan Dayma, Elmer De Camps, Charles de Leeuw, Xavier De Pena, Christopher DeAngelo, Elliott Delgado, Anthony DeMarinis, Stacy Diaz, Evan Dishion, Nicola Helen Duzak, Emily Edwards, Alec Egan, Mohamed Elashwal, Osama El-Gabalawy, Matthew Eli, Awab Elnaeem, Sally El Sammak, Dylan Erwin, Stephanie Estevez-Marin, Gray Evans, Najat Fadlallah, Aria Fariborzi, Richard Ferro, Adam Fletcher, Kimberly A. Foley, Kyle Fratta, Samantha Friday, Nikhila Gandrakota, Siva Garapati, Nicolas Curi Gawlinski, Joanna Georgakas, Beth Anne George, Ashley Ghaemi, E. Sophia Gonzalez, Justin Graff, Gabriel Graham, Donovan Griggs, David Gruen, Gursewak Hadday, Jacqueline Hairston, Hunter Harrison, Gull Shahmir Hasnat, Maximillan Hawkins, Grecia Haymee, Briana Hernandez, Robin Hilder, Tammy Hua, Derrek Humphries, Audrey Hunt, Nanki Hura, Danny Ibrahim, Jyothik Varun Inampudi, Hnin Ingyin, Maham Irfan, Mina Iskandar, Kritika Iyer, Christina Jacobs, Arpit Jain, Neil K. Jain, Ala Jamal, Natalie Jansen, Jordan Jay, Mohammad Jmasi, Colton Junod, Talia Kamdjou, Filip Kaniski, Lydia Kaoutzani, Panagiotis Kaparaliotis, Srikrishna Karnatapu, Patrick Keller, Olivia Keller-Baruch, Cameron Kerl, Ahmed Ali Khan, Sara Khan, Shaima Khandaker, Samir Khouzam, Sonya Klein, Elana Kleinman, Andrew Ko, Soheil Kooraki, Anna Kukharchuk, Dennis Vu Kulp, Anil A. Kumar, Julie Kurek, Chloe Lahoud, Mike Lawandy, Ramy Lawandy, Jessica Lazar, Andrea Leal-Lopez, Lynda Lee, Chime Lhatso, Christine Lin, Benjamin Lodge, Soon Khai Low, Estefanía Henríquez Luthje, Lisa-Qiao MacDonald, Divya Madhavarapu, Mahir Mameledzija, Keerer Mann, Rajver Mann, Nadeen Mansour, Yusra Mansour, Bridget Martinez, Ahmad Mashlah, Rick Mathews, Amy McGregor, Alexandra & Joshua Medeiros & Fowler, Viviana Medina, Areeka Memon, Pedro G. R. Menicucci, Ben Meyers, Stephan A. Miller, Fatima Mirza, Murli Mishra, Elana Molcho, Guarina Molina, John Moon, Nayla Mroueh, Neha Mylarapu, Behnam Nabavizadeh, Moeko Nagatsuka, Ghazal Naghibzadeh, Alice Nassar, Nadya Nee, Lucas Nelson, Zach Nelson, Monica Nemat, Kenneth Nguyen, Michael Nguyen, Christian

FAS1_2019_00_Frontmatter.indd 15 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Nieves, Nyia Njamfa, Ahmed Noor, Kyle Nyugen, Ahamd Obeidat, Gerald Olayan, Anndres Olson, Hasaan Omar, Daniel Ortiz, Michael O'Shea, Zonghao Pan, Vasilis Sebastian Paraschos, Christopher Parrino, Janak Patel, Vanisha Patel, Cyril Patra, Rita Paulis, Dmytro Pavlenko, Nancy A. Pina, Alexander Polyak, Jackeline Porto, Shannon D. Powell, Jacob Pruett, Laith Rahabneh, Kamleshun Ramphul, Janhvi Rana, Nidaa Rasheed, Abdul Sattar Raslan, Tomas Ream, Rashelle Ripa, Amanda Michelle Ritchie, Helio Manuel Grullón Rodríguez, Sarah Rohrig, Gessel Romero, Alexander Rose, Rachel Rose, Erica Rubin, Areesha Saati, Jeffrey Sackey, Raza H. Sagarwala, Chhavi Saini, Sergii Sakhno, Allie Sakowicz, Shadia Saleh, Roshun Sangani, Dhruv Sarwal, Abeer Sarwar, M. Sathyanarayanan, Neetu Scariya, Tonio Felix Schaffert, Melissa Schechter, Kathryn Scheinberg, Emma Schnuckle, Emma Schulte, Taylor Schweigert, Lee Seifert, Sheila Serin, Deeksha Seth, Omid Shafaat, Nirav Shah, Samir K. Shah, Wasif Nauman Shah, Muhanad Shaib, Ahmed Shakir, Purnima Sharma, Tina Sharma, Kayla Sheehan, Dr. Priya Shenwai, Sami Shoura, Kris Sifeldeen, Akhand Singh, Manik Inder Singh, Ramzi Y. Skaik, Samantha A. Smith, Timothy Smith, Emilie Song, Hang Song, Shichen Song, Luke Sorensen, Charles Starling, Jonathan Andrew Stone, Nathan Stumpf, Johnny Su, Bahaa Eddine Succar, Saranya Sundaram, Steven Svoboda, Clara Sze, Olive Tang, Brian Tanksley, Omar Tayh, Joshua Taylor, Valerie Teano, Warren Teltser, Steffanie Camilo Tertulien, Roger Torres, Michael Trainer, Andrew Trinh, Aalap K. Trivedi, Georgeanna Tsoumas, Elizabeth Tsui, Cem Turam, Methma Udawatta, Daramfon Udofia, Adaku Ume, Rio Varghese, Judith Vásquez, Earl Vialpando, Sagar Vinayak, Phuong Vo, Habiba Wada, Jason Wang, Tiffany Wang, Zoe Warczak, Mitchell Waters, Rachel Watson, Elizabeth Douglas Weigel, Rabbi Michael Weingarten, Kaystin Weisenberger, Aidan Woodthorpe, Mattia Wruble, Angela Wu, Catherine Xie, Rebecca Xu, Nicholas Yeisley, Sammy Yeroushalmi, Melissas Yuan, Sahil Zaveri, and Yolanda Zhang.

FAS1_2019_00_Frontmatter.indd 16 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

How to Contribute

This version of *First Aid for the USMLE Step 1* incorporates thousands of contributions and improvements suggested by student and faculty advisors. We invite you to participate in this process. Please send us your suggestions for:

- Study and test-taking strategies for the USMLE Step 1
- New facts, mnemonics, diagrams, and clinical images
- High-yield topics that may appear on future Step 1 exams
- Personal ratings and comments on review books, question banks, apps, videos, and courses

For each new entry incorporated into the next edition, you will receive **up to a $20 Amazon.com gift card** as well as personal acknowledgment in the next edition. Significant contributions will be compensated at the discretion of the authors. Also, let us know about material in this edition that you feel is low yield and should be deleted.

All submissions including potential errata should ideally be supported with hyperlinks to a dynamically updated Web resource such as UpToDate, AccessMedicine, and ClinicalKey.

We welcome potential errata on grammar and style if the change improves readability. Please note that *First Aid* style is somewhat unique; for example, we have fully adopted the *AMA Manual of Style* recommendations on eponyms ("We recommend that the possessive form be omitted in eponymous terms") and on abbreviations (no periods with eg, ie, etc). We also avoid periods in tables unless required for full sentences. Kindly refrain from submitting "style errata" unless you find specific inconsistencies with the *AMA Manual of Style*.

The preferred way to submit new entries, clarifications, mnemonics, or potential corrections with a valid, authoritative reference is via our website: **www.firstaidteam.com**.

This website will be continuously updated with validated errata, new high-yield content, and a new online platform to contribute suggestions, mnemonics, diagrams, clinical images, and potential errata.

Alternatively, you can email us at: **firstaid@scholarrx.com.**

Contributions submitted by **May 15, 2020**, receive priority consideration for the 2021 edition of *First Aid for the USMLE Step 1*. We thank you for taking the time to share your experience and apologize in advance that we cannot individually respond to all contributors as we receive thousands of contributions each year.

FAS1_2019_00_Frontmatter.indd 17 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

NOTE TO CONTRIBUTORS

All contributions become property of the authors and are subject to editing and reviewing. Please verify all data and spellings carefully. Contributions should be supported by at least two high-quality references.

Check our website first to avoid duplicate submissions. In the event that similar or duplicate entries are received, only the first complete entry received with valid, authoritative references will be credited. Please follow the style, punctuation, and format of this edition as much as possible.

JOIN THE FIRST AID TEAM

The *First Aid* author team is pleased to offer part-time and full-time paid internships in medical education and publishing to motivated medical students and physicians. Internships range from a few months (eg, a summer) up to a full year. Participants will have an opportunity to author, edit, and earn academic credit on a wide variety of projects, including the popular *First Aid* series.

For 2020, we are actively seeking passionate medical students and graduates with a specific interest in improving our medical illustrations, expanding our database of medical photographs, and developing the software that supports our crowdsourcing platform. We welcome people with prior experience and talent in these areas. Relevant skills include clinical imaging, digital photography, digital asset management, information design, medical illustration, graphic design, tutoring, and software development.

Please email us at **firstaid@scholarrx.com** with a CV and summary of your interest or sample work.

FAS1_2019_00_Frontmatter.indd 18 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

How to Use This Book

CONGRATULATIONS: You now possess the book that has guided nearly two million students to USMLE success for 30 years. With appropriate care, the binding should last the useful life of the book. Keep in mind that putting excessive flattening pressure on any binding will accelerate its failure. If you purchased a book that you believe is defective, please **immediately** return it to the place of purchase. If you encounter ongoing issues, you can also contact Customer Service at our publisher, McGraw-Hill Education, at https://www.mheducation.com/contact.html.

START EARLY: Use this book as early as possible while learning the basic medical sciences. The first semester of your first year is not too early! Devise a study plan by reading Section I: Guide to Efficient Exam Preparation, and make an early decision on resources to use by checking Section IV: Top-Rated Review Resources. Note that *First Aid* is neither a textbook nor a comprehensive review book, and it is not a panacea for inadequate preparation.

CONSIDER *FIRST AID* YOUR ANNOTATION HUB: Annotate material from other resources, such as class notes or comprehensive textbooks, into your book. This will keep all the high-yield information you need in one place. Other tips on keeping yourself organized:

- For best results, use fine-tipped ballpoint pens (eg, BIC Pro+, Uni-Ball Jetstream Sports, Pilot Drawing Pen, Zebra F-301). If you like gel pens, try Pentel Slicci, and for markers that dry almost immediately, consider Staedtler Triplus Fineliner, Pilot Drawing Pen, and Sharpies.
- Consider using pens with different colors of ink to indicate different sources of information (eg, blue for USMLE-Rx Step 1 Qmax, green for UWorld Step 1 Qbank).
- Choose highlighters that are bright and dry quickly to minimize smudging and bleeding through the page (eg, Tombow Kei Coat, Sharpie Gel).
- Many students de-spine their book and get it 3-hole-punched. This will allow you to insert materials from other sources, including curricular materials.

INTEGRATE STUDY WITH CASES, FLASH CARDS, AND QUESTIONS: To broaden your learning strategy, consider integrating your *First Aid* study with case-based reviews (eg, *First Aid Cases for the USMLE Step 1*), flash cards (eg, First Aid Flash Facts), and practice questions (eg, the USMLE-Rx Step 1 Qmax). Read the chapter in the book, then test your comprehension by using cases, flash cards, and questions that cover the same topics. Maintain access to more comprehensive resources (eg, *First Aid for the Basic Sciences: General Principles* and *Organ Systems* and First Aid Express videos) for deeper review as needed.

PRIME YOUR MEMORY: Return to your annotated Sections II and III several days before taking the USMLE Step 1. The book can serve as a useful way of retaining key associations and keeping high-yield facts fresh in your memory just prior to the exam. The Rapid Review section includes high-yield topics to help guide your studying.

CONTRIBUTE TO FIRST AID: Reviewing the book immediately after your exam can help us improve the next edition. Decide what was truly high and low yield and send us your comments. Feel free to send us scanned images from your annotated *First Aid* book as additional support. Of course, always remember that **all examinees are under agreement with the NBME to not disclose the specific details of copyrighted test material.**

FAS1_2019_00_Frontmatter.indd 19 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Selected USMLE Laboratory Values

* = Included in the Biochemical Profile (SMA-12)

Blood, Plasma, Serum	Reference Range	SI Reference Intervals
*Alanine aminotransferase (ALT, GPT at 30°C)	8–20 U/L	8–20 U/L
Amylase, serum	25–125 U/L	25–125 U/L
*Aspartate aminotransferase (AST, GOT at 30°C)	8–20 U/L	8–20 U/L
Bilirubin, serum (adult)		
Total // Direct	0.1–1.0 mg/dL // 0.0–0.3 mg/dL	2–17 μmol/L // 0–5 μmol/L
*Calcium, serum (Total)	8.4–10.2 mg/dL	2.1–2.8 mmol/L
*Cholesterol, serum (Total)	Rec: < 200 mg/dL	< 5.2 mmol/L
*Creatinine, serum (Total)	0.6–1.2 mg/dL	53–106 μmol/L
Electrolytes, serum		
Sodium (Na^+)	136–145 mEq/L	136–145 mmol/L
Chloride (Cl^-)	95–105 mEq/L	95–105 mmol/L
* Potassium (K^+)	3.5–5.0 mEq/L	3.5–5.0 mmol/L
Bicarbonate (HCO_3^-)	22–28 mEq/L	22–28 mmol/L
Magnesium (Mg^{2+})	1.5–2 mEq/L	0.75–1.0 mmol/L
Gases, arterial blood (room air)		
P_{O_2}	75–105 mm Hg	10.0–14.0 kPa
P_{CO_2}	33–45 mm Hg	4.4–5.9 kPa
pH	7.35–7.45	$[H^+]$ 36–44 nmol/L
*Glucose, serum	Fasting: 70–110 mg/dL 2-h postprandial: < 120 mg/dL	3.8–6.1 mmol/L < 6.6 mmol/L
Growth hormone – arginine stimulation	Fasting: < 5 ng/mL provocative stimuli: > 7 ng/mL	< 5 μg/L > 7 μg/L
Osmolality, serum	275–295 mOsm/kg	275–295 mOsm/kg
*Phosphatase (alkaline), serum (p-NPP at 30°C)	20–70 U/L	20–70 U/L
*Phosphorus (inorganic), serum	3.0–4.5 mg/dL	1.0–1.5 mmol/L
Prolactin, serum (hPRL)	< 20 ng/mL	< 20 μg/L
*Proteins, serum		
Total (recumbent)	6.0–7.8 g/dL	60–78 g/L
Albumin	3.5–5.5 g/dL	35–55 g/L
Globulins	2.3–3.5 g/dL	23–35 g/L
Thyroid-stimulating hormone, serum or plasma	0.5–5.0 μU/mL	0.5–5.0 mU/L
*Urea nitrogen, serum (BUN)	7–18 mg/dL	1.2–3.0 mmol/L
*Uric acid, serum	3.0–8.2 mg/dL	0.18–0.48 mmol/L

(continues)

FAS1_2019_00_Frontmatter.indd 20 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

Cerebrospinal Fluid	Reference Range	SI Reference Intervals
Glucose	40–70 mg/dL	2.2–3.9 mmol/L
Hematologic		
Erythrocyte count	Male: 4.3–5.9 million/mm^3 Female: 3.5–5.5 million/mm^3	4.3–5.9 × 10^{12}/L 3.5–5.5 × 10^{12}/L
Erythrocyte sedimentation rate (Westergen)	Male: 0–15 mm/h Female: 0–20 mm/h	0–15 mm/h 0–20 mm/h
Hematocrit	Male: 41–53% Female: 36–46%	0.41–0.53 0.36–0.46
Hemoglobin, blood	Male: 13.5–17.5 g/dL Female: 12.0–16.0 g/dL	2.09–2.71 mmol/L 1.86–2.48 mmol/L
Hemoglobin, plasma	1–4 mg/dL	0.16–0.62 µmol/L
Leukocyte count and differential		
Leukocyte count	4,500–11,000/mm^3	4.5–11.0 × 10^9/L
Segmented neutrophils	54–62%	0.54–0.62
Band forms	3–5%	0.03–0.05
Eosinophils	1–3%	0.01–0.03
Basophils	0–0.75%	0–0.0075
Lymphocytes	25–33%	0.25–0.33
Monocytes	3–7%	0.03–0.07
Mean corpuscular hemoglobin	25.4–34.6 pg/cell	0.39–0.54 fmol/cell
Mean corpuscular volume	80–100 μm^3	80–100 fL
Partial thromboplastin time (activated)	25–40 seconds	25–40 seconds
Platelet count	150,000–400,000/mm^3	150–400 × 10^9/L
Prothrombin time	11–15 seconds	11–15 seconds
Reticulocyte count	0.5–1.5% of red cells	0.005–0.015
Sweat		
Chloride	0–35 mmol/L	0–35 mmol/L
Urine		
Creatinine clearance	Male: 97–137 mL/min Female: 88–128 mL/min	
Osmolality	50–1,400 mOsmol/kg H_2O	
Proteins, total	< 150 mg/24 h	< 0.15 g/24 h

FAS1_2019_00_Frontmatter.indd 21 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

First Aid Checklist for the USMLE Step 1

This is an example of how you might use the information in Section I to prepare for the USMLE Step 1. Refer to corresponding topics in Section I for more details.

Years Prior

- ☐ Use top-rated review resources for first-year medical school courses.
- ☐ Ask for advice from those who have recently taken the USMLE Step 1.

Months Prior

- ☐ Review computer test format and registration information.
- ☐ Register six months in advance.
- ☐ Carefully verify name and address printed on scheduling permit. Make sure the name on scheduling permit matches the name printed on your photo ID.
- ☐ Go online for test date ASAP.
- ☐ Define your exam goals (pass comfortably, beat the mean, ace the test)
- ☐ Set up a realistic timeline for study. Cover less crammable subjects first.
- ☐ Evaluate and choose study materials (review books, question banks).
- ☐ Use a question bank to simulate the USMLE Step 1 to pinpoint strengths and weaknesses in knowledge and test-taking skills.

Weeks Prior

- ☐ Do another test simulation in a question bank.
- ☐ Assess how close you are to your goal.
- ☐ Pinpoint remaining weaknesses. Stay healthy (exercise, sleep).
- ☐ Verify information on admission ticket (eg, location, date).

One Week Prior

- ☐ Remember comfort measures (loose clothing, earplugs, etc).
- ☐ Work out test site logistics (eg, location, transportation, parking, lunch).
- ☐ Print or download your Scheduling Permit and Scheduling Confirmation to your phone.

One Day Prior

- ☐ Relax.
- ☐ Lightly review short-term material if necessary. Skim high-yield facts.
- ☐ Get a good night's sleep.

Day of Exam

- ☐ Relax.
- ☐ Eat breakfast.
- ☐ Minimize bathroom breaks during exam by avoiding excessive morning caffeine.

After Exam

- ☐ Celebrate, regardless of how well you feel you did.
- ☐ Send feedback to us on our website at **www.firstaidteam.com**.

FAS1_2019_00_Frontmatter.indd 22 11/14/19 4:35 PM

(c) ketabton.com: The Digital Library

SECTION I

Guide to Efficient Exam Preparation

"I don't love studying. I hate studying. I like learning. Learning is beautiful."

—Natalie Portman

"Finally, from so little sleeping and so much reading, his brain dried up and he went completely out of his mind."

—Miguel de Cervantes Saavedra, *Don Quixote*

"Sometimes the questions are complicated and the answers are simple."

—Dr. Seuss

"He who knows all the answers has not been asked all the questions."

—Confucius

"The expert in anything was once a beginner."

—Helen Hayes

"It always seems impossible until it's done."

—Nelson Mandela

- Introduction 2
- USMLE Step 1—The Basics 2
- Defining Your Goal 12
- Learning Strategies 13
- Timeline for Study 16
- Study Materials 20
- Test-Taking Strategies 22
- Clinical Vignette Strategies 23
- If You Think You Failed 24
- Testing Agencies 24
- References 25

FAS1_2019_00_Section_I.indd 1 11/7/19 2:48 PM

(c) ketabton.com: The Digital Library

INTRODUCTION

Relax.

This section is intended to make your exam preparation easier, not harder. Our goal is to reduce your level of anxiety and help you make the most of your efforts by helping you understand more about the United States Medical Licensing Examination, Step 1 (USMLE Step 1). As a medical student, you are no doubt familiar with taking standardized examinations and quickly absorbing large amounts of material. When you first confront the USMLE Step 1, however, you may find it all too easy to become sidetracked from your goal of studying with maximal effectiveness. Common mistakes that students make when studying for Step 1 include the following:

- Starting to study (including *First Aid*) too late
- Starting to study intensely too early and burning out
- Starting to prepare for boards before creating a knowledge foundation
- Using inefficient or inappropriate study methods
- Buying the wrong resources or buying too many resources
- Buying only one publisher's review series for all subjects
- Not using practice examinations to maximum benefit
- Not understanding how scoring is performed or what the score means
- Not using review books along with your classes
- Not analyzing and improving your test-taking strategies
- Getting bogged down by reviewing difficult topics excessively
- Studying material that is rarely tested on the USMLE Step 1
- Failing to master certain high-yield subjects owing to overconfidence
- Using *First Aid* as your sole study resource
- Trying to prepare for it all alone

The test at a glance:

- *8-hour exam*
- *Up to a total of 280 multiple choice items*
- *7 test blocks (60 min/block)*
- *Up to 40 test items per block*
- *45 minutes of break time, plus another 15 if you skip the tutorial*

In this section, we offer advice to help you avoid these pitfalls and be more productive in your studies.

USMLE STEP 1—THE BASICS

The USMLE Step 1 is the first of three examinations that you must pass in order to become a licensed physician in the United States. The USMLE is a joint endeavor of the National Board of Medical Examiners (NBME) and the Federation of State Medical Boards (FSMB). The USMLE serves as the single examination system for US medical students and international medical graduates (IMGs) seeking medical licensure in the United States.

(c) ketabton.com: The Digital Library

The Step 1 exam includes test items that can be grouped by the organizational constructs outlined in Table 1 (in order of tested frequency).

TABLE 1. Frequency of Various Constructs Tested on the USMLE Step 1.*

Competency	Range, %
Medical knowledge: applying foundational science concepts	52–62
Patient care: diagnosis	20–30
Patient care: management	7–12
Practice-based learning & improvement	5–7
Communication/professionalism	2–5
Discipline	**Range, %**
Pathology	45–52
Physiology	26–34
Pharmacology	16–23
Biochemistry & nutrition	14–24
Microbiology & immunology	15–22
Gross anatomy & embryology	11–15
Histology & cell biology	9–13
Behavioral sciences	8–12
Genetics	5–9

System	Range, %
General principles	13–17
Behavioral health & nervous systems/special senses	9–13
Respiratory & renal/urinary systems	9–13
Reproductive & endocrine systems	9–13
Blood & lymphoreticular/immune systems	7–11
Multisystem processes & disorders	7–11
Musculoskeletal, skin & subcutaneous tissue	6–10
Cardiovascular system	6–10
Gastrointestinal system	5–9
Biostatistics & epidemiology/population health	5–7
Social sciences: communication skills/ethics	3–5

*Percentages are subject to change at any time. www.usmle.org

How Is the Computer-Based Test (CBT) Structured?

The CBT Step 1 exam consists of one "optional" tutorial/simulation block and seven "real" question blocks of up to 40 questions per block with no more than 280 questions in total, timed at 60 minutes per block. A short 11-question survey follows the last question block. The computer begins the survey with a prompt to proceed to the next block of questions.

Once an examinee finishes a particular question block on the CBT, he or she must click on a screen icon to continue to the next block. Examinees **cannot** go back and change their answers to questions from any previously completed block. However, changing answers is allowed **within** a block of questions as long as the block has not been ended and if time permits.

What Is the CBT Like?

Given the unique environment of the CBT, it's important that you become familiar ahead of time with what your test-day conditions will be like. You can access a 15-minute tutorial and practice blocks at http://orientation.nbme.org/Launch/USMLE/STPF1. This tutorial interface is very similar to the one you will use in the exam; learn it now and you can skip taking it during the exam, giving you up to 15 extra minutes of break time. You can gain experience with the CBT format by taking the 120 practice questions (3 blocks with 40 questions each) available online or by signing up for a practice session at a test center for a fee.

(c) ketabton.com: The Digital Library

For security reasons, examinees are not allowed to bring any personal electronic equipment into the testing area. This includes both digital and analog watches, iPods, tablets, calculators, cell phones, and electronic paging devices. Examinees are also prohibited from carrying in their books, notes, pens/pencils, and scratch paper. Food and beverages are also prohibited in the testing area. The testing centers are monitored by audio and video surveillance equipment. However, most testing centers allot each examinee a small locker outside the testing area in which he or she can store snacks, beverages, and personal items.

- *Keyboard shortcuts:*
  - *A, B, etc—letter choices*
  - *Enter or spacebar—move to next question*
  - *Esc—exit pop-up Calculator and Notes windows*

Questions are typically presented in multiple choice format, with 4–5 possible answer options. There is a countdown timer on the lower left corner of the screen as well. There is also a button that allows the examinee to mark a question for review. If a given question happens to be longer than the screen (which occurs very rarely), a scroll bar will appear on the right, allowing the examinee to see the rest of the question. Regardless of whether the examinee clicks on an answer choice or leaves it blank, he or she must click the "Next" button to advance to the next question.

- *Heart sounds are tested via media questions. Make sure you know how different heart diseases sound on auscultation.*

The USMLE features a small number of media clips in the form of audio and/or video. There may even be a question with a multimedia heart sound simulation. In these questions, a digital image of a torso appears on the screen, and the examinee directs a digital stethoscope to various auscultation points to listen for heart and breath sounds. The USMLE orientation materials include several practice questions in these formats. During the exam tutorial, examinees are given an opportunity to ensure that both the audio headphones and the volume are functioning properly. If you are already familiar with the tutorial and planning on skipping it, first skip ahead to the section where you can test your headphones. After you are sure the headphones are working properly, proceed to the exam.

- *Be sure to test your headphones during the tutorial.*

- *Familiarize yourself with the commonly tested lab values (eg, Hgb, WBC, platelets, Na^+, K^+).*

The examinee can call up a window displaying normal laboratory values. In order to do so, he or she must click the "Lab" icon on the top part of the screen. Afterward, the examinee will have the option to choose between "Blood," "Cerebrospinal," "Hematologic," or "Sweat and Urine." The normal values screen may obscure the question if it is expanded. The examinee may have to scroll down to search for the needed lab values. You might want to memorize some common lab values so you spend less time on questions that require you to analyze these.

- *Illustrations on the test include:*
  - *Gross specimen photos*
  - *Histology slides*
  - *Medical imaging (eg, x-ray, CT, MRI)*
  - *Electron micrographs*
  - *Line drawings*

The CBT interface provides a running list of questions on the left part of the screen at all times. The software also permits examinees to highlight or cross out information by using their mouse. There is a "Notes" icon on the top part of the screen that allows students to write notes to themselves for review at a later time. Finally, the USMLE has recently added new functionality including text magnification and reverse color (white text on black background). Being familiar with these features can save time and may help you better view and organize the information you need to answer a question.

- *Ctrl-Alt-Delete are the keys of death during the exam. Don't touch them at the same time!*

(c) ketabton.com: The Digital Library

For those who feel they might benefit, the USMLE offers an opportunity to take a simulated test, or "CBT Practice Session" at a Prometric center. Students are eligible to register for this three-and-one-half-hour practice session after they have received their scheduling permit.

You can take a shortened CBT practice test at a Prometric center.

The same USMLE Step 1 sample test items (120 questions) available on the USMLE website, www.usmle.org, are used at these sessions. **No new items will be presented.** The practice session is available at a cost of $75 (or more if taken outside of the US and Canada) and is divided into a short tutorial and three 1-hour blocks of ~40 test items each. Students receive a printed percent-correct score after completing the session. **No explanations of questions are provided.**

You may register for a practice session online at www.usmle.org. A separate scheduling permit is issued for the practice session. Students should allow two weeks for receipt of this permit.

How Do I Register to Take the Exam?

The Prometric website will display a calendar with open test dates.

Prometric test centers offer Step 1 on a year-round basis, except for the first two weeks in January and major holidays. The exam is given every day except Sunday at most centers. Some schools administer the exam on their own campuses. Check with the test center you want to use before making your exam plans.

US students can apply to take Step 1 at the NBME website. This application allows you to select one of 12 overlapping three-month blocks in which to be tested (eg, April–May–June, June–July–August). Choose your three-month eligibility period wisely. If you need to reschedule outside your initial three-month period, you can request a one-time extension of eligibility for the next contiguous three-month period, and pay a rescheduling fee. The application also includes a photo ID form that must be certified by an official at your medical school to verify your enrollment. After the NBME processes your application, it will send you a scheduling permit.

The scheduling permit you receive from the NBME will contain your USMLE identification number, the eligibility period in which you may take the exam, and two additional numbers. The first of these is known as your "scheduling number." You must have this number in order to make your exam appointment with Prometric. The second number is known as the "candidate identification number," or CIN. Examinees must enter their CINs at the Prometric workstation in order to access their exams. However, you will not be allowed to bring your permit into the exam and will be asked to copy your CIN onto your scratch paper. Prometric has no access to the codes. **Make sure to bring a paper or electronic copy of your permit with you to the exam!** Also bring an unexpired, government-issued photo ID that includes your signature (such as a driver's license or passport). Make sure the name on your photo ID exactly matches the name that appears on your scheduling permit.

(c) ketabton.com: The Digital Library

The confirmation emails that Prometric and NBME send are not the same as the scheduling permit.

Once you receive your scheduling permit, you may access the Prometric website or call Prometric's toll-free number to arrange a time to take the exam. You may contact Prometric two weeks before the test date if you want to confirm identification requirements. Although requests for taking the exam may be completed more than six months before the test date, examinees will not receive their scheduling permits earlier than six months before the eligibility period. The eligibility period is the three-month period you have chosen to take the exam. Most medical students choose the April–June or June–August period. Because exams are scheduled on a "first-come, first-served" basis, it is recommended that you book an exam date on the Prometric website as soon as you receive your permit. Prometric will provide appointment confirmation on a print-out and by email. Be sure to read the latest *USMLE Bulletin of Information* for further details.

Test scheduling is done on a "first-come, first-served" basis. It's important to schedule an exam date as soon as you receive your scheduling permit.

What If I Need to Reschedule the Exam?

You can change your test date and/or center by contacting Prometric at 1-800-MED-EXAM (1-800-633-3926) or www.prometric.com. Make sure to have your CIN when rescheduling. If you are rescheduling by phone, you must speak with a Prometric representative; leaving a voicemail message will not suffice. To avoid a rescheduling fee, you will need to request a change at least 31 calendar days before your appointment. Please note that your rescheduled test date must fall within your assigned three-month eligibility period.

Register six months in advance for seating and scheduling preference.

When Should I Register for the Exam?

You should plan to register as far in advance as possible ahead of your desired test date (eg, six months), but, depending on your particular test center, new dates and times may open closer to the date. Scheduling early will guarantee that you will get either your test center of choice or one within a 50-mile radius of your first choice. For most US medical students, the desired testing window is in June, since most medical school curricula for the second year end in May or June. Thus, US medical students should plan to register before January in anticipation of a June test date. The timing of the exam is more flexible for IMGs, as it is related only to when they finish exam preparation. Talk with upperclassmen who have already taken the test so you have real-life experience from students who went through a similar curriculum, then formulate your own strategy.

Where Can I Take the Exam?

Your testing location is arranged with Prometric when you book your test date (after you receive your scheduling permit). For a list of Prometric locations nearest you, visit www.prometric.com.

(c) ketabton.com: The Digital Library

How Long Will I Have to Wait Before I Get My Scores?

The USMLE reports scores in three to four weeks, unless there are delays in score processing. Examinees will be notified via email when their scores are available. By following the online instructions, examinees will be able to view, download, and print their score report online for ~120 days after score notification, after which scores can only be obtained through requesting an official USMLE transcript. Additional information about score timetables and accessibility is available on the official USMLE website.

What About Time?

Gain extra break time by skipping the tutorial or finishing a block early.

Time is of special interest on the CBT exam. Here's a breakdown of the exam schedule:

15 minutes	Tutorial (skip if familiar with test format and features)
7 hours	Seven 60-minute question blocks
45 minutes	Break time (includes time for lunch)

The computer will keep track of how much time has elapsed on the exam. However, the computer will show you only how much time you have remaining in a given block. Therefore, it is up to you to determine if you are pacing yourself properly (at a rate of approximately one question per 90 seconds).

The computer does not warn you if you are spending more than your allotted time for a break. You should therefore budget your time so that you can take a short break when you need one and have time to eat. You must be especially careful not to spend too much time in between blocks (you should keep track of how much time elapses from the time you finish a block of questions to the time you start the next block). After you finish one question block, you'll need to click to proceed to the next block of questions. If you do not click within 30 seconds, you will automatically be entered into a break period.

Be careful to watch the clock on your break time.

Break time for the day is 45 minutes, but you are not required to use all of it, nor are you required to use any of it. You can gain extra break time (but not extra time for the question blocks) by skipping the tutorial or by finishing a block ahead of the allotted time. Any time remaining on the clock when you finish a block gets added to your remaining break time. Once a new question block has been started, you may not take a break until you have reached the end of that block. If you do so, this will be recorded as an "unauthorized break" and will be reported on your final score report.

Finally, be aware that it may take a few minutes of your break time to "check out" of the secure resting room and then "check in" again to resume testing, so plan accordingly. The "check-in" process may include fingerprints, pocket checks, and metal detector scanning. Some students recommend pocketless clothing on exam day to streamline the process.

(c) ketabton.com: The Digital Library

If I Freak Out and Leave, What Happens to My Score?

Your scheduling permit shows a CIN that you will need to enter to start your exam. Entering the CIN is the same as breaking the seal on a test book, and you are considered to have started the exam when you do so. However, no score will be reported if you do not complete the exam. In fact, if you leave at any time from the start of the test to the last block, no score will be reported. The fact that you started but did not complete the exam, however, will appear on your USMLE score transcript. Even though a score is not posted for incomplete tests, examinees may still get an option to request that their scores be calculated and reported if they desire; unanswered questions will be scored as incorrect.

The exam ends when all question blocks have been completed or when their time has expired. As you leave the testing center, you will receive a printed test-completion notice to document your completion of the exam. To receive an official score, you must finish the entire exam.

What Types of Questions Are Asked?

Nearly three fourths of Step 1 questions begin with a description of a patient.

All questions on the exam are **one-best-answer multiple choice items.** Most questions consist of a clinical scenario or a direct question followed by a list of five or more options. You are required to select the single best answer among the options given. There are no "except," "not," or matching questions on the exam. A number of options may be partially correct, in which case you must select the option that best answers the question or completes the statement. Additionally, keep in mind that experimental questions may appear on the exam, which do not affect your score.

How Is the Test Scored?

Each Step 1 examinee receives an electronic score report that includes the examinee's pass/fail status, a three-digit test score, a bar chart comparing the examinee's performance to that of other examinees', and a graphic depiction of the examinee's performance by physician task, discipline and organ system.

The USMLE score report highlights the examinee's strength and weaknesses by providing an overview of their performance by physician task, discipline and organ system compared to their overall performance on the exam. Each of the questions (minus experimental questions) is tagged according to any or all relevant content areas. Yellow-colored boxes (lower, same, higher) on your score report indicate your performance in each specific content area relative to your overall performance on the exam. This is often a direct consequence of the total number of questions for each physician task, discipline or system, which is indicated by percentage range after each specified content area on the score report (see Figure 1).

(c) ketabton.com: The Digital Library

FIGURE 1. Samples from the USMLE Step 1 Performance Profile.



The NBME provides a three-digit test score based on the total number of items answered correctly on the examination, which corresponds to a particular percentile (see Figure 2). Your three-digit score will be qualified by the mean and standard deviation of US and Canadian medical school first-time examinees.

The mean Step 1 score for US medical students continues to rise, from 200 in 1991 to 230 in 2018.

Since some questions may be experimental and are not counted, it is possible to get different scores for the same number of correct answers. In 2018, the mean score was 230 with a standard deviation of 19.

The passing score for Step 1 is 194. The NBME does not report the minimum number of correct responses needed to pass, but estimates that it is roughly 60–70%. The NBME may adjust the minimum passing score in the future, so please check the USMLE website or www.firstaidteam.com for updates.

According to the USMLE, medical schools receive a listing of total scores and pass/fail results plus group summaries by discipline and organ system. Students can withhold their scores from their medical school if they wish. Official USMLE transcripts, which can be sent on request to residency programs, include only total scores, not performance profiles.

(c) ketabton.com: The Digital Library

FIGURE 2. Score and Percentile for First-time Step 1 Takers.



N=72,473 including US and Canadian medical school students testing between January 1, 2016–December 31, 2018. www.usmle.org.

Consult the USMLE website or your medical school for the most current and accurate information regarding the examination.

What Does My Score Mean?

The most important point with the Step 1 score is passing versus failing. Passing essentially means, "Hey, you're on your way to becoming a fully licensed doc." As Table 2 shows, the majority of students pass the exam, so remember, we told you to relax.

TABLE 2. Passing Rates for the 2017–2018 USMLE Step 1.[2]

	2017		2018	
	No. Tested	% Passing	No. Tested	% Passing
Allopathic 1st takers	20,353	96%	20,670	96%
Repeaters	1,029	67%	941	67%
Allopathic total	21,382	94%	21,611	95%
Osteopathic 1st takers	3,786	95%	4,092	96%
Repeaters	49	76%	44	73%
Osteopathic total	3,835	95%	4,136	96%
Total US/Canadian	**25,217**	**94%**	**25,747**	**94%**
IMG 1st takers	14,900	78%	14,332	80%
Repeaters	2,303	41%	2,111	44%
IMG total	**17,203**	73%	**16,443**	**75%**
Total Step 1 examinees	**42,420**	**85%**	**42,190**	**86%**

(c) ketabton.com: The Digital Library

Beyond that, the main point of having a quantitative score is to give you a sense of how well you've done on the exam and to help schools and residencies rank their students and applicants, respectively.

Official NBME/USMLE Resources

The NBME offers a Comprehensive Basic Science Examination (CBSE) for practice that is a shorter version of the Step 1. The CBSE contains four blocks of 50 questions each and covers material that is typically learned during the basic science years. Scores range from 45 to 95 and correlate with a Step 1 equivalent (see Table 3). The standard error of measurement is approximately 3 points, meaning a score of 80 would estimate the student's proficiency is somewhere between 77 and 83. In other words, the actual Step 1 score could be predicted to be between 218 and 232. Of course, these values do not correlate exactly, and they do not reflect different test preparation methods. Many schools use this test to gauge whether a student is expected to pass Step 1. If this test is offered by your school, it is usually conducted at the end of regular didactic time before any dedicated Step 1 preparation. If you do not encounter the CBSE before your dedicated study time, you need not worry about taking it. Use the information to help set realistic goals and timetables for your success.

The NBME also offers six forms of Comprehensive Basic Science Self-Assessment (CBSSA). Students who prepared for the exam using this web-based tool reported that they found the format and content highly indicative of questions tested on the actual exam. In addition, the CBSSA is a fair predictor of USMLE performance (see Table 4). The test interface, however, does not match the actual USMLE test interface, so practicing with these forms alone is not advised.

The CBSSA exists in two formats: standard-paced and self-paced, both of which consist of four sections of 50 questions each (for a total of 200 multiple choice items). The standard-paced format allows the user up to 75 minutes to complete each section, reflecting time limits similar to the actual exam. By contrast, the self-paced format places a 5-hour time limit on answering all multiple choice questions. Every few years, a new form is released and an older one is retired, reflecting changes in exam content. Therefore, the newer exams tend to be more similar to the actual Step 1, and scores from these exams tend to provide a better estimation of exam day performance.

Keep in mind that this bank of questions is available only on the web. The NBME requires that users start and complete the exam within 90 days of purchase. Once the assessment has begun, users are required to complete the sections within 20 days. Following completion of the questions, the CBSSA provides a performance profile indicating the user's relative strengths and weaknesses, much like the report profile for the USMLE Step 1 exam. The profile is scaled with an average score of 500 and a standard deviation of 100. In addition to the performance profile, examinees will be informed of the number of questions answered incorrectly. You will have the ability to review the text of the incorrect question with the correct answer.

TABLE 3. CBSE to USMLE Score Prediction.

CBSE Score	Step 1 Equivalent
≥ 94	≥ 260
92	255
90	250
88	245
86	240
84	235
82	230
80	225
78	220
76	215
74	210
72	205
70	200
68	195
66	190
64	185
62	180
60	175
58	170
56	165
54	160
52	155
50	150
48	145
46	140
≤ 44	≤ 135

▸ *Practice questions may be easier than the actual exam.*

(c) ketabton.com: The Digital Library

TABLE 4. CBSSA to USMLE Score Prediction.

CBSSA Score	Approximate USMLE Step 1 Score
150	155
200	165
250	175
300	186
350	196
400	207
450	217
500	228
550	238
600	248
650	259
700	269
750	280
800	290

Explanations for the correct answer, however, will not be provided. The NBME charges $60 for assessments with expanded feedback. The fees are payable by credit card or money order. For more information regarding the CBSE and the CBSSA, visit the NBME's website at www.nbme.org.

The NBME scoring system is weighted for each assessment exam. While some exams seem more difficult than others, the score reported takes into account these inter-test differences when predicting Step 1 performance. Also, while many students report seeing Step 1 questions "word-for-word" out of the assessments, the NBME makes special note that no live USMLE questions are shown on any NBME assessment.

Lastly, the International Foundations of Medicine (IFOM) offers a Basic Science Examination (BSE) practice exam at participating Prometric test centers for $200. Students may also take the self-assessment test online for $35 through the NBME's website. The IFOM BSE is intended to determine an examinee's relative areas of strength and weakness in general areas of basic science—not to predict performance on the USMLE Step 1 exam—and the content covered by the two examinations is somewhat different. However, because there is substantial overlap in content coverage and many IFOM items were previously used on the USMLE Step 1, it is possible to roughly project IFOM performance onto the USMLE Step 1 score scale. More information is available at http://www.nbme.org/ifom/.

DEFINING YOUR GOAL

It is useful to define your own personal performance goal when approaching the USMLE Step 1. Your style and intensity of preparation can then be matched to your goal. Furthermore, your goal may depend on your school's requirements, your specialty choice, your grades to date, and your personal assessment of the test's importance. Do your best to define your goals early so that you can prepare accordingly.

Some competitive residency programs place more weight on Step 1 scores when choosing candidates to interview.

The value of the USMLE Step 1 score in selecting residency applicants remains controversial, and some have called for less emphasis to be placed on the score when selecting or screening applicants.[3] For the time being, however, it continues to be an important part of the residency application, and it is not uncommon for some specialties to implement filters that screen out applicants who score below a certain cutoff. This is more likely to be seen in competitive specialties (eg, orthopedic surgery, ophthalmology, dermatology, otolaryngology). Independent of your career goals, you can maximize your future options by doing your best to obtain the highest score possible (see Figure 3). At the same time, your Step 1 score is only one of a number of factors that are assessed when you apply for residency. In fact, many residency programs value other criteria such as letters of recommendation, third-year clerkship grades, honors, and research experience more than a high score on Step 1. Fourth-year medical students who have recently completed the residency application process can be a valuable resource in this regard.

Fourth-year medical students have the best feel for how Step 1 scores factor into the residency application process.

(c) ketabton.com: The Digital Library

FIGURE 3. Median USMLE Step 1 Score by Specialty for Matched US Seniors.[a,b]



[a]Vertical lines show interquartile range.
[b]Data from National Resident Matching Program. Charting outcomes in the match. http://www.nrmp.org/wp-content/uploads/2018/06/Charting-Outcomes-in-the-Match-2018-Seniors.pdf. Published July 2018.

LEARNING STRATEGIES

Many students feel overwhelmed during the preclinical years and struggle to find an effective learning strategy. Table 5 lists several learning strategies you can try and their estimated effectiveness for Step 1 preparation based on the literature (see References). These are merely suggestions, and it's important to take your learning preferences into account. Your comprehensive learning approach will contain a combination of strategies (eg, elaborative interrogation followed by practice testing, mnemonics review using spaced repetition, etc). Regardless of your choice, the foundation of knowledge you build during your basic science years is the most important resource for success on the USMLE Step 1.

The foundation of knowledge you build during your basic science years is the most important resource for success on the USMLE Step 1.

HIGH EFFICACY

Practice Testing

Also called "retrieval practice," practice testing has both direct and indirect benefits to the learner.[4] Effortful retrieval of answers does not only identify weak spots—it directly strengthens long-term retention of material.[5] The more effortful the recall, the better the long-term retention. This advantage has been shown to result in higher test scores and GPAs.[6] In fact, research has shown a positive correlation between the number of boards-style practice questions completed and Step 1 scores among medical students.[7]

Research has shown a positive correlation between the number of boards-style practice questions completed and Step 1 scores among medical students.

Practice testing should be done with "interleaving" (mixing of questions from different topics in a single session). Question banks often allow you to intermingle topics. Interleaved practice helps learners develop their ability to focus on the relevant concept when faced with many possibilities. Practicing topics in massed fashion (eg, all cardiology, then all dermatology) may seem intuitive, but there is strong evidence that interleaving correlates with longer-

(c) ketabton.com: The Digital Library

TABLE 5. Effective Learning Strategies.

EFFICACY	STRATEGY	EXAMPLE RESOURCES
High efficacy	Practice testing (retrieval practice)	UWorld Qbank NBME Self-Assessments USMLE-Rx QMax Kaplan Qbank
	Distributed practice	USMLE-Rx Flash Facts Anki Firecracker Memorang Osmosis
Moderate efficacy	Mnemonics	*Pre-made:* SketchyMedical Picmonic *Self-made:* Mullen Memory
	Elaborative interrogation/ self-explanation	
	Concept mapping	Coggle FreeMind XMind MindNode
Low efficacy	Rereading	
	Highlighting/underlining	
	Summarization	

term retention and increased student achievement, especially on tasks that involve problem solving.[5]

In addition to using question banks, you can test yourself by arranging your notes in a question-answer format (eg, via flash cards). Testing these Q&As in random order allows you to reap the benefit of interleaved practice. Bear in mind that the utility of practice testing comes from the practice of information retrieval, so simply reading through Q&As will attenuate this benefit.

Distributed Practice

Also called "spaced repetition," distributed practice is the opposite of massed practice or "cramming." Learners review material at increasingly spaced out intervals (days to weeks to months). Massed learning may produce more short-term gains and satisfaction, but learners who use distributed practice have better mastery and retention over the long term.[5,9]

Flash cards are a simple way to incorporate both distributed practice and practice testing. Studies have linked spaced repetition learning with flash

(c) ketabton.com: The Digital Library

cards to improved long-term knowledge retention and higher exam scores.[6,8,10] Apps with automated spaced-repetition software (SRS) for flash cards exist for smartphones and tablets, so the cards are accessible anywhere. Proceed with caution: there is an art to making and reviewing cards. The ease of quickly downloading or creating digital cards can lead to flash card overload (it is unsustainable to make 50 flash cards per lecture!). Even at a modest pace, the thousands upon thousands of cards are too overwhelming for Step 1 preparation. Unless you have specific high-yield cards (and have checked the content with high-yield resources), stick to pre-made cards by reputable sources that curate the vast amount of knowledge for you.

Studies have linked spaced repetition learning with flash cards to improved long-term knowledge retention and higher exam scores.

If you prefer pen and paper, consider using a planner or spreadsheet to organize your study material over time. Distributed practice allows for some forgetting of information, and the added effort of recall over time strengthens the learning.

MODERATE EFFICACY

Mnemonics

A "mnemonic" refers to any device that assists memory, such as acronyms, mental imagery (eg, keywords with or without memory palaces), etc. Keyword mnemonics have been shown to produce superior knowledge retention when compared with rote memorization in many scenarios. However, they are generally more effective when applied to memorization-heavy, keyword-friendly topics and may not be broadly suitable.[5] Keyword mnemonics may not produce long-term retention, so consider combining mnemonics with distributed, retrieval-based practice (eg, via flash cards with SRS).

Self-made mnemonics may have an advantage when material is simple and keyword friendly. If you can create your own mnemonic that accurately represents the material, this will be more memorable. When topics are complex and accurate mnemonics are challenging to create, pre-made mnemonics may be more effective, especially if you are inexperienced at creating mnemonics.[11]

Elaborative Interrogation/Self-Explanation

Elaborative interrogation ("why" questions) and self-explanation (general questioning) prompt learners to generate explanations for facts. When reading passages of discrete facts, consider using these techniques, which have been shown to be more effective than rereading (eg, improved recall and better problem-solving/diagnostic performance).[5,12,13]

Elaborative interrogation and self-explanation prompt learners to generate explanations for facts, which improves recall and problem solving.

Concept Mapping

Concept mapping is a method for graphically organizing knowledge, with concepts enclosed in boxes and lines drawn between related concepts.

(c) ketabton.com: The Digital Library

Creating or studying concept maps may be more effective than other activities (eg, writing or reading summaries/outlines). However, studies have reached mixed conclusions about its utility, and the small size of this effect raises doubts about its authenticity and pedagogic significance.[14]

LOW EFFICACY

Rereading

While the most commonly used method among surveyed students, rereading has not been shown to correlate with grade point average.[9] Due to its popularity, rereading is often a comparator in studies on learning. Other strategies that we have discussed (eg, practice testing) have been shown to be significantly more effective than rereading.

Highlighting/Underlining

Because this method is passive, it tends to be of minimal value for learning and recall. In fact, lower-performing students are more likely to use these techniques.[9] Students who highlight and underline do not learn how to actively recall learned information and thus find it difficult to apply knowledge to exam questions.

Summarization

While more useful for improving performance on generative measures (eg, free recall or essays), summarization is less useful for exams that depend on recognition (eg, multiple choice). Findings on the overall efficacy of this method have been mixed.[5]

TIMELINE FOR STUDY

Before Starting

Your preparation for the USMLE Step 1 should begin when you enter medical school. Organize and commit to studying from the beginning so that when the time comes to prepare for the USMLE, you will be ready with a strong foundation.

Make a Schedule

After you have defined your goals, map out a study schedule that is consistent with your objectives, your vacation time, the difficulty of your ongoing coursework, and your family and social commitments (see Figure 4). Determine whether you want to spread out your study time or concentrate it into 14-hour study days in the final weeks. Then factor in your own history in

(c) ketabton.com: The Digital Library

FIGURE 4. Typical Timeline for the USMLE Step 1.



▸ *Customize your schedule. Tackle your weakest section first.*

preparing for standardized examinations (eg, SAT, MCAT). Talk to students at your school who have recently taken Step 1. Ask them for their study schedules, especially those who have study habits and goals similar to yours. Sample schedules can be found at https://firstaidteam.com/schedules/.

Typically, US medical schools allot between four and eight weeks for dedicated Step 1 preparation. The time you dedicate to exam preparation will depend on your target score as well as your success in preparing yourself during the first two years of medical school. Some students reserve about a week at the end of their study period for final review; others save just a few days. When you have scheduled your exam date, do your best to adhere to it. Studies show that a later testing date does not translate into a higher score, so avoid pushing back your test date without good reason.[15]

Make your schedule realistic, and set achievable goals. Many students make the mistake of studying at a level of detail that requires too much time for a comprehensive review—reading *Gray's Anatomy* in a couple of days is not a realistic goal! Have one catch-up day per week of studying. No matter how well you stick to your schedule, unexpected events happen. But don't let yourself procrastinate because you have catch-up days; stick to your schedule as closely as possible and revise it regularly on the basis of your actual progress. Be careful not to lose focus. Beware of feelings of inadequacy when comparing study schedules and progress with your peers. **Avoid others who stress you out.** Focus on a few top-rated resources that suit your learning style—not on some obscure books your friends may pass down to you. Accept the fact that you cannot learn it all.

▸ *Avoid burnout. Maintain proper diet, exercise, and sleep habits.*

You will need time for uninterrupted and focused study. Plan your personal affairs to minimize crisis situations near the date of the test. Allot an adequate number of breaks in your study schedule to avoid burnout. Maintain a healthy lifestyle with proper diet, exercise, and sleep.

Another important aspect of your preparation is your studying environment. **Study where you have always been comfortable studying.** Be sure to include everything you need close by (review books, notes, coffee, snacks, etc). If you're the kind of person who cannot study alone, form a study group with other students taking the exam. The main point here is to create a comfortable environment with minimal distractions.

(c) ketabton.com: The Digital Library

Year(s) Prior

The knowledge you gained during your first two years of medical school and even during your undergraduate years should provide the groundwork on which to base your test preparation. Student scores on NBME subject tests (commonly known as "shelf exams") have been shown to be highly correlated with subsequent Step 1 scores.[16] Moreover, undergraduate science GPAs as well as MCAT scores are strong predictors of performance on the Step 1 exam.[17]

▸ *Buy review books early (first year) and use while studying for courses.*

We also recommend that you buy highly rated review books early in your first year of medical school and use them as you study throughout the two years. When Step 1 comes along, these books will be familiar and personalized to the way in which you learn. It is risky and intimidating to use unfamiliar review books in the final two or three weeks preceding the exam. Some students find it helpful to personalize and annotate *First Aid* throughout the curriculum.

Months Prior

Review test dates and the application procedure. Testing for the USMLE Step 1 is done on a year-round basis. If you have disabilities or special circumstances, contact the NBME as early as possible to discuss test accommodations (see the Section I Supplement at www.firstaidteam.com/bonus).

▸ *Simulate the USMLE Step 1 under "real" conditions before beginning your studies.*

Use this time to finalize your ideal schedule. Consider upcoming breaks and whether you want to relax or study. Work backward from your test date to make sure you finish at least one question bank. Also add time to redo missed or flagged questions (which may be half the bank). This is the time to build a structured plan with enough flexibility for the realities of life.

Begin doing blocks of questions from reputable question banks under "real" conditions. Don't use tutor mode until you're sure you can finish blocks in the allotted time. It is important to continue balancing success in your normal studies with the Step 1 test preparation process.

Weeks Prior (Dedicated Preparation)

▸ *In the final two weeks, focus on review, practice questions, and endurance. Stay confident!*

Your dedicated prep time may be one week or two months. You should have a working plan as you go into this period. Finish your schoolwork strong, take a day off, and then get to work. Start by simulating a full-length USMLE Step 1 if you haven't yet done so. Consider doing one NBME CBSSA and the free questions from the NBME website. Alternatively, you could choose 7 blocks of randomized questions from a commercial question bank. Make sure you get feedback on your strengths and weaknesses and adjust your studying accordingly. Many students study from review sources or comprehensive programs for part of the day, then do question blocks. Also, keep in mind that reviewing a question block can take upward of two hours. Feedback from CBSSA exams and question banks will help you focus on your weaknesses.

(c) ketabton.com: The Digital Library

One Week Prior

Make sure you have your CIN (found on your scheduling permit) as well as other items necessary for the day of the examination, including a current driver's license or another form of photo ID with your signature (make sure the name on your **ID exactly** matches that on your scheduling permit). Confirm the Prometric testing center location and test time. Work out how you will get to the testing center and what parking and traffic problems you might encounter. Drive separately from other students taking the test on the same day, and exchange cell phone numbers in case of emergencies. If possible, visit the testing site to get a better idea of the testing conditions you will face. Determine what you will do for lunch. Make sure you have everything you need to ensure that you will be comfortable and alert at the test site. It may be beneficial to adjust your schedule to start waking up at the same time that you will on your test day. And of course, make sure to maintain a healthy lifestyle and get enough sleep.

- *One week before the test:*
  - *Sleep according to the same schedule you'll use on test day*
  - *Review the CBT tutorial one last time*
  - *Call Prometric to confirm test date and time*

One Day Prior

Try your best to relax and rest the night before the test. Double-check your admissions and test-taking materials as well as the comfort measures discussed earlier so that you will not have to deal with such details on the morning of the exam. At this point it will be more effective to review short-term memory material that you're already familiar with than to try to learn new material. The Rapid Review section at the end of this book is high yield for last-minute studying. Remember that regardless of how hard you have studied, you cannot know everything. There will be things on the exam that you have never even seen before, so do not panic. Do not underestimate your abilities.

Many students report difficulty sleeping the night prior to the exam. This is often exacerbated by going to bed much earlier than usual. Do whatever it takes to ensure a good night's sleep (eg, massage, exercise, warm milk, no back-lit screens at night). Do not change your daily routine prior to the exam. Exam day is not the day for a caffeine-withdrawal headache.

Morning of the Exam

- *No notes, books, calculators, pagers, cell phones, recording devices, or watches of any kind are allowed in the testing area, but they are allowed in lockers.*

On the morning of the Step 1 exam, wake up at your regular time and eat a normal breakfast. If you think it will help you, have a close friend or family member check to make sure you get out of bed. Make sure you have your scheduling permit admission ticket, test-taking materials, and comfort measures as discussed earlier. Wear loose, comfortable clothing. Plan for a variable temperature in the testing center. Arrive at the test site 30 minutes before the time designated on the admission ticket; however, do not come too early, as doing so may intensify your anxiety. When you arrive at the test site, the proctor should give you a USMLE information sheet that will explain critical factors such as the proper use of break time. Seating may be assigned, but ask to be reseated if necessary; you need to be seated in an area

(c) ketabton.com: The Digital Library

Arrive at the testing center 30 minutes before your scheduled exam time. If you arrive more than half an hour late, you will not be allowed to take the test.

that will allow you to remain comfortable and to concentrate. Get to know your testing station, especially if you have never been in a Prometric testing center before. Listen to your proctors regarding any changes in instructions or testing procedures that may apply to your test site.

Finally, remember that it is natural (and even beneficial) to be a little nervous. Focus on being mentally clear and alert. Avoid panic. When you are asked to begin the exam, take a deep breath, focus on the screen, and then begin. Keep an eye on the timer. Take advantage of breaks between blocks to stretch, maybe do some jumping jacks, and relax for a moment with deep breathing or stretching.

After the Test

After you have completed the exam, be sure to have fun and relax regardless of how you may feel. Taking the test is an achievement in itself. Remember, you are much more likely to have passed than not. Enjoy the free time you have before your clerkships. Expect to experience some "reentry" phenomena as you try to regain a real life. Once you have recovered sufficiently from the test (or from partying), we invite you to send us your feedback, corrections, and suggestions for entries, facts, mnemonics, strategies, resource ratings, and the like (see p. xvii, How to Contribute). Sharing your experience will benefit fellow medical students and IMGs.

STUDY MATERIALS

Quality Considerations

Although an ever-increasing number of review books and software are now available on the market, the quality of such material is highly variable. Some common problems are as follows:

- Certain review books are too detailed to allow for review in a reasonable amount of time or cover subtopics that are not emphasized on the exam.
- Many sample question books were originally written years ago and have not been adequately updated to reflect recent trends.
- Some question banks test to a level of detail that you will not find on the exam.

If a given review book is not working for you, stop using it no matter how highly rated it may be or how much it costs.

Review Books

In selecting review books, be sure to weigh different opinions against each other, read the reviews and ratings in Section IV of this guide, examine the books closely in the bookstore, and choose carefully. You are investing not only money but also your limited study time. Do not worry about finding the "perfect" book, as many subjects simply do not have one, and different students prefer different formats. Supplement your chosen books with personal notes from other sources, including what you learn from question banks.

(c) ketabton.com: The Digital Library

There are two types of review books: those that are stand-alone titles and those that are part of a series. Books in a series generally have the same style, and you must decide if that style works for you. However, a given style is not optimal for every subject.

Charts and diagrams may be the best approach for physiology and biochemistry, whereas tables and outlines may be preferable for microbiology.

You should also find out which books are up to date. Some recent editions reflect major improvements, whereas others contain only cursory changes. Take into consideration how a book reflects the format of the USMLE Step 1.

Apps

With the explosion of smartphones and tablets, apps are an increasingly popular way to review for the Step 1 exam. The majority of apps are compatible with both iOS and Android. Many popular Step 1 review resources (eg, UWorld, USMLE-Rx) have apps that are compatible with their software. Many popular web references (eg, UpToDate) also now offer app versions. All of these apps offer flexibility, allowing you to study while away from a computer (eg, while traveling).

Practice Tests

Taking practice tests provides valuable information about potential strengths and weaknesses in your fund of knowledge and test-taking skills. Some students use practice examinations simply as a means of breaking up the monotony of studying and adding variety to their study schedule, whereas other students rely almost solely on practice. You should also subscribe to one or more high-quality question banks. In addition, students report that many current practice-exam books have questions that are, on average, shorter and less clinically oriented than those on the current USMLE Step 1.

Most practice exams are shorter and less clinical than the real thing.

Additionally, some students preparing for the Step 1 exam have started to incorporate case-based books intended primarily for clinical students on the wards or studying for the Step 2 CK exam. *First Aid Cases for the USMLE Step 1* aims to directly address this need.

After taking a practice test, spend time on each question and each answer choice whether you were right or wrong. There are important teaching points in each explanation. Knowing why a wrong answer choice is incorrect is just as important as knowing why the right answer is correct. Do not panic if your practice scores are low as many questions try to trick or distract you to highlight a certain point. Use the questions you missed or were unsure about to develop focused plans during your scheduled catch-up time.

Use practice tests to identify concepts and areas of weakness, not just facts that you missed.

Textbooks and Course Syllabi

Limit your use of textbooks and course syllabi for Step 1 review. Many textbooks are too detailed for high-yield review and include material that is generally not tested on the USMLE Step 1 (eg, drug dosages, complex chemical structures). Syllabi, although familiar, are inconsistent across

(c) ketabton.com: The Digital Library

medical schools and frequently reflect the emphasis of individual faculty, which often does not correspond to that of the USMLE Step 1. Syllabi also tend to be less organized than top-rated books and generally contain fewer diagrams and study questions.

TEST-TAKING STRATEGIES

Practice! Develop your test-taking skills and strategies well before the test date.

Your test performance will be influenced by both your knowledge and your test-taking skills. You can strengthen your performance by considering each of these factors. Test-taking skills and strategies should be developed and perfected well in advance of the test date so that you can concentrate on the test itself. We suggest that you try the following strategies to see if they might work for you.

Pacing

Time management is an important skill for exam success.

You have seven hours to complete up to 280 questions. Note that each one-hour block contains up to 40 questions. This works out to approximately 90 seconds per question. We recommend following the "1 minute rule" to pace yourself. Spend no more than 1 minute on each question. If you are still unsure about the answer after this time, mark the question, make an educated guess, and move on. Following this rule, you should have approximately 20 minutes left after all questions are answered, which you can use to revisit all of your marked questions. Remember that some questions may be experimental and do not count for points (and reassure yourself that these experimental questions are the ones that are stumping you). In the past, pacing errors have been detrimental to the performance of even highly prepared examinees. The bottom line is to keep one eye on the clock at all times!

Dealing with Each Question

There are several established techniques for efficiently approaching multiple choice questions; find what works for you. One technique begins with identifying each question as easy, workable, or impossible. Your goal should be to answer all easy questions, resolve all workable questions in a reasonable amount of time, and make quick and intelligent guesses on all impossible questions. Most students read the stem, think of the answer, and turn immediately to the choices. A second technique is to first skim the answer choices to get a context, then read the last sentence of the question (the lead-in), and then read through the passage quickly, extracting only information relevant to answering the question. This can be particularly helpful for questions with long clinical vignettes. Try a variety of techniques on practice exams and see what works best for you. If you get overwhelmed, remember that a 30-second time out to refocus may get you back on track.

(c) ketabton.com: The Digital Library

Guessing

There is **no penalty** for wrong answers. Thus, **no test block should be left with unanswered questions.** A hunch is probably better than a random guess. If you have to guess, we suggest selecting an answer you recognize over one with which you are totally unfamiliar.

Changing Your Answer

The conventional wisdom is not to change answers that you have already marked unless there is a convincing and logical reason to do so—in other words, go with your "first hunch." Many question banks tell you how many questions you changed from right to wrong, wrong to wrong, and wrong to right. Use this feedback to judge how good a second-guesser you are. If you have extra time, reread the question stem and make sure you didn't misinterpret the question.

Go with your first hunch, unless you are certain that you are a good second-guesser.

CLINICAL VIGNETTE STRATEGIES

In recent years, the USMLE Step 1 has become increasingly clinically oriented. This change mirrors the trend in medical education toward introducing students to clinical problem solving during the basic science years. The increasing clinical emphasis on Step 1 may be challenging to those students who attend schools with a more traditional curriculum.

Be prepared to read fast and think on your feet!

What Is a Clinical Vignette?

A clinical vignette is a short (usually paragraph-long) description of a patient, including demographics, presenting symptoms, signs, and other information concerning the patient. Sometimes this paragraph is followed by a brief listing of important physical findings and/or laboratory results. The task of assimilating all this information and answering the associated question in the span of one minute can be intimidating. So be prepared to read quickly and think on your feet. Remember that the question is often indirectly asking something you already know.

Practice questions that include case histories or descriptive vignettes are critical for Step 1 preparation.

Strategy

Remember that Step 1 vignettes usually describe diseases or disorders in their most classic presentation. So look for cardinal signs (eg, malar rash for SLE or nuchal rigidity for meningitis) in the narrative history. Be aware that the question will contain classic signs and symptoms instead of buzzwords. Sometimes the data from labs and the physical exam will help you confirm or reject possible diagnoses, thereby helping you rule answer choices in or out. In some cases, they will be a dead giveaway for the diagnosis.

Step 1 vignettes usually describe diseases or disorders in their most classic presentation.

(c) ketabton.com: The Digital Library

Making a diagnosis from the history and data is often not the final answer. Not infrequently, the diagnosis is divulged at the end of the vignette, after you have just struggled through the narrative to come up with a diagnosis of your own. The question might then ask about a related aspect of the diagnosed disease. Consider skimming the answer choices and lead-in before diving into a long stem. However, be careful with skimming the answer choices; going too fast may warp your perception of what the vignette is asking.

IF YOU THINK YOU FAILED

After the test, many examinees feel that they have failed, and most are at the very least unsure of their pass/fail status. There are several sensible steps you can take to plan for the future in the event that you do not achieve a passing score. First, save and organize all your study materials, including review books, practice tests, and notes. Familiarize yourself with the reapplication procedures for Step 1, including application deadlines and upcoming test dates.

If you pass Step 1 (score of 194 or above), you are not allowed to retake the exam.

Make sure you know both your school's and the NBME's policies regarding retakes. The NBME allows a maximum of six attempts to pass each Step examination.[18] You may take Step 1 no more than three times within a 12-month period. Your fourth and subsequent attempts must be at least 12 months after your first attempt at that exam and at least six months after your most recent attempt at that exam.

The performance profiles on the back of the USMLE Step 1 score report provide valuable feedback concerning your relative strengths and weaknesses. Study these profiles closely. Set up a study timeline to strengthen gaps in your knowledge as well as to maintain and improve what you already know. Do not neglect high-yield subjects. It is normal to feel somewhat anxious about retaking the test, but if anxiety becomes a problem, seek appropriate counseling.

TESTING AGENCIES

- **National Board of Medical Examiners (NBME) / USMLE Secretariat**
  Department of Licensing Examination Services
  3750 Market Street
  Philadelphia, PA 19104-3102
  (215) 590-9500 (operator) or
  (215) 590-9700 (automated information line)
  Email: webmail@nbme.org
  www.nbme.org

(c) ketabton.com: The Digital Library

- **Educational Commission for Foreign Medical Graduates (ECFMG)**
  3624 Market Street
  Philadelphia, PA 19104-2685
  (215) 386-5900
  Email: info@ecfmg.org
  www.ecfmg.org

REFERENCES

1. United States Medical Licensing Examination. Available from: https://www.usmle.org/pdfs/step-1/content_step1.pdf. Accessed October 17, 2019.
2. United States Medical Licensing Examination. 2018 Performance Data. Available from: https://www.usmle.org/performance-data/default.aspx#2018_step-1. Accessed October 17, 2019.
3. Prober CG, Kolars JC, First LR, et al. A plea to reassess the role of United States Medical Licensing Examination Step 1 scores in residency selection. *Acad Med.* 2016;91(1):12–15.
4. Roediger HL, Butler AC. The critical role of retrieval practice in long-term retention. *Trends Cogn Sci.* 2011;15(1):20–27.
5. Dunlosky J, Rawson KA, Marsh EJ, et al. Improving students' learning with effective learning techniques: promising directions from cognitive and educational psychology. *Psychol Sci Publ Int.* 2013;14(1):4–58.
6. Larsen DP, Butler AC, Lawson AL, et al. The importance of seeing the patient: test-enhanced learning with standardized patients and written tests improves clinical application of knowledge. *Adv Health Sci Educ.* 2013;18(3):409–425.
7. Panus PC, Stewart DW, Hagemeier NE, et al. A subgroup analysis of the impact of self-testing frequency on examination scores in a pathophysiology course. *Am J Pharm Educ.* 2014;78(9):165.
8. Deng F, Gluckstein JA, Larsen DP. Student-directed retrieval practice is a predictor of medical licensing examination performance. *Perspect Med Educ.* 2015;4(6):308–313.
9. McAndrew M, Morrow CS, Atiyeh L, et al. Dental student study strategies: are self-testing and scheduling related to academic performance? *J Dent Educ.* 2016;80(5):542–552.
10. Augustin M. How to learn effectively in medical school: test yourself, learn actively, and repeat in intervals. *Yale J Biol Med.* 2014;87(2):207–212.
11. Bellezza FS. Mnemonic devices: classification, characteristics, and criteria. *Rev Educ Res.* 1981;51(2):247–275.
12. Dyer J-O, Hudon A, Montpetit-Tourangeau K, et al. Example-based learning: comparing the effects of additionally providing three different integrative learning activities on physiotherapy intervention knowledge. *BMC Med Educ.* 2015;15:37.
13. Chamberland M, Mamede S, St-Onge C, et al. Self-explanation in learning clinical reasoning: the added value of examples and prompts. *Med Educ.* 2015;49(2):193–202.
14. Nesbit JC, Adesope OO. Learning with concept and knowledge maps: a meta-analysis. *Rev Educ Res.* 2006;76(3):413–448.

(c) ketabton.com: The Digital Library

15. Pohl CA, Robeson MR, Hojat M, et al. Sooner or later? USMLE Step 1 performance and test administration date at the end of the second year. *Acad Med.* 2002;77(10):S17–S19.
16. Holtman MC, Swanson DB, Ripkey DR, et al. Using basic science subject tests to identify students at risk for failing Step 1. *Acad Med.* 2001;76(10):S48–S51.
17. Basco WT, Way DP, Gilbert GE, et al. Undergraduate institutional MCAT scores as predictors of USMLE Step 1 performance. *Acad Med.* 2002;77(10):S13–S16.
18. United States Medical Licensing Examination. 2019 USMLE Bulletin of Information. Available from: https://www.usmle.org/pdfs/bulletin/2019bulletin.pdf. Accessed July 23, 2018.

(c) ketabton.com: The Digital Library

SECTION I SUPPLEMENT

Special Situations

Please visit **www.firstaidteam.com/bonus/** to view this section.

- First Aid for the International Medical Graduate 2
- First Aid for the Osteopathic Medical Student 13
- First Aid for the Podiatric Medical Student 17
- First Aid for the Student Requiring Test Accommodations 20

FAS1_2019_00_Section_I.indd 27 11/7/19 2:48 PM

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

SECTION I SUPPLEMENT

Special Situations

- First Aid for the International Medical Graduate 2
- First Aid for the Osteopathic Medical Student 13
- First Aid for the Podiatric Medical Student 17
- First Aid for the Student Requiring Test Accommodations 20

FAS1_2019_00_Section_I_Suppl_ONLINE.indd 1 11/8/19 7:43 AM

(c) ketabton.com: The Digital Library

FIRST AID FOR THE INTERNATIONAL MEDICAL GRADUATE

▸ *IMGs make up approximately 25% of the US physician population.*

"International medical graduate" (IMG) is the term used to describe any student or graduate of a non-US, non-Canadian, non-Puerto Rican medical school, regardless of whether he or she is a US citizen/resident or not.

IMG's Steps to Licensure in the United States

▸ *More detailed information can be found in the ECFMG* Information Booklet, *available at www.ecfmg.org/pubshome.html.*

To be eligible to take the USMLE Steps, you (the applicant) must be officially enrolled in a medical school located outside the United States and Canada that is listed in the World Directory of Medical Schools (WDOMS; www.wdoms.org) and meet the ECFMG eligibility requirements, both at the time you apply for examination and on your test day. In addition, your "Graduation Year" must be listed as "Current" at the time you apply and on your test day.

If you are an IMG, you must go through the following steps (not necessarily in this order) to apply for residency programs and become licensed to practice in the United States. You must complete these steps even if you are already a practicing physician and have completed a residency program in your own country.

▸ *Applicants may apply online for USMLE Step 1, Step 2 CK, or Step 2 CS at www.ecfmg.org.*

- Pass USMLE Step 1, Step 2 CK, and Step 2 CS, as well as obtain a medical school diploma (not necessarily in this order). All three exams can be taken during medical school. If you have already graduated prior to taking any of the Steps, then you will need to verify your academic credentials (confirmation of enrollment and medical degree) prior to applying for any Step exam.
- You will be certified electronically by the Educational Commission for Foreign Medical Graduates (ECFMG) after above steps are successfully completed. You should receive your formal ECFMG certificate in the mail within the next 1–2 weeks. The ECFMG will not issue a certificate (even if all the USMLE scores are submitted) until it verifies your medical diploma with your medical school.
- You must have a valid ECFMG certificate before entering an accredited residency program in the United States, although you can begin the Electronic Residency Application Service (ERAS) application and interviews before you receive the certificate.
- Apply for residency positions in your fields of interest, either directly or through the ERAS and the National Residency Matching Program (NRMP), otherwise known as "the Match." To be entered into the Match, you need to have passed all the examinations necessary for ECFMG certification (ie, Step 1, Step 2 CK, and Step 2 CS) by the rank order list deadline (usually in late February before the Match). If you do not pass these exams by the deadline, you will be withdrawn from the Match.

(c) ketabton.com: The Digital Library

- If you are not a US citizen or green-card holder (permanent resident), you will need to obtain a visa that will allow you to enter and work in the United States after you have matched successfully.
- Sign up to receive the ECFMG and ERAS email newsletter to keep up to date with their most current policies and deadlines.
- If required by the state in which your residency program is located, obtain an educational/training/limited medical license. Your residency program may assist you with this application. Note that medical licensing is the prerogative of each individual state, not of the federal government, and that states vary with respect to their laws about licensing.
- Once you have the ECFMG certification, take the USMLE Step 3 during your residency, and then obtain a full medical license. Once you have a state-issued license, you are permitted to practice in federal institutions such as Veterans Affairs (VA) hospitals and Indian Health Service facilities in any state. This can open the door to "moonlighting" opportunities and possibilities for an H1B visa application if relevant. For details on individual state rules, write to the licensing board in the state in question or contact the Federation of State Medical Boards (FSMB). If you need to apply for an H1B visa for starting residency, you need to first take and pass the USMLE Step 3 exam, preferably before you Match. However, you will be able to apply for and take the USMLE Step 3 exam only after you graduate from medical school.
- Complete your residency and then take the appropriate specialty board exams if you wish to become board certified (eg, in internal medicine or surgery). If you already have a specialty certification in another country, some specialty boards may grant you six months' or one year's credit toward your total residency time.
- Currently, most residency programs are accepting applications through ERAS. For more information, see *First Aid for the Match* or contact:

  ECFMG/ERAS Program
  3624 Market Street
  Philadelphia, PA 19104-2685 USA
  (215) 386-5900
  Email: eras-support@ecfmg.org
  www.ecfmg.org/eras

- For detailed information on the USMLE Steps, visit the USMLE website at http://www.usmle.org.

Keep informed by signing up for the ECFMG email newsletter at www.ecfmg.org/resources.

The USMLE and the IMG

The USMLE is a series of standardized exams that give IMGs and US medical graduates a level playing field. The passing marks for IMGs for Step 1, Step 2 CK, and Step 2 CS are determined by a statistical distribution that is based on the scores of US medical school students. For example, to pass Step 1, you will probably have to score higher than the bottom 8–10% of US and Canadian graduates.

(c) ketabton.com: The Digital Library

IMGs have a maximum of six attempts to pass any USMLE Step, and must pass the USMLE Steps required for ECFMG certification within a seven-year period.

Under USMLE program rules, a maximum of six attempts will be permitted to pass any USMLE Step or component exam. There is a limit of three attempts within a 12-month period for any of the USMLE Steps.

Timing of the USMLE

For an IMG, the timing of a complete application is critical. It is extremely important that you send in your application early if you are to obtain the maximum number of interviews. Complete all exam requirements by August of the year in which you wish to apply. Check the ECFMG website for deadlines to take and pass the various Step exams to be eligible for the NRMP Match.

IMG applicants must pass the USMLE Steps required for ECFMG certification (Step 1, Steps 2 CK and 2 CS) within a seven-year period. The USMLE program recommends, although not all jurisdictions impose, a seven-year limit for completion of the three-step USMLE program.

In terms of USMLE exam order, arguments can be made for taking the Step 1 or the Step 2 CK exam first. For example, you may consider taking the Step 2 CK exam first if you have just graduated from medical school and the clinical topics are still fresh in your mind. However, keep in mind that there is substantial overlap between Step 1 and Step 2 CK topics in areas such as pharmacology, pathophysiology, and biostatistics. You might therefore consider taking the Step 1 and Step 2 CK exams close together to take advantage of this overlap in your test preparation.

USMLE Step 1 and the IMG

Significance of the Test. Step 1 is one of the three exams required for the ECFMG certification. Since most US graduates apply to residency with their Step 1 scores only, it may be the only objective tool available with which to compare IMGs with US graduates.

Signing Up. We advise that you read the FAQ section on the ECFMG website carefully. Most of the services you will need to use involve either IWA or OASIS. If you have not yet completed medical school, follow these steps to sign up for Step 1:

- Apply and pay for an ECFMG/USMLE ID number on the ECFMG website.
- After receiving an email with your ID number, log in to IWA/OASIS, enter your details, and complete the "On-Line part of your USMLE Step 1 application." Choose your test center location and 3-month eligibility period. Additional fees apply if you need to change your eligibility period.
- Pay the Step 1 fee plus any international test surcharges that may apply.
- Access and complete Form 186 (Certification of Identity Form) from IWA as part of the Application for ECFMG Certification.

(c) ketabton.com: The Digital Library

- Follow the instructions on the form to notarize Form 186 using the online service NotaryCam.com. The fee for this service is included in the ECFMG application fee.
- Once notarized by NotaryCam.com and submitted, Form 186 will remain valid indefinitely. A valid, previously completed Form 186 will remain valid for five years from the date it was accepted.
- After receiving a confirmation email from the ECFMG, you may book an exam date and location on www.prometric.com.

Eligibility Period. A three-month period of your choice.

Fee. The fee for Step 1 is $940 plus an international test delivery surcharge (if you choose a testing region other than the United States or Canada).

Statistics. In 2018–2019, 80% of IMG examinees passed Step 1 on their first attempt, compared with 96% of MD degree examinees from the United States and Canada.

Tips. Although few if any students feel totally prepared to take Step 1, IMGs in particular require serious study and preparation in order to reach their full potential on this exam. It is also imperative that IMGs do their best on Step 1, as a poor score on Step 1 is a distinct disadvantage in applying for most residencies. Remember that if you pass Step 1, you cannot retake it in an attempt to improve your score. Your goal should thus be to beat the mean, because you can then assert with confidence that you have done better than average for US students (see Table 1). Higher Step 1 scores will also

A higher Step 1 score will improve your chances of getting into a highly competitive specialty.

TABLE 1. USMLE Step 1 Mean Score of Matched Applicants in 2018.

Specialty	US Graduates	US IMGs	Non-US IMGs
All specialties	233	222	234
Anesthesiology	232	231	240
Dermatology	249	—	238
Diagnostic radiology	240	239	241
Emergency medicine	233	232	229
Family medicine	220	211	220
General surgery	236	237	242
Internal medicine	233	225	236
Neurology	231	227	236
Obstetrics and gynecology	230	229	231
Pathology	233	226	230
Pediatrics	227	221	230
Physical medicine and rehabilitation	225	226	238
Psychiatry	226	214	222

Source: www.nrmp.org.

(c) ketabton.com: The Digital Library

lend credibility to your residency application and help you get into highly competitive specialties such as radiology, orthopedics, and dermatology.

Commercial Review Courses. Do commercial review courses help improve your scores? Reports vary, and such courses can be expensive. For some students these programs can provide a more structured learning environment with professional support. However, review courses consume a significant chunk of time away from independent study. Many IMGs participate in review courses as they typically need higher scores to compete effectively with US and Canadian candidates for residency positions. (For more information on review courses, see Section IV in the book.)

USMLE Step 2 CK and the IMG

What Is the Step 2 CK? It is a computerized test of the clinical sciences consisting of up to 318 multiple-choice questions divided into eight blocks. Each block contains a maximum of 40 questions and needs to be completed within 60 minutes. It can be taken at Prometric centers in the United States and several other countries.

The areas tested on the Step 2 CK relate to the clerkships provided at US medical schools.

Content. The Step 2 CK includes test items in the following content areas:

- Internal medicine
- Obstetrics and gynecology
- Pediatrics
- Preventive medicine
- Psychiatry
- Surgery
- Other areas relevant to the provision of care under supervision

Significance of the Test. The Step 2 CK is required for the ECFMG certificate. It reflects the level of clinical knowledge of the applicant. It tests clinical subjects, primarily internal medicine. Other areas tested are orthopedics, ENT, ophthalmology, safety science, epidemiology, professionalism, and ethics.

Eligibility. Students and graduates from medical schools that are listed in WDOMS and meet the ECFMG eligibility requirement to take the Step 2 CK. Students must have completed at least two years of medical school. This means that students must have completed the basic medical science component of the medical school curriculum by the beginning of the eligibility period selected.

Eligibility Period. A three-month period of your choice.

Fee. The fee for the Step 2 CK is $940 plus an international test delivery surcharge (if you choose a testing region other than the United States or Canada).

(c) ketabton.com: The Digital Library

Statistics. In 2017–2018, 83% of ECFMG candidates passed the Step 2 CK on their first attempt, compared with 97% of MD degree examinees from US and Canadian schools.

Tips. It's better to take the Step 2 CK after your internal medicine rotation because most of the questions on the exam give clinical scenarios and ask you to make medical diagnoses and clinical decisions. In addition, because this is a clinical sciences exam, cultural and geographic considerations play a greater role than is the case with Step 1. For example, if your medical education gave you ample exposure to malaria, brucellosis, and malnutrition but little to alcohol withdrawal, child abuse, and cholesterol screening, you must work to familiarize yourself with topics that are more heavily emphasized in US medicine. You must also have a basic understanding of the legal and social aspects of US medicine, because you will be asked questions about communicating with and advising patients.

Be familiar with topics that are heavily emphasized in US medicine, such as cholesterol screening.

USMLE Step 2 CS and the IMG

What Is the Step 2 CS? The Step 2 CS is a test of clinical and communication skills administered as a one-day, eight-hour exam. It includes 12 encounters with standardized patients (15 minutes each, with 10 minutes to write a note after each encounter).

Content. The Step 2 CS tests the ability to communicate in English as well as interpersonal skills, data-gathering skills, the ability to perform a physical exam, and the ability to formulate a brief note, a differential diagnosis, and a list of diagnostic tests. The areas that are covered in the exam are as follows:

- Internal medicine
- Surgery
- Obstetrics and gynecology
- Pediatrics
- Psychiatry
- Family medicine

Unlike the USMLE Step 1, Step 2 CK, or Step 3, **there are no numerical grades for the Step 2 CS**—it's simply either a "pass" or a "fail." To pass, a candidate must attain a passing performance in **each** of the following three components:

The Step 2 CS is graded as pass/fail.

- Integrated Clinical Encounter (ICE): includes Data Gathering, Physical Exam, and the electronic Patient Note
- Spoken English Proficiency (SEP)
- Communication and Interpersonal Skills (CIS)

According to the NBME, the most commonly failed component for IMGs is the CIS.

(c) ketabton.com: The Digital Library

Significance of the Test. The Step 2 CS assesses spoken English language proficiency and is required for the ECFMG certificate. The Test of English as a Foreign Language (TOEFL) is no longer required.

Eligibility. Students must have completed at least two years of medical school in order to take the test. That means students must have completed the basic medical science component of the medical school curriculum at the time they apply for the exam.

Fee. The fee for the Step 2 CS is $1580.

Statistics. In 2017–2018, 75% of ECFMG candidates passed the Step 2 CS on their first attempt, compared with 95% of MD degree examinees from US and Canadian schools.

▸ *Try to take the Step 2 CS the year before you plan to Match.*

Scheduling. You must schedule the Step 2 CS within **four months** of the date indicated on your notification of registration. You must take the exam within 12 months of the date indicated on your notification of registration. It is generally advisable to take the Step 2 CS as soon as possible in the year before your Match, as often the results either come in late or arrive too late to allow you to retake the test and pass it before the Match.

Test Site Locations. The Step 2 CS is currently administered at the following five locations:

- Philadelphia, PA
- Atlanta, GA
- Los Angeles, CA
- Chicago, IL
- Houston, TX

For more information about the Step 2 CS exam, please refer to *First Aid for the Step 2 CS*.

USMLE Step 3 and the IMG

What Is the USMLE Step 3? It is a two-day computerized test in clinical medicine consisting of 413 multiple-choice questions and 13 computer-based case simulations (CCS). The exam aims to test your knowledge and its application to patient care and clinical decision making (ie, this exam tests if you can safely practice medicine independently and without supervision). Please go to the USMLE website to learn more about recent changes to the exam.

▸ *Complete the Step 3 exam before you apply for an H1B visa.*

Significance of the Test. Taking Step 3 before residency is critical for IMGs seeking an H1B visa and is also a bonus that can be added to the residency application. Step 3 is also required to obtain a full medical license in the United States and can be taken during residency for this purpose.

Fee. The fee for Step 3 is $895.

(c) ketabton.com: The Digital Library

Eligibility. Examinees are no longer required to apply to the Step 3 exam under the eligibility requirements of a specific medical licensing authority. Those wishing to sit for the Step 3 exam, independent of the place of residence, must meet the following requirements:

- Have completed an MD or DO degree from an LCME- or AOA-accredited US or Canadian medical school, or from a medical school outside the US and Canada listed in the World Directory of Medical Schools.
- Have taken and passed the Step 1, Step 2 CK, and Step 2 CS exams.
- If an IMG, be certified by the ECFMG.

The Step 3 exam is not available outside the United States. Applications can be found online at www.fsmb.org and must be submitted to the FSMB.

Statistics. In 2018, 90% of IMG candidates passed the Step 3 on their first attempt, compared with 98% of MD degree examinees from US and Canadian schools.

Residencies and the IMG

In the Match, the number of US-citizen IMG applications has grown over the past few years, while the percentage accepted has remained constant (see Table 2). More information about residency programs can be obtained at www.ama-assn.org.

The Match and the IMG

Given the growing number of IMG candidates with strong applications, you should bear in mind that good USMLE scores are not the only way to gain a competitive edge. However, USMLE Step 1 and Step 2 CK scores continue to be used as the initial screening mechanism when candidates are being considered for interviews.

TABLE 2. IMGs in the Match.

Applicants	2016	2017	2018	2019
US-citizen IMGs	5,323	5,069	5,075	5,080
% US-citizen IMGs accepted	53.9	54.8	57.1	59
Non-US-citizen IMGs	7,460	7,284	7,067	6,869
% non-US-citizen IMGs accepted	50.5	52.4	56.1	58.6
US seniors (non-IMGs)	18,187	18,539	18,818	18,925
% US seniors accepted	93.8	94.3	94.3	93.9
DO graduates		3,590	4,617	6,001
% DO graduates accepted		81.7	81.7	84.6

Source: www.nrmp.org.

(c) ketabton.com: The Digital Library

Based on accumulated IMG Match experiences over recent years, here are a few pointers to help IMGs maximize their chances for a residency interview:

- **Apply early.** Programs offer a limited number of interviews and often select candidates on a first-come, first-served basis. Because of this, you should aim to complete the entire process of applying for the ERAS token, registering with the Association of American Medical Colleges (AAMC), mailing necessary documents to ERAS, and completing the ERAS application by mid-September (see Figure 1). Community programs usually send out interview offers earlier than do university and university-affiliated programs.

All US hospitals allow externship only when the applicant is actively enrolled in a medical school, so plan ahead.

- **US clinical experience helps.** Externships and observerships in a US hospital setting have emerged as an important credential on an IMG application. Externships are like short-term medical school internships and offer hands-on clinical experience. Observerships, also called "shadowing," involve following a physician and observing how he or she manages patients. Some programs require students to have participated in an externship or observership before applying. It is best to gain such an experience before or at the time you apply to various programs so that you can mention it on your

FIGURE 1. IMG Timeline for Application.



(c) ketabton.com: The Digital Library

ERAS application. If such an experience or opportunity comes up after you apply, be sure to inform the programs accordingly.

- **Clinical research helps.** University programs are attracted to candidates who show a strong interest in clinical research and academics. They may even relax their application criteria for individuals with unique backgrounds and strong research experience. Publications in well-known journals are an added bonus.
- **Time the Step 2 CS well.** ECFMG has published the new Step 2 CS score-reporting schedule for 2019–2020 at http://www.ecfmg.org. Most program directors would like to see a passing score on the Step 1, Step 2 CK, and Step 2 CS exams before they rank an IMG on their rank order list in mid-February. There have been many instances in which candidates have lost a potential Match—either because of delayed CS results or because they have been unable to retake the exam on time following a failure. It is difficult to predict a result on the Step 2 CS, since the grading process is not very transparent. Therefore, it is advisable to take the Step 2 CS as early as possible in the application year.
- **US letters of recommendation help.** Letters of recommendation from clinicians practicing in the United States carry more weight than recommendations from home countries.
- **Step up the Step 3.** If H1B visa sponsorship is desired, aim to have Step 3 results by January of the Match year. In addition to the visa advantage you will gain, an early and good Step 3 score may benefit IMGs who have been away from clinical medicine for a while as well as those who have low scores on Step 1 and the Step 2 CK. Note that the Step 3 can be taken only after medical school graduation.

A good score on the Step 3 may help offset poorer scores on the Step 1 or 2 CK exams.

- **Verify medical credentials in a timely manner.** Do not overlook the medical school credential verification process. The ECFMG certificate arrives only after credentials have been verified and after you have passed Step 1, the Step 2 CK, and the Step 2 CS, so you should keep track of the process and check their application status online using IWA/OASIS.
- **Don't count on a pre-Match.** Programs participating in NRMP Match can no longer offer a pre-Match.

What if You Do Not Match?

For applicants who do not Match into a residency program, there's SOAP (Supplemental Offer and Acceptance Program). Under SOAP, unmatched applicants will have access to the list of unfilled programs at noon Eastern time on the Monday of Match week. The unfilled programs electing to participate in SOAP will offer positions to unmatched applicants through the Registration, Ranking, and Results (R3) system. A series of "rounds" will begin at noon Eastern time on Wednesday of Match week until 5:00 PM Eastern time on Friday of Match week. Detailed information about SOAP can be found at the NRMP website at http://www.nrmp.org.

(c) ketabton.com: The Digital Library

Resources for the IMG

- **Educational Commission for Foreign Medical Graduates (ECFMG)**
  3624 Market Street
  Philadelphia, PA 19104-2685
  (215) 386-5900
  Fax: (215) 386-9196
  Email: info@ecfmg.org
  www.ecfmg.org

  The ECFMG telephone number is answered only between 9:00 AM–5:00 PM Monday through Friday EST. The ECFMG often takes a long time to answer the phone, which is frequently busy at peak times of the year, and then gives you a long voice-mail message—so it is better to email early than to rely on a last-minute phone call. When contacting the ECFMG by email, include your USMLE/ECFMG Identification Number and use the email address that you registered with the ECFMG. Do not contact the NBME, as all IMG exam matters are conducted by the ECFMG. The ECFMG also publishes an information booklet on ECFMG certification and the USMLE program, which gives details on the dates and locations of forthcoming Step tests for IMGs together with application forms. The *Information Booklet* is available to view and download on the ECFMG's website at www.ecfmg.org, where they also have a complete list of fees for certification posted (see Table 3).

TABLE 3. Estimated Costs for IMGs (as of 2019).

Exams and Services	Fee(s)
USMLE Step 1	$940 + international surcharge (eg, $195 in all European countries offering the exam)
USMLE Step 2 CK	$940 + international surcharge (eg, $220 in all European countries offering the exam)
USMLE Step 2 CS	$1580
USMLE Step 3	$895
ERAS	$130 registration fee (ECFMG token fee) $80 USMLE transcript assessment $99 for programs 1–10 $15 each for programs 11–20 $19 each for programs 21–30 $26 each for programs 31+
NRMP	$85 registration fee (for ranking 20 programs) $30 per additional program ranked $35 per partner (couples match only) $50 late registration fee (sign up before November 30 to avoid paying this fee)
J-1 visa application fee	$160 visa application fee $340 annual ECFMG application fee $220 payable to Homeland Security (SEVIS fee)

(c) ketabton.com: The Digital Library

- **Federation of State Medical Boards (FSMB)**
  400 Fuller Wiser Road, Suite 300
  Euless, TX 76039-3856
  (817) 868-4041
  Email: usmle@fsmb.org
  www.fsmb.org

  The FSMB has a number of publications available, including free policy documents. All of these documents are available to view and download for free on the FSMB's website at www.fsmb.org. For Step 3 inquiries, the telephone number is (817) 868-4041.

The AMA has dedicated a portion of its website to information on IMG demographics, residencies, immigration, and the like. This information can be found at www.ama-assn.org.

FIRST AID FOR THE OSTEOPATHIC MEDICAL STUDENT

What Is the COMLEX-USA Level 1?

The National Board of Osteopathic Medical Examiners (NBOME) administers the Comprehensive Osteopathic Medical Licensing Examination, or COMLEX-USA. Like the USMLE, the COMLEX-USA is administered over three levels.

The COMLEX-USA series assesses osteopathic medical knowledge and clinical skills using clinical presentations and physician tasks. A description of the COMLEX-USA Written Examination Blueprints for each level, which outline the various clinical presentations and physician tasks that examinees will encounter, is given on the NBOME website. Another stated goal of the COMLEX-USA Level 1 is to create a more primary care–oriented exam that integrates osteopathic principles into clinical situations.

To be eligible to take the COMLEX-USA Level 1, you must be on track to satisfactorily complete your first two years in an AOA-accredited medical school. The office of the dean at each school informs the NBOME that the student will complete the first two years of medical school and is in good standing. At this point, the NBOME sends out an email with detailed instructions on how to register for the exam.

For all three levels of the COMLEX-USA, raw scores are converted to a percentile score and a score ranging from 5 to 800. For Levels 1 and 2, a score of 400 is required to pass; for Level 3, a score of 350 is needed. COMLEX-USA scores are posted at the NBOME website 4–6 weeks after the test and usually mailed within 8 weeks after the test. The mean score is always 500.

(c) ketabton.com: The Digital Library

If you pass a COMLEX-USA examination, you are not allowed to retake it to improve your grade. Currently, if you fail, there is no specific limit to the number of times you can retake it in order to pass. However, a student may not take the exam more than four times in one year. Levels 2 and 3 exams must be passed in sequential order within seven years of passing Level 1.

Note that candidates taking COMLEX-USA examinations will be limited to a total of six attempts for each examination.

What Is the Structure of the COMLEX-USA Level 1?

The COMLEX-USA Level 1 is a computer-based examination consisting of 400 questions over an eight-hour period in a single day (nine hours counting breaks). Most of the questions are in one-best-answer format, but a small number are matching-type questions. Some one-best-answer questions are bundled together around a common question stem that usually takes the form of a clinical scenario. Every section of the COMLEX-USA Level 1 ends with either matching questions, multiple questions around a single stem, or both. New question formats may gradually be introduced, but candidates will be notified if this occurs. Multimedia questions are also included on the exam.

Questions are grouped into eight subsections of 50 questions each in a manner similar to that of the USMLE. The individual subsections are not timed, but the exam is divided into two blocks consisting of four subsections. Each subsection consists of 200 questions to be completed within four hours. Reviewing and changing answers may be done only in the current subsection. A "review page" is presented for each subsection in order to advise test takers of questions completed, questions marked for further review, and incomplete questions for which no answer has been given.

Breaks are even more structured with COMLEX-USA than they are with the USMLE. Students are allowed to take an optional 10-minute break at the end of the second and sixth subsections. After subsection 4, students are given a 40-minute lunch break. These are the only times a student is permitted a break. Any unused break time will not be added to the time allotted for taking the examination. More information about the computer-based COMLEX-USA examinations can be obtained from www.nbome.org.

What Is the Difference Between the USMLE and the COMLEX-USA?

According to the NBOME, the COMLEX-USA Level 1 focuses broadly on the following categories, with osteopathic principles and practices integrated into each section:

- Health promotion and disease prevention
- The history and physical
- Diagnostic technologies

(c) ketabton.com: The Digital Library

- Management
- Scientific understanding of mechanisms
- Health care delivery

The test interface for the COMLEX-USA Level 1 is not the same as the USMLE Step 1 interface.

Although the COMLEX-USA and the USMLE are similar in scope, content, and emphasis, some differences are worth noting. For example, the interface is different; you cannot search for lab values. Instead, lab values and reference ranges (where appropriate) are included directly in the clinical vignette or test question. Fewer details are given about a patient's condition, so a savvy student needs to know how to differentiate between similar pathologies. Also, age, gender, and race are key factors for diagnosis on the COMLEX-USA. Images or videos are embedded in the question stem and the examinee has to click an attachment button to see the image. If you don't read the question carefully, the attachment buttons are very easy to miss. A standard calculator feature is embedded in the examination interface.

COMLEX-USA Level 1 tests osteopathic principles in addition to basic science materials but does not emphasize lab techniques. Although both exams often require that you apply and integrate knowledge over several areas of basic science to answer a given question, many students who took both tests reported that the questions differed somewhat in style. Students reported, for example, that USMLE questions generally required that the test taker reason and draw from the information given (often a two-step process), whereas those on the COMLEX-USA exam tended to be more straightforward and that multiple different questions are asked pertaining to one question stem.

COMLEX-USA test takers can expect to have only a few questions on biochemistry, molecular biology, or lab technique. On the other hand, microbiology is very heavily tested by clinical presentation and by lab identification. The COMLEX-USA exam also focuses more on disease management, specific legal principles (eg, Tarasoff case and the Emergency Treatment Act) and more detailed ethical principles (eg, res ipsa loquitur) than the USMLE Step 1. Another main difference is that the COMLEX-USA exam stresses osteopathic manipulative medicine. Therefore, question banks specific to the USMLE will not be adequate, and supplementation with a question bank specific to the COMLEX-USA is highly recommended. The most commonly used are COMBANK or COMQUEST.

Students also commented that the COMLEX-USA utilized "buzzwords," although limited in their use (eg, "rose spots" in typhoid fever), whereas the USMLE avoided buzzwords in favor of descriptions of clinical findings or symptoms (eg, rose-colored papules on the abdomen rather than rose spots). Finally, USMLE appeared to have more photographs than did the COMLEX-USA. In general, the overall impression was that the USMLE was a more "thought-provoking" exam, while the COMLEX-USA was more of a "knowledge-based" exam.

(c) ketabton.com: The Digital Library

Who Should Take Both the USMLE and the COMLEX-USA?

Aside from facing the COMLEX-USA Level 1, you must decide if you will also take the USMLE Step 1. We recommend that you consider taking both the USMLE and the COMLEX-USA under the following circumstances:

- **If you are applying to allopathic residencies.** Although there is growing acceptance of COMLEX-USA certification on the part of allopathic residencies, some allopathic programs prefer or even require passage of the USMLE Step 1. These include many academic programs, programs in competitive specialties (eg, orthopedics, ophthalmology, or dermatology), and programs in competitive geographic areas (eg, Vermont, Utah, and California). Fourth-year osteopathic medical students who have already Matched may be a good source of information about which programs and specialties look for USMLE scores. It is also a good idea to contact program directors at the institutions you are interested in to ask about their policy regarding the COMLEX-USA versus the USMLE.
- **If you are unsure about your postgraduate training plans.** Successful passage of both the COMLEX-USA Level 1 and the USMLE Step 1 is certain to provide you with the greatest possible range of options when you are applying for internship and residency training.

If you're not sure whether you need to take either the COMLEX-USA Level 1 or the USMLE Step 1, consider taking both to keep your Match options open.

In addition, the COMLEX-USA Level 1 has in recent years placed increasing emphasis on questions related to primary care medicine and prevention. Having a strong background in family or primary care medicine can help test takers when they face questions on prevention.

How Do I Prepare for the COMLEX-USA Level 1?

Student experience suggests that you should start studying for the COMLEX-USA four to six months before the test is given, as an early start will allow you to spend up to a month on each subject. The recommendations made in Section I regarding study and testing methods, strategies, and resources, as well as the books suggested in Section IV for the USMLE Step 1, hold true for the COMLEX-USA as well.

Another important source of information is in the *Examination Guidelines and Sample Exam*, a booklet that discusses the breakdown of each subject while also providing sample questions and corresponding answers. Many students, however, felt that this breakdown provided only a general guideline and was not representative of the level of difficulty of the actual COMLEX-USA. The sample questions did not provide examples of clinical vignettes, which made up approximately 25% of the exam. You will receive this publication with registration materials for the COMLEX-USA Level 1, but you can also receive a copy and additional information by writing:

NBOME
8765 W. Higgins Road, Suite 200
Chicago, IL 60631-4174
(773) 714-0622
www.nbome.org

(c) ketabton.com: The Digital Library

The NBOME developed the Comprehensive Osteopathic Medical Self-Assessment Examination (COMSAE) series to fill the need for self-assessment on the part of osteopathic medical students. Many students take the COMSAE exam before the COMLEX-USA in addition to using test-bank questions and board review books. Students can purchase a copy of this exam at www.nbome.org/comsae.asp.

In recent years, students have reported an emphasis in certain areas. For example:

- There was an increased emphasis on upper limb anatomy/brachial plexus.
- Specific topics were repeatedly tested on the exam. These included cardiovascular physiology and pathology, acid-base physiology, diabetes, benign prostatic hyperplasia, sexually transmitted diseases, measles, and rubella. Thyroid and adrenal function, neurology (head injury), specific drug treatments for bacterial infection, migraines/cluster headaches, and drug mechanisms also received heavy emphasis.
- Behavioral science questions were based on psychiatry.
- High-yield osteopathic manipulative technique (OMT) topics included an emphasis on the sympathetic and parasympathetic innervations of viscera and nerve roots, rib mechanics/diagnosis, and basic craniosacral theory. Students who spend time reviewing basic anatomy, studying nerve and dermatome innervations, and understanding how to perform basic OMT techniques (eg, muscle energy or counterstrain) can improve their scores.

You must know the Chapman reflex points and the obscure names of physical exam signs.

COMLEX is heavy on "bugs and drugs."

The COMLEX-USA Level 1 also includes multimedia-based questions. Such questions test the student's ability to perform a good physical exam and to elicit various physical diagnostic signs (eg, Murphy sign).

FIRST AID FOR THE PODIATRIC MEDICAL STUDENT

The National Board of Podiatric Medical Examiners (NBPME) offers the American Podiatric Medical Licensing Examinations (APMLE), which are designed to assess whether a candidate possesses the knowledge required to practice as a minimally competent entry-level podiatric surgeon. The APMLE is used as part of the licensing process governing the practice of podiatric medicine and surgery. The APMLE is recognized by all 50 states and the District of Columbia, the US Army, the US Navy, and the Canadian provinces of Alberta, British Columbia, and Ontario. Individual states use the examination scores differently; therefore, doctor of podiatric medicine (DPM) candidates should refer to the *NBPME Part I and Part II Information Bulletin 2019*.

(c) ketabton.com: The Digital Library

- *Areas tested on the NBPME Part I:*
  - *General anatomy*
  - *Lower extremity anatomy*
  - *Biochemistry*
  - *Physiology*
  - *Medical microbiology & immunology*
  - *Pathology*
  - *Pharmacology*

The APMLE Part I is generally taken after the completion of the second year of podiatric medical education. Unlike the USMLE Step 1, there is no behavioral science section, nor is biomechanics tested. The exam samples seven basic science disciplines: general anatomy (13%); lower extremity anatomy (25%); biochemistry (10%); physiology (13%); microbiology and immunology (13%); pathology (13%); and pharmacology (13%). A detailed outline of topics and subtopics covered on the exam can be found in the *Candidate Information Bulletin Part I Examination*, available at www.apmle.org.

Your APMLE Appointment

Applicants have to register for the exam online at www.prometric.com/NBPME. Once registration is completed, you will receive an Authorization to Test (ATT) email notification that allows you to schedule your exam online. This should be done promptly to secure the testing location and exam date of your choice. The exam will be offered at an independent Prometric testing facility. Test centers within a 50-mile radius of a podiatric medicine school specifically reserve a number of seats on each APMLE Part I exam date. You may take the exam at any Prometric site regardless of which school you attend. Specific instructions about exam dates and registration deadlines can be found in the *Candidate Information Bulletin*.

Exam Format

The APMLE Part I is a written exam consisting of 205 questions. The test consists exclusively of one-best-answer multiple choice questions with four options per question. A review screen showing all answered, unanswered, and marked questions will be available at the end. Students are encouraged to mark questions and return to these for review at the end of the exam if time allows. Examinees have four hours in which to complete the exam and are given scratch paper that must be turned in at the end of the exam. Some questions on the exam will be "trial questions." These questions are evaluated as future board questions but are not counted in your score.

Interpreting Your Score

Exam results are emailed to examinees approximately four weeks after the exam date, and are also available online via the Prometric dashboard. APMLE scores are reported as pass/fail, with a scaled score of at least 75 needed to pass. Historically, 85% of first-time test takers pass the APMLE Part I. Failing candidates receive a report with a score between 55 and 74 in addition to diagnostic messages intended to help identify strengths or weaknesses in specific content areas. If you fail the APMLE Part I, you must retake the entire examination at a later date. There is no limit to the number of times you can retake the exam.

(c) ketabton.com: The Digital Library

Preparation for the APMLE Part I

Begin studying for the APMLE Part I at least three months prior to the test date. The suggestions made in Section I regarding study and testing methods for the USMLE Step 1 can be applied to the APMLE as well. This book should, however, be used as a supplement and not as the sole source of information. Neither you nor your school or future residency will ever see your actual passing numerical score. Competing with colleagues should not be an issue, and study groups are beneficial to many.

A study method that helps many students is to copy the outline of the material to be tested from the *Candidate Information Bulletin*. Check off each topic during your study, because doing so will ensure that you have engaged each topic. If you are pressed for time, prioritize subjects on the basis of their weight on the exam. A full 25% of the APMLE Part I focuses on lower extremity anatomy. In this area, students should rely on the notes and material that they received from their class. Remember, lower extremity anatomy is the podiatric physician's specialty—so everything about it is important. Do not forget to study osteology. Keep your old tests and look through old lower extremity class exams, since each of the podiatric colleges submits questions from its faculty. This strategy will give you an understanding of the types of questions that may be asked. On the APMLE Part I, you will see some of the same classic lower extremity anatomy questions you were tested on in school.

▸ *Know the anatomy of the lower extremity!*

The APMLE, like the USMLE, requires that you apply and integrate knowledge over several areas of basic science in order to answer exam questions. Students report that many questions emphasize clinical presentations; however, the facts in this book are very useful in helping students recall the various diseases and organisms. DPM candidates should expand on the high-yield pharmacology section and study antifungal drugs and treatments for *Pseudomonas*, methicillin-resistant *S aureus*, candidiasis, and erythrasma. The high-yield section focusing on pathology is very useful; however, additional emphasis on diabetes mellitus and all its secondary manifestations, particularly peripheral neuropathy, should not be overlooked. Students should also focus on renal physiology and drug elimination, the biochemistry of gout, and neurophysiology, all of which have been noted to be important topics on the APMLE Part I exam.

A sample set of questions is found on the APMLE website www.apmle.org. These samples are somewhat similar in difficulty to actual board questions. If you have any questions regarding registration, fees, test centers, authorization forms, or score reports, please contact your college registrar or:

Prometric
877-302-8952
Email: nbpmeinquiry@prometric.com
www.prometric.com

(c) ketabton.com: The Digital Library

FIRST AID FOR THE STUDENT REQUIRING TEST ACCOMMODATIONS

The USMLE provides accommodations for students with documented disabilities. The basis for such accommodations is the Americans with Disabilities Act (ADA) of 1990. The ADA defines a disability as "a significant limitation in one or more major life activities." This includes both "observable/physical" disabilities (eg, blindness, hearing loss, narcolepsy) and "hidden/mental disabilities" (eg, attention-deficit hyperactivity disorder, chronic fatigue syndrome, learning disabilities).

US students seeking ADA-compliant accommodations must contact the NBME directly; IMGs, contact the ECFMG.

To provide appropriate support, the administrators of the USMLE must be informed of both the nature and the severity of an examinee's disability. Such documentation is required for an examinee to receive testing accommodations. Accommodations include extra time on tests, low-stimulation environments, extra or extended breaks, and zoom text.

Who Can Apply for Accommodations?

Students or graduates of a school in the United States or Canada that is accredited by the Liaison Committee on Medical Education (LCME) or the AOA may apply for test accommodations directly from the NBME. Requests are granted only if they meet the ADA definition of a disability. If you are a disabled student or a disabled graduate of a foreign medical school, you must contact the ECFMG (see the following page).

Who Is Not Eligible for Accommodations?

Individuals who do not meet the ADA definition of disabled are not eligible for test accommodations. Difficulties not eligible for test accommodations include test anxiety, slow reading without an identified underlying cognitive deficit, English as a second language, and learning difficulties that have not been diagnosed as a medically recognized disability.

Understanding the Need for Documentation

Although most learning-disabled medical students are all too familiar with the often exhausting process of providing documentation of their disability, you should realize that **applying for USMLE accommodation is different from these previous experiences.** This is because the NBME determines whether an individual is disabled solely on the basis of the guidelines set by the ADA. **Previous accommodation does not in itself justify provision of an accommodation for the USMLE,** so be sure to review the NBME guidelines carefully.

(c) ketabton.com: The Digital Library

Getting the Information

The first step in applying for USMLE special accommodations is to contact the NBME and obtain a guidelines and questionnaire booklet. For the Step 1, Step 2 CK, and Step 2 CS exams, this can be obtained by calling or writing to:

Disability Services
National Board of Medical Examiners
3750 Market Street
Philadelphia, PA 19104-3102
(215) 590-9509
Email: disabilityservices@nbme.org
www.usmle.org/test-accommodations

Internet access to this information is also available at www.nbme.org. This information is also relevant for IMGs, since the information is the same as that sent by the ECFMG.

Foreign graduates should contact the ECFMG to obtain information on special accommodations by calling or writing to:

ECFMG
3624 Market Street
Philadelphia, PA 19104-2685
(215) 386-5900
www.ecfmg.org

When you get this information, take some time to read it carefully. The guidelines are clear and explicit about what you need to do to obtain accommodations.

(c) ketabton.com: The Digital Library

SECTION II

High-Yield General Principles

"There comes a time when for every addition of knowledge you forget something that you knew before. It is of the highest importance, therefore, not to have useless facts elbowing out the useful ones."

—Sir Arthur Conan Doyle, *A Study in Scarlet*

"Never regard study as a duty, but as the enviable opportunity to learn."

—Albert Einstein

"Live as if you were to die tomorrow. Learn as if you were to live forever."

—Gandhi

- How to Use the Database 30
- Biochemistry 33
- Immunology 95
- Microbiology 123
- Pathology 205
- Pharmacology 229
- Public Health Sciences 255

FAS1_2019_01-Biochem.indd 29 11/7/19 3:16 PM

(c) ketabton.com: The Digital Library

HOW TO USE THE DATABASE

The 2020 edition of *First Aid for the USMLE Step 1* contains a revised and expanded database of basic science material that students, student authors, and faculty authors have identified as high yield for board review. The information is presented in a partially organ-based format. Hence, Section II is devoted to the foundational principles of biochemistry, microbiology, immunology, basic pathology, basic pharmacology, and public health sciences. Section III focuses on organ systems, with subsections covering the embryology, anatomy and histology, physiology, clinical pathology, and clinical pharmacology relevant to each. Each subsection is then divided into smaller topic areas containing related facts. Individual facts are generally presented in a three-column format, with the **Title** of the fact in the first column, the **Description** of the fact in the second column, and the **Mnemonic** or **Special Note** in the third column. Some facts do not have a mnemonic and are presented in a two-column format. Others are presented in list or tabular form in order to emphasize key associations.

The database structure used in Sections II and III is useful for reviewing material already learned. These sections are **not** ideal for learning complex or highly conceptual material for the first time.

The database of high-yield facts is not comprehensive. Use it to complement your core study material and not as your primary study source. The facts and notes have been condensed and edited to emphasize the essential material, and as a result, each entry is "incomplete" and arguably "over-simplified." Often, the more you research a topic, the more complex it becomes, with certain topics resisting simplification. Work with the material, add your own notes and mnemonics, and recognize that not all memory techniques work for all students.

We update the database of high-yield facts annually to keep current with new trends in boards emphasis, including clinical relevance. However, we must note that inevitably many other high-yield topics are not yet included in our database.

We actively encourage medical students and faculty to submit high-yield topics, well-written entries, diagrams, clinical images, and useful mnemonics so that we may enhance the database for future students. We also solicit recommendations of alternate tools for study that may be useful in preparing for the examination, such as charts, flash cards, apps, and online resources (see How to Contribute, p. xix).

(c) ketabton.com: The Digital Library

Image Acknowledgments

All images and diagrams marked with Rx are © USMLE-Rx.com (MedIQ Learning, LLC) and reproduced here by special permission. All images marked with RU are © Dr. Richard P. Usatine, author of *The Color Atlas of Family Medicine*, *The Color Atlas of Internal Medicine*, and *The Color Atlas of Pediatrics*, and are reproduced here by special permission (www.usatinemedia.com). Images and diagrams marked with * are adapted or reproduced with permission of other sources as listed on page 727. Images and diagrams with no acknowledgment are part of this book.

Disclaimer

The entries in this section reflect student opinions of what is high yield. Because of the diverse sources of material, no attempt has been made to trace or reference the origins of entries individually. We have regarded mnemonics as essentially in the public domain. Errata will gladly be corrected if brought to the attention of the authors, either through our online errata submission form at www.firstaidteam.com or directly by email to firstaid@scholarrx.com.

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

HIGH-YIELD PRINCIPLES IN

Biochemistry

"Biochemistry is the study of carbon compounds that crawl."

—Mike Adams

"We think we have found the basic mechanism by which life comes from life."

—Francis H. C. Crick

"The biochemistry and biophysics are the notes required for life; they conspire, collectively, to generate the real unit of life, the organism."

—Ursula Goodenough

Molecular 34

Cellular 46

Laboratory Techniques 52

Genetics 56

Nutrition 65

Metabolism 72

This high-yield material includes molecular biology, genetics, cell biology, and principles of metabolism (especially vitamins, cofactors, minerals, and single-enzyme-deficiency diseases). When studying metabolic pathways, emphasize important regulatory steps and enzyme deficiencies that result in disease, as well as reactions targeted by pharmacologic interventions. For example, understanding the defect in Lesch-Nyhan syndrome and its clinical consequences is higher yield than memorizing every intermediate in the purine salvage pathway.

Do not spend time learning details of organic chemistry, mechanisms, or physical chemistry. Detailed chemical structures are infrequently tested; however, many structures have been included here to help students learn reactions and the important enzymes involved. Familiarity with the biochemical techniques that have medical relevance—such as ELISA, immunoelectrophoresis, Southern blotting, and PCR—is useful. Review the related biochemistry when studying pharmacology or genetic diseases as a way to reinforce and integrate the material.

FAS1_2019_01-Biochem.indd 33 11/7/19 3:16 PM

(c) ketabton.com: The Digital Library

▶ BIOCHEMISTRY—MOLECULAR

Chromatin structure



DNA exists in the condensed, chromatin form to fit into the nucleus. DNA loops twice around a histone octamer to form a nucleosome ("**beads on a string**"). H1 binds to the nucleosome and to "linker DNA," thereby stabilizing the chromatin fiber.

Phosphate groups give DNA a ⊖ charge. Lysine and arginine give histones a ⊕ charge.

In mitosis, DNA condenses to form chromosomes. DNA and histone synthesis occurs during S phase.

Mitochondria have their own DNA, which is circular and does not utilize histones.

Heterochromatin 	Condensed, appears darker on EM (labeled H in **A**; Nu, nucleolus). Sterically inaccessible, thus transcriptionally inactive. ↑ methylation, ↓ acetylation.	**H**etero**C**hromatin = **H**ighly **C**ondensed. Barr bodies (inactive X chromosomes) may be visible on the periphery of nucleus.
Euchromatin	Less condensed, appears lighter on EM (labeled E in **A**). Transcriptionally active, sterically accessible.	*Eu* = true, "truly transcribed." **E**uchromatin is **E**xpressed.
DNA methylation	Changes the expression of a DNA segment without changing the sequence. Involved with aging, carcinogenesis, genomic imprinting, transposable element repression, and inactivation of the X chromosome.	DNA is methylated in imprinting. Methylation within gene promoter (CpG islands) typically represses (silences) gene transcription. CpG **M**ethylation **M**akes DNA **M**ute.
Histone methylation	Usually causes reversible transcriptional suppression, but can also cause activation depending on location of methyl groups.	Histone **M**ethylation **M**ostly **M**akes DNA **M**ute.
Histone acetylation	Removal of histone's ⊕ charge → relaxed DNA coiling → ↑ transcription.	Histone **A**cetylation makes DNA **A**ctive.
Histone deacetylation	Removal of acetyl groups → tightened DNA coiling → ↓ transcription.	

(c) ketabton.com: The Digital Library

Nucleotides

NucleoSide = base + (deoxy)ribose (Sugar).
NucleoTide = base + (deoxy)ribose + phosphaTe; linked by 3′-5′ phosphodiester bond.

5′ end of incoming nucleotide bears the triphosphate (energy source for the bond). Triphosphate bond is target of 3′ hydroxyl attack.

PURines (A,G)—2 rings.
PYrimidines (C,U,T)—1 ring.

PURe As Gold.
CUT the PY (pie).
Thymine has a methyl.

Deamination reactions:
Cytosine → uracil
Adenine → hypoxanthine
Guanine → xanthine
5-methylcytosine → thymine

C-G bond (3 H bonds) stronger than A-T bond (2 H bonds). ↑ C-G content → ↑ melting temperature of DNA. "C-G bonds are like Crazy Glue."

Uracil found in RNA; thymine in DNA. Methylation of uracil makes thymine.

Amino acids necessary for purine synthesis (cats purr until they GAG):
Glycine
Aspartate
Glutamine



(c) ketabton.com: The Digital Library

De novo pyrimidine and purine synthesis

Various immunosuppressive, antineoplastic, and antibiotic drugs function by interfering with nucleotide synthesis:



Pyrimidine synthesis:

- Leflunomide: inhibits dihydroorotate dehydrogenase
- 5-fluorouracil (5-FU) and its prodrug capecitabine: form 5-F-dUMP, which inhibits thymidylate synthase (↓ dTMP)

Purine synthesis:

- 6-mercaptopurine (6-MP) and its prodrug azathioprine: inhibit de novo purine synthesis
- Mycophenolate and ribavirin: inhibit inosine monophosphate dehydrogenase

Purine and pyrimidine synthesis:

- Hydroxyurea: inhibits ribonucleotide reductase
- Methotrexate (MTX), trimethoprim (TMP), and pyrimethamine: inhibit dihydrofolate reductase (↓ deoxythymidine monophosphate [dTMP]) in humans, bacteria, and protozoa, respectively

CPS**1** = m**1**tochondria (urea cycle)
CPS**2** = cy**TWO**sol

(c) ketabton.com: The Digital Library

Purine salvage deficiencies



ADA, adenosine deaminase; APRT, adenine phosphoribosyltransferase; HGPRT, hypoxanthine guanine phosphoribosyltransferase; XO, xanthine oxidase.

Adenosine deaminase deficiency	ADA is required for degradation of adenosine and deoxyadenosine. ↓ ADA → ↑ dATP → ↓ ribonucleotide reductase activity → lymphotoxicity.	One of the major causes of autosomal recessive SCID.
Lesch-Nyhan syndrome	Defective purine salvage due to absent **HGPRT**, which converts hypoxanthine to IMP and guanine to GMP. Results in excess uric acid production and de novo purine synthesis. X-linked recessive. Findings: intellectual disability, self-mutilation, aggression, hyperuricemia (orange "sand" [sodium urate crystals] in diaper), gout, dystonia, macrocytosis. Treatment: allopurinol or febuxostat (2nd line).	**HGPRT**: **H**yperuricemia **G**out **P**issed off (aggression, self-mutilation) **R**etardation (intellectual disability) Dys**T**onia

Genetic code features

Unambiguous	Each codon specifies only 1 amino acid.	
Degenerate/ redundant	Most amino acids are coded by multiple codons. **Wobble**—codons that differ in 3rd ("wobble") position may code for the same tRNA/amino acid. Specific base pairing is usually required only in the first 2 nucleotide positions of mRNA codon.	Exceptions: methionine (AUG) and tryptophan (UGG) encoded by only 1 codon.
Commaless, nonoverlapping	Read from a fixed starting point as a continuous sequence of bases.	Exceptions: some viruses.
Universal	Genetic code is conserved throughout evolution.	Exception in humans: mitochondria.

(c) ketabton.com: The Digital Library

DNA replication	Eukaryotic DNA replication is more complex than in prokaryotes but uses many enzymes analogous to those listed below. In both prokaryotes and eukaryotes, DNA replication is semiconservative, involves continuous and discontinuous (Okazaki fragment) synthesis, and occurs in the 5′ → 3′ direction.	
Origin of replication A	Particular consensus sequence in genome where DNA replication begins. May be single (prokaryotes) or multiple (eukaryotes).	AT-rich sequences (such as TATA box regions) are found in promoters and origins of replication.
Replication fork B	Y-shaped region along DNA template where leading and lagging strands are synthesized.	
Helicase C	Unwinds DNA template at replication fork.	Helicase Halves DNA. Deficient in Bloom syndrome (*BLM* gene mutation).
Single-stranded binding proteins D	Prevent strands from reannealing.	
DNA topoisomerases E	Create a single- or double-stranded break in the helix to add or remove supercoils.	In eukaryotes: irinotecan/topotecan inhibit topoisomerase (TOP) I, etoposide/teniposide inhibit TOP II. In prokaryotes: fluoroquinolones inhibit TOP II (DNA gyrase) and TOP IV.
Primase F	Makes an RNA primer on which DNA polymerase III can initiate replication.	
DNA polymerase III G	Prokaryotes only. Elongates leading strand by adding deoxynucleotides to the 3′ end. Elongates lagging strand until it reaches primer of preceding fragment.	DNA polymerase III has 5′ → 3′ synthesis and proofreads with 3′ → 5′ exonuclease. Drugs blocking DNA replication often have a modified 3′ OH, thereby preventing addition of the next nucleotide ("chain termination").
DNA polymerase I H	Prokaryotes only. Degrades RNA primer; replaces it with DNA.	Same functions as DNA polymerase III, also excises RNA primer with 5′ → 3′ exonuclease.
DNA ligase I	Catalyzes the formation of a phosphodiester bond within a strand of double-stranded DNA.	Joins Okazaki fragments. Ligase Links DNA.
Telomerase	Eukaryotes only. A reverse transcriptase (RNA-dependent DNA polymerase) that adds DNA (TTAGGG) to 3′ ends of chromosomes to avoid loss of genetic material with every duplication.	Often dysregulated in cancer cells, allowing unlimited replication. Telomerase TAGs for Greatness and Glory.



(c) ketabton.com: The Digital Library

Mutations in DNA

Severity of damage: silent << missense < nonsense < frameshift.

Types of single nucleotide (point) mutations:

- Transition—purine to purine (eg, A to G) or pyrimidine to pyrimidine (eg, C to T).
- Transversion—purine to pyrimidine (eg, A to T) or pyrimidine to purine (eg, C to G).

Single nucleotide substitutions

Silent mutation

Nucleotide substitution codes for same (synonymous) amino acid; often base change in 3rd position of codon (tRNA wobble).

Missense mutation

Nucleotide substitution results in changed amino acid (called conservative if new amino acid has similar chemical structure).

Examples include sickle cell disease (substitution of glutamic acid with valine).

Nonsense mutation

Nucleotide substitution results in early stop codon (UGA, UAA, UAG). Usually results in nonfunctional protein. Stop the nonsense!

Other mutations

Frameshift mutation

Deletion or insertion of a number of nucleotides not divisible by 3 → misreading of all nucleotides downstream. Protein may be shorter or longer, and its function may be disrupted or altered.

Examples include Duchenne muscular dystrophy, Tay-Sachs disease.

Splice site mutation

Retained intron in mRNA → protein with impaired or altered function.

Examples include rare causes of cancers, dementia, epilepsy, some types of β-thalassemia, Gaucher disease, Marfan syndrome.



(c) ketabton.com: The Digital Library

Lac operon

Classic example of a genetic response to an environmental change. Glucose is the preferred metabolic substrate in *E coli*, but when glucose is absent and lactose is available, the *lac* operon is activated to switch to lactose metabolism. Mechanism of shift:

- Low glucose → ↑ adenylate cyclase activity → ↑ generation of cAMP from ATP → activation of catabolite activator protein (CAP) → ↑ transcription.
- High lactose → unbinds repressor protein from repressor/operator site → ↑ transcription.



DNA repair

Single strand		
Nucleotide excision repair	Specific endonucleases release the oligonucleotides containing damaged bases; DNA polymerase and ligase fill and reseal the gap, respectively. Repairs bulky helix-distorting lesions. Occurs in G_1 phase of cell cycle.	Defective in xeroderma pigmentosum (inability to repair DNA pyrimidine dimers caused by UV exposure). Findings: dry skin, extreme light sensitivity, skin cancer.
Base excision repair	Base-specific **G**lycosylase removes altered base and creates AP site (apurinic/apyrimidinic). One or more nucleotides are removed by AP-**E**ndonuclease, which cleaves 5′ end. AP-**L**yase cleaves 3′ end. DNA **P**olymerase-β fills the gap and DNA **L**igase seals it. Occurs throughout cell cycle.	Important in repair of spontaneous/toxic deamination. "**GEL PL**ease"
Mismatch repair	Mismatched nucleotides in newly synthesized (unmethylated) strand are removed and gap is filled and resealed. Occurs predominantly in S phase of cell cycle.	Defective in Lynch syndrome (hereditary nonpolyposis colorectal cancer [HNPCC]).
Double strand		
Nonhomologous end joining	Brings together 2 ends of DNA fragments to repair double-stranded breaks.	Defective in ataxia-telangiectasia. No requirement for homology. Some DNA may be lost.
Homologous recombination	Requires 2 homologous DNA duplexes. A strand from damaged dsDNA is repaired using a complementary strand from intact homologous dsDNA as a template.	Defective in breast/ovarian cancers with *BRCA1* mutation and in Fanconi anemia. Restores duplexes accurately without loss of nucleotides.

(c) ketabton.com: The Digital Library

Functional organization of a eukaryotic gene



Regulation of gene expression

Promoter	Site where RNA polymerase II and multiple other transcription factors bind to DNA upstream from gene locus (AT-rich upstream sequence with TATA and CAAT boxes).	Promoter mutation commonly results in dramatic ↓ in level of gene transcription.
Enhancer	DNA locus where regulatory proteins ("**activators**") bind, **increasing** expression of a gene on the same chromosome.	Enhancers and silencers may be located close to, far from, or even within (in an intron) the gene whose expression they regulate.
Silencer	DNA locus where regulatory proteins ("**repressors**") bind, **decreasing** expression of a gene on the same chromosome.	

RNA processing (eukaryotes)



Initial transcript is called heterogeneous nuclear RNA (hnRNA). hnRNA is then modified and becomes mRNA.

The following processes occur in the nucleus:

- Capping of 5′ end (addition of 7-methylguanosine cap)
- Polyadenylation of 3′ end (≈ 200 As)
- Splicing out of introns

Capped, tailed, and spliced transcript is called mRNA.

mRNA is transported out of nucleus to be translated in cytosol.

mRNA quality control occurs at cytoplasmic processing bodies (P-bodies), which contain exonucleases, decapping enzymes, and microRNAs; mRNAs may be degraded or stored in P-bodies for future translation.

Poly-A polymerase does not require a template.

AAUAAA = polyadenylation signal.

(c) ketabton.com: The Digital Library

RNA polymerases

Eukaryotes	RNA polymerase I makes **r**RNA, the most common (**r**ampant) type; present only in nucleolus. RNA polymerase II makes **m**RNA (**m**assive), **mi**croRNA (**mi**RNA), and **s**mall **n**uclear RNA (**sn**RNA). RNA polymerase III makes 5S rRNA, **t**RNA (**t**iny). No proofreading function, but can initiate chains. RNA polymerase II opens DNA at promoter site.	I, II, and III are numbered in the same order that their products are used in protein synthesis: rRNA, mRNA, then tRNA. α-amanitin, found in *Amanita phalloides* (death cap mushrooms), inhibits RNA polymerase II. Causes severe hepatotoxicity if ingested. Actinomycin D, also called dactinomycin, inhibits RNA polymerase in both prokaryotes and eukaryotes.
Prokaryotes	1 RNA polymerase (multisubunit complex) makes all 3 kinds of RNA.	Rifampin inhibits DNA-dependent RNA polymerase in prokaryotes.

Splicing of pre-mRNA Part of process by which precursor mRNA (pre-mRNA) is transformed into mature mRNA. Alterations in snRNP assembly can cause clinical disease; eg, in spinal muscular atrophy, snRNP assembly is affected due to ↓ SMN protein → congenital degeneration of anterior horns of spinal cord → symmetric weakness (hypotonia, or "floppy baby syndrome").



(c) ketabton.com: The Digital Library

Introns vs exons

Exons contain the actual genetic information coding for protein.

Introns do not code for protein, but are important in regulation of gene expression.

Different exons are frequently combined by alternative splicing to produce a larger number of unique proteins.

Alternative splicing can produce a variety of protein products from a single hnRNA sequence (eg, transmembrane vs secreted Ig, tropomyosin variants in muscle, dopamine receptors in the brain).

Introns are **in**tervening sequences and stay **in** the nucleus, whereas **ex**ons **ex**it and are **ex**pressed.



(c) ketabton.com: The Digital Library

tRNA

Structure	75–90 nucleotides, 2° structure, cloverleaf form, anticodon end is opposite 3′ aminoacyl end. All tRNAs, both eukaryotic and prokaryotic, have CCA at 3′ end along with a high percentage of chemically modified bases. The amino acid is covalently bound to the 3′ end of the tRNA. **CCA** **C**an **C**arry **A**mino acids. T-arm: contains the TΨC (ribothymidine, pseudouridine, cytidine) sequence necessary for tRNA-ribosome binding. **T**-arm **T**ethers tRNA molecule to ribosome. D-arm: contains **D**ihydrouridine residues necessary for tRNA recognition by the correct aminoacyl-tRNA synthetase. **D**-arm allows **D**etection of the tRNA by aminoacyl-tRNA synthetase. Attachment site: the 5′-CCA-3′ is the amino acid acceptor site.
Charging	Aminoacyl-tRNA synthetase (uses ATP; 1 unique enzyme per respective amino acid) and binding of charged tRNA to the codon are responsible for the accuracy of amino acid selection. Aminoacyl-tRNA synthetase matches an amino acid to the tRNA by scrutinizing the amino acid before and after it binds to tRNA. If an incorrect amino acid is attached, the bond is hydrolyzed. A mischarged tRNA reads the usual codon but inserts the wrong amino acid.



Start and stop codons

mRNA start codons	AUG (or rarely GUG).	**AUG** in**AUG**urates protein synthesis.
Eukaryotes	Codes for methionine, which may be removed before translation is completed.	
Prokaryotes	Codes for *N*-formylmethionine (fMet).	fMet stimulates neutrophil chemotaxis.
mRNA stop codons	UGA, UAA, UAG.	**UGA** = **U** **G**o **A**way. **UAA** = **U** **A**re **A**way. **UAG** = **U** **A**re **G**one.

(c) ketabton.com: The Digital Library

Protein synthesis

Initiation

1. Eukaryotic initiation factors (eIFs) identify the 5′ cap.
2. eIFs help assemble the 40S ribosomal subunit with the initiator tRNA.
3. eIFs released when the mRNA and the ribosomal 60S subunit assemble with the complex. Requires GTP.

Eukaryotes: 40S + 60S → 80S (Even).
Prokaryotes: 30S + 50S → 70S (Prime).
Synthesis occurs from N-terminus to C-terminus.

ATP—tRNA Activation (charging).
GTP—tRNA Gripping and Going places (translocation).

Elongation

❶ Aminoacyl-tRNA binds to A site (except for initiator methionine, which binds the P site), requires an elongation factor and GTP.
❷ rRNA ("ribozyme") catalyzes peptide bond formation, transfers growing polypeptide to amino acid in A site.
❸ Ribosome advances 3 nucleotides toward 3′ end of mRNA, moving peptidyl tRNA to P site (translocation).

Think of "going APE":
A site = incoming Aminoacyl-tRNA.
P site = accommodates growing Peptide.
E site = holds Empty tRNA as it Exits.

Termination

Eukaryotic release factors (eRFs) recognize the stop codon and halt translation → completed polypeptide is released from ribosome. Requires GTP.



Posttranslational modifications

Trimming	Removal of N- or C-terminal propeptides from zymogen to generate mature protein (eg, trypsinogen to trypsin).
Covalent alterations	Phosphorylation, glycosylation, hydroxylation, methylation, acetylation, and ubiquitination.

Chaperone protein — Intracellular protein involved in facilitating and maintaining protein folding. In yeast, heat shock proteins (eg, HSP60) are expressed at high temperatures to prevent protein denaturing/misfolding.

(c) ketabton.com: The Digital Library

▸ BIOCHEMISTRY—CELLULAR

Cell cycle phases

Checkpoints control transitions between phases of cell cycle. This process is regulated by cyclins, cyclin-dependent kinases (CDKs), and tumor suppressors. M phase (shortest phase of cell cycle) includes mitosis (prophase, prometaphase, metaphase, anaphase, telophase) and cytokinesis (cytoplasm splits in two). G_1 and G_0 are of variable duration.

REGULATION OF CELL CYCLE

Cyclin-dependent kinases	Constitutively expressed but inactive when not bound to cyclin.
Cyclins	Regulatory proteins that control cell cycle events; phase specific; activate CDKs.
Cyclin-CDK complexes	Phosphorylate other proteins to coordinate cell cycle progression; must be activated and inactivated at appropriate times for cell cycle to progress.
Tumor suppressors	p53 → p21 induction → CDK inhibition → Rb hypophosphorylation (activation) → G_1-S progression inhibition. Mutations in tumor suppressor genes can result in unrestrained cell division (eg, Li-Fraumeni syndrome).
	Growth factors (eg, insulin, PDGF, EPO, EGF) bind tyrosine kinase receptors to transition the cell from G_1 to S phase.



CELL TYPES

Permanent	Remain in G_0, regenerate from stem cells.	Neurons, skeletal and cardiac muscle, RBCs.
Stable (quiescent)	Enter G_1 from G_0 when stimulated.	Hepatocytes, lymphocytes, PCT, periosteal cells.
Labile	Never go to G_0, divide rapidly with a short G_1. Most affected by chemotherapy.	Bone marrow, gut epithelium, skin, hair follicles, germ cells.

Rough endoplasmic reticulum	Site of synthesis of secretory (exported) proteins and of N-linked oligosaccharide addition to lysosomal and other proteins. Nissl bodies (RER in neurons)—synthesize peptide neurotransmitters for secretion. Free ribosomes—unattached to any membrane; site of synthesis of cytosolic, peroxisomal, and mitochondrial proteins.	Mucus-secreting goblet cells of the small intestine and antibody-secreting plasma cells are rich in RER. Proteins within organelles (eg, ER, Golgi bodies, lysosomes) are formed in RER.
Smooth endoplasmic reticulum	Site of steroid synthesis and detoxification of drugs and poisons. Lacks surface ribosomes. Location of glucose-6-phosphatase (last step of glycogenolysis).	Liver hepatocytes and steroid hormone–producing cells of the adrenal cortex and gonads are rich in SER.

(c) ketabton.com: The Digital Library

Cell trafficking

Golgi is distribution center for proteins and lipids from ER to vesicles and plasma membrane. Posttranslational events in Golgi include modifying N-oligosaccharides on asparagine, adding O-oligosaccharides on serine and threonine, and adding mannose-6-phosphate to proteins for lysosomal trafficking.

Endosomes are sorting centers for material from outside the cell or from the Golgi, sending it to lysosomes for destruction or back to the membrane/Golgi for further use.

I-cell disease (inclusion cell disease/mucolipidosis type II)—inherited lysosomal storage disorder (autosomal recessive); defect in *N*-acetylglucosaminyl-1-phosphotransferase → failure of the Golgi to phosphorylate mannose residues (↓ mannose-6-phosphate) on glycoproteins → proteins are secreted extracellularly rather than delivered to lysosomes. Results in coarse facial features, gingival hyperplasia, clouded corneas, restricted joint movements, claw hand deformities, kyphoscoliosis, and high plasma levels of lysosomal enzymes. Often fatal in childhood.



Signal recognition particle (SRP)

Abundant, cytosolic ribonucleoprotein that traffics polypeptide-ribosome complex from the cytosol to the RER. Absent or dysfunctional SRP → accumulation of protein in cytosol.

Vesicular trafficking proteins

COPI: Golgi → Golgi (retrograde); *cis*-Golgi → ER.

COPII: ER → *cis*-Golgi (anterograde).

"Two (COPII) steps forward (anterograde); one (COPI) step back (retrograde)."

Clathrin: *trans*-Golgi → lysosomes; plasma membrane → endosomes (receptor-mediated endocytosis [eg, LDL receptor activity]).

Peroxisome

Membrane-enclosed organelle involved in:

- β-oxidation of very-long-chain fatty acids (VLCFA) (strictly peroxisomal process)
- α-oxidation of branched-chain fatty acids (strictly peroxisomal process)
- Catabolism of amino acids and ethanol
- Synthesis of cholesterol, bile acids, and plasmalogens (important membrane phospholipid, especially in white matter of brain)

Zellweger syndrome—autosomal recessive disorder of peroxisome biogenesis due to mutated *PEX* genes. Hypotonia, seizures, hepatomegaly, early death.

Refsum disease—autosomal recessive disorder of α-oxidation → phytanic acid not metabolized to pristanic acid. Scaly skin, ataxia, cataracts/night blindness, shortening of 4th toe, epiphyseal dysplasia. Treatment: diet, plasmapheresis.

Adrenoleukodystrophy—X-linked recessive disorder of β-oxidation due to mutation in *ABCD1* gene → VLCFA buildup in adrenal glands, white (leuko) matter of brain, testes. Progressive disease that can lead to adrenal gland crisis, coma, and death.

(c) ketabton.com: The Digital Library

Proteasome Barrel-shaped protein complex that degrades damaged or ubiquitin-tagged proteins. Defects in the ubiquitin-proteasome system have been implicated in some cases of Parkinson disease.

Cytoskeletal elements A network of protein fibers within the cytoplasm that supports cell structure, cell and organelle movement, and cell division.

TYPE OF FILAMENT	PREDOMINANT FUNCTION	EXAMPLES
Microfilaments	Muscle contraction, cytokinesis	Actin, microvilli.
Intermediate filaments	Maintain cell structure	Vimentin, desmin, cytokeratin, lamins, glial fibrillary acidic protein (GFAP), neurofilaments.
Microtubules	Movement, cell division	Cilia, flagella, mitotic spindle, axonal trafficking, centrioles.

Microtubule



Cylindrical outer structure composed of a helical array of polymerized heterodimers of α- and β-tubulin. Each dimer has 2 GTP bound. Incorporated into flagella, cilia, mitotic spindles. Grows slowly, collapses quickly. Also involved in slow axoplasmic transport in neurons.

Molecular motor proteins—transport cellular cargo toward opposite ends of microtubule.
- **RE**trograde to microtubule (+ → –)—**DY**nein.
- **A**nterograde to microtubule (– → +)—**K**inesin.

Clostridium tetani, herpes simplex virus, poliovirus, and rabies virus use dynein for retrograde transport to the neuronal cell body.

Drugs that act on microtubules (**M**icrotubules **G**et **C**onstructed **V**ery **P**oorly):
- **M**ebendazole (antihelminthic)
- **G**riseofulvin (antifungal)
- **C**olchicine (antigout)
- **V**incristine/**V**inblastine (anticancer)
- **P**aclitaxel (anticancer)

Negative end **N**ear **N**ucleus.
Positive end **P**oints to **P**eriphery.

REa**DY**? **A**ttac**K**!

(c) ketabton.com: The Digital Library

Cilia structure

9 doublet + 2 singlet arrangement of microtubules A.

Basal body (base of cilium below cell membrane) consists of 9 microtubule triplets B with no central microtubules.

Axonemal dynein—ATPase that links peripheral 9 doublets and causes bending of cilium by differential sliding of doublets.

Gap junctions enable coordinated ciliary movement.

Kartagener syndrome (1° ciliary dyskinesia)—immotile cilia due to a dynein arm defect. Autosomal recessive. Results in ↓ male and female fertility due to immotile sperm and dysfunctional fallopian tube cilia, respectively; ↑ risk of ectopic pregnancy. Can cause bronchiectasis, recurrent sinusitis, chronic ear infections, conductive hearing loss, and situs inversus (eg, dextrocardia on CXR C). ↓ nasal nitric oxide (used as screening test). (Kartagener's restaurant: take-out only; there's no dynein "dine-in".)



Sodium-potassium pump

Na^+-K^+ ATPase is located in the plasma membrane with ATP site on cytosolic side. For each ATP consumed, 3 Na^+ leave the cell (pump phosphorylated) and 2 K^+ enter the cell (pump dephosphorylated).

Plasma membrane is an asymmetric lipid bilayer containing cholesterol, phospholipids, sphingolipids, glycolipids, and proteins.

Pumpkin = pump K^+ in.

Ouabain (a cardiac glycoside) inhibits by binding to K^+ site.

Cardiac glycosides (digoxin and digitoxin) directly inhibit the Na^+-K^+ ATPase, which leads to indirect inhibition of Na^+/Ca^{2+} exchange → ↑ $[Ca^{2+}]_i$ → ↑ cardiac contractility.



(c) ketabton.com: The Digital Library

Collagen	Most abundant protein in the human body. Extensively modified by posttranslational modification. Organizes and strengthens extracellular matrix.	Be So Totally Cool, Read Books.
Type I	Most common (90%)—Bone (made by osteoblasts), Skin, Tendon, dentin, fascia, cornea, **late** wound repair.	Type I: bone. ↓ production in osteogenesis imperfecta type I.
Type II	Cartilage (including hyaline), vitreous body, nucleus pulposus.	Type II: cartwolage.
Type III	Reticulin—skin, blood vessels, uterus, fetal tissue, **early** wound repair.	Type III: deficient in the uncommon, vascular type of Ehlers-Danlos syndrome (ThreE D).
Type IV	Basement membrane (basal lamina), lens.	Type IV: under the floor (basement membrane). Defective in Alport syndrome; targeted by autoantibodies in Goodpasture syndrome.

Collagen synthesis and structure



1. **Synthesis**—translation of collagen α chains (preprocollagen)—usually Gly-X-Y (X and Y are proline or lysine). Collagen is ⅓ glycine; glycine content of collagen is less variable than that of lysine and proline. Hydroxyproline is used for lab quantification of collagen.
2. **Hydroxylation**—hydroxylation of specific proline and lysine residues. Requires vitamin C; deficiency → scurvy.
3. **Glycosylation**—glycosylation of pro-α-chain hydroxylysine residues and formation of procollagen via hydrogen and disulfide bonds (triple helix of 3 collagen α chains). Problems forming triple helix → osteogenesis imperfecta.
4. **Exocytosis**—exocytosis of procollagen into extracellular space.
5. **Proteolytic processing**—cleavage of disulfide-rich terminal regions of procollagen → insoluble tropocollagen.
6. **Cross-linking**—reinforcement of many staggered tropocollagen molecules by covalent lysine-hydroxylysine cross-linkage (by copper-containing lysyl oxidase) to make collagen fibrils. Problems with cross-linking → Menkes disease.

(c) ketabton.com: The Digital Library

Osteogenesis imperfecta

Genetic bone disorder (brittle bone disease) caused by a variety of gene defects (most commonly *COL1A1* and *COL1A2*).

Most common form is autosomal dominant with ↓ production of otherwise normal type I collagen. Manifestations include:

- Multiple fractures and bone deformities after minimal trauma (eg, during birth)
- Blue sclerae B due to the translucent connective tissue over choroidal veins
- Some forms have tooth abnormalities, including opalescent teeth that wear easily due to lack of dentin (dentinogenesis imperfecta)
- Conductive hearing loss (abnormal ossicles)

May be confused with child abuse.

Treat with bisphosphonates to ↓ fracture risk.

Patients can't **BITE**:

Bones = multiple fractures
I (eye) = blue sclerae
Teeth = dental imperfections
Ear = hearing loss



Ehlers-Danlos syndrome

Faulty collagen synthesis causing hyperextensible skin A, hypermobile joints B, and tendency to bleed (easy bruising).

Multiple types. Inheritance and severity vary. Can be autosomal dominant or recessive. May be associated with joint dislocation, berry and aortic aneurysms, organ rupture.

Hypermobility type (joint instability): most common type.

Classical type (joint and skin symptoms): caused by a mutation in type V collagen (eg, *COL5A1*, *COL5A2*).

Vascular type (fragile tissues including vessels [eg, aorta], muscles, and organs that are prone to rupture [eg, gravid uterus]): mutations in type III procollagen (eg, *COL3A1*).



Menkes disease

X-linked recessive connective tissue disease caused by impaired copper absorption and transport due to defective Menkes protein (*ATP7A*, vs *ATP7B* in Wilson disease). Low copper levels (vs high levels in Wilson disease). Leads to ↓ activity of lysyl oxidase (copper is a necessary cofactor) → defective collagen. Results in brittle, "kinky" hair, growth retardation, hypotonia, ↑ risk of cerebral aneurysms.

(c) ketabton.com: The Digital Library

Elastin



Stretchy protein within skin, lungs, large arteries, elastic ligaments, vocal cords, ligamenta flava (connect vertebrae → relaxed and stretched conformations).
Rich in nonhydroxylated proline, glycine, and lysine residues, vs the hydroxylated residues of collagen.
Tropoelastin with fibrillin scaffolding.
Cross-linking takes place extracellularly and gives elastin its elastic properties.
Broken down by elastase, which is normally inhibited by α_1-antitrypsin.
α_1-Antitrypsin deficiency results in unopposed elastase activity, which can cause COPD.
Changes with aging: ↓ dermal collagen and elastin, ↓ synthesis of collagen fibrils; cross-linking remains normal.

Marfan syndrome—autosomal dominant (with variable expression) connective tissue disorder affecting skeleton, heart, and eyes. *FBN1* gene mutation on chromosome 15 (fifteen) results in defective fibrillin, a glycoprotein that forms a sheath around elastin. Findings: tall with long extremities; pectus carinatum (more specific) or pectus excavatum A; hypermobile joints; long, tapering fingers and toes (arachnodactyly); cystic medial necrosis of aorta; aortic root aneurysm rupture or dissection (most common cause of death); mitral valve prolapse. Subluxation of lenses, typically upward and temporally (vs downward and medially in homocystinuria).

▸ BIOCHEMISTRY—LABORATORY TECHNIQUES

Polymerase chain reaction

Molecular biology lab procedure used to amplify a desired fragment of DNA. Useful as a diagnostic tool (eg, neonatal HIV, herpes encephalitis).



1. **Denaturation**—DNA is heated to ~95°C to separate the strands.
2. **Annealing**—Sample is cooled to ~55°C. DNA primers, a heat-stable DNA polymerase *(Taq)*, and deoxynucleotide triphosphates (dNTPs) are added. DNA primers anneal to the specific sequence to be amplified on each strand.
3. **Elongation**—Temperature is increased to ~72°C. DNA polymerase attaches dNTPs to the strand to replicate the sequence after each primer.

Heating and cooling cycles continue until the DNA sample size is sufficient.

Reverse transcriptase polymerase chain reaction

Detects and quantifies mRNA levels in a sample. Uses reverse transcription to create a complementary DNA template that is amplified via standard PCR procedure.

(c) ketabton.com: The Digital Library

CRISPR/Cas9

A genome editing tool derived from bacteria. Consists of a guide RNA (gRNA) ❶, which is complementary to a target DNA sequence, and an endonuclease (Cas9), which makes a single- or double-strand break at the target site ❷. Break imperfectly repaired by nonhomologous end joining (NHEJ) → accidental frameshift mutations ("knock-out") 3A, or a donor DNA sequence can be added to fill in the gap using homology-directed repair (HDR) 3B.

Not used clinically. Potential applications include removing virulence factors from pathogens, replacing disease-causing alleles of genes with healthy variants, and specifically targeting tumor cells.



Blotting procedures

Southern blot

1. **DNA** sample is enzymatically cleaved into smaller pieces, which are separated on a gel by electrophoresis, and then transferred to a filter.
2. Filter is exposed to radiolabeled DNA probe that recognizes and anneals to its complementary strand.
3. Resulting double-stranded, labeled piece of DNA is visualized when filter is exposed to film.



Northern blot

Similar to Southern blot, except that an **RNA** sample is electrophoresed. Useful for studying mRNA levels, which are reflective of gene expression.

Western blot

Sample protein is separated via gel electrophoresis and transferred to a membrane. Labeled antibody is used to bind to relevant **protein.**

Southwestern blot

Identifies **DNA-binding proteins** (eg, c-Jun, c-Fos [leucine zipper motif]) using labeled double-stranded DNA probes.

SNoW DRoP:
Southern = DNA
Northern = RNA
Western = Protein

(c) ketabton.com: The Digital Library

Flow cytometry

Laboratory technique to assess size, granularity, and protein expression (immunophenotype) of individual cells in a sample.

Commonly used in workup of hematologic abnormalities (eg, leukemia, paroxysmal nocturnal hemoglobinuria, fetal RBCs in mother's blood) and immunodeficiencies (eg, $CD4^+$ cell count in HIV).

Cells are tagged with antibodies specific to surface or intracellular proteins. Antibodies are then tagged with a unique fluorescent dye. Sample is analyzed one cell at a time by focusing a laser on the cell and measuring light scatter and intensity of fluorescence.



Data are plotted either as histogram (one measure) or scatter plot (any two measures, as shown). In illustration:

- Cells in left lower quadrant ⊖ for both CD8 and CD3.
- Cells in right lower quadrant ⊕ for CD8 and ⊖ for CD3. In this example, right lower quadrant is empty because all CD8-expressing cells also express CD3.
- Cells in left upper quadrant ⊕ for CD3 and ⊖ for CD8.
- Cells in right upper quadrant ⊕ for both CD8 and CD3.



Microarrays

Thousands of nucleic acid sequences are arranged in grids on glass or silicon. DNA or RNA probes are hybridized to the chip, and a scanner detects the relative amounts of complementary binding. Used to profile gene expression levels of thousands of genes simultaneously to study certain diseases and treatments. Able to detect single nucleotide polymorphisms (SNPs) and copy number variations (CNVs) for a variety of applications including genotyping, clinical genetic testing, forensic analysis, cancer mutations, and genetic linkage analysis.

Enzyme-linked immunosorbent assay

Immunologic test used to detect the presence of either a specific antigen or antibody in a patient's blood sample. Detection involves the use of an antibody linked to an enzyme. Added substrate reacts with enzyme, producing a detectable signal. Can have high sensitivity and specificity, but is less specific than Western blot.

(c) ketabton.com: The Digital Library

Karyotyping

Colchicine is added to cultured cells to halt chromosomes in metaphase. Chromosomes are stained, ordered, and numbered according to morphology, size, arm-length ratio, and banding pattern (arrows in A point to extensive abnormalities in a cancer cell).

Can be performed on a sample of blood, bone marrow, amniotic fluid, or placental tissue.

Used to diagnose chromosomal imbalances (eg, autosomal trisomies, sex chromosome disorders).



Fluorescence in situ hybridization

Fluorescent DNA or RNA probe binds to specific gene site of interest on chromosomes (arrows in A point to abnormalities in a cancer cell, whose karyotype is seen above; each fluorescent color represents a chromosome-specific probe).

Used for specific localization of genes and direct visualization of chromosomal anomalies at the molecular level.

- Microdeletion—no fluorescence on a chromosome compared to fluorescence at the same locus on the second copy of that chromosome.
- Translocation—fluorescence signal that corresponds to one chromosome is found in a different chromosome (two white arrows in A show fragments of chromosome 17 that have translocated to chromosome 19).
- Duplication—a second copy of a chromosome, resulting in a trisomy or tetrasomy (two blue arrows show duplicated chromosomes 8, resulting in a tetrasomy).

Molecular cloning

Production of a recombinant DNA molecule in a bacterial host.

Steps:

1. Isolate eukaryotic mRNA (post-RNA processing) of interest.
2. Add reverse transcriptase (an RNA-dependent DNA polymerase) to produce complementary DNA (cDNA, lacks introns).
3. Insert cDNA fragments into bacterial plasmids containing antibiotic resistance genes.
4. Transform (insert) recombinant plasmid into bacteria.
5. Surviving bacteria on antibiotic medium produce cloned DNA (copies of cDNA).

(c) ketabton.com: The Digital Library

Gene expression modifications	Transgenic strategies in mice involve: ▪ Random insertion of gene into mouse genome ▪ Targeted insertion or deletion of gene through homologous recombination with mouse gene	Knock-**out** = removing a gene, taking it **out**. Knock-**in** = **in**serting a gene. Random insertion—constitutive expression. Targeted insertion—conditional expression.
Cre-lox system	Can inducibly manipulate genes at specific developmental points (eg, to study a gene whose deletion causes embryonic death).	
RNA interference	Process whereby small non-coding RNA molecules target mRNAs to inhibit gene expression.	
MicroRNA (miRNA)	Naturally produced by the cell as hairpin structures. Loose nucleotide pairing allows broader targeting of related mRNAs, blocking translation and accelerating mRNA degradation.	Abnormal expression of miRNAs contributes to certain malignancies (eg, by silencing an mRNA from a tumor suppressor gene).
Small interfering RNA (siRNA)	Usually derived from exogenous dsRNA source (eg, virus). Once inside a cell, siRNA requires complete nucleotide pairing, leading to highly specific mRNA targeting. Results in mRNA cleavage prior to translation.	Can be produced by in vitro transcription for gene "knockdown" experiments.

▸ BIOCHEMISTRY—GENETICS

Genetic terms

TERM	DEFINITION	EXAMPLE
Codominance	Both alleles contribute to the phenotype of the heterozygote.	Blood groups A, B, AB; α_1-antitrypsin deficiency; HLA groups.
Variable expressivity	Patients with the same genotype have varying phenotypes.	2 patients with neurofibromatosis type 1 (NF1) may have varying disease severity.
Incomplete penetrance	Not all individuals with a mutant genotype show the mutant phenotype. % penetrance × probability of inheriting genotype = risk of expressing phenotype.	*BRCA1* gene mutations do not always result in breast or ovarian cancer.
Pleiotropy	One gene contributes to multiple phenotypic effects.	Untreated phenylketonuria (PKU) manifests with light skin, intellectual disability, and musty body odor.
Anticipation	Increased severity or earlier onset of disease in succeeding generations.	Trinucleotide repeat diseases (eg, Huntington disease).
Loss of heterozygosity	If a patient inherits or develops a mutation in a tumor suppressor gene, the complementary allele must be deleted/mutated before cancer develops. This is not true of oncogenes.	Retinoblastoma and the "two-hit hypothesis," Lynch syndrome (HNPCC), Li-Fraumeni syndrome.

(c) ketabton.com: The Digital Library

Genetic terms *(continued)*

TERM	DEFINITION	EXAMPLE
Dominant negative mutation	Exerts a dominant effect. A heterozygote produces a nonfunctional altered protein that also prevents the normal gene product from functioning.	A single mutated *p53* tumor suppressor gene results in a protein that is able to bind DNA and block the nonmutated p53 from binding to the promoter.
Linkage disequilibrium	Tendency for certain alleles at 2 linked loci to occur together more or less often than expected by chance. Measured in a population, not in a family, and often varies in different populations.	
Mosaicism A	Presence of genetically distinct cell lines in the same individual. Somatic mosaicism—mutation arises from mitotic errors after fertilization and propagates through multiple tissues or organs. Gonadal mosaicism—mutation only in egg or sperm cells. If parents and relatives do not have the disease, suspect gonadal (or germline) mosaicism.	**McCune-Albright syndrome**—due to G_s-protein activating mutation. Presents with unilateral café-au-lait spots A with ragged edges, polyostotic fibrous dysplasia (bone is replaced by collagen and fibroblasts), and at least one endocrinopathy (eg, precocious puberty). Lethal if mutation occurs before fertilization (affecting all cells), but survivable in patients with mosaicism.
Locus heterogeneity	Mutations at different loci can produce a similar phenotype.	Albinism.
Allelic heterogeneity	Different mutations in the same locus produce the same phenotype.	β-thalassemia.
Heteroplasmy	Presence of both normal and mutated mtDNA, resulting in variable expression in mitochondrially inherited disease.	mtDNA passed from mother to all children.
Uniparental disomy	Offspring receives 2 copies of a chromosome from 1 parent and no copies from the other parent. HeterodIsomy (heterozygous) indicates a meiosis I error. IsodIsomy (homozygous) indicates a meiosis II error or postzygotic chromosomal duplication of one of a pair of chromosomes, and loss of the other of the original pair.	Uniparental is euploid (correct number of chromosomes). Most occurrences of uniparental disomy (UPD) → normal phenotype. Consider isodisomy in an individual manifesting a recessive disorder when only one parent is a carrier. Examples: Prader-Willi and Angelman syndromes.

Hardy-Weinberg population genetics

	A (p)	a (q)
A (p)	AA (p^2)	Aa (pq)
a (q)	Aa (pq)	aa (q^2)

If p and q represent the frequencies of alleles A and a, respectively, in a population, then $p + q = 1$:

- p^2 = frequency of homozygosity for allele A
- q^2 = frequency of homozygosity for allele a
- $2pq$ = frequency of heterozygosity (carrier frequency, if an autosomal recessive disease)

Therefore, the sum of the frequencies of these genotypes is $p^2 + 2pq + q^2 = 1$.

The frequency of an X-linked recessive disease in males = q and in females = q^2.

Hardy-Weinberg law assumptions include:

- No mutation occurring at the locus
- Natural selection is not occurring
- Completely random mating
- No net migration
- Large population

If a population is in Hardy-Weinberg equilibrium, then the values of p and q remain constant from generation to generation.

(c) ketabton.com: The Digital Library

Disorders of imprinting

Imprinting—one gene copy is silenced by methylation, and only the other copy is expressed → parent-of-origin effects.

	Prader-Willi syndrome	AngelMan syndrome
WHICH GENE IS SILENT?	Maternally derived genes are silenced Disease occurs when the **P**aternal allele is deleted or mutated	Paternally derived *UBE3A* is silenced Disease occurs when the **M**aternal allele is deleted or mutated
SIGNS AND SYMPTOMS	Hyperphagia, obesity, intellectual disability, hypogonadism, hypotonia	**S**eizures, **A**taxia, severe **I**ntellectual disability, inappropriate **L**aughter ("happy puppet") Set **SAIL** for Angel Island
CHROMOSOMES INVOLVED	Chromosome 15 of paternal origin	*UBE3A* on maternal copy of chromosome 15
NOTES	25% of cases are due to maternal uniparental disomy	5% of cases are due to paternal uniparental disomy
	Prader has no **D**ad (**P**aternal **D**eletion)	**MD**s are angels (**M**aternal **D**eletion)

(c) ketabton.com: The Digital Library

Modes of inheritance

Autosomal dominant



	A	a
a	Aa	aa
a	Aa	aa

Often due to defects in structural genes. Many generations, both males and females are affected.

Often pleiotropic (multiple apparently unrelated effects) and variably expressive (different between individuals). Family history crucial to diagnosis. With one affected (heterozygous) parent, on average, ½ of children affected.

Autosomal recessive



	A	a
A	AA	Aa
a	Aa	aa

With 2 carrier (heterozygous) parents, on average: ¼ of children will be affected (homozygous), ½ of children will be carriers, and ¼ of children will be neither affected nor carriers.

Often due to enzyme deficiencies. Usually seen in only 1 generation. Commonly more severe than dominant disorders; patients often present in childhood.

↑ risk in consanguineous families.

Unaffected individual with affected sibling has 2/3 probability of being a carrier.

X-linked recessive



	X	X
X	XX	XX
Y	XY	XY

	X	X
X	XX	XX
Y	XY	XY

Sons of heterozygous mothers have a 50% chance of being affected. No male-to-male transmission. Skips generations.

Commonly more severe in males. Females usually must be homozygous to be affected.

X-linked dominant



	X	X
X	XX	XX
Y	XY	XY

	X	X
X	XX	XX
Y	XY	XY

Transmitted through both parents. Mothers transmit to 50% of daughters and sons; fathers transmit to all daughters but no sons.

Examples: fragile X syndrome, Alport syndrome, hypophosphatemic rickets (also called X-linked hypophosphatemia)—phosphate wasting at proximal tubule → rickets-like presentation.

Mitochondrial inheritance



Transmitted only through the mother. All offspring of affected females may show signs of disease.

Variable expression in a population or even within a family due to heteroplasmy.

Mitochondrial myopathies—rare disorders; often present with myopathy, lactic acidosis, and CNS disease, eg, MELAS syndrome (mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes). 2° to failure in oxidative phosphorylation. Muscle biopsy often shows "ragged red fibers" (due to accumulation of diseased mitochondria in the subsarcolemma of the muscle fiber).

Leber hereditary optic neuropathy—cell death in optic nerve neurons → subacute bilateral vision loss in teens/young adults, 90% males. Usually permanent.

□ = unaffected male; ■ = affected male; ○ = unaffected female; ● = affected female.

(c) ketabton.com: The Digital Library

Autosomal dominant diseases

Achondroplasia, autosomal dominant polycystic kidney disease, familial adenomatous polyposis, familial hypercholesterolemia, hereditary hemorrhagic telangiectasia (Osler-Weber-Rendu syndrome), hereditary spherocytosis, Huntington disease, Li-Fraumeni syndrome, Marfan syndrome, multiple endocrine neoplasias, myotonic muscular dystrophy, neurofibromatosis type 1 (von Recklinghausen disease), neurofibromatosis type 2, tuberous sclerosis, von Hippel-Lindau disease.

Autosomal recessive diseases

Oculocutaneous albinism, autosomal recessive polycystic kidney disease (ARPKD), cystic fibrosis, Friedreich ataxia, glycogen storage diseases, hemochromatosis, Kartagener syndrome, mucopolysaccharidoses (except Hunter syndrome), phenylketonuria, sickle cell anemia, sphingolipidoses (except Fabry disease), thalassemias, Wilson disease.

Cystic fibrosis

GENETICS	Autosomal recessive; defect in *CFTR* gene on chromosome 7; commonly a deletion of Phe508. Most common lethal genetic disease in Caucasian population.
PATHOPHYSIOLOGY	*CFTR* encodes an ATP-gated Cl^- channel that secretes Cl^- in lungs and GI tract, and reabsorbs Cl^- in sweat glands. Most common mutation → misfolded protein → protein retained in RER and not transported to cell membrane, causing ↓ Cl^- (and H_2O) secretion; ↑ intracellular Cl^- results in compensatory ↑ Na^+ reabsorption via epithelial Na^+ channels (ENaC) → ↑ H_2O reabsorption → abnormally thick mucus secreted into lungs and GI tract. ↑ Na^+ reabsorption also causes more negative transepithelial potential difference.
DIAGNOSIS	↑ Cl^- concentration in pilocarpine-induced sweat test is diagnostic. Can present with contraction alkalosis and hypokalemia (ECF effects analogous to a patient taking a loop diuretic) because of ECF H_2O/Na^+ losses via sweating and concomitant renal K^+/H^+ wasting. ↑ immunoreactive trypsinogen (newborn screening).
COMPLICATIONS	Recurrent pulmonary infections (eg, *S aureus* [infancy and early childhood], *P aeruginosa* [adulthood], allergic bronchopulmonary aspergillosis [ABPA]), chronic bronchitis and bronchiectasis → reticulonodular pattern on CXR, opacification of sinuses. Pancreatic insufficiency, malabsorption with steatorrhea, fat-soluble vitamin deficiencies (A, D, E, K), biliary cirrhosis, liver disease. Meconium ileus in newborns. Infertility in men (absence of vas deferens, spermatogenesis may be unaffected) and subfertility in women (amenorrhea, abnormally thick cervical mucus). Nasal polyps, clubbing of nails.
TREATMENT	Multifactorial: chest physiotherapy, albuterol, aerosolized dornase alfa (DNase), and hypertonic saline facilitate mucus clearance. Azithromycin used as anti-inflammatory agent. Ibuprofen slows disease progression. Pancreatic enzyme replacement therapy for pancreatic insufficiency. In patients with Phe508 deletion: combination of lumacaftor (corrects misfolded proteins and improves their transport to cell surface) and ivacaftor (opens Cl^- channels → improved chloride transport).



FAS1_2019_01-Biochem.indd 60 11/7/19 3:16 PM

(c) ketabton.com: The Digital Library

X-linked recessive disorders	Ornithine transcarbamylase deficiency, Fabry disease, Wiskott-Aldrich syndrome, Ocular albinism, G6PD deficiency, Hunter syndrome, Bruton agammaglobulinemia, Hemophilia A and B, Lesch-Nyhan syndrome, Duchenne (and Becker) muscular dystrophy. X-inactivation (lyonization)—one copy of female X chromosome forms a transcriptionally inactive Barr body. Female carriers variably affected depending on the pattern of inactivation of the X chromosome carrying the mutant vs normal gene.	Oblivious Female Will Often Give Her Boys Her x-Linked Disorders Females with Turner syndrome (45,XO) are more likely to have an X-linked recessive disorder.

Muscular dystrophies

Duchenne  	X-linked disorder typically due to **frameshift** deletions or nonsense mutations → truncated or absent dystrophin protein → progressive myofiber damage. Weakness begins in pelvic girdle muscles and progresses superiorly. Pseudohypertrophy of calf muscles due to fibrofatty replacement of muscle A. Waddling gait. Onset before 5 years of age. Dilated cardiomyopathy is common cause of death. Gowers sign—patient uses upper extremities to help stand up. Classically seen in Duchenne muscular dystrophy, but also seen in other muscular dystrophies and inflammatory myopathies (eg, polymyositis).	Duchenne = deleted dystrophin. Dystrophin gene (*DMD*) is the largest protein-coding human gene → ↑ chance of spontaneous mutation. Dystrophin helps anchor muscle fibers, primarily in skeletal and cardiac muscle. It connects the intracellular cytoskeleton (actin) to the transmembrane proteins α- and β-dystroglycan, which are connected to the extracellular matrix (ECM). Loss of dystrophin → myonecrosis. ↑ CK and aldolase; genetic testing confirms diagnosis. 
Becker	X-linked disorder typically due to **non-frameshift** deletions in dystrophin gene (partially functional instead of truncated). Less severe than Duchenne (Becker is better). Onset in adolescence or early adulthood.	Deletions can cause both Duchenne and Becker muscular dystrophies. ⅔ of cases have large deletions spanning one or more exons.
Myotonic dystrophy	Autosomal dominant. CTG trinucleotide repeat expansion in the *DMPK* gene → abnormal expression of myotonin protein kinase → myotonia (eg, difficulty releasing hand from handshake), muscle wasting, cataracts, testicular atrophy, frontal balding, arrhythmia.	Cataracts, Toupee (early balding in men), Gonadal atrophy.

(c) ketabton.com: The Digital Library

Rett syndrome

Sporadic disorder seen almost exclusively in girls (affected males die in utero or shortly after birth). Most cases are caused by de novo mutation of *MECP2* on X chromosome. Symptoms of **Rett** syndrome usually appear between ages 1–4 and are characterized by regression (**Rett**urn) in motor, verbal, and cognitive abilities; ataxia; seizures; growth failure; and stereotyped hand-wringing.

Fragile X syndrome

X-linked dominant inheritance. Trinucleotide repeat in *FMR1* gene → hypermethylation → ↓ expression. Most common inherited cause of intellectual disability (Down syndrome is the most common genetic cause, but most cases occur sporadically).

Findings: post-pubertal macroorchidism (enlarged testes), long face with a large jaw, large everted ears, autism, mitral valve prolapse, hypermobile joints.

Trinucleotide repeat expansion [$(CGG)_n$] occurs during oogenesis.

Trinucleotide repeat expansion diseases

Huntington disease, **my**otonic dystrophy, **fragile X** syndrome, and **Frie**dreich ataxia.

May show genetic anticipation (disease severity ↑ and age of onset ↓ in successive generations).

Try (**tri**nucleotide) **hunting** for **my fragile** cage-**free** eggs (**X**).

DISEASE	TRINUCLEOTIDE REPEAT	MODE OF INHERITANCE	MNEMONIC
Huntington disease	$(\mathbf{CAG})_n$	AD	**C**audate has ↓ **A**Ch and **G**ABA
Myotonic dystrophy	$(\mathbf{CTG})_n$	AD	**C**ataracts, **T**oupee (early balding in men), **G**onadal atrophy in men, reduced fertility in women
Fragile X syndrome	$(\mathbf{CGG})_n$	XD	**C**hin (protruding), **G**iant **G**onads
Friedreich ataxia	$(\mathbf{GAA})_n$	AR	Ataxic **GAA**it

(c) ketabton.com: The Digital Library

Autosomal trisomies

Down syndrome (trisomy 21)

Single palmar crease Rx

Findings: intellectual disability, flat facies, prominent epicanthal folds, single palmar crease, incurved 5th finger, gap between 1st 2 toes, duodenal atresia, Hirschsprung disease, congenital heart disease (eg, ASD), Brushfield spots. Associated with early-onset Alzheimer disease (chromosome 21 codes for amyloid precursor protein), ↑ risk of AML/ALL.

95% of cases due to meiotic nondisjunction (↑ with advanced maternal age; from 1:1500 in women < 20 to 1:25 in women > 45 years old).

4% of cases due to unbalanced Robertsonian translocation, most typically between chromosomes 14 and 21. Only 1% of cases are due to postfertilization mitotic error.

Incidence 1:700.

Drinking age (21).

Most common viable chromosomal disorder and most common cause of genetic intellectual disability.

First-trimester ultrasound commonly shows ↑ nuchal translucency and hypoplastic nasal bone. Markers for Down syndrome are HI up: ↑ hCG, ↑ inhibin.

The 5 A's of Down syndrome:

- Advanced maternal age
- Atresia (duodenal)
- Atrioventricular septal defect
- Alzheimer disease (early onset)
- AML/ALL

Edwards syndrome (trisomy 18)

Overlapping fingers Rx

Findings: PRINCE Edward—Prominent occiput, Rocker-bottom feet, Intellectual disability, Nondisjunction, Clenched fists with overlapping fingers, low-set Ears, micrognathia (small jaw), congenital heart disease, omphalocele, myelomeningocele. Death usually occurs by age 1 year.

Incidence 1:8000.

Election age (18).

2nd most common autosomal trisomy resulting in live birth (most common is Down syndrome).

Patau syndrome (trisomy 13)

Cutis aplasia Rx

Findings: severe intellectual disability, rocker-bottom feet, microphthalmia, microcephaly, cleft liP/Palate, holoProsencephaly, Polydactyly, cutis aPlasia, congenital heart (Pump) disease, Polycystic kidney disease, omphalocele. Death usually occurs by age 1.

Incidence 1:15,000.

Puberty (13).

Defect in fusion of prechordal mesoderm → midline defects.



Rx

1st trimester screening		
Trisomy	β-hCG	PAPP-A
21	↑	↓
18	↓	↓
13	↓	↓

2nd trimester screening				
Trisomy	β-hCG	Inhibin A	Estriol	AFP
21	↑	↑	↓	↓
18	↓	— or ↓	↓	↓
13	—	—	—	—

(c) ketabton.com: The Digital Library

Genetic disorders by chromosome

CHROMOSOME	SELECTED EXAMPLES
3	von Hippel-Lindau disease, renal cell carcinoma
4	ADPKD (*PKD2*), achondroplasia, Huntington disease
5	Cri-du-chat syndrome, familial adenomatous polyposis
6	Hemochromatosis (*HFE*)
7	Williams syndrome, cystic fibrosis
9	Friedreich ataxia, tuberous sclerosis *(TSC1)*
11	Wilms tumor, β-globin gene defects (eg, sickle cell disease, β-thalassemia), MEN1
13	Patau syndrome, Wilson disease, retinoblastoma (*RB1*), *BRCA2*
15	Prader-Willi syndrome, Angelman syndrome, Marfan syndrome
16	ADPKD (*PKD1*), α-globin gene defects (eg, α-thalassemia), tuberous sclerosis *(TSC2)*
17	Neurofibromatosis type 1, *BRCA1*, *TP53*
18	Edwards syndrome
21	Down syndrome
22	Neurofibromatosis type 2, DiGeorge syndrome (22q11)
X	Fragile X syndrome, X-linked agammaglobulinemia, Klinefelter syndrome (XXY)

Robertsonian translocation

Chromosomal translocation that commonly involves chromosome pairs 21, 22, 13, 14, and 15.
One of the most common types of translocation. Occurs when the long arms of 2 acrocentric chromosomes (chromosomes with centromeres near their ends) fuse at the centromere and the 2 short arms are lost.
Balanced translocations normally do not cause any abnormal phenotype. Unbalanced translocations can result in miscarriage, stillbirth, and chromosomal imbalance (eg, Down syndrome, Patau syndrome).

Cri-du-chat syndrome

Cri du chat = **cry** of the **cat**. Congenital deletion on short arm of chromosome 5 (46,XX or XY, 5p−).
Findings: microcephaly, moderate to severe intellectual disability, high-pitched **cry**ing/**meow**ing, epicanthal folds, cardiac abnormalities (VSD).

Williams syndrome

A

Congenital microdeletion of long arm of chromosome 7 (deleted region includes elastin gene).
Findings: distinctive "**elf**in" facies A, intellectual disability, hypercalcemia, well-developed verbal skills, extreme friendliness with strangers, cardiovascular problems (eg, supravalvular aortic stenosis, renal artery stenosis). Think **Will** Ferrell in **Elf**.

(c) ketabton.com: The Digital Library

▸ BIOCHEMISTRY—NUTRITION

Vitamins: fat soluble	A, D, E, K. Absorption dependent on ileum and pancreas. Toxicity more common than for water-soluble vitamins because fat-soluble vitamins accumulate in fat.	Malabsorption syndromes with steatorrhea (eg, cystic fibrosis and celiac disease) or mineral oil intake can cause fat-soluble vitamin deficiencies.
Vitamins: water soluble	B_1 (thiamine: TPP) B_2 (riboflavin: FAD, FMN) B_3 (niacin: NAD^+) B_5 (pantothenic acid: CoA) B_6 (pyridoxine: PLP) B_7 (biotin) B_9 (folate) B_{12} (cobalamin) C (ascorbic acid)	All wash out easily from body except B_{12} and B_9 (folate). B_{12} stored in liver for ~ 3–4 years. B_9 stored in liver for ~ 3–4 months. B-complex deficiencies often result in dermatitis, glossitis, and diarrhea. Can be coenzymes (eg, ascorbic acid) or precursors to coenzymes (eg, FAD, NAD^+).

(c) ketabton.com: The Digital Library

Vitamin A

Includes retinal, retinol, retinoic acid.

FUNCTION

Antioxidant; constituent of visual pigments (**retina**l); essential for normal differentiation of epithelial cells into specialized tissue (pancreatic cells, mucus-secreting cells); prevents squamous metaplasia. Used to treat measles and acute promyelocytic leukemia (APL).

Retinol is vitamin **A**, so think **retin-A** (used topically for wrinkles and **A**cne).
Found in liver and leafy vegetables.
Supplementation in vitamin A-deficient measles patients may improve outcomes.
Use oral isotretinoin to treat severe cystic acne.
Use *all*-trans retinoic acid to treat acute promyelocytic leukemia.

DEFICIENCY



Night blindness (nyctalopia); dry, scaly skin (xerosis cutis); corneal squamous metaplasia → Bitot spots (keratin debris; foamy appearance on conjunctiva A); corneal degeneration (keratomalacia); immunosuppression.

EXCESS

Acute toxicity—nausea, vomiting, vertigo, and blurred vision.
Chronic toxicity—alopecia, dry skin (eg, scaliness), hepatic toxicity and enlargement, arthralgias, and idiopathic intracranial hypertension.
Teratogenic (cleft palate, cardiac abnormalities), therefore a ⊖ pregnancy test and two forms of contraception are required before isotretinoin (vitamin A derivative) is prescribed.

Isotretinoin **is terat**ogenic.

Vitamin B_1

Also called thiamine.

FUNCTION

In thiamine pyrophosphate (TPP), a cofactor for several dehydrogenase enzyme reactions:
- **B**ranched-chain ketoacid dehydrogenase
- **α**-ketoglutarate dehydrogenase (TCA cycle)
- **P**yruvate dehydrogenase (links glycolysis to TCA cycle)
- **T**ransketolase (HMP shunt)

DEFICIENCY

Impaired glucose breakdown → ATP depletion worsened by glucose infusion; highly aerobic tissues (eg, brain, heart) are affected first.
In alcoholic or malnourished patients, give thiamine before dextrose to ↓ risk of precipitating Wernicke encephalopathy.
Diagnosis made by ↑ in RBC transketolase activity following vitamin B_1 administration.

Be APT.
Spell beriberi as **B**er1**B**er1 to remember vitamin B_1.
Wernicke encephalopathy—acute, life-threatening, neurologic condition; classic triad of confusion, ophthalmoplegia, ataxia.
Korsakoff syndrome—amnestic disorder due to chronic alcohol consumption; presents with confabulation, personality changes, memory loss (permanent).
Wernicke-Korsakoff syndrome—damage to medial dorsal nucleus of thalamus, mammillary bodies. Presentation is combination of Wernicke encephalopathy and Korsakoff syndrome.
Dry beriberi—polyneuropathy, symmetric muscle wasting.
Wet beriberi—high-output cardiac failure (dilated cardiomyopathy), edema.

(c) ketabton.com: The Digital Library

Vitamin B_2	Also called riboflavin.	
FUNCTION	Component of flavins FAD and FMN, used as cofactors in redox reactions, eg, the succinate dehydrogenase reaction in the TCA cycle.	FAD and FMN are derived from riboFlavin ($B_2 \approx 2$ ATP).
DEFICIENCY	Cheilosis (inflammation of lips, scaling and fissures at the corners of the mouth), Corneal vascularization.	The 2 C's of B_2.

Vitamin B_3	Also called niacin, nicotinic acid.	
FUNCTION	Constituent of NAD^+, $NADP^+$ (used in redox reactions). Derived from tryptophan. Synthesis requires vitamins B_2 and B_6. Used to treat dyslipidemia; lowers levels of VLDL and raises levels of HDL.	NAD derived from Niacin ($B_3 \approx 3$ ATP).
DEFICIENCY A	Glossitis. Severe deficiency leads to pellagra, which can also be caused by Hartnup disease, malignant carcinoid syndrome (↑ tryptophan metabolism), and isoniazid (↓ vitamin B_6). Symptoms of pellagra: Diarrhea, Dementia (also hallucinations), Dermatitis (C3/C4 dermatome circumferential "broad collar" rash [Casal necklace], hyperpigmentation of sun-exposed limbs A).	The 3 D's of B_3. **Hartnup disease**—autosomal recessive. Deficiency of neutral amino acid (eg, tryptophan) transporters in proximal renal tubular cells and on enterocytes → neutral aminoaciduria and ↓ absorption from the gut → ↓ tryptophan for conversion to niacin → pellagra-like symptoms. Treat with high-protein diet and nicotinic acid. Deficiency of vitamin B_3 → **pell**agra.
EXCESS	Facial flushing (induced by prostaglandin, not histamine; can avoid by taking aspirin with niacin), hyperglycemia, hyperuricemia.	Excess of vitamin B_3 → **pod**agra.

Vitamin B_5	Also called pantothenic acid.	
FUNCTION	Essential component of coenzyme A (CoA, a cofactor for acyl transfers) and fatty acid synthase.	B_5 is "pento"thenic acid.
DEFICIENCY	Dermatitis, enteritis, alopecia, adrenal insufficiency.	

Vitamin B_6	Also called pyridoxine.
FUNCTION	Converted to pyridoxal phosphate (PLP), a cofactor used in transamination (eg, ALT and AST), decarboxylation reactions, glycogen phosphorylase. Synthesis of glutathione, cystathionine, heme, niacin, histamine, and neurotransmitters including serotonin, epinephrine, norepinephrine (NE), dopamine, and GABA.
DEFICIENCY	Convulsions, hyperirritability, peripheral neuropathy (deficiency inducible by isoniazid and oral contraceptives), sideroblastic anemia (due to impaired hemoglobin synthesis and iron excess).

(c) ketabton.com: The Digital Library

Vitamin B_7	Also called biotin.	
FUNCTION	Cofactor for carboxylation enzymes (which add a 1-carbon group): ▪ Pyruvate carboxylase: pyruvate (3C) → oxaloacetate (4C) ▪ Acetyl-CoA carboxylase: acetyl-CoA (2C) → malonyl-CoA (3C) ▪ Propionyl-CoA carboxylase: propionyl-CoA (3C) → methylmalonyl-CoA (4C)	
DEFICIENCY	Relatively rare. Dermatitis, enteritis, alopecia. Caused by long-term antibiotic use or excessive ingestion of raw egg whites.	"**Avid**in in egg whites **avid**ly binds biotin."

Vitamin B_9	Also called folate.	
FUNCTION	Converted to tetrahydrofolic acid (THF), a coenzyme for 1-carbon transfer/methylation reactions. Important for the synthesis of nitrogenous bases in DNA and RNA.	Found in leafy green vegetables. Absorbed in jejunum. **Fol**ate from **fol**iage. Small reserve pool stored primarily in the liver.
DEFICIENCY	Macrocytic, megaloblastic anemia; hypersegmented polymorphonuclear cells (PMNs); glossitis; no neurologic symptoms (as opposed to vitamin B_{12} deficiency). Labs: ↑ homocysteine, normal methylmalonic acid levels. Seen in alcoholism and pregnancy.	Deficiency can be caused by several drugs (eg, phenytoin, sulfonamides, methotrexate). Supplemental maternal folic acid at least 1 month prior to conception and during early pregnancy to ↓ risk of neural tube defects. Give vitamin B_9 for the 9 months of pregnancy.

(c) ketabton.com: The Digital Library

Vitamin B_{12}

Also called cobalamin.

FUNCTION

Cofactor for methionine synthase (transfers CH_3 groups as methylcobalamin) and methylmalonyl-CoA mutase. Important for DNA synthesis.

DEFICIENCY

Macrocytic, megaloblastic anemia; hypersegmented PMNs; paresthesias and subacute combined degeneration (degeneration of dorsal columns, lateral corticospinal tracts, and spinocerebellar tracts) due to abnormal myelin. Associated with ↑ serum homocysteine and methylmalonic acid levels, along with 2° folate deficiency. Prolonged deficiency → irreversible nerve damage.

Found in animal products.

Synthesized only by microorganisms. Very large reserve pool (several years) stored primarily in the liver. Deficiency caused by malabsorption (eg, sprue, enteritis, *Diphyllobothrium latum*, achlorhydria, bacterial overgrowth, alcohol excess), lack of intrinsic factor (eg, pernicious anemia, gastric bypass surgery), absence of terminal ileum (surgical resection, eg, for Crohn disease), certain drugs (eg, metformin), or insufficient intake (eg, veganism).

Anti-intrinsic factor antibodies diagnostic for pernicious anemia.

Folate supplementation can mask the hematologic symptoms of B_{12} deficiency, but not the neurologic symptoms.



Vitamin C

Also called ascorbic acid.

FUNCTION

Antioxidant; also facilitates iron absorption by reducing it to Fe^{2+} state. Necessary for hydroxylation of proline and lysine in collagen synthesis. Necessary for dopamine β-hydroxylase, which converts dopamine to NE.

Found in fruits and vegetables.

Pronounce "**absorb**ic" acid.

Ancillary treatment for methemoglobinemia by reducing Fe^{3+} to Fe^{2+}.

DEFICIENCY

Scurvy—swollen gums, easy bruising, petechiae, hemarthrosis, anemia, poor wound healing, perifollicular and subperiosteal hemorrhages, "corkscrew" hair.

Weakened immune response.

Vitamin **C** deficiency causes s**C**urvy due to a **C**ollagen synthesis defect.

EXCESS

Nausea, vomiting, diarrhea, fatigue, calcium oxalate nephrolithiasis. Can ↑ iron toxicity in predisposed individuals by increasing dietary iron absorption (ie, can worsen hereditary hemochromatosis or transfusion-related iron overload).

(c) ketabton.com: The Digital Library

Vitamin D	D_3 (cholecalciferol) from exposure of skin (stratum basale) to sun, ingestion of fish, milk, plants. D_2 (ergocalciferol) from ingestion of plants, fungi, yeasts. Both converted to 25-OH D_3 (storage form) in liver and to the active form 1,25-$(OH)_2$ D_3 (calcitriol) in kidney.
FUNCTION	↑ intestinal absorption of Ca^{2+} and PO_4^{3-}. ↑ bone mineralization at low levels. ↑ bone resorption at higher levels.
REGULATION	↑ PTH, ↓ Ca^{2+}, ↓ PO_4^{3-} → ↑ 1,25-$(OH)_2D_3$ production. 1,25-$(OH)_2D_3$ feedback inhibits its own production. ↑ PTH → ↑ Ca^{2+} reabsorption and ↓ PO_4^{3-} reabsorption in the kidney.
DEFICIENCY A	Rickets in children (deformity, such as genu varum "bowlegs" A), osteomalacia in adults (bone pain and muscle weakness), hypocalcemic tetany. Caused by malabsorption, ↓ sun exposure, poor diet, chronic kidney disease (CKD), advanced liver disease. Give oral vitamin D to breastfed infants. Deficiency is exacerbated by pigmented skin, premature birth.
EXCESS	Hypercalcemia, hypercalciuria, loss of appetite, stupor. Seen in granulomatous diseases (↑ activation of vitamin D by epithelioid macrophages).

Vitamin E	Includes tocopherol, tocotrienol.	
FUNCTION	Antioxidant (protects RBCs and membranes from free radical damage).	
DEFICIENCY	Hemolytic anemia, acanthocytosis, muscle weakness, demyelination of posterior columns (↓ position and vibration sensation) and spinocerebellar tract (ataxia).	Neurologic presentation may appear similar to vitamin B_{12} deficiency, but without megaloblastic anemia, hypersegmented neutrophils, or ↑ serum methylmalonic acid levels.
EXCESS	Risk of enterocolitis in infants.	High-dose supplementation may alter metabolism of vitamin K → enhanced anticoagulant effects of warfarin.

(c) ketabton.com: The Digital Library

Vitamin K

Includes phytomenadione, phylloquinone, phytonadione, menaquinone.

FUNCTION

Activated by epoxide reductase to the reduced form, which is a cofactor for the γ-carboxylation of glutamic acid residues on various proteins required for blood clotting. Synthesized by intestinal flora.

K is for **K**oagulation. Necessary for the maturation of clotting factors II, VII, IX, X, and proteins C and S. Warfarin inhibits vitamin K–dependent synthesis of these factors and proteins.

DEFICIENCY

Neonatal hemorrhage with ↑ PT and ↑ aPTT but normal bleeding time (neonates have sterile intestines and are unable to synthesize vitamin K). Can also occur after prolonged use of broad-spectrum antibiotics.

Not in breast milk; neonates are given vitamin K injection at birth to prevent hemorrhagic disease of the newborn.

Zinc

FUNCTION

Mineral essential for the activity of 100+ enzymes. Important in the formation of zinc fingers (transcription factor motif).

DEFICIENCY

Delayed wound healing, suppressed immunity, male hypogonadism, ↓ adult hair (axillary, facial, pubic), dysgeusia, anosmia. Associated with acrodermatitis enteropathica (A, defect in intestinal zinc absorption). May predispose to alcoholic cirrhosis.

Protein-energy malnutrition

Kwashiorkor

Protein malnutrition resulting in skin lesions, edema due to ↓ plasma oncotic pressure, liver malfunction (fatty change due to ↓ apolipoprotein synthesis). Clinical picture is small child with swollen abdomen A.

Kwashiorkor results from protein-deficient **MEALS**:

- **M**alnutrition
- **E**dema
- **A**nemia
- **L**iver (fatty)
- **S**kin lesions (eg, hyperkeratosis, dyspigmentation)

Marasmus

Malnutrition not causing edema. Diet is deficient in calories but no nutrients are entirely absent.

Marasmus results in **M**uscle wasting B.

(c) ketabton.com: The Digital Library

Ethanol metabolism



Fomepizole—blocks alcohol DH; antidote **F**or **O**verdoses of **M**ethanol or **E**thylene glycol.
Disulfiram— blocks acetaldehyde dehydrogenase → ↑ acetaldehyde → ↑ hangover symptoms → **dis**couraging drinking.
NAD^+ is the limiting reagent.
Alcohol dehydrogenase operates via zero-order kinetics.



Ethanol metabolism ↑ NADH/NAD^+ ratio in liver, causing:
1. Lactic acidosis—↑ pyruvate conversion to lactate
2. Fasting hypoglycemia—↓ gluconeogenesis due to ↑ conversion of OAA to malate
3. Ketoacidosis—diversion of acetyl-CoA into ketogenesis rather than TCA cycle
4. Hepatosteatosis— ↑ conversion of DHAP to glycerol-3-P (4A); acetyl-CoA diverges into fatty acid synthesis (4B), which combines with glycerol-3-P to synthesize triglycerides

↑ NADH/NAD^+ ratio inhibits TCA cycle → ↑ acetyl-CoA used in ketogenesis (→ ketoacidosis), lipogenesis (→ hepatosteatosis).

▶ BIOCHEMISTRY—METABOLISM

Metabolism sites

Mitochondria	Fatty acid oxidation (β-oxidation), acetyl-CoA production, TCA cycle, oxidative phosphorylation, ketogenesis.	
Cytoplasm	Glycolysis, HMP shunt, and synthesis of cholesterol (SER), proteins (ribosomes, RER), fatty acids, and nucleotides.	
Both	**H**eme synthesis, **U**rea cycle, **G**luconeogenesis.	**HUG**s take **two** (both).

(c) ketabton.com: The Digital Library

Enzyme terminology	An enzyme's name often describes its function. For example, glucokinase is an enzyme that catalyzes the phosphorylation of glucose using a molecule of ATP. The following are commonly used enzyme descriptors.
Kinase	Catalyzes transfer of a phosphate group from a high-energy molecule (usually ATP) to a substrate (eg, phosphofructokinase).
Phosphorylase	Adds inorganic phosphate onto substrate without using ATP (eg, glycogen phosphorylase).
Phosphatase	Removes phosphate group from substrate (eg, fructose-1,6-bisphosphatase).
Dehydrogenase	Catalyzes oxidation-reduction reactions (eg, pyruvate dehydrogenase).
Hydroxylase	Adds hydroxyl group (–OH) onto substrate (eg, tyrosine hydroxylase).
Carboxylase	Transfers CO_2 groups with the help of biotin (eg, pyruvate carboxylase).
Mutase	Relocates a functional group within a molecule (eg, vitamin B_{12}–dependent methylmalonyl-CoA mutase).
Synthase/synthetase	Joins two molecules together using a source of energy (eg, ATP, acetyl-CoA, nucleotide sugar).

Rate-determining enzymes of metabolic processes

PROCESS	ENZYME	REGULATORS
Glycolysis	Phosphofructokinase-1 (PFK-1)	AMP ⊕, fructose-2,6-bisphosphate ⊕ ATP ⊖, citrate ⊖
Gluconeogenesis	Fructose-1,6-bisphosphatase	AMP ⊖, fructose-2,6-bisphosphate ⊖
TCA cycle	Isocitrate dehydrogenase	ADP ⊕ ATP ⊖, NADH ⊖
Glycogenesis	Glycogen synthase	Glucose-6-phosphate ⊕, insulin ⊕, cortisol ⊕ Epinephrine ⊖, glucagon ⊖
Glycogenolysis	Glycogen phosphorylase	Epinephrine ⊕, glucagon ⊕, AMP ⊕ Glucose-6-phosphate ⊖, insulin ⊖, ATP ⊖
HMP shunt	Glucose-6-phosphate dehydrogenase (G6PD)	$NADP^+$ ⊕ NADPH ⊖
De novo pyrimidine synthesis	Carbamoyl phosphate synthetase II	ATP ⊕, PRPP ⊕ UTP ⊖
De novo purine synthesis	Glutamine-phosphoribosylpyrophosphate (PRPP) amidotransferase	AMP ⊖, inosine monophosphate (IMP) ⊖, GMP ⊖
Urea cycle	Carbamoyl phosphate synthetase I	N-acetylglutamate ⊕
Fatty acid synthesis	Acetyl-CoA carboxylase (ACC)	Insulin ⊕, citrate ⊕ Glucagon ⊖, palmitoyl-CoA ⊖
Fatty acid oxidation	Carnitine acyltransferase I	Malonyl-CoA ⊖
Ketogenesis	HMG-CoA synthase	
Cholesterol synthesis	HMG-CoA reductase	Insulin ⊕, thyroxine ⊕, estrogen ⊕ Glucagon ⊖, cholesterol ⊖

(c) ketabton.com: The Digital Library

Summary of pathways

1. Galactokinase *(mild galactosemia)*
2. Galactose-1-phosphate uridyltransferase *(severe galactosemia)*
3. Hexokinase/glucokinase
4. Glucose-6-phosphatase *(von Gierke disease)*
5. Glucose-6-phosphate dehydrogenase
6. Transketolase
7. Phosphofructokinase-1
8. Fructose-1,6-bisphosphatase
9. Fructokinase *(essential fructosuria)*
10. Aldolase B *(fructose intolerance)*
11. Aldolase B *(liver)*, A *(muscle)*
12. Triose phosphate isomerase
13. Pyruvate kinase
14. Pyruvate dehydrogenase
15. Pyruvate carboxylase
16. PEP carboxykinase
17. Citrate synthase
18. Isocitrate dehydrogenase
19. α-ketoglutarate dehydrogenase
20. Carbamoyl phosphate synthetase I
21. Ornithine transcarbamylase
22. Propionyl-CoA carboxylase
23. HMG-CoA reductase



ATP production

Aerobic metabolism of one glucose molecule produces 32 net ATP via malate-aspartate shuttle (heart and liver), 30 net ATP via glycerol-3-phosphate shuttle (muscle).

Anaerobic glycolysis produces only 2 net ATP per glucose molecule.

ATP hydrolysis can be coupled to energetically unfavorable reactions.

Arsenic causes glycolysis to produce zero net ATP.

(c) ketabton.com: The Digital Library

Activated carriers

CARRIER MOLECULE	CARRIED IN ACTIVATED FORM
ATP	Phosphoryl groups
NADH, NADPH, $FADH_2$	Electrons
CoA, lipoamide	Acyl groups
Biotin	CO_2
Tetrahydrofolates	1-carbon units
S-adenosylmethionine (SAM)	CH_3 groups
TPP	Aldehydes

Universal electron acceptors

Nicotinamides (NAD^+, $NADP^+$ from vitamin B_3) and flavin nucleotides (FAD from vitamin B_2).

NAD^+ is generally used in **catabolic** processes to carry reducing equivalents away as NADH.

NADPH is used in **anabolic** processes (eg, steroid and fatty acid synthesis) as a supply of reducing equivalents.

NADPH is a product of the HMP shunt.

NADPH is used in:

- Anabolic processes
- Respiratory burst
- Cytochrome P-450 system
- Glutathione reductase

Hexokinase vs glucokinase

Phosphorylation of glucose to yield glucose-6-phosphate is catalyzed by glucokinase in the liver and hexokinase in other tissues. Hexokinase sequesters glucose in tissues, where it is used even when glucose concentrations are low. At high glucose concentrations, glucokinase helps to store glucose in liver.

	Hexokinase	Glucokinase
Location	Most tissues, except liver and pancreatic β cells	Liver, β cells of pancreas
K_m	Lower (↑ affinity)	Higher (↓ affinity)
V_{max}	Lower (↓ capacity)	Higher (↑ capacity)
Induced by insulin	No	Yes
Feedback inhibition by	Glucose-6-phosphate	Fructose-6-phosphate

(c) ketabton.com: The Digital Library

Glycolysis regulation, key enzymes

Net glycolysis (cytoplasm):
Glucose + 2 P_i + 2 ADP + 2 NAD^+ → 2 pyruvate + 2 ATP + 2 NADH + 2 H^+ + 2 H_2O.

Equation not balanced chemically, and exact balanced equation depends on ionization state of reactants and products.

REQUIRE ATP

Glucose —(Hexokinase/glucokinase)→ Glucose-6-P

Glucose-6-P ⊖ hexokinase.
Fructose-6-P ⊖ glucokinase.

Fructose-6-P —(Phosphofructokinase-1 (rate-limiting step))→ Fructose-1,6-BP

AMP ⊕, fructose-2,6-bisphosphate ⊕.
ATP ⊖, citrate ⊖.

PRODUCE ATP

1,3-BPG ⟷ (Phosphoglycerate kinase) 3-PG

Phosphoenolpyruvate —(Pyruvate kinase)→ Pyruvate

Fructose-1,6-bisphosphate ⊕.
ATP ⊖, alanine ⊖.

Regulation by fructose-2,6-bisphosphate

Fructose bisphosphatase-2 (FBPase-2) and phosphofructokinase-2 (PFK-2) are the same bifunctional enzyme whose function is reversed by phosphorylation by protein kinase A.



Fasting state: ↑ glucagon → ↑ cAMP → ↑ protein kinase A → ↑ FBPase-2, ↓ PFK-2, less glycolysis, more gluconeogenesis.

Fa**B**ian the **P**easant (**FBP**) has to work hard when starving.

Fed state: ↑ insulin → ↓ cAMP → ↓ protein kinase A → ↓ FBPase-2, ↑ PFK-2, more glycolysis, less gluconeogenesis.

Prince **F**rederic**K** (**PFK**) works only when fed.

Pyruvate dehydrogenase complex

Mitochondrial enzyme complex linking glycolysis and TCA cycle. Differentially regulated in fed (active)/fasting (inactive) states.
Reaction: pyruvate + NAD^+ + CoA → acetyl-CoA + CO_2 + NADH.
Contains 3 enzymes requiring 5 cofactors:

1. **T**hiamine pyrophosphate (B_1)
2. **L**ipoic acid
3. **C**oA (B_5, pantothenic acid)
4. **F**AD (B_2, riboflavin)
5. **N**AD^+ (B_3, niacin)

Activated by: ↑ NAD^+/NADH ratio, ↑ ADP ↑ Ca^{2+}.

The complex is similar to the α-ketoglutarate dehydrogenase complex (same cofactors, similar substrate and action), which converts α-ketoglutarate → succinyl-CoA (TCA cycle).

The **L**ovely **C**oenzymes **F**or **N**erds.
Arsenic inhibits lipoic acid. Arsenic poisoning clinical findings: imagine a vampire (pigmentary skin changes, skin cancer), vomiting and having diarrhea, running away from a cutie (QT prolongation) with garlic breath.

(c) ketabton.com: The Digital Library

Pyruvate dehydrogenase complex deficiency	Causes a buildup of pyruvate that gets shunted to lactate (via LDH) and alanine (via ALT). X-linked.
FINDINGS	Neurologic defects, lactic acidosis, ↑ serum alanine starting in infancy.
TREATMENT	↑ intake of ketogenic nutrients (eg, high fat content or ↑ lysine and leucine).

Pyruvate metabolism



Functions of different pyruvate metabolic pathways (and their associated cofactors):

1. Alanine aminotransferase (B_6): alanine carries amino groups to the liver from muscle
2. Pyruvate carboxylase (biotin): oxaloacetate can replenish TCA cycle or be used in gluconeogenesis
3. Pyruvate dehydrogenase (B_1, B_2, B_3, B_5, lipoic acid): transition from glycolysis to the TCA cycle
4. Lactic acid dehydrogenase (B_3): end of anaerobic glycolysis (major pathway in RBCs, WBCs, kidney medulla, lens, testes, and cornea)

TCA cycle



Also called Krebs cycle. Pyruvate → acetyl-CoA produces 1 NADH, 1 CO_2.

The TCA cycle produces 3 NADH, 1 $FADH_2$, 2 CO_2, 1 GTP per acetyl-CoA = 10 ATP/acetyl-CoA (2× everything per glucose). TCA cycle reactions occur in the mitochondria.

α-ketoglutarate dehydrogenase complex requires the same cofactors as the pyruvate dehydrogenase complex (vitamins B_1, B_2, B_3, B_5, lipoic acid).

Citrate **I**s **K**rebs' **S**tarting **S**ubstrate **F**or **M**aking **O**xaloacetate.

(c) ketabton.com: The Digital Library

Electron transport chain and oxidative phosphorylation

NADH electrons from glycolysis enter mitochondria via the malate-aspartate or glycerol-3-phosphate shuttle. $FADH_2$ electrons are transferred to complex II (at a lower energy level than NADH). The passage of electrons results in the formation of a proton gradient that, coupled to oxidative phosphorylation, drives the production of ATP.



ATP PRODUCED VIA ATP SYNTHASE		
	1 NADH → 2.5 ATP; 1 $FADH_2$ → 1.5 ATP.	
OXIDATIVE PHOSPHORYLATION POISONS		
Electron transport inhibitors	Directly inhibit electron transport, causing a ↓ proton gradient and block of ATP synthesis.	Roten**one**: complex **one** inhibitor. "An-**3**-mycin" (antimycin) A: complex **3** inhibitor. Cyan**ide**, carbon monox**ide**, az**ide** (the **-ides**, 4 letters) inhibit complex **IV**.
ATP synthase inhibitors	Directly inhibit mitochondrial ATP synthase, causing an ↑ proton gradient. No ATP is produced because electron transport stops.	Oligomycin.
Uncoupling agents	↑ permeability of membrane, causing a ↓ proton gradient and ↑ O_2 consumption. ATP synthesis stops, but electron transport continues. Produces heat.	2,4-Dinitrophenol (used illicitly for weight loss), aspirin (fevers often occur after overdose), thermogenin in brown fat (has more mitochondria than white fat).

Gluconeogenesis, irreversible enzymes		**P**athway **P**roduces **F**resh **G**lucose.
Pyruvate carboxylase	In mitochondria. Pyruvate → oxaloacetate.	Requires biotin, ATP. Activated by acetyl-CoA.
Phosphoenolpyruvate carboxykinase	In cytosol. Oxaloacetate → phosphoenolpyruvate.	Requires GTP.
Fructose-1,6-bisphosphatase	In cytosol. Fructose-1,6-bisphosphate → fructose-6-phosphate.	Citrate ⊕, AMP ⊖, fructose 2,6-bisphosphate ⊖.
Glucose-6-phosphatase	In ER. Glucose-6-phosphate → glucose.	
	Occurs primarily in liver; serves to maintain euglycemia during fasting. Enzymes also found in kidney, intestinal epithelium. Deficiency of the key gluconeogenic enzymes causes hypoglycemia. (Muscle cannot participate in gluconeogenesis because it lacks glucose-6-phosphatase). Odd-chain fatty acids yield 1 propionyl-CoA during metabolism, which can enter the TCA cycle (as succinyl-CoA), undergo gluconeogenesis, and serve as a glucose source. Even-chain fatty acids cannot produce new glucose, since they yield only acetyl-CoA equivalents.	

(c) ketabton.com: The Digital Library

Pentose phosphate pathway

Also called HMP shunt. Provides a source of NADPH from abundantly available glucose-6-P (NADPH is required for reductive reactions, eg, glutathione reduction inside RBCs, fatty acid and cholesterol biosynthesis). Additionally, this pathway yields ribose for nucleotide synthesis. Two distinct phases (oxidative and nonoxidative), both of which occur in the cytoplasm. No ATP is used or produced.

Sites: lactating mammary glands, liver, adrenal cortex (sites of fatty acid or steroid synthesis), RBCs.

REACTIONS

Oxidative (irreversible)

Nonoxidative (reversible)



Glucose-6-phosphate dehydrogenase deficiency

NADPH is necessary to keep glutathione reduced, which in turn detoxifies free radicals and peroxides. ↓ NADPH in RBCs leads to hemolytic anemia due to poor RBC defense against oxidizing agents (eg, fava beans, sulfonamides, nitrofurantoin, primaquine/chloroquine, antituberculosis drugs). Infection (most common cause) can also precipitate hemolysis; inflammatory response produces free radicals that diffuse into RBCs, causing oxidative damage.

X-linked recessive disorder; most common human enzyme deficiency; more prevalent among African Americans. ↑ malarial resistance.

Heinz bodies—denatured globin chains precipitate within RBCs due to oxidative stress.

Bite cells—result from the phagocytic removal of **Heinz** bodies by splenic macrophages. Think, "**Bite** into some **Heinz** ketchup."



(c) ketabton.com: The Digital Library

Disorders of fructose metabolism

Essential fructosuria

Involves a defect in **fructokinase**. Autosomal recessive. A benign, asymptomatic condition (fructo**kin**ase deficiency is **kin**der), since fructose is not trapped in cells. Hexokinase becomes 1° pathway for converting fructose to fructose-6-phosphate.

Symptoms: fructose appears in blood and urine.

Disorders of fructose metabolism cause milder symptoms than analogous disorders of galactose metabolism.

Hereditary fructose intolerance

Hereditary deficiency of **aldolase B**. Autosomal recessive. Fructose-1-phosphate accumulates, causing a ↓ in available phosphate, which results in inhibition of glycogenolysis and gluconeogenesis. Symptoms present following consumption of fruit, juice, or honey. Urine dipstick will be ⊖ (tests for glucose only); reducing sugar can be detected in the urine (nonspecific test for inborn errors of carbohydrate metabolism).

Symptoms: hypoglycemia, jaundice, cirrhosis, vomiting.

Treatment: ↓ intake of fructose, sucrose (glucose + fructose), and sorbitol (metabolized to fructose).



Disorders of galactose metabolism

Galactokinase deficiency

Hereditary deficiency of **galactokinase**. Galactitol accumulates if galactose is present in diet. Relatively mild condition. Autosomal recessive.

Symptoms: galactose appears in blood (galactosemia) and urine (galactosuria); infantile cataracts. May present as failure to track objects or to develop a social smile. Galacto**kin**ase deficiency is **kin**der (benign condition).

Classic galactosemia

Absence of **galactose-1-phosphate uridyltransferase**. Autosomal recessive. Damage is caused by accumulation of toxic substances (including galactitol, which accumulates in the lens of the eye).

Symptoms develop when infant begins feeding (lactose present in breast milk and routine formula) and include failure to thrive, jaundice, hepatomegaly, infantile cataracts, intellectual disability. Can predispose to *E coli* sepsis in neonates.

Treatment: exclude galactose and lactose (galactose + glucose) from diet.



Fructose is to **A**ldolase **B** as **G**alactose is to **U**ridyl**T**ransferase (**FAB GUT**).

The more serious defects lead to PO_4^{3-} depletion.

(c) ketabton.com: The Digital Library

Sorbitol

An alternative method of trapping glucose in the cell is to convert it to its alcohol counterpart, sorbitol, via aldose reductase. Some tissues then convert sorbitol to fructose using sorbitol dehydrogenase; tissues with an insufficient amount/activity of this enzyme are at risk of intracellular sorbitol accumulation, causing osmotic damage (eg, cataracts, retinopathy, and peripheral neuropathy seen with chronic hyperglycemia in diabetes).

High blood levels of galactose also result in conversion to the osmotically active galactitol via aldose reductase.

Liver, **O**varies, and **Se**minal vesicles have both enzymes (they **LOSe** sorbitol).

Lens has primarily aldose reductase. **R**etina, **K**idneys, and **S**chwann cells have only aldose reductase (**LuRKS**).

Lactase deficiency

Insufficient lactase enzyme → dietary lactose intolerance. Lactase functions on the intestinal brush border to digest lactose (in milk and milk products) into glucose and galactose.

Primary: age-dependent decline after childhood (absence of lactase-persistent allele), common in people of Asian, African, or Native American descent.

Secondary: loss of intestinal brush border due to gastroenteritis (eg, rotavirus), autoimmune disease.

Congenital lactase deficiency: rare, due to defective gene.

Stool demonstrates ↓ pH and breath shows ↑ hydrogen content with lactose hydrogen breath test. Intestinal biopsy reveals normal mucosa in patients with hereditary lactose intolerance.

FINDINGS	Bloating, cramps, flatulence, osmotic diarrhea.
TREATMENT	Avoid dairy products or add lactase pills to diet; lactose-free milk.

Amino acids

Only L-amino acids are found in proteins.

Essential	**PVT TIM HaLL**: **P**henylalanine, **V**aline, **T**ryptophan, **T**hreonine, **I**soleucine, **M**ethionine, **H**istidine, **L**eucine, **L**ysine. Glucogenic: **Met**hionine, **his**tidine, **val**ine. We **met his val**entine, she is so **sweet** (**gluco**genic). Glucogenic/ketogenic: Isoleucine, phenylalanine, threonine, tryptophan. Ketogenic: **L**eucine, **L**ysine. The on**Ly** pure**Ly** ketogenic amino acids.
Acidic	Aspartic **acid**, glutamic **acid**. Negatively charged at body pH.
Basic	**Ar**ginine, **his**tidine, **lys**ine. Arginine is most **basic**. Histidine has no charge at body pH. Arginine and histidine are required during periods of growth. Arginine and lysine are ↑ in histones which bind negatively charged DNA. **His lys** (lies) **are basic**.

(c) ketabton.com: The Digital Library

Urea cycle

Amino acid catabolism results in the formation of common metabolites (eg, pyruvate, acetyl-CoA), which serve as metabolic fuels. Excess nitrogen generated by this process is converted to urea and excreted by the kidneys.

Ordinarily, Careless Crappers Are Also Frivolous About Urination.





Transport of ammonia by alanine



Hyperammonemia



Can be acquired (eg, liver disease) or hereditary (eg, urea cycle enzyme deficiencies).

Presents with flapping tremor (eg, asterixis), slurring of speech, somnolence, vomiting, cerebral edema, blurring of vision.

↑ NH_3 changes relative amounts of α-ketoglutarate, glutamate, GABA, and glutamine to favor ↑ glutamine. CNS toxicity may involve ↓ GABA, ↓ α-ketoglutarate, TCA cycle inhibition, and cerebral edema due to glutamine-induced osmotic shifts.

Treatment: limit protein in diet.

May be given to ↓ ammonia levels:

- Lactulose to acidify GI tract and trap NH_4^+ for excretion.
- Antibiotics (eg, rifaximin, neomycin) to ↓ ammoniagenic bacteria.
- Benzoate, phenylacetate, or phenylbutyrate react with glycine or glutamine, forming products that are excreted renally.



(c) ketabton.com: The Digital Library

Ornithine transcarbamylase deficiency

Most common urea cycle disorder. X-linked recessive (vs other urea cycle enzyme deficiencies, which are autosomal recessive). Interferes with the body's ability to eliminate ammonia. Often evident in the first few days of life, but may present later. Excess carbamoyl phosphate is converted to orotic acid (part of the pyrimidine synthesis pathway).

Findings: ↑ orotic acid in blood and urine, ↓ BUN, symptoms of hyperammonemia. No megaloblastic anemia (vs orotic aciduria).

Amino acid derivatives



BH_4 = tetrahydrobiopterin

Catecholamine synthesis/tyrosine catabolism



(c) ketabton.com: The Digital Library

Phenylketonuria

Due to ↓ phenylalanine hydroxylase or ↓ tetrahydrobiopterin (BH_4) cofactor (malignant PKU). Tyrosine becomes essential. ↑ phenylalanine → ↑ phenyl ketones in urine.
Findings: intellectual disability, growth retardation, seizures, fair complexion, eczema, musty body odor.
Treatment: ↓ phenylalanine and ↑ tyrosine in diet, tetrahydrobiopterin supplementation.

Maternal PKU—lack of proper dietary therapy during pregnancy. Findings in infant: microcephaly, intellectual disability, growth retardation, congenital heart defects.

Autosomal recessive. Incidence ≈ 1:10,000.
Screening occurs 2–3 days after birth (normal at birth because of maternal enzyme during fetal life).
Phenyl ketones—phenylacetate, phenyllactate, and phenylpyruvate.
Disorder of **aromatic** amino acid metabolism → musty body **odor.**
PKU patients must avoid the artificial sweetener aspartame, which contains phenylalanine.

Maple syrup urine disease

Blocked degradation of **branched** amino acids (**I**soleucine, **L**eucine, **V**aline) due to ↓ branched-chain α-ketoacid dehydrogenase (B_1). Causes ↑ α-ketoacids in the blood, especially those of leucine.
Treatment: restriction of isoleucine, leucine, valine in diet, and thiamine supplementation.

Autosomal recessive.
Presentation: vomiting, poor feeding, urine smells like maple syrup/burnt sugar. Causes severe CNS defects, intellectual disability, death.
I Love **V**ermont **maple syrup** from maple trees (with **B_1ranches**).

Alkaptonuria

Congenital deficiency of homogentisate oxidase in the degradative pathway of tyrosine to fumarate → pigment-forming homogentisic acid builds up in tissue A. Autosomal recessive. Usually benign.
Findings: bluish-black connective tissue, ear cartilage, and sclerae (ochronosis); urine turns black on prolonged exposure to air. May have debilitating arthralgias (homogentisic acid toxic to cartilage).

Homocystinuria

Causes (all autosomal recessive):
- Cystathionine synthase deficiency (treatment: ↓ methionine, ↑ cysteine, ↑ B_6, B_{12}, and folate in diet)
- ↓ affinity of cystathionine synthase for pyridoxal phosphate (treatment: ↑↑ B_6 and ↑ cysteine in diet)
- Methionine synthase (homocysteine methyltransferase) deficiency (treatment: ↑ methionine in diet)
- Methylenetetrahydrofolate reductase (MTHFR) deficiency (treatment: ↑ folate in diet)

All forms result in excess homocysteine.
HOMOCYstinuria: ↑↑ **H**omocysteine in urine, **O**steoporosis, **M**arfanoid habitus, **O**cular changes (downward and inward lens subluxation), **C**ardiovascular effects (thrombosis and atherosclerosis → stroke and MI), k**Y**phosis, intellectual disability, fair complexion. In homocystinuria, lens subluxes "down and in" (vs Marfan, "up and fans out").



(c) ketabton.com: The Digital Library

Cystinuria

Hereditary defect of renal PCT and intestinal amino acid transporter that prevents reabsorption of **C**ystine, **O**rnithine, **L**ysine, and **A**rginine (**COLA**).

Excess cystine in the urine can lead to recurrent precipitation of hexagonal cystine stones A.

Treatment: urinary alkalinization (eg, potassium citrate, acetazolamide) and chelating agents (eg, penicillamine) ↑ solubility of cystine stones; good hydration.

Autosomal recessive. Common (1:7000). Urinary cyanide-nitroprusside test is diagnostic.

Cystine is made of 2 cysteines connected by a disulfide bond.

Organic acidemias

Most commonly present in infancy with poor feeding, vomiting, hypotonia, high anion gap metabolic acidosis, hepatomegaly, seizures. Organic acid accumulation:

- Inhibits gluconeogenesis → ↓ fasting blood glucose levels, ↑ ketoacidosis → high anion gap metabolic acidosis
- Inhibits urea cycle → hyperammonemia

Propionic acidemia	Deficiency of propionyl-CoA carboxylase → ↑ propionyl-CoA, ↓ methylmalonic acid.	Treatment: low-protein diet limited in substances that metabolize into propionyl-CoA: **V**aline, **O**dd-chain fatty acids, **M**ethionine, **I**soleucine, **T**hreonine (**VOMIT**).
Methylmalonic acidemia	Deficiency of methylmalonyl-CoA mutase or vitamin B_{12}.	

Glycogen regulation by insulin and glucagon/epinephrine



(c) ketabton.com: The Digital Library

Glycogen	Branches have α-(1,6) bonds; linkages have α-(1,4) bonds.
Skeletal muscle	Glycogen undergoes glycogenolysis → glucose-1-phosphate → glucose-6-phosphate, which is rapidly metabolized during exercise.
Hepatocytes	Glycogen is stored and undergoes glycogenolysis to maintain blood sugar at appropriate levels. Glycogen phosphorylase ❹ liberates glucose-1-phosphate residues off branched glycogen until 4 glucose units remain on a branch. Then 4-α-D-glucanotransferase (debranching enzyme ❺) moves 3 of the 4 glucose units from the branch to the linkage. Then α-1,6-glucosidase (debranching enzyme ❻) cleaves off the last residue, liberating glucose. "Limit dextrin" refers to the two to four residues remaining on a branch after glycogen phosphorylase has already shortened it.



Glycogen storage disease type

- I Von Gierke disease
- II Pompe disease
- III Cori disease
- V McArdle disease

Glycogen enzymes

1. UDP-glucose pyrophosphorylase
2. Glycogen synthase
3. Branching enzyme
4. Glycogen phosphorylase
5. Debranching enzyme (4-α-D-glucanotransferase)
6. Debranching enzyme (α-1,6-glucosidase)
7. α-1,4-glucosidase Rx

Note: A small amount of glycogen is degraded in lysosomes by ❼ α-1,4-glucosidase (acid maltase).

(c) ketabton.com: The Digital Library

Glycogen storage diseases

At least 15 types have been identified, all resulting in abnormal glycogen metabolism and an accumulation of glycogen within cells. Periodic acid–Schiff stain identifies glycogen and is useful in identifying these diseases.

Very Poor Carbohydrate Metabolism.
Types I, II, III, and V are autosomal recessive.

DISEASE	FINDINGS	DEFICIENT ENZYME	COMMENTS
Von Gierke disease (type I)	Severe fasting hypoglycemia, ↑↑ Glycogen in liver and kidneys, ↑ blood lactate, ↑ triglycerides, ↑ uric acid (Gout), and hepatomegaly, renomegaly. Liver does not regulate blood glucose.	Glucose-6-phosphatase	Treatment: frequent oral glucose/cornstarch; avoidance of fructose and galactose Impaired gluconeogenesis and glycogenolysis
Pompe disease (type II)	Cardiomegaly, hypertrophic cardiomyopathy, hypotonia, exercise intolerance, and systemic findings lead to early death.	Lysosomal acid α-1,4-glucosidase (acid maltase) with α-1,6-glucosidase activity	PomPe trashes the PumP (1st and 4th letter; heart, liver, and muscle)
Cori disease (type III)	Similar to von Gierke disease, but milder symptoms and normal blood lactate levels. Can lead to cardiomyopathy. Limit dextrin–like structures accumulate in cytosol.	Debranching enzymes (α-1,6-glucosidase and 4-α-D-glucanotransferase)	Gluconeogenesis is intact
McArdle disease (type V)	↑ glycogen in muscle, but muscle cannot break it down → painful Muscle cramps, Myoglobinuria (red urine) with strenuous exercise, and arrhythmia from electrolyte abnormalities. Second-wind phenomenon noted during exercise due to ↑ muscular blood flow.	Skeletal muscle glycogen phosphorylase (Myophosphorylase) Characterized by a flat venous lactate curve with normal rise in ammonia levels during exercise	Blood glucose levels typically unaffected McArdle = Muscle

(c) ketabton.com: The Digital Library

Lysosomal storage diseases

Each is caused by a deficiency in one of the many lysosomal enzymes. Results in an accumulation of abnormal metabolic products.

DISEASE	FINDINGS	DEFICIENT ENZYME	ACCUMULATED SUBSTRATE	INHERITANCE
Sphingolipidoses				
Tay-Sachs disease 	Progressive neurodegeneration, developmental delay, hyperreflexia, hyperacusis, "cherry-red" spot on macula A, lysosomes with onion skin, no hepatosplenomegaly (vs Niemann-Pick).	❶ HeXosaminidase A ("TAy-SaX")	GM_2 ganglioside	AR
Fabry disease	Early: triad of episodic peripheral neuropathy, angiokeratomas B, hypohidrosis. Late: progressive renal failure, cardiovascular disease.	❷ α-galactosidase A	Ceramide trihexoside (globotriaosylceramide)	XR
Metachromatic leukodystrophy	Central and peripheral demyelination with ataxia, dementia.	❸ Arylsulfatase A	Cerebroside sulfate	AR
Krabbe disease	Peripheral neuropathy, destruction of oligodendrocytes, developmental delay, optic atrophy, globoid cells.	❹ Galactocerebrosidase (galactosylceramidase)	Galactocerebroside, psychosine	AR
Gaucher disease	Most common. Hepatosplenomegaly, pancytopenia, osteoporosis, avascular necrosis of femur, bone crises, Gaucher cells C (lipid-laden macrophages resembling crumpled tissue paper).	❺ Glucocerebrosidase (β-glucosidase); treat with recombinant glucocerebrosidase	Glucocerebroside	AR
Niemann-Pick disease	Progressive neurodegeneration, hepatosplenomegaly, foam cells (lipid-laden macrophages) D, "cherry-red" spot on macula A.	❻ Sphingomyelinase	Sphingomyelin	AR
Mucopolysaccharidoses				
Hurler syndrome	Developmental delay, gargoylism, airway obstruction, corneal clouding, hepatosplenomegaly.	α-L-iduronidase	Heparan sulfate, dermatan sulfate	AR
Hunter syndrome	Mild Hurler + aggressive behavior, no corneal clouding.	Iduronate-2-sulfatase	Heparan sulfate, dermatan sulfate	XR



No man picks (**Niemann-Pick**) his nose with his **sphing**er (**sphing**omyelinase).
Hunters see clearly (no corneal clouding) and aggressively aim for the **X** (**X**-linked recessive).
↑ incidence of Tay-Sachs, Niemann-Pick, some forms of Gaucher disease in Ashkenazi Jews.

(c) ketabton.com: The Digital Library

Fatty acid metabolism



Fatty acid synthesis requires transport of citrate from mitochondria to cytosol. Predominantly occurs in liver, lactating mammary glands, and adipose tissue.

Long-chain fatty acid (LCFA) degradation requires carnitine-dependent transport into the mitochondrial matrix.

"SYtrate" = SYnthesis.
CARnitine = CARnage of fatty acids.

Systemic 1° carnitine deficiency—no cellular uptake of carnitine → no transport of LCFAs into mitochondria → toxic accumulation of LCFAs in the cytosol. Causes weakness, hypotonia, hypoketotic hypoglycemia, dilated cardiomyopathy.

Medium-chain acyl-CoA dehydrogenase deficiency—↓ ability to break down fatty acids into acetyl-CoA → accumulation of fatty acyl carnitines in the blood with hypoketotic hypoglycemia. Causes vomiting, lethargy, seizures, coma, liver dysfunction, hyperammonemia. Can lead to sudden death in infants or children. Treat by avoiding fasting.

(c) ketabton.com: The Digital Library

Ketone bodies

In the liver, fatty acids and amino acids are metabolized to acetoacetate and β-hydroxybutyrate (to be used in muscle and brain).

In prolonged starvation and diabetic ketoacidosis, oxaloacetate is depleted for gluconeogenesis. In alcoholism, excess NADH shunts oxaloacetate to malate. All of these processes lead to a buildup of acetyl-CoA, which is shunted into ketone body synthesis.

Ketone bodies: acetone, acetoacetate, β-hydroxybutyrate.

Breath smells like acetone (fruity odor).

Urine test for ketones can detect acetoacetate, but not β-hydroxybutyrate.

RBCs cannot utilize ketones; they strictly use glucose.

HMG-CoA lyase for ketone production.

HMG-CoA reductase for cholesterol synthesis.



(c) ketabton.com: The Digital Library

Metabolic fuel use



1g **carb**/protein = **4** kcal
1g **alcohol** = **7** kcal
1g **fatty acid** = **9** kcal
(# letters = # kcal)

Fasting and starvation	Priorities are to supply sufficient glucose to the brain and RBCs and to preserve protein.	
Fed state (after a meal)	Glycolysis and aerobic respiration.	Insulin stimulates storage of lipids, proteins, and glycogen.
Fasting (between meals)	Hepatic glycogenolysis (major); hepatic gluconeogenesis, adipose release of FFA (minor).	Glucagon and epinephrine stimulate use of fuel reserves.
Starvation days 1–3	Blood glucose levels maintained by: ▪ Hepatic glycogenolysis ▪ Adipose release of FFA ▪ Muscle and liver, which shift fuel use from glucose to FFA ▪ Hepatic gluconeogenesis from peripheral tissue lactate and alanine, and from adipose tissue glycerol and propionyl-CoA (from odd-chain FFA—the only triacylglycerol components that contribute to gluconeogenesis)	Glycogen reserves depleted after day 1. RBCs lack mitochondria and therefore cannot use ketones. 
Starvation after day 3	Adipose stores (ketone bodies become the main source of energy for the brain). After these are depleted, vital protein degradation accelerates, leading to organ failure and death. Amount of excess stores determines survival time.	

(c) ketabton.com: The Digital Library

Lipid transport



(c) ketabton.com: The Digital Library

Key enzymes in lipid transport

Cholesteryl ester transfer protein	Mediates transfer of cholesteryl esters to other lipoprotein particles.
Hepatic lipase	Degrades TGs remaining in IDL.
Hormone-sensitive lipase	Degrades TGs stored in adipocytes.
Lecithin-cholesterol acyltransferase	Catalyzes esterification of ⅔ of plasma cholesterol.
Lipoprotein lipase	Degrades TGs in circulating chylomicrons.
Pancreatic lipase	Degrades dietary TGs in small intestine.
PCSK9	Degrades LDL receptor → ↑ serum LDL. Inhibition → ↑ LDL receptor recycling → ↓ serum LDL.



Major apolipoproteins

Apolipoprotein	Function	Chylomicron	Chylomicron remnant	VLDL	IDL	LDL	HDL
E	Mediates remnant uptake (everything except LDL)	✓	✓	✓	✓		✓
A-I	Found only on alpha-lipoproteins (HDL), activates LCAT						✓
C-II	Lipoprotein lipase cofactor that catalyzes cleavage.	✓		✓	✓		✓
B-48	Mediates chylomicron secretion into lymphatics Only on particles originating from the intestines	✓	✓				
B-100	Binds LDL receptor Only on particles originating from the liver			✓	✓	✓	

(c) ketabton.com: The Digital Library

Lipoprotein functions	Lipoproteins are composed of varying proportions of cholesterol, TGs, and phospholipids. LDL and HDL carry the most cholesterol. Cholesterol is needed to maintain cell membrane integrity and synthesize bile acids, steroids, and vitamin D.
Chylomicron	Delivers dietary TGs to peripheral tissues. Delivers cholesterol to liver in the form of chylomicron remnants, which are mostly depleted of their TGs. Secreted by intestinal epithelial cells.
VLDL	Delivers hepatic TGs to peripheral tissue. Secreted by liver.
IDL	Delivers TGs and cholesterol to liver. Formed from degradation of VLDL.
LDL	Delivers hepatic cholesterol to peripheral tissues. Formed by hepatic lipase modification of IDL in the liver and peripheral tissue. Taken up by target cells via receptor-mediated endocytosis. **L**DL is **L**ethal.
HDL	Mediates reverse cholesterol transport from peripheral tissues to liver. Acts as a repository for apolipoproteins C and E (which are needed for chylomicron and VLDL metabolism). Secreted from both liver and intestine. Alcohol ↑ synthesis. **H**DL is **H**ealthy.

Abetalipoproteinemia	Autosomal recessive. Mutation in gene that encodes microsomal transfer protein (*MTP*). Chylomicrons, VLDL, LDL absent. Deficiency in ApoB-48, ApoB-100. Affected infants present with severe fat malabsorption, steatorrhea, failure to thrive. Later manifestations include retinitis pigmentosa, spinocerebellar degeneration due to vitamin E deficiency, progressive ataxia, acanthocytosis. Intestinal biopsy shows lipid-laden enterocytes. Treatment: restriction of long-chain fatty acids, large doses of oral vitamin E.

Familial dyslipidemias

TYPE	INHERITANCE	PATHOGENESIS	↑ BLOOD LEVEL	CLINICAL
I—Hyper-chylomicronemia	AR	Lipoprotein lipase or apolipoprotein C-II deficiency	Chylomicrons, TG, cholesterol	Pancreatitis, hepatosplenomegaly, and eruptive/pruritic xanthomas (no ↑ risk for atherosclerosis). Creamy layer in supernatant.
II—Familial hyper-cholesterolemia	AD	Absent or defective LDL receptors, or defective ApoB-100	IIa: LDL, cholesterol IIb: LDL, cholesterol, VLDL	Heterozygotes (1:500) have cholesterol ≈ 300 mg/dL; homozygotes (very rare) have cholesterol ≥ 700 mg/dL. Accelerated atherosclerosis (may have MI before age 20), tendon (Achilles) xanthomas, and corneal arcus.
III—Dysbeta-lipoproteinemia	AR	Defective ApoE	Chylomicrons, VLDL	Premature atherosclerosis, tuberoeruptive and palmar xanthomas.
IV—Hyper-triglyceridemia	AD	Hepatic overproduction of VLDL	VLDL, TG	Hypertriglyceridemia (> 1000 mg/dL) can cause acute pancreatitis. Related to insulin resistance.

(c) ketabton.com: The Digital Library

HIGH-YIELD PRINCIPLES IN

Immunology

"I hate to disappoint you, but my rubber lips are immune to your charms."
—Batman & Robin

"The fully engaged heart is the antibody for the infection of violence."
—Mark Nepo

Learning the components of the immune system and their roles in host defense at the cellular level is essential for both the understanding of disease pathophysiology and clinical practice. Know the immune mechanisms of responses to vaccines. Both congenital and acquired immunodeficiencies are very testable. Cell surface markers are high yield for understanding immune cell interactions and for laboratory diagnosis. Know the roles and functions of major cytokines and chemokines.

Lymphoid Structures 96
Cellular Components 99
Immune Responses 104
Immunosuppressants 120

FAS1_2019_02-Immunology.indd 95 11/7/19 3:24 PM

(c) ketabton.com: The Digital Library

▸ IMMUNOLOGY—LYMPHOID STRUCTURES

Immune system organs

1° organs:
- Bone marrow—immune cell production, B cell maturation
- Thymus—T cell maturation

2° organs:
- Spleen, lymph nodes, tonsils, Peyer patches
- Allow immune cells to interact with antigen

Lymph node

A 2° lymphoid organ that has many afferents, 1 or more efferents. Encapsulated, with trabeculae A B. Functions are nonspecific filtration by macrophages, circulation of B and T cells, and immune response activation.

Follicle

Site of B-cell localization and proliferation. In outer cortex. 1° follicles are dense and quiescent. 2° follicles have pale central germinal centers and are active.

Medulla

Consists of medullary cords (closely packed lymphocytes and plasma cells) and medullary sinuses. Medullary sinuses communicate with efferent lymphatics and contain reticular cells and macrophages.

Paracortex

Contains T cells. Region of cortex between follicles and medulla. Contains high endothelial venules through which T and B cells enter from blood. Not well developed in patients with DiGeorge syndrome.

Paracortex enlarges in an extreme cellular immune response (eg, EBV and other viral infections → paracortical hyperplasia → lymphadenopathy).



(c) ketabton.com: The Digital Library

Lymphatic drainage associations



Lymph node cluster	Area of body drained	Associated pathology
Cervical, supraclavicular	Head and neck	Upper respiratory tract infection Infectious mononucleosis Kawasaki disease
Mediastinal	Trachea and esophagus	Pulmonary TB Sarcoidosis (bilateral) 1° lung cancer Granulomatous disease
Hilar	Lungs	
Axillary	Upper limb, breast, skin above umbilicus	Mastitis Metastasis (especially breast cancer)
Celiac	Liver, stomach, spleen, pancreas, upper duodenum	Mesenteric lymphadenitis Typhoid fever Ulcerative colitis Celiac disease
Superior mesenteric	Lower duodenum, jejunum, ileum, colon to splenic flexure	
Inferior mesenteric	Colon from splenic flexure to upper rectum	
Para-aortic	Testes, ovaries, kidneys, uterus	Metastasis
External iliac	Cervix, superior bladder, and body of uterus	Sexually transmitted infections Medial foot/leg cellulitis (superficial inguinal)
Internal iliac	Lower rectum to anal canal (above pectinate line), bladder, vagina (middle third), cervix, prostate	
Superficial inguinal	Anal canal (below pectinate line), skin below umbilicus (except popliteal area), scrotum, vulva	
Popliteal	Dorsolateral foot, posterior calf	Lateral foot/leg cellulitis

Right lymphatic duct drains right side of body above diaphragm into junction of the right subclavian and internal jugular vein

Thoracic duct drains below the diaphragm and left thorax and upper limb into junction of left subclavian and internal jugular veins (rupture of thoracic duct can cause chylothorax) ℞

(c) ketabton.com: The Digital Library

Spleen



Located in LUQ of abdomen, anterolateral to left kidney, protected by 9th-11th ribs.

Sinusoids are long, vascular channels in red pulp (red arrows in A) with fenestrated "barrel hoop" basement membrane.

- T cells are found in the periarteriolar lymphatic sheath (PALS) within the white pulp (white arrows in A).
- B cells are found in follicles within the white pulp.
- The marginal zone, in between the red pulp and white pulp, contains macrophages and specialized B cells, and is where antigen-presenting cells (APCs) capture blood-borne antigens for recognition by lymphocytes.

Splenic macrophages remove encapsulated bacteria.

Splenic dysfunction (eg, postsplenectomy state, sickle cell disease autosplenectomy): ↓ IgM → ↓ complement activation → ↓ C3b opsonization → ↑ susceptibility to encapsulated organisms.

Postsplenectomy blood findings:

- Howell-Jolly bodies (nuclear remnants)
- Target cells
- Thrombocytosis (loss of sequestration and removal)
- Lymphocytosis (loss of sequestration)

Vaccinate patients undergoing splenectomy or with splenic dysfunction against encapsulated organisms (pneumococci, Hib, meningococci).



Thymus





Located in the anterosuperior mediastinum. Site of T-cell differentiation and maturation. Encapsulated. **Th**ymus epithelium is derived from **Th**ird pharyngeal pouch (endoderm), whereas thymic lymphocytes are of mesodermal origin. Cortex is dense with immature T cells; **M**edulla is pale with **M**ature T cells and Hassall corpuscles A containing epithelial reticular cells.

Normal neonatal thymus "sail-shaped" on CXR B, involutes by age 3 years.

T cells = **T**hymus
B cells = **B**one marrow

Absent thymic shadow or hypoplastic thymus seen in some immunodeficiencies (eg, SCID, DiGeorge syndrome).

Thymoma—neoplasm of thymus. Associated with myasthenia gravis, superior vena cava syndrome, pure red cell aplasia, Good syndrome.

(c) ketabton.com: The Digital Library

▸ IMMUNOLOGY—CELLULAR COMPONENTS

Innate vs adaptive immunity

	Innate immunity	Adaptive immunity
COMPONENTS	Neutrophils, macrophages, monocytes, dendritic cells, natural killer (NK) cells (lymphoid origin), complement, physical epithelial barriers, secreted enzymes	T cells, B cells, circulating antibodies
MECHANISM	Germline encoded	Variation through V(D)J recombination during lymphocyte development
RESISTANCE	Resistance persists through generations; does not change within an organism's lifetime	Microbial resistance not heritable
RESPONSE TO PATHOGENS	Nonspecific Occurs rapidly (minutes to hours) No memory response	Highly specific, refined over time Develops over long periods; memory response is faster and more robust
SECRETED PROTEINS	Lysozyme, complement, C-reactive protein (CRP), defensins, cytokines	Immunoglobulins
KEY FEATURES IN PATHOGEN RECOGNITION	Toll-like receptors (TLRs): pattern recognition receptors that recognize pathogen-associated molecular patterns (PAMPs) and lead to activation of NF-κB. Examples of PAMPs include LPS (gram ⊖ bacteria), flagellin (bacteria), nucleic acids (viruses)	Memory cells: activated B and T cells; subsequent exposure to a previously encountered antigen → stronger, quicker immune response

(c) ketabton.com: The Digital Library

Major histocompatibility complex I and II

MHC encoded by HLA genes. Present antigen fragments to T cells and bind T-cell receptors (TCRs).

	MHC I	MHC II
LOCI	HLA-A, HLA-B, HLA-C MHC I loci have 1 letter	HLA-DP, HLA-DQ, HLA-DR MHC II loci have 2 letters
BINDING	TCR and CD8	TCR and CD4
STRUCTURE	1 long chain, 1 short chain	2 equal-length chains (2 α, 2 β)
EXPRESSION	All nucleated cells, APCs, platelets (except RBCs)	APCs
FUNCTION	Present endogenous antigens (eg, viral or cytosolic proteins) to CD8+ cytotoxic T cells	Present exogenous antigens (eg, bacterial proteins) to CD4+ helper T cells
ANTIGEN LOADING	Antigen peptides loaded onto MHC I in RER after delivery via TAP (transporter associated with antigen processing)	Antigen loaded following release of invariant chain in an acidified endosome
ASSOCIATED PROTEINS	β_2-microglobulin	Invariant chain
STRUCTURE		

HLA subtypes associated with diseases

HLA SUBTYPE	DISEASE	MNEMONIC
A3	Hemochromatosis	HA3mochromatosis
B8	Addison disease, myasthenia gravis, Graves disease	Don't Be late(8), Dr. Addison, or else you'll send my patient to the grave
B27	Psoriatic arthritis, Ankylosing spondylitis, IBD-associated arthritis, Reactive arthritis	PAIR. Also called seronegative arthropathies
C	Psoriasis	
DQ2/DQ8	Celiac disease	I ate (8) too (2) much gluten at Dairy Queen
DR2	Multiple sclerosis, hay fever, SLE, Goodpasture syndrome	DRive 2 multiple hay pastures
DR3	DM type 1, SLE, Graves disease, Hashimoto thyroiditis, Addison disease	2-3, S-L-E
DR4	Rheumatoid arthritis, DM type 1, Addison disease	There are 4 walls in 1 "rheum" (room)
DR5	Hashimoto thyroiditis	Hashimoto is an odd Dr (DR3, DR5)

(c) ketabton.com: The Digital Library

Functions of natural killer cells	Lymphocyte member of innate immune system. Use perforin and granzymes to induce apoptosis of virally infected cells and tumor cells. Activity enhanced by IL-2, IL-12, IFN-α, and IFN-β. Induced to kill when exposed to a nonspecific activation signal on target cell and/or to an absence of MHC I on target cell surface. Also kills via antibody-dependent cell-mediated cytotoxicity (CD16 binds Fc region of bound IgG, activating the NK cell).

Major functions of B and T cells

B cells	Humoral immunity. Recognize and present antigen—undergo somatic hypermutation to optimize antigen specificity. Produce antibody—differentiate into plasma cells to secrete specific immunoglobulins. Maintain immunologic memory—memory B cells persist and accelerate future response to antigen.
T cells	Cell-mediated immunity. CD4+ T cells help B cells make antibodies and produce cytokines to recruit phagocytes and activate other leukocytes. CD8+ T cells directly kill virus-infected and tumor cells via perforin and granzymes (similar to NK cells). Delayed cell-mediated hypersensitivity (type IV). Acute and chronic cellular organ rejection. Rule of 8: MHC II × CD4 = 8; MHC I × CD8 = 8.

(c) ketabton.com: The Digital Library

Differentiation of T cells



Positive selection
Thymic cortex. T cells expressing TCRs capable of binding self-MHC on cortical epithelial cells survive.

Negative selection
Thymic medulla. T cells expressing TCRs with high affinity for self antigens undergo apoptosis or become regulatory T cells. Tissue-restricted self-antigens are expressed in the thymus due to the action of autoimmune regulator (AIRE); deficiency leads to autoimmune polyendocrine syndrome-1 (Chronic mucocutaneous candidiasis, Hypoparathyroidism, Adrenal insufficiency, Recurrent *Candida* infections). "Without AIRE, your body will CHAR".

Macrophage-lymphocyte interaction
Th1 cells secrete IFN-γ, which enhances the ability of monocytes and macrophages to kill microbes they ingest. This function is also enhanced by interaction of T cell CD40L with CD40 on macrophages. Macrophages also activate lymphocytes via antigen presentation.

Cytotoxic T cells
Kill virus-infected, neoplastic, and donor graft cells by inducing apoptosis.
Release cytotoxic granules containing preformed proteins (eg, perforin, granzyme B).
Cytotoxic T cells have CD8, which binds to MHC I on virus-infected cells.

Regulatory T cells
Help maintain specific immune tolerance by suppressing $CD4^+$ and $CD8^+$ T-cell effector functions.
Identified by expression of CD3, CD4, CD25, and FOXP3.
Activated regulatory T cells (Tregs) produce anti-inflammatory cytokines (eg, IL-10, TGF-β).

IPEX (Immune dysregulation, Polyendocrinopathy, Enteropathy, X-linked) syndrome—genetic deficiency of FOXP3 → autoimmunity. Characterized by enteropathy, endocrinopathy, nail dystrophy, dermatitis, and/or other autoimmune dermatologic conditions. Associated with diabetes in male infants.

(c) ketabton.com: The Digital Library

T- and B-cell activation

APCs: B cells, dendritic cells, Langerhans cells, macrophages.
Two signals are required for T-cell activation, B-cell activation, and class switching.

T-cell activation

1. Dendritic cell (specialized APC) samples and processes antigen, then migrates to the draining lymph node.
2. T-cell activation (signal 1): exogenous antigen is presented on MHC II and recognized by TCR on Th (CD4+) cell. Endogenous or cross-presented antigen is presented on MHC I to Tc (CD8+) cell.
3. Proliferation and survival (signal 2): costimulatory signal via interaction of B7 protein (CD80/86) on dendritic cell and CD28 on naïve T cell.
4. Activated Th cell produces cytokines. Tc cell able to recognize and kill virus-infected cell.



B-cell activation and class switching

1. Th-cell activation as above.
2. B-cell receptor–mediated endocytosis.
3. Exogenous antigen is presented on MHC II and recognized by TCR on Th cell.
4. CD40 receptor on B cell binds CD40 ligand (CD40L) on Th cell.
5. Th cells secrete cytokines that determine Ig class switching of B cells.
6. B cells are activated, undergo class switching and affinity maturation, and begin producing antibodies.



(c) ketabton.com: The Digital Library

▸ IMMUNOLOGY—IMMUNE RESPONSES

Antibody structure and function

Fab (containing the variable/hypervariable regions) consisting of light (L) and heavy (H) chains recognizes antigens. Fc region of IgM and IgG fixes complement. Heavy chain contributes to Fc and Fab regions. Light chain contributes only to Fab region.





Fab:
- **F**ragment, **a**ntigen **b**inding
- Determines idiotype: unique antigen-binding pocket; only 1 antigenic specificity expressed per B cell

Fc (5 C's):
- **C**onstant
- **C**arboxy terminal
- **C**omplement binding
- **C**arbohydrate side chains
- **C**onfers (determines) isotype (IgM, IgD, etc)

Generation of antibody diversity (antigen independent)
1. Random recombination of VJ (light-chain) or V(D)J (heavy-chain) genes
2. Random addition of nucleotides to DNA during recombination by terminal deoxynucleotidyl transferase (TdT)
3. Random combination of heavy chains with light chains

Generation of antibody specificity (antigen dependent)
4. Somatic hypermutation and affinity maturation (variable region)
5. Isotype switching (constant region)

(c) ketabton.com: The Digital Library

Immunoglobulin isotypes

All isotypes can exist as monomers. Mature, naïve B cells prior to activation express IgM and IgD on their surfaces. They may differentiate in germinal centers of lymph nodes by isotype switching (gene rearrangement; induced by cytokines and CD40L) into plasma cells that secrete IgA, IgE, or IgG.

Affinity refers to the individual antibody-antigen interaction, while avidity describes the cumulative binding strength of all antibody-antigen interactions in a multivalent molecule.

IgG

Main antibody in 2° response to an antigen. Most abundant isotype in serum. Fixes complement, opsonizes bacteria, neutralizes bacterial toxins and viruses. Only isotype that crosses the placenta (provides infants with passive immunity that starts to wane after birth). "IgG Greets the Growing fetus."

IgA



Prevents attachment of bacteria and viruses to mucous membranes; does not fix complement. Monomer (in circulation) or dimer (with J chain when secreted). Crosses epithelial cells by transcytosis. Produced in GI tract (eg, by Peyer patches) and protects against gut infections (eg, *Giardia*). Most produced antibody overall, but has lower serum concentrations. Released into secretions (tears, saliva, mucus) and breast milk. Picks up secretory component from epithelial cells, which protects the Fc portion from luminal proteases.

IgM



Produced in the 1° (immediate) response to an antigen. Fixes complement. Antigen receptor on the surface of B cells. Monomer on B cell, pentamer with J chain when secreted. Pentamer enables avid binding to antigen while humoral response evolves.

IgD

Unclear function. Found on surface of many B cells and in serum.

IgE



Binds mast cells and basophils; cross-links when exposed to allergen, mediating immediate (type I) hypersensitivity through release of inflammatory mediators such as histamine. Contributes to immunity to parasites by activating eosinophils.

Antigen type and memory

Thymus-independent antigens

Antigens lacking a peptide component (eg, lipopolysaccharides from gram ⊖ bacteria); cannot be presented by MHC to T cells. Weakly immunogenic; vaccines often require boosters and adjuvants (eg, capsular polysaccharide subunit of *Streptococcus pneumoniae* PPSV23 vaccine).

Thymus-dependent antigens

Antigens containing a protein component (eg, *Streptococcus pneumoniae* PCV13 vaccine, polysaccharides conjugated to diphtheria toxin-like protein). Class switching and immunologic memory occur as a result of direct contact of B cells with Th cells.

(c) ketabton.com: The Digital Library

Complement

System of hepatically synthesized plasma proteins that play a role in innate immunity and inflammation. Membrane attack complex (MAC) defends against gram ⊖ bacteria. The CH_{50} test is used to screen for activation of the classical complement pathway.

ACTIVATION PATHWAYS

Classic—IgG or IgM mediated.
Alternative—microbe surface molecules.
Lectin—mannose or other sugars on microbe surface.

GM makes **classic** cars.

FUNCTIONS

C3b—opsonization.
C3a, C4a, C5a—anaphylaxis.
C5a—neutrophil chemotaxis.
C5b-9 (MAC)—cytolysis.

C3b binds to lipopolysaccharides on bacteria.

Opsonins—C3b and IgG are the two 1° opsonins in bacterial defense; enhance phagocytosis. C3b also helps clear immune complexes.

Opsonin (Greek) = to prepare for eating.

Inhibitors—decay-accelerating factor (DAF, aka CD55) and C1 esterase inhibitor help prevent complement activation on self cells (eg, RBCs).



(c) ketabton.com: The Digital Library

Complement disorders

Complement protein deficiencies	
Early complement deficiencies (C1-C4)	Increased risk of severe, recurrent pyogenic sinus and respiratory tract infections. Increased risk of SLE.
Terminal complement deficiencies (C5–C9)	Increased susceptibility to recurrent *Neisseria* bacteremia.
Complement regulatory protein deficiencies	
C1 esterase inhibitor deficiency	Causes hereditary angioedema due to unregulated activation of kallikrein → ↑ bradykinin. Characterized by ↓ C4 levels. ACE inhibitors are contraindicated (also ↑ bradykinin).
Paroxysmal nocturnal hemoglobinuria 	A defect in the *PIGA* gene preventing the formation of glycosylphosphatidylinositol (GPI) anchors for complement inhibitors, such as decay-accelerating factor (DAF/CD55) and membrane inhibitor of reactive lysis (MIRL/CD59). Causes complement-mediated intravascular hemolysis → ↓ haptoglobin, dark urine A.

(c) ketabton.com: The Digital Library

Important cytokines	Acute (IL-1, IL-6, TNF-α), then recruit (IL-8, IL-12).	
SECRETED BY MACROPHAGES		
Interleukin-1	Causes fever, acute inflammation. Activates endothelium to express adhesion molecules. Induces chemokine secretion to recruit WBCs. Also called osteoclast-activating factor.	"**Hot T-bone** st**EAK**": IL-1: fever (**hot**). IL-2: stimulates **T** cells. IL-3: stimulates **bone** marrow. IL-4: stimulates Ig**E** production. IL-5: stimulates Ig**A** production. IL-6: stimulates a**K**ute-phase protein production.
Interleukin-6	Causes fever and stimulates production of acute-phase proteins.	
Tumor necrosis factor-α	Activates endothelium. Causes WBC recruitment, vascular leak.	Causes cachexia in malignancy. Maintains granulomas in TB. IL-1, IL-6, TNF-α can mediate fever and sepsis.
Interleukin-8	Major chemotactic factor for neutrophils.	"**Clean up** on **aisle 8**." Neutrophils are recruited by **IL-8** to **clear** infections.
Interleukin-12	Induces differentiation of T cells into Th1 cells. Activates NK cells.	
SECRETED BY ALL T CELLS		
Interleukin-2	Stimulates growth of helper, cytotoxic, and regulatory T cells, and NK cells.	
Interleukin-3	Supports growth and differentiation of bone marrow stem cells. Functions like GM-CSF.	
FROM Th1 CELLS		
Interferon-γ	Secreted by NK cells and T cells in response to antigen or IL-12 from macrophages; stimulates macrophages to kill phagocytosed pathogens. Inhibits differentiation of Th2 cells.	Also activates NK cells to kill virus-infected cells. Increases MHC expression and antigen presentation by all cells. Activates macrophages to induce granuloma formation.
FROM Th2 CELLS		
Interleukin-4	Induces differentiation of T cells into Th (**help**er) **2** cells. Promotes growth of **B** cells. Enhances class switching to Ig**E** and Ig**G**.	Ain't too proud **2 BEG 4 help**.
Interleukin-5	Promotes growth and differentiation of B cells. Enhances class switching to IgA. Stimulates growth and differentiation of eosinophils.	
Interleukin-10	Attenuates inflammatory response. Decreases expression of MHC class II and Th1 cytokines. Inhibits activated macrophages and dendritic cells. Also secreted by regulatory T cells.	TGF-β and IL-**10** **b**oth at**ten**uate the immune response.

(c) ketabton.com: The Digital Library

Respiratory burst

Also called oxidative burst. Involves the activation of the phagocyte NADPH oxidase complex (eg, in neutrophils, monocytes), which utilizes O_2 as a substrate. Plays an important role in the immune response → rapid release of reactive oxygen species (ROS). NADPH plays a role in both the creation and neutralization of ROS. Myeloperoxidase contains a blue-green, heme-containing pigment that gives sputum its color.



Phagocytes of patients with CGD can utilize H_2O_2 generated by invading organisms and convert it to ROS. Patients are at ↑ risk for infection by catalase ⊕ species (eg, *S aureus*, *Aspergillus*) capable of neutralizing their own H_2O_2, leaving phagocytes without ROS for fighting infections.

Pyocyanin of *P aeruginosa* generates ROS to kill competing pathogens. Oxidative burst also leads to K^+ influx, which releases lysosomal enzymes. Lactoferrin is a protein found in secretory fluids and neutrophils that inhibits microbial growth via iron chelation.

Interferons

IFN-α, IFN-β, IFN-γ

MECHANISM

A part of innate host defense, **interfer**ons **interfer**e with both RNA and DNA viruses. Cells infected with a virus synthesize these glycoproteins, which act on local cells, priming them for viral defense by downregulating protein synthesis to resist potential viral replication and by upregulating MHC expression to facilitate recognition of infected cells. Also play a major role in activating antitumor immunity.

CLINICAL USE

Chronic HBV, Kaposi sarcoma, hairy cell leukemia, condyloma acuminatum, renal cell carcinoma, malignant melanoma, multiple sclerosis, chronic granulomatous disease.

ADVERSE EFFECTS

Flu-like symptoms, depression, neutropenia, myopathy, interferon-induced autoimmunity.

(c) ketabton.com: The Digital Library

Cell surface proteins

T cells	TCR (binds antigen-MHC complex) CD3 (associated with TCR for signal transduction) CD28 (binds B7 on APC)	
Helper T cells	CD4, CD40L, CXCR4/CCR5 (co-receptors for HIV)	
Cytotoxic T cells	CD8	
Regulatory T cells	CD4, CD25	
B cells	Ig (binds antigen) CD19, CD20, CD21 (receptor for Epstein-Barr virus), CD40 MHC II, B7	Must be 21 to drink Beer in a Barr
Macrophages	CD14 (receptor for PAMPs, eg, LPS), CD40 CCR5 MHC II, B7 (CD80/86) Fc and C3b receptors (enhanced phagocytosis)	
NK cells	CD16 (binds Fc of IgG), CD56 (suggestive marker for NK)	
Hematopoietic stem cells	CD34	

Anergy — State during which a cell cannot become activated by exposure to its antigen. T and B cells become anergic when exposed to their antigen without costimulatory signal (signal 2). Another mechanism of self-tolerance.

Passive vs active immunity

	Passive	Active
MEANS OF ACQUISITION	Receiving preformed antibodies	Exposure to exogenous antigens
ONSET	Rapid	Slow
DURATION	Short span of antibodies (half-life = 3 weeks)	Long-lasting protection (memory)
EXAMPLES	IgA in breast milk, maternal IgG crossing placenta, antitoxin, humanized monoclonal antibody	Natural infection, vaccines, toxoid
NOTES	After exposure to Tetanus toxin, Botulinum toxin, HBV, Varicella, Rabies virus, or Diphtheria toxin, unvaccinated patients are given preformed antibodies (passive)—"To Be Healed Very Rapidly before Dying"	Combined passive and active immunizations can be given for hepatitis B or rabies exposure

(c) ketabton.com: The Digital Library

Vaccination

Induces an active immune response (humoral and/or cellular) to specific pathogens.

VACCINE TYPE	DESCRIPTION	PROS/CONS	EXAMPLES
Live attenuated vaccine	Microorganism loses its pathogenicity but retains capacity for transient growth within inoculated host. Induces **cellular and humoral responses.** MMR and varicella vaccines can be given to HIV ⊕ patients without evidence of immunity if CD4 cell count ≥ 200 cells/mm^3.	Pros: induces strong, often lifelong immunity. Cons: may revert to virulent form. Often contraindicated in pregnancy and immunodeficiency.	**A**denovirus (nonattenuated, given to military recruits), **T**yphoid (Ty21a, oral), **P**olio (Sabin), **V**aricella (chickenpox), **Small**pox, **B**CG, **Y**ellow fever, **Inf**luenza (intranasal), **MMR**, **R**otavirus "**A**ttention **T**eachers! **P**lease **V**accinate **Small**, **B**eautiful **Y**oung **Inf**ants with **MMR** **R**egularly!"
Killed or inactivated vaccine	Pathogen is inactivated by heat or chemicals. Maintaining epitope structure on surface antigens is important for immune response. Mainly induces a **humoral response.**	Pros: safer than live vaccines. Cons: weaker immune response; booster shots usually required.	Hepatitis **A**, **T**yphoid (Vi polysaccharide, intramuscular), **R**abies, **I**nfluenza, **P**olio (Sal**K**) **A TRIP** could **K**ill you
Subunit	Includes only the antigens that best stimulate the immune system.	Pros: lower chance of adverse reactions. Cons: expensive, weaker immune response.	HBV (antigen = HBsAg), HPV (types 6, 11, 16, and 18), acellular pertussis (aP), *Neisseria meningitidis* (various strains), *Streptococcus pneumoniae*, *Haemophilus influenzae* type b.
Toxoid	Denatured bacterial toxin with an intact receptor binding site. Stimulates the immune system to make antibodies without potential for causing disease.	Pros: protects against the bacterial toxins. Cons: antitoxin levels decrease with time, may require a booster.	*Clostridium tetani*, *Corynebacterium diphtheriae*

(c) ketabton.com: The Digital Library

Hypersensitivity types Four types (**ABCD**): **A**naphylactic and **A**topic (type I), Anti**B**ody-mediated (type II), Immune **C**omplex (type III), **D**elayed (cell-mediated, type IV). Types I, II, and III are all antibody-mediated.

Type I hypersensitivity



Anaphylactic and atopic—two phases:

- Immediate (minutes): antigen crosslinks preformed IgE on presensitized mast cells → immediate degranulation → release of histamine (a vasoactive amine) and tryptase (a marker of mast cell activation).
- Late (hours): chemokines (attract inflammatory cells, eg, eosinophils) and other mediators (eg, leukotrienes) from mast cells → inflammation and tissue damage.

First (type) and **F**ast (anaphylaxis).

Test: skin test or blood test (ELISA) for allergen-specific IgE.

Example:

- Anaphylaxis (eg, food, drug, or bee sting allergies)
- Allergic asthma

Type II hypersensitivity



Antibody-dependent cellular cytotoxicity

Antibodies bind to cell-surface antigens → cellular destruction, inflammation, and cellular dysfunction.

Direct Coombs test—detects antibodies attached **direct**ly to the RBC surface.

Indirect Coombs test—detects presence of unbound antibodies in the serum

Cellular destruction—cell is opsonized (coated) by antibodies, leading to either:

- Phagocytosis and/or activation of complement system.
- NK cell killing (antibody-dependent cellular cytotoxicity).

Examples:

- Autoimmune-hemolytic anemia
- Immune thrombocytopenia
- Transfusion reactions
- Hemolytic disease of the newborn

Inflammation—binding of antibodies to cell surfaces → activation of complement system and Fc receptor-mediated inflammation.

Examples:

- Goodpasture syndrome
- Rheumatic fever
- Hyperacute transplant rejection

Cellular dysfunction—antibodies bind to cell surface receptors → abnormal blockade or activation of downstream process.

Examples:

- Myasthenia gravis
- Graves disease
- Pemphigus vulgaris

(c) ketabton.com: The Digital Library

Hypersensitivity types *(continued)*

Type III hypersensitivity



Immune complex—antigen-antibody (mostly IgG) complexes activate complement, which attracts neutrophils; neutrophils release lysosomal enzymes.

Can be associated with vasculitis and systemic manifestations.

Serum sickness—the prototypic immune complex disease. Antibodies to foreign proteins are produced and 1–2 weeks later, antibody-antigen complexes form and deposit in tissues → complement activation → inflammation and tissue damage.

Arthus reaction—a local subacute immune complex-mediated hypersensitivity reaction. Intradermal injection of antigen into a presensitized (has circulating IgG) individual leads to immune complex formation in the skin (eg, enhanced local reaction to a booster vaccination). Characterized by edema, necrosis, and activation of complement.

In type **III** reaction, imagine an immune complex as **3** things stuck together: antigen-antibody-complement.

Examples:

- SLE
- Polyarteritis nodosa
- Poststreptococcal glomerulonephritis

Fever, urticaria, arthralgia, proteinuria, lymphadenopathy occur 1–2 weeks after antigen exposure. Serum sickness-like reactions are associated with some drugs (may act as haptens, eg, penicillin) and infections (eg, hepatitis B).

Type IV hypersensitivity



Two mechanisms, each involving T cells:

1. Direct cell cytotoxicity: CD8+ cytotoxic T cells kill targeted cells.
2. Inflammatory reaction: effector CD4+ T cells recognize antigen and release inflammation-inducing cytokines (shown in illustration).

Response does not involve antibodies (vs types I, II, and III).

Examples: contact dermatitis (eg, poison ivy, nickel allergy) and graft-versus-host disease.

Tests: PPD for TB infection; patch test for contact dermatitis; *Candida* skin test for T cell immune function.

4**T**'s: **T** cells, **T**ransplant rejections, **T**B skin tests, **T**ouching (contact dermatitis).

Fourth (type) and **last** (delayed).

(c) ketabton.com: The Digital Library

Blood transfusion reactions

TYPE	PATHOGENESIS	TIMING	CLINICAL PRESENTATION	DONOR BLOOD	HOST BLOOD
Allergic/ anaphylactic reaction	Type I hypersensitivity reaction against plasma proteins in transfused blood IgA-deficient individuals should receive blood products without IgA	Within minutes to 2-3 hr (due to release of preformed inflammatory mediators in degranulating mast cells)	Allergies: urticaria, pruritus Anaphylaxis: wheezing, hypotension, respiratory arrest, shock		
Acute hemolytic transfusion reaction	Type II hypersensitivity reaction Typically causes intravascular hemolysis (ABO blood group incompatibility)	During transfusion or within 24 hr (due to preformed antibodies)	Fever, hypotension, tachypnea, tachycardia, flank pain, hemoglobinuria (intravascular), jaundice (extravascular)		
Febrile nonhemolytic transfusion reaction	Cytokines created by donor WBCs accumulate during storage of blood products Reactions prevented by leukoreduction of blood products	Within 1-6 hr (due to preformed cytokines)	Fever, headaches, chills, flushing More common in children		
Transfusion-related acute lung injury	Two-hit mechanism: ▪ Neutrophils are sequestered and primed in pulmonary vasculature due to recipient risk factors ▪ Neutrophils are activated by a product (eg, antileukocyte antibodies) in the transfused blood and release inflammatory mediators → ↑ capillary permeability → pulmonary edema	Within minutes to 6 hr	Respiratory distress, noncardiogenic pulmonary edema		
Delayed hemolytic transfusion reaction	Anamnestic response to a foreign antigen on donor RBCs (most commonly Rh or other minor blood group antigens) previously encountered by recipient Typically causes extravascular hemolysis	Onset over 24 hr Usually presents within 1-2 wk (due to slow destruction by reticuloendothelial system)	Generally self limited and clinically silent Mild fever, hyperbilirubinemia		

(c) ketabton.com: The Digital Library

Autoantibodies

AUTOANTIBODY	ASSOCIATED DISORDER
Anti-postsynaptic ACh receptor	Myasthenia gravis
Anti-presynaptic voltage-gated calcium channel	Lambert-Eaton myasthenic syndrome
Anti-β_2 glycoprotein I	Antiphospholipid syndrome
Antinuclear (ANA)	Nonspecific screening antibody, often associated with SLE
Anticardiolipin, lupus anticoagulant	SLE, antiphospholipid syndrome
Anti-dsDNA, anti-Smith	SLE
Antihistone	Drug-induced lupus
Anti-U1 RNP (ribonucleoprotein)	Mixed connective tissue disease
Rheumatoid factor (IgM antibody against IgG Fc region), anti-CCP (more specific)	Rheumatoid arthritis
Anti-Ro/SSA, anti-La/SSB	Sjögren syndrome
Anti-Scl-70 (anti-DNA topoisomerase I)	Scleroderma (diffuse)
Anticentromere	Limited scleroderma (CREST syndrome)
Antisynthetase (eg, anti-Jo-1), anti-SRP, anti-helicase (anti-Mi-2)	Polymyositis, dermatomyositis
Antimitochondrial	1° biliary cholangitis
Anti-smooth muscle	Autoimmune hepatitis type 1
MPO-ANCA/p-ANCA	Microscopic polyangiitis, eosinophilic granulomatosis with polyangiitis (Churg-Strauss syndrome), ulcerative colitis
PR3-ANCA/c-ANCA	Granulomatosis with polyangiitis (Wegener)
Anti-phospholipase A_2 receptor	1° membranous nephropathy
Anti-hemidesmosome	Bullous pemphigoid
Anti-desmoglein (anti-desmosome)	Pemphigus vulgaris
Antithyroglobulin, antithyroid peroxidase (antimicrosomal)	Hashimoto thyroiditis
Anti-TSH receptor	Graves disease
IgA anti-endomysial, IgA anti-tissue transglutaminase, IgA and IgG deamidated gliadin peptide	Celiac disease
Anti-glutamic acid decarboxylase, islet cell cytoplasmic antibodies	Type 1 diabetes mellitus
Antiparietal cell, anti-intrinsic factor	Pernicious anemia
Anti-glomerular basement membrane	Goodpasture syndrome

(c) ketabton.com: The Digital Library

Immunodeficiencies

DISEASE	DEFECT	PRESENTATION	FINDINGS
B-cell disorders			
X-linked (Bruton) agammaglobulinemia	Defect in *BTK*, a tyrosine kinase gene → no B-cell maturation; X-linked recessive (↑ in Boys)	Recurrent bacterial and enteroviral infections after 6 months (↓ maternal IgG)	Absent B cells in peripheral blood, ↓ Ig of all classes. Absent/scanty lymph nodes and tonsils (1° follicles and germinal centers absent) → live vaccines contraindicated
Selective IgA deficiency	Cause unknown Most common 1° immunodeficiency	Majority Asymptomatic Can see Airway and GI infections, Autoimmune disease, Atopy, Anaphylaxis to IgA-containing products	↓ IgA with normal IgG, IgM levels ↑ susceptibility to giardiasis Can cause false-positive β-hCG test
Common variable immunodeficiency	Defect in B-cell differentiation. Cause unknown in most cases	May present in childhood but usually diagnosed after puberty ↑ risk of autoimmune disease, bronchiectasis, lymphoma, sinopulmonary infections	↓ plasma cells, ↓ immunoglobulins
T-cell disorders			
Thymic aplasia	22q11 microdeletion; failure to develop 3rd and 4th pharyngeal pouches → absent thymus and parathyroids DiGeorge syndrome—thymic, parathyroid, cardiac defects Velocardiofacial syndrome—palate, facial, cardiac defects	CATCH-22: Cardiac defects (conotruncal abnormalities [eg, tetralogy of Fallot, truncus arteriosus]), Abnormal facies, Thymic hypoplasia → T-cell deficiency (recurrent viral/fungal infections), Cleft palate, Hypocalcemia 2° to parathyroid aplasia → tetany	↓ T cells, ↓ PTH, ↓ Ca^{2+} Thymic shadow absent on CXR
IL-12 receptor deficiency	↓ Th1 response; autosomal recessive	Disseminated mycobacterial and fungal infections; may present after administration of BCG vaccine	↓ IFN-γ Most common cause of Mendelian susceptibility to mycobacterial diseases (MSMD)
Autosomal dominant hyper-IgE syndrome (Job syndrome)	Deficiency of Th17 cells due to *STAT3* mutation → impaired recruitment of neutrophils to sites of infection	Cold (noninflamed) staphylococcal Abscesses, retained Baby teeth, Coarse facies, Dermatologic problems (eczema), ↑ IgE, bone Fractures from minor trauma	↑ IgE ↑ eosinophils Learn the ABCDEF's to get a Job!
Chronic mucocutaneous candidiasis	T-cell dysfunction Impaired cell-mediated immunity against *Candida* sp Classic form caused by defects in *AIRE*	Persistent noninvasive *Candida albicans* infections of skin and mucous membranes	Absent in vitro T-cell proliferation in response to *Candida* antigens Absent cutaneous reaction to *Candida* antigens

(c) ketabton.com: The Digital Library

Immunodeficiencies *(continued)*

DISEASE	DEFECT	PRESENTATION	FINDINGS
B- and T-cell disorders			
Severe combined immunodeficiency	Several types including defective IL-2R gamma chain (most common, X-linked recessive); adenosine deaminase deficiency (autosomal recessive); *RAG* mutation → VDJ recombination defect	Failure to thrive, chronic diarrhea, thrush Recurrent viral, bacterial, fungal, and protozoal infections	↓ T-cell receptor excision circles (TRECs) Absence of thymic shadow (CXR), germinal centers (lymph node biopsy), and T cells (flow cytometry)
Ataxia-telangiectasia 	Defects in *ATM* gene → failure to detect DNA damage → failure to halt progression of cell cycle → mutations accumulate; autosomal recessive	Triad: cerebellar defects (Ataxia), spider Angiomas (telangiectasia A), IgA deficiency ↑↑ sensitivity to radiation (limit x-ray exposure)	↑ AFP ↓ IgA, IgG, and IgE Lymphopenia, cerebellar atrophy ↑ risk of lymphoma and leukemia
Hyper-IgM syndrome	Most commonly due to defective CD40L on Th cells → class switching defect; X-linked recessive	Severe pyogenic infections early in life; opportunistic infection with *Pneumocystis*, *Cryptosporidium*, CMV	Normal or ↑ IgM ↓↓ IgG, IgA, IgE Failure to make germinal centers
Wiskott-Aldrich syndrome	Mutation in *WAS* gene; leukocytes and platelets unable to reorganize actin cytoskeleton → defective antigen presentation; X-linked recessive	**WATER**: **W**iskott-**A**ldrich: **T**hrombocytopenia, **E**czema, **R**ecurrent (pyogenic) infections ↑ risk of autoimmune disease and malignancy	↓ to normal IgG, IgM ↑ IgE, IgA Fewer and smaller platelets
Phagocyte dysfunction			
Leukocyte adhesion deficiency (type 1)	Defect in LFA-1 integrin (CD18) protein on phagocytes; impaired migration and chemotaxis; autosomal recessive	**L**ate separation (>30 days) of umbilical cord, **a**bsent pus, **d**ysfunctional neutrophils → recurrent skin and mucosal bacterial infections	↑ neutrophils in blood Absence of neutrophils at infection sites → impaired wound healing
Chédiak-Higashi syndrome 	Defect in lysosomal trafficking regulator gene (*LYST*) Microtubule dysfunction in phagosome-lysosome fusion; autosomal recessive	**PLAIN**: **P**rogressive neurodegeneration, **L**ymphohistiocytosis, **A**lbinism (partial), recurrent pyogenic **I**nfections, peripheral **N**europathy	Giant granules (B, arrows) in granulocytes and platelets. Pancytopenia Mild coagulation defects
Chronic granulomatous disease	Defect of NADPH oxidase → ↓ reactive oxygen species (eg, superoxide) and ↓ respiratory burst in neutrophils; X-linked form most common	↑ susceptibility to catalase ⊕ organisms	Abnormal dihydrorhodamine (flow cytometry) test (↓ green fluorescence) Nitroblue tetrazolium dye reduction test (obsolete) fails to turn blue

(c) ketabton.com: The Digital Library

Infections in immunodeficiency

PATHOGEN	↓ T CELLS	↓ B CELLS	↓ GRANULOCYTES	↓ COMPLEMENT
Bacteria	Sepsis	Encapsulated (**P**lease **SHINE** my **SK**i**S**): *P**seudomonas aeruginosa***, *S**treptococcus pneumoniae***, *H**aemophilus I**nfluenzae*** type b, *N**eisseria meningitidis***, *E**scherichia coli***, *S**almonella***, *K**lebsiella pneumoniae***, Group B *S**treptococcus***	**S**ome **B**acteria **P**roduce **N**o **S**erious granules: *S**taphylococcus***, *B**urkholderia cepacia***, *P**seudomonas aeruginosa***, *N**ocardia***, *S**erratia***	Encapsulated species with early complement deficiencies *Neisseria* with late complement (C5–C9) deficiencies
Viruses	CMV, EBV, JC virus, VZV, chronic infection with respiratory/GI viruses	Enteroviral encephalitis, poliovirus (live vaccine contraindicated)	N/A	N/A
Fungi/parasites	*Candida* (local), PCP, *Cryptococcus*	GI giardiasis (no IgA)	*Candida* (systemic), *Aspergillus*, *Mucor*	N/A

Note: **B**-cell deficiencies tend to produce recurrent **b**acterial infections, whereas T-cell deficiencies produce more fungal and viral infections.

(c) ketabton.com: The Digital Library

Transplant rejection

TYPE OF REJECTION	ONSET	PATHOGENESIS	FEATURES
Hyperacute 	Within minutes	Pre-existing recipient antibodies react to donor antigen (type II hypersensitivity reaction), activate complement	Widespread thrombosis of graft vessels (arrows within glomerulus A) → ischemia/necrosis Graft must be removed
Acute 	Weeks to months	Cellular: CD8+ T cells and/or CD4+ T cells activated against donor MHCs (type IV hypersensitivity reaction) Humoral: similar to hyperacute, except antibodies develop after transplant	Vasculitis of graft vessels with dense interstitial lymphocytic infiltrate B Prevent/reverse with immunosuppressants
Chronic 	Months to years	CD4+ T cells respond to recipient APCs presenting donor peptides, including allogeneic MHC Both cellular and humoral components (type II and IV hypersensitivity reactions)	Recipient T cells react and secrete cytokines → proliferation of vascular smooth muscle, parenchymal atrophy, interstitial fibrosis Dominated by arteriosclerosis C Organ-specific examples: ▪ Chronic allograft nephropathy ▪ Bronchiolitis obliterans ▪ Accelerated atherosclerosis (heart) ▪ Vanishing bile duct syndrome
Graft-versus-host disease	Varies	Grafted immunocompetent T cells proliferate in the immunocompromised host and reject host cells with "foreign" proteins → severe organ dysfunction Type IV hypersensitivity reaction	Maculopapular rash, jaundice, diarrhea, hepatosplenomegaly Usually in bone marrow and liver transplants (rich in lymphocytes) Potentially beneficial in bone marrow transplant for leukemia (graft-versus-tumor effect) For immunocompromised patients, irradiate blood products prior to transfusion to prevent GVHD

(c) ketabton.com: The Digital Library

▸ IMMUNOLOGY—IMMUNOSUPPRESSANTS

Immunosuppressants Agents that block lymphocyte activation and proliferation. Reduce acute transplant rejection by suppressing cellular immunity (used as prophylaxis). Frequently combined to achieve greater efficacy with ↓ toxicity. Chronic suppression ↑ risk of infection and malignancy.

DRUG	MECHANISM	INDICATIONS	TOXICITY	NOTES
Cyclosporine	Calcineurin inhibitor; binds cyclophilin Blocks T-cell activation by **preventing IL-2 transcription**	Psoriasis, rheumatoid arthritis	**Nephrotoxicity**, hypertension, hyperlipidemia, neurotoxicity, gingival hyperplasia, hirsutism	Both calcineurin inhibitors are highly nephrotoxic, especially in higher doses or in patients with decreased renal function
Tacrolimus (FK506)	Calcineurin inhibitor; binds FK506 binding protein (FKBP) Blocks T-cell activation by **preventing IL-2 transcription**		Similar to cyclosporine, ↑ risk of diabetes and neurotoxicity; no gingival hyperplasia or hirsutism	
Sirolimus (Rapamycin)	mTOR inhibitor; binds FKBP Blocks T-cell activation and B-cell differentiation by **preventing response to IL-2**	Kidney transplant rejection prophylaxis specifically Sir Basil's kidney transplant	"PanSirtopenia" (pancytopenia), insulin resistance, hyperlipidemia; **not nephrotoxic**	Kidney "sir-vives." Synergistic with cyclosporine Also used in drug-eluting stents
Basiliximab	Monoclonal antibody; blocks IL-2R		Edema, hypertension, tremor	
Azathioprine	Antimetabolite precursor of 6-mercaptopurine Inhibits lymphocyte proliferation by blocking nucleotide synthesis	Rheumatoid arthritis, Crohn disease, glomerulonephritis, other autoimmune conditions	Pancytopenia	6-MP degraded by xanthine oxidase; toxicity ↑ by allopurinol Pronounce "azathio-purine"
Mycophenolate Mofetil	Reversibly inhibits IMP dehydrogenase, preventing purine synthesis of B and T cells	Lupus nephritis	GI upset, pancytopenia, hypertension, hyperglycemia Less nephrotoxic and neurotoxic	Associated with invasive CMV infection
Glucocorticoids	Inhibit NF-κB Suppress both B- and T-cell function by ↓ transcription of many cytokines Induce T cell apoptosis	Many autoimmune and inflammatory disorders, adrenal insufficiency, asthma, CLL, non-Hodgkin lymphoma	Cushing syndrome, osteoporosis, hyperglycemia, diabetes, amenorrhea, adrenocortical atrophy, peptic ulcers, psychosis, cataracts, avascular necrosis (femoral head)	Demargination of WBCs causes artificial leukocytosis Adrenal insufficiency may develop if drug is stopped abruptly after chronic use

(c) ketabton.com: The Digital Library

Immunosuppression targets



Recombinant cytokines and clinical uses

CYTOKINE	AGENT	CLINICAL USES
Bone marrow stimulation		
Erythropoietin	Epoetin alfa (EPO analog)	Anemias (especially in renal failure)
Colony stimulating factors	Filgrastim (G-CSF), Sargramostim (GM-CSF)	Leukopenia; recovery of granulocyte and monocyte counts
Thrombopoietin	Romiplostim (TPO analog), eltrombopag (TPO receptor agonist)	Autoimmune thrombocytopenia Platelet stimulator
Immunotherapy		
Interleukin-2	Aldesleukin	Renal cell carcinoma, metastatic melanoma
Interferons	IFN-α	Chronic hepatitis C (not preferred) and B, renal cell carcinoma
	IFN-β	Multiple sclerosis
	IFN-γ	Chronic granulomatous disease

(c) ketabton.com: The Digital Library

Therapeutic antibodies

AGENT	TARGET	CLINICAL USE	NOTES
Cancer therapy			
Alemtuzumab	CD52	CLL, multiple sclerosis	"Alymtuzumab"—chronic lymphocytic leukemia
Bevacizumab	VEGF	Colorectal cancer, renal cell carcinoma, non-small cell lung cancer	Also used for neovascular age-related macular degeneration, proliferative diabetic retinopathy, and macular edema
Rituximab	CD20	B-cell non-Hodgkin lymphoma, CLL, rheumatoid arthritis, ITP, multiple sclerosis	Risk of PML in patients with JC virus CD20—"ri2ximab"
Trastuzumab	HER2	Breast cancer, gastric cancer	HER2—"tras2zumab"
Autoimmune disease therapy			
Adalimumab, infliximab	Soluble TNF-α	IBD, rheumatoid arthritis, ankylosing spondylitis, psoriasis	Etanercept is a decoy TNF-α receptor and not a monoclonal antibody
Eculizumab	Complement protein C5	Paroxysmal nocturnal hemoglobinuria	
Ixekizumab, secukinumab	IL-17A	Psoriasis, psoriatic arthritis	
Natalizumab	α4-integrin	Multiple sclerosis, Crohn disease	α4-integrin: WBC adhesion Risk of PML in patients with JC virus
Ustekinumab	IL-12/IL-23	Psoriasis, psoriatic arthritis	
Other applications			
Abciximab	Platelet glycoproteins IIb/IIIa	Antiplatelet agent for prevention of ischemic complications in patients undergoing percutaneous coronary intervention	ABC is as easy as 123
Denosumab	RANKL	Osteoporosis; inhibits osteoclast maturation (mimics osteoprotegerin)	Denosumab helps make dense bones
Omalizumab	IgE	Refractory allergic asthma; prevents IgE binding to FcεRI	
Palivizumab	RSV F protein	RSV prophylaxis for high-risk infants	PaliVIzumab—VIrus

(c) ketabton.com: The Digital Library

HIGH-YIELD PRINCIPLES IN

Microbiology

"Support bacteria. They're the only culture some people have."

—Steven Wright

"What lies behind us and what lies ahead of us are tiny matters compared to what lies within us."

—Henry S. Haskins

"Infectious disease is merely a disagreeable instance of a widely prevalent tendency of all living creatures to save themselves the bother of building, by their own efforts, the things they require."

—Hans Zinsser

Microbiology questions on the Step 1 exam often require two (or more) steps: Given a certain clinical presentation, you will first need to identify the most likely causative organism, and you will then need to provide an answer regarding some features of that organism or relevant antimicrobial agents. For example, a description of a child with fever and a petechial rash will be followed by a question that reads, "From what site does the responsible organism usually enter the blood?"

This section therefore presents organisms in two major ways: in individual microbial "profiles" and in the context of the systems they infect and the clinical presentations they produce. You should become familiar with both formats. When reviewing the systems approach, remind yourself of the features of each microbe by returning to the individual profiles. Also be sure to memorize the laboratory characteristics that allow you to identify microbes.

Basic Bacteriology 124
Clinical Bacteriology 134
Mycology 151
Parasitology 155
Virology 162
Systems 178
Antimicrobials 187

FAS1_2019_03-Microbiology.indd 123 11/14/19 12:19 PM

(c) ketabton.com: The Digital Library

▸ MICROBIOLOGY—BASIC BACTERIOLOGY

Bacterial structures

STRUCTURE	CHEMICAL COMPOSITION	FUNCTION
Appendages		
Flagellum	Proteins	Motility
Pilus/fimbria	Glycoprotein	Mediate adherence of bacteria to cell surface; sex pilus forms during conjugation
Specialized structures		
Spore	Keratin-like coat; dipicolinic acid; peptidoglycan, DNA	Gram ⊕ only Survival: resist dehydration, heat, chemicals
Cell envelope		
Capsule	Discrete layer usually made of polysaccharides (and rarely proteins)	Protects against phagocytosis
Slime (S) layer	Loose network of polysaccharides	Mediates adherence to surfaces, especially foreign surfaces (eg, indwelling catheters)
Outer membrane	Outer leaflet: contains endotoxin (LPS/LOS) Embedded proteins: porins and other outer membrane proteins (OMPs) Inner leaflet: phospholipids	Gram ⊖ only Endotoxin: lipid A induces TNF and IL-1; antigenic O polysaccharide component Most OMPs are antigenic Porins: transport across outer membrane
Periplasm	Space between cytoplasmic membrane and outer membrane in gram ⊖ bacterial (peptidoglycan in middle)	Accumulates components exiting gram ⊖ cells, including hydrolytic enzymes (eg, β-lactamases)
Cell wall	Peptidoglycan is a sugar backbone with peptide side chains cross-linked by transpeptidase	Net-like structure gives rigid support, protects against osmotic pressure damage
Cytoplasmic membrane	Phospholipid bilayer sac with embedded proteins (eg, penicillin-binding proteins [PBPs]) and other enzymes Lipoteichoic acids (gram positive) only extend from membrane to exterior	Site of oxidative and transport enzymes; PBPs involved in cell wall synthesis Lipoteichoic acids induce TNF-α and IL-1

Cell envelope



(c) ketabton.com: The Digital Library

Stains

Gram stain	First-line lab test in bacterial identification. Bacteria with thick peptidoglycan layer retain crystal violet dye (gram ⊕); bacteria with thin peptidoglycan layer turn red or pink (gram ⊖) with counterstain. These bugs do not Gram stain well (**T**hese **L**ittle **M**icrobes **M**ay **U**nfortunately **L**ack **R**eal **C**olor **B**ut **A**re **E**verywhere):	
	Treponema, ***L**eptospira*	Too thin to be visualized
	Mycobacteria	Cell wall has high lipid content
	***M**ycoplasma*, ***U**reaplasma*	No cell wall
	***L**egionella*, ***R**ickettsia*, ***C**hlamydia*, ***B**artonella*, ***A**naplasma*, ***E**hrlichia*	Primarily intracellular; also, *Chlamydia* lack classic peptidoglycan because of ↓ muramic acid
Giemsa stain	***Rick**ettsia*, ***Chlamydia***, **Try**panosomes A, ***P**lasmodium*, ***B**orrelia*, ***H**elicobacter pylori*	**Rick**y got ***Chlamydia*** as he **Tri**ed to **Pl**ease the **B**ored **H**ot "**Geisha**"
Periodic acid–Schiff stain	Stains **glycogen**, mucopolysaccharides; used to diagnose Whipple disease (*Tropheryma whipplei* B)	**PaS**s the **sugar**
Ziehl-Neelsen stain (carbol fuchsin)	Acid-fast bacteria (eg, *Mycobacteria* C, *Nocardia*; stains mycolic acid in cell wall); protozoa (eg, *Cryptosporidium* oocysts)	Auramine-rhodamine stain is more often used for screening (inexpensive, more sensitive)
India ink stain	*Cryptococcus neoformans* D; mucicarmine can also be used to stain thick polysaccharide capsule red	
Silver stain	Fungi (eg, *Coccidioides* E, *Pneumocystis jirovecii*), *Legionella*, *Helicobacter pylori*	
Fluorescent antibody stain	Used to identify many bacteria, viruses, *Pneumocystis jirovecii*, *Giardia*, and *Cryptosporidium*	Example is FTA-ABS for syphilis



(c) ketabton.com: The Digital Library

Properties of growth media	The same type of media can possess both (or neither) of these properties.
Selective media	Favors the growth of particular organism while preventing growth of other organisms. Example: Thayer-Martin agar contains antibiotics that allow the selective growth of *Neisseria* by inhibiting the growth of other sensitive organisms.
Indicator (differential) media	Yields a color change in response to the metabolism of certain organisms. Example: MacConkey agar contains a pH indicator; a lactose fermenter like *E coli* will convert lactose to acidic metabolites → color change to pink.

Special culture requirements

BUG	MEDIA USED FOR ISOLATION	MEDIA CONTENTS/OTHER
H influenzae	Chocolate agar	Factors V (NAD^+) and X (hematin)
N gonorrhoeae, N meningitidis	Thayer-Martin agar	Selectively favors growth of *Neisseria* by inhibiting growth of gram ⊕ organisms with **V**ancomycin, gram ⊖ organisms except *Neisseria* with **T**rimethoprim and **C**olistin, and fungi with **N**ystatin **V**ery **T**ypically **C**ultures ***N**eisseria*
B pertussis	Bordet-Gengou agar (**Bordet** for ***Bordet**ella*) Regan-Lowe medium	Potato extract Charcoal, blood, and antibiotic
C diphtheriae	Tellurite agar, Löffler medium	
M tuberculosis	Löwenstein-Jensen medium, Middlebrook medium, rapid automated broth cultures	
M pneumoniae	Eaton agar	Requires cholesterol
Lactose-fermenting enterics	MacConkey agar	Fermentation produces acid, causing colonies to turn pink
E coli	Eosin–methylene blue (EMB) agar	Colonies with green metallic sheen
Brucella, Francisella, Legionella, Pasteurella	**Charcoal** yeast extract agar buffered with **cysteine** and **iron**	The **Ella** siblings, **Bruce**, **Francis**, a **legion**naire, and a **pasteur** (pastor), built the Sistine (**cysteine**) chapel out of **charcoal** and **iron**.
Fungi	**Sab**ouraud agar	"**Sab**'s a **fun guy**!"

Aerobes	Use an O_2-dependent system to generate ATP. Examples include *Nocardia*, *Pseudomonas aeruginosa*, *Mycobacterium tuberculosis*, and *Bordetella pertussis*. Reactivation of *M tuberculosis* (eg, after immunocompromise or TNF-α inhibitor use) has a predilection for the apices of the lung.

(c) ketabton.com: The Digital Library

Anaerobes	Examples include *Clostridium*, *Bacteroides*, *Fusobacterium*, and *Actinomyces israelii*. They lack catalase and/or superoxide dismutase and are thus susceptible to oxidative damage. Generally foul smelling (short-chain fatty acids), are difficult to culture, and produce gas in tissue (CO_2 and H_2).	Anaerobes Can't Breathe Fresh Air. Anaerobes are normal flora in GI tract, typically pathogenic elsewhere. AminO_2glycosides are ineffective against anaerobes because these antibiotics require O_2 to enter into bacterial cell.
Facultative anaerobes	May use O_2 as a terminal electron acceptor to generate ATP, but can also use fermentation and other O_2-independent pathways.	Streptococci, staphylococci, and enteric gram ⊖ bacteria.

Intracellular bacteria

Obligate intracellular	*Rickettsia*, *Chlamydia*, *Coxiella* Rely on host ATP	Stay inside (cells) when it is Really Chilly and Cold
Facultative intracellular	*Salmonella*, *Neisseria*, *Brucella*, *Mycobacterium*, *Listeria*, *Francisella*, *Legionella*, *Yersinia pestis*	Some Nasty Bugs May Live FacultativeLY

Encapsulated bacteria 	Examples are *Pseudomonas aeruginosa*, *Streptococcus pneumoniae* A, *Haemophilus influenzae* type b, *Neisseria meningitidis*, *Escherichia coli*, *Salmonella*, *Klebsiella pneumoniae*, and group B Strep. Their capsules serve as an antiphagocytic virulence factor. Capsular polysaccharide + protein conjugate serves as an antigen in vaccines.	Please SHiNE my SKiS. Are opsonized, and then cleared by spleen. Asplenics (No Spleen Here) have ↓ opsonizing ability and thus ↑ risk for severe infections; need vaccines to protect against: ▪ *N meningitidis* ▪ *S pneumoniae* ▪ *H influenzae*
Encapsulated bacteria vaccines	Some vaccines containing polysaccharide capsule antigens are conjugated to a carrier protein, enhancing immunogenicity by promoting T-cell activation and subsequent class switching. A polysaccharide antigen alone cannot be presented to T cells.	Pneumococcal vaccines: PCV13 (pneumococcal conjugate vaccine), PPSV23 (pneumococcal polysaccharide vaccine with no conjugated protein). *H influenzae* type b (conjugate vaccine). Meningococcal vaccine (conjugate vaccine).
Urease-positive organisms	*Proteus*, *Cryptococcus*, *H pylori*, *Ureaplasma*, *Nocardia*, *Klebsiella*, *S epidermidis*, *S saprophyticus*. Urease hydrolyzes urea to release ammonia and CO_2 → ↑ pH. Predisposes to struvite (ammonium magnesium phosphate) stones, particularly *Proteus*.	Pee CHUNKSS.

(c) ketabton.com: The Digital Library

Catalase-positive organisms



Catalase degrades H_2O_2 into H_2O and bubbles of O_2 **A** before it can be converted to microbicidal products by the enzyme myeloperoxidase. People with chronic granulomatous disease (NADPH oxidase deficiency) have recurrent infections with certain catalase ⊕ organisms.

Examples: *Nocardia*, *Staphylococci*, *Serratia*, *Candida*, *Listeria*, *E coli*, *Burkholderia cepacia*, *Pseudomonas*, *Aspergillus*, *Helicobacter pylori*, *Bordetella pertussis*.

Pigment-producing bacteria

Actinomyces israelii—yellow "sulfur" granules, which are composed of filaments of bacteria	Israel has yellow sand
S aureus—yellow pigment	*Aureus* (Latin) = gold
P aeruginosa—blue-green pigment (pyocyanin and pyoverdin)	Aerugula is green
Serratia marcescens—red pigment	Think red Sriracha hot sauce

In vivo biofilm-producing bacteria

S epidermidis	Catheter and prosthetic device infections
Viridans streptococci *(S mutans, S sanguinis)*	Dental plaques, infective endocarditis
P aeruginosa	Respiratory tree colonization in patients with cystic fibrosis, ventilator-associated pneumonia Contact lens–associated keratitis
Nontypeable (unencapsulated) *H influenzae*	Otitis media

(c) ketabton.com: The Digital Library

Spore-forming bacteria



Some gram ⊕ bacteria can form spores A when nutrients are limited. Spores lack metabolic activity and are highly resistant to heat and chemicals. Core contains dipicolinic acid. Must autoclave to kill spores (as is done to surgical equipment) by steaming at 121°C for 15 minutes.

Examples: *B anthracis* (anthrax), *B cereus* (food poisoning), *C botulinum* (botulism), *C difficile* (pseudomembranous colitis), *C perfringens* (gas gangrene), *C tetani* (tetanus).

Bacterial virulence factors	These promote evasion of host immune response.
Protein A	Binds Fc region of IgG. Prevents opsonization and phagocytosis. Expressed by S *a*ureus.
IgA protease	Enzyme that cleaves IgA, allowing bacteria to adhere to and colonize mucous membranes. Secreted by S *p*neumoniae, H *i*nfluenzae type b, and N*e*isseria (SHiN).
M protein	Helps prevent phagocytosis. Expressed by group A streptococci. Shares similar epitopes to human cellular proteins (molecular mimicry); possibly underlies the autoimmune response seen in acute rheumatic fever.

(c) ketabton.com: The Digital Library

Bacterial genetics

Transformation

Competent bacteria can bind and import short pieces of environmental naked bacterial chromosomal DNA (from bacterial cell lysis). The transfer and expression of newly transferred genes is called transformation. A feature of many bacteria, especially ***S** pneumoniae*, ***H** influenzae* type b, and **N***eisseria* (**SHiN**).

Adding deoxyribonuclease degrades naked DNA, preventing transformation.



Conjugation

$F^+ \times F^-$

F^+ plasmid contains genes required for sex pilus and conjugation. Bacteria without this plasmid are termed F^-. Sex pilus on F^+ bacterium contacts F^- bacterium. A single strand of plasmid DNA is transferred across the conjugal bridge ("mating bridge"). No transfer of chromosomal DNA.



Hfr × F^-

F^+ plasmid can become incorporated into bacterial chromosomal DNA, termed high-frequency recombination (Hfr) cell. Transfer of leading part of plasmid and a few flanking chromosomal genes. High-frequency recombination may integrate some of those bacterial genes. Recipient cell remains F^- but now may have new bacterial genes.



Transduction

Generalized

A packaging "error." Lytic phage infects bacterium, leading to cleavage of bacterial DNA. Parts of bacterial chromosomal DNA may become packaged in phage capsid. Phage infects another bacterium, transferring these genes.



Specialized

An "excision" event. Lysogenic phage infects bacterium; viral DNA incorporates into bacterial chromosome. When phage DNA is excised, flanking bacterial genes may be excised with it. DNA is packaged into phage capsid and can infect another bacterium.

Genes for the following 5 bacterial toxins are encoded in a lysogenic phage (**ABCD'S**): Group **A** strep erythrogenic toxin, **B**otulinum toxin, **C**holera toxin, **D**iphtheria toxin, **S**higa toxin.



(c) ketabton.com: The Digital Library

Bacterial genetics *(continued)*

Transposition

A "jumping" process involving a transposon (specialized segment of DNA), which can copy and excise itself and then insert into the same DNA molecule or an unrelated DNA (eg, plasmid or chromosome). Critical in creating plasmids with multiple drug resistance and transfer across species lines (eg, Tn*1546* with *van*A from *Enterococcus* to *S aureus*).



Main features of exotoxins and endotoxins

	Exotoxins	Endotoxins
SOURCE	Certain species of gram ⊕ and gram ⊖ bacteria	Outer cell membrane of most gram ⊖ bacteria
SECRETED FROM CELL	Yes	No
CHEMISTRY	Polypeptide	Lipid A component of LPS (structural part of bacteria; released when lysed)
LOCATION OF GENES	Plasmid or bacteriophage	Bacterial chromosome
ADVERSE EFFECTS	High (fatal dose on the order of 1 µg)	Low (fatal dose on the order of hundreds of micrograms)
CLINICAL EFFECTS	Various effects (see following pages)	Fever, shock (hypotension), DIC
MODE OF ACTION	Various modes (see following pages)	Induces TNF, IL-1, and IL-6
ANTIGENICITY	Induces high-titer antibodies called antitoxins	Poorly antigenic
VACCINES	Toxoids used as vaccines	No toxoids formed and no vaccine available
HEAT STABILITY	Destroyed rapidly at 60°C (except staphylococcal enterotoxin and *E coli* heat-stable toxin)	Stable at 100°C for 1 hr
TYPICAL DISEASES	Tetanus, botulism, diphtheria, cholera	Meningococcemia; sepsis by gram ⊖ rods



FAS1_2019_03-Microbiology.indd 131 11/14/19 12:20 PM

(c) ketabton.com: The Digital Library

Bacteria with exotoxins

BACTERIA	TOXIN	MECHANISM	MANIFESTATION
Inhibit protein synthesis			
Corynebacterium diphtheriae	Diphtheria toxin[a]	Inactivate elongation factor (EF-2)	Pharyngitis with pseudomembranes in throat and severe lymphadenopathy (bull neck), myocarditis
Pseudomonas aeruginosa	Exotoxin A[a]		Host cell death
***Shigella* spp**	Shiga toxin (ST)[a]	Inactivate 60S ribosome by removing adenine from rRNA	GI mucosal damage → dysentery; ST also enhances cytokine release, causing hemolytic-uremic syndrome (HUS)
Enterohemorrhagic *E coli*	Shiga-like toxin (SLT)[a]		SLT enhances cytokine release, causing HUS (prototypically in EHEC serotype O157:H7) Unlike *Shigella*, EHEC does not invade host cells
Increase fluid secretion			
Enterotoxigenic *E coli*	Heat-**labile** toxin (LT)[a]	Overactivates adenylate cyclase (↑ c**A**MP) → ↑ Cl^- secretion in gut and H_2O efflux	Watery diarrhea: "**labile** in the **A**ir (**A**denylate cyclase), **stable** on the **G**round (**G**uanylate cyclase)"
	Heat-**stable** toxin (ST)	Overactivates guanylate cyclase (↑ c**G**MP) → ↓ resorption of NaCl and H_2O in gut	
Bacillus anthracis	Anthrax toxin[a]	Mimics adenylate cyclase (↑ cAMP)	Likely responsible for characteristic edematous borders of black eschar in cutaneous anthrax
Vibrio cholerae	Cholera toxin[a]	Overactivates adenylate cyclase (↑ cAMP) by permanently activating G_s → ↑ Cl^- secretion in gut and H_2O efflux	Voluminous "rice-water" diarrhea
Inhibit phagocytic ability			
Bordetella pertussis	Pertussis toxin[a]	Inactivates inhibitory G subunit (G_i) → activation of adenylate cyclase → ↑ cAMP	Whooping cough—child coughs on expiration and "whoops" on inspiration (toxin may not actually be a cause of cough; can cause "100-day cough" in adults)
Inhibit release of neurotransmitter			
Clostridium tetani	Tetanospasmin[a]	Both are proteases that cleave SNARE (soluble NSF attachment protein receptor), a set of proteins required for neurotransmitter release via vesicular fusion	Toxin prevents release of **inhibitory** (GABA and glycine) neurotransmitters from Renshaw cells in spinal cord → spastic paralysis, risus sardonicus, trismus (lockjaw)
Clostridium botulinum	Botulinum toxin[a]		Toxin prevents release of **stimulatory** (ACh) signals at neuromuscular junction → flaccid paralysis (floppy baby)

[a]An AB toxin (aka, two-component toxin [or three for anthrax]) with **B** enabling **b**inding and triggering uptake (endocytosis) of the **a**ctive **A** component. The A components are usually ADP ribosyltransferases; others have enzymatic activities as listed in chart.

(c) ketabton.com: The Digital Library

Bacteria with exotoxins *(continued)*

BACTERIA	TOXIN	MECHANISM	MANIFESTATION
Lyse cell membranes			
Clostridium perfringens	Alpha toxin	Phospholipase (lecithinase) that degrades tissue and cell membranes	Degradation of phospholipids → myonecrosis ("gas gangrene") and hemolysis ("double zone" of hemolysis on blood agar)
Streptococcus pyogenes	Streptolysin O	Protein that degrades cell membrane	Lyses RBCs; contributes to β-hemolysis; host antibodies against toxin (ASO) used to diagnose rheumatic fever (do not confuse with immune complexes of poststreptococcal glomerulonephritis)
Superantigens causing shock			
Staphylococcus aureus	Toxic shock syndrome toxin (TSST-1)	Cross-links β region of TCR to MHC class II on APCs outside of the antigen binding site → overwhelming release of IL-1, IL-2, IFN-γ, and TNF-α → shock	Toxic shock syndrome: fever, rash, shock; other toxins cause scalded skin syndrome (exfoliative toxin) and food poisoning (heat-stable enterotoxin)
Streptococcus pyogenes	Erythrogenic exotoxin A		Toxic shock–like syndrome: fever, rash, shock; scarlet fever

Endotoxin

LPS found in outer membrane of gram ⊖ bacteria (both cocci and rods). Composed of O antigen + core polysaccharide + lipid A (the toxic component).

Released upon cell lysis or by living cells by blebs detaching from outer surface membrane (vs exotoxin, which is actively secreted).

Three main effects: macrophage activation (TLR4/CD14), complement activation, and tissue factor activation.

ENDOTOXINS:
Edema
Nitric oxide
DIC/**D**eath
Outer membrane
TNF-α
O-antigen + core polysaccharide + lipid A
e**X**tremely heat stable
IL-1 and **I**L-6
Neutrophil chemotaxis
Shock



(c) ketabton.com: The Digital Library

▸ MICROBIOLOGY—CLINICAL BACTERIOLOGY

Gram-positive lab algorithm



Important **tests** are in **bold.** Important ***pathogens*** are in ***bold italics.***
Note: Enterococcus is either α- or γ-hemolytic.

Gram-positive cocci antibiotic tests

Staphylococci	Novobiocin—*Saprophyticus* is Resistant; *Epidermidis* is Sensitive	On the office's "staph" retreat, there was no stress
Streptococci	Optochin—*Viridans* is Resistant; *Pneumoniae* is Sensitive	OVRPS (overpass)
	Bacitracin—group B strep are Resistant; group A strep are Sensitive	B-BRAS

(c) ketabton.com: The Digital Library

α-hemolytic bacteria



Gram ⊕ cocci. Partial oxidation of hemoglobin causes greenish or brownish color without clearing around growth on blood agar A. Include the following organisms:

- *Streptococcus pneumoniae* (catalase ⊖ and optochin sensitive)
- Viridans streptococci (catalase ⊖ and optochin resistant)

β-hemolytic bacteria



Gram ⊕ cocci. Complete lysis of RBCs → pale/clear area surrounding colony on blood agar A. Include the following organisms:

- *Staphylococcus aureus* (catalase and coagulase ⊕)
- *Streptococcus pyogenes*—group A strep (catalase ⊖ and bacitracin sensitive)
- *Streptococcus agalactiae*—group *B* strep (catalase ⊖ and bacitracin resistant)

Staphylococcus aureus



Gram ⊕, β-hemolytic, catalase ⊕, coagulase ⊕ cocci in clusters A. Protein A (virulence factor) binds Fc-IgG, inhibiting complement activation and phagocytosis. Commonly colonizes the nares, ears, axilla, and groin.

Causes:

- Inflammatory disease—skin infections, organ abscesses, pneumonia (often after influenza virus infection), endocarditis, septic arthritis, and osteomyelitis.
- Toxin-mediated disease—toxic shock syndrome (TSST-1), scalded skin syndrome (exfoliative toxin), rapid-onset food poisoning (enterotoxins).

MRSA (methicillin-resistant *S aureus*)—important cause of serious nosocomial and community-acquired infections; resistance due to altered penicillin-binding protein. *mec*A gene from staphylococcal chromosomal cassette involved in penicillin resistance.

TSST-1 is a superantigen that binds to MHC II and T-cell receptor, resulting in polyclonal T-cell activation and cytokine release.

Staphylococcal toxic shock syndrome (TSS)—fever, vomiting, rash, desquamation, shock, end-organ failure. TSS results in ↑ AST, ↑ ALT, ↑ bilirubin. Associated with prolonged use of vaginal tampons or nasal packing.

Compare with *Streptococcus pyogenes* TSS (a toxic shock–like syndrome associated with painful skin infection).

S aureus food poisoning due to ingestion of preformed toxin → short incubation period (2–6 hr) followed by nonbloody diarrhea and emesis. Enterotoxin is heat stable → not destroyed by cooking.

S aureus makes coagulase and toxins. Forms fibrin clot around itself → abscess.

Staphylococcus epidermidis

Gram ⊕, catalase ⊕, coagulase ⊖, urease ⊕ cocci in clusters. Novobiocin sensitive. Does not ferment mannitol (vs *S aureus*).

Normal flora of skin; contaminates blood cultures.

Infects prosthetic devices (eg, hip implant, heart valve) and IV catheters by producing adherent biofilms.

(c) ketabton.com: The Digital Library

Staphylococcus saprophyticus

Gram ⊕, catalase ⊕, coagulase ⊖, urease ⊕ cocci in clusters. Novobiocin resistant.
Normal flora of female genital tract and perineum.
Second most common cause of uncomplicated UTI in young women (most common is *E coli*).

Streptococcus pneumoniae

A



Gram ⊕, α-hemolytic, lancet-shaped diplococci A.
Encapsulated. IgA protease. Optochin sensitive and bile soluble. Most commonly causes:
- Meningitis
- Otitis media (in children)
- Pneumonia
- Sinusitis

Pneumococcus is associated with "rusty" sputum, sepsis in patients with sickle cell disease, and asplenic patients.
No virulence without capsule.

Viridans group streptococci

Gram ⊕, α-hemolytic cocci. Optochin resistant and bile insoluble. Normal flora of the oropharynx.
Streptococcus mutans and *S mitis* cause dental caries.
S sanguinis makes dextrans that bind to fibrin-platelet aggregates on damaged **heart** valves, causing subacute bacterial endocarditis.

Viridans group strep live in the mouth, because they are not afraid **of-the-chin** (**op-to-chin** resistant).
Sanguinis = **blood**. Think, "there is lots of **blood** in the **heart**" (endocarditis).

Streptococcus pyogenes (group A streptococci)

A



Gram ⊕ cocci in chains A. Group A strep cause:
- Pyogenic—pharyngitis, cellulitis, impetigo ("honey-crusted" lesions), erysipelas
- Toxigenic—scarlet fever, toxic shock–like syndrome, necrotizing fasciitis
- Immunologic—rheumatic fever, glomerulonephritis

Bacitracin sensitive, β-hemolytic, pyrrolidonyl arylamidase (PYR) ⊕. Hyaluronic acid capsule and M protein inhibit phagocytosis. Antibodies to M protein enhance host defenses against *S pyogenes* but can give rise to rheumatic fever.
ASO titer or anti-DNase B antibodies indicate recent *S pyogenes* infection.

"**Ph**"yogenes **ph**aryngitis can result in rheumatic "**ph**ever" and glomerulone**ph**ritis.
Strains causing impetigo can induce glomerulonephritis.
Scarlet fever—blanching, sandpaper-like body rash, strawberry tongue, and circumoral pallor in the setting of group A streptococcal pharyngitis (erythrogenic toxin ⊕).

(c) ketabton.com: The Digital Library

***Streptococcus agalactiae* (group B streptococci)**	Gram ⊕ cocci, bacitracin resistant, β-hemolytic, colonizes vagina; causes pneumonia, meningitis, and sepsis, mainly in babies. Produces CAMP factor, which enlarges the area of hemolysis formed by *S aureus*. (Note: CAMP stands for the authors of the test, not cyclic AMP.) Hippurate test ⊕. PYR ⊖. Screen pregnant women at 35–37 weeks of gestation with rectal and vaginal swabs. Patients with ⊕ culture receive intrapartum penicillin/ampicillin prophylaxis.	Group B for Babies!
Streptococcus bovis	Gram ⊕ cocci, colonizes the gut. *S gallolyticus* (*S bovis* biotype 1) can cause bacteremia and subacute endocarditis and is associated with colon cancer.	Bovis in the blood = cancer in the colon.
Enterococci	Gram ⊕ cocci. Enterococci (*E faecalis* and *E faecium*) are normal colonic flora that are penicillin G resistant and cause UTI, biliary tract infections, and subacute endocarditis (following GI/GU procedures). Catalase ⊖, PYR ⊕, variable hemolysis. VRE (vancomycin-resistant enterococci) are an important cause of nosocomial infection.	Enterococci are more resilient than streptococci, can grow in 6.5% NaCl and bile (lab test). *Entero* = intestine, *faecalis* = feces, *strepto* = twisted (chains), *coccus* = berry.
Bacillus anthracis	Gram ⊕, spore-forming rod that produces anthrax toxin (an exotoxin consisting of protective antigen, lethal factor, and edema factor). Has a polypeptide capsule (poly D-glutamate). Colonies show a halo of projections, sometimes referred to as "medusa head" appearance.	
Cutaneous anthrax 	Painless papule surrounded by vesicles → ulcer with black eschar A (painless, necrotic) → uncommonly progresses to bacteremia and death.	
Pulmonary anthrax	Inhalation of spores, most commonly from contaminated animals or animal products, although also a potential bioweapon → flu-like symptoms that rapidly progress to fever, pulmonary hemorrhage, mediastinitis (CXR may show widened mediastinum), and shock. Also called woolsorter's disease.	

(c) ketabton.com: The Digital Library

Bacillus cereus	Gram ⊕ rod. Causes food poisoning. Spores survive cooking rice (reheated rice syndrome). Keeping rice warm results in germination of spores and enterotoxin formation. Emetic type causes nausea and vomiting within 1–5 hours. Caused by cereulide, a preformed toxin. Diarrheal type causes watery, nonbloody diarrhea and GI pain within 8–18 hours. Management: supportive care (antibiotics are ineffective against toxins).

Clostridia

Gram ⊕, spore-forming, obligate anaerobic rods. Tetanus toxin and botulinum toxin are proteases that cleave SNARE proteins involved in neurotransmission.

Clostridium tetani	Produces tetanospasmin, an exotoxin causing tetanus. Tetanospasmin blocks release of GABA and glycine from Renshaw cells in spinal cord. Causes spastic paralysis, trismus (lockjaw), risus sardonicus (raised eyebrows and open grin), opisthotonos (spasms of spinal extensors). Prevent with tetanus vaccine. Treat with antitoxin +/– vaccine booster, antibiotics, diazepam (for muscle spasms), and wound debridement.	Tetanus is tetanic paralysis.
Clostridium botulinum	Produces a heat-labile toxin that inhibits ACh release at the neuromuscular junction, causing botulism. In adults, disease is caused by ingestion of preformed toxin. In babies, ingestion of spores (eg, in honey) leads to disease (floppy baby syndrome). Treat with human botulinum immunoglobulin.	Symptoms of botulism (the 4 D's): Diplopia, Dysarthria, Dysphagia, Dyspnea. *Botulinum* is from bad bottles of food, juice, and honey (causes a descending flaccid paralysis). Local botulinum toxin A (Botox) injections used to treat focal dystonia, hyperhidrosis, muscle spasms, and cosmetic reduction of facial wrinkles.
Clostridium perfringens A	Produces α-toxin (lecithinase, a phospholipase) that can cause myonecrosis (gas gangrene A; presents as soft tissue crepitus) and hemolysis. If heavily spore-contaminated food is cooked but left standing too long at < 60°C, spores germinate → vegetative bacteria → heat-labile enterotoxin → food poisoning symptoms in 10-12 hours, resolution in 24 hours.	*Perfringens* perforates a gangrenous leg.
Clostridioides difficile  B	Produces toxins A and B, which damage enterocytes. Both toxins lead to watery diarrhea → pseudomembranous colitis B. Often 2° to antibiotic use, especially clindamycin or ampicillin; associated with PPIs. Diagnosed by PCR or antigen detection of one or both toxins in stool. Complications: toxic megacolon.	*Difficile* causes diarrhea. Treatment: oral vancomycin, metronidazole, or fidaxomicin. For recurrent cases, consider repeating prior regimen or fecal microbiota transplant.

(c) ketabton.com: The Digital Library

Corynebacterium diphtheriae



Gram ⊕ rods occurring in angular arrangements; transmitted via respiratory droplets. Causes diphtheria via exotoxin encoded by β-prophage. Potent exotoxin inhibits protein synthesis via ADP-ribosylation of EF-2, leading to possible necrosis in pharynx, cardiac, and CNS tissue.

Symptoms include pseudomembranous pharyngitis (grayish-white membrane A) with lymphadenopathy. Toxin dissemination may cause myocarditis, arrhythmias, neuropathies.

Lab diagnosis based on gram ⊕ rods with metachromatic (blue and red) granules and ⊕ Elek test for toxin.

Toxoid vaccine prevents diphtheria.

Coryne = club shaped (metachromatic granules on Löffler media).

Black colonies on cystine-tellurite agar.

ABCDEFG:
- **A**DP-ribosylation
- β-prophage
- ***C**orynebacterium*
- ***D**iphtheriae*
- **E**longation **F**actor 2
- **G**ranules

Treatment: antibiotic therapy +/– diphtheria antitoxin.

Listeria monocytogenes



Gram ⊕, facultative intracellular rod; acquired by ingestion of unpasteurized dairy products and cold deli meats, transplacental transmission, by vaginal transmission during birth. Grows well at refrigeration temperatures ("cold enrichment").

Forms "rocket tails" (red in A) via actin polymerization that allow intracellular movement and cell-to-cell spread across cell membranes, thereby avoiding antibody. Characteristic tumbling motility in broth.

Can cause amnionitis, septicemia, and spontaneous abortion in pregnant women; granulomatosis infantiseptica; meningitis in immunocompromised patients, neonates, and older adults; mild, self-limited gastroenteritis in healthy individuals.

Treatment: ampicillin.

Nocardia vs *Actinomyces*





Both are gram ⊕ and form long, branching filaments resembling fungi.

Nocardia	*Actinomyces*
Aerobe	Anaerobe
Acid fast (weak) A	Not acid fast B
Found in soil	Normal oral, reproductive, and GI flora
Causes pulmonary infections in immunocompromised (can mimic TB but with ⊖ PPD); cutaneous infections after trauma in immunocompetent; can spread to CNS	Causes oral/facial abscesses that drain through sinus tracts; often associated with dental caries/extraction and other maxillofacial trauma; forms yellow "sulfur granules"; can also cause PID with IUDs
Treat with sulfonamides (TMP-SMX)	Treat with penicillin

Treatment is a **SNAP**: **S**ulfonamides—***N**ocardia*; ***A**ctinomyces*—**P**enicillin

---

(c) ketabton.com: The Digital Library

Mycobacteria



Gram ⊕ acid fast rods (pink rods, arrows in A).

Mycobacterium tuberculosis (TB, often resistant to multiple drugs).

M avium–intracellulare (causes disseminated, non-TB disease in AIDS; often resistant to multiple drugs). Prophylaxis with azithromycin when CD4+ count < 50 cells/mm^3.

M scrofulaceum (cervical lymphadenitis in children).

M marinum (hand infection in aquarium handlers).

TB symptoms include fever, night sweats, weight loss, cough (nonproductive or productive), hemoptysis.

Cord factor creates a "serpentine cord" appearance in virulent *M tuberculosis* strains; activates macrophages (promoting granuloma formation) and induces release of TNF-α. Sulfatides (surface glycolipids) inhibit phagolysosomal fusion.

Tuberculosis



Interferon-γ release assay (IGRA) has fewer false positives from BCG vaccination.

PPD ⊕ if current infection or past exposure.

PPD ⊖ if no infection and in sarcoidosis or HIV infection (especially with low CD4+ cell count).

Caseating granulomas with central necrosis and Langhans giant cell (single example in A) are characteristic of 2° tuberculosis.



(c) ketabton.com: The Digital Library

Leprosy

Also called Hansen disease. Caused by *Mycobacterium leprae*, an acid-fast bacillus that likes cool temperatures (infects skin and superficial nerves—"glove and stocking" loss of sensation A) and cannot be grown in vitro. Diagnosed via skin biopsy or tissue PCR. Reservoir in United States: armadillos.

Leprosy has 2 forms (many cases fall temporarily between two extremes):

- Lepromatous—presents diffusely over the skin, with Leonine (Lion-like) facies B, and is communicable (high bacterial load); characterized by low cell-mediated immunity with a largely Th2 response. Lepromatous form can be Lethal.
- Tuberculoid—limited to a few hypoesthetic, hairless skin plaques; characterized by high cell-mediated immunity with a largely Th1-type response and low bacterial load.

Treatment: dapsone and rifampin for tuberculoid form; clofazimine is added for lepromatous form.

Gram-negative lab algorithm



Important **tests** are in **bold.** Important ***pathogens*** are in ***bold italics.***
[a]Pleomorphic rod/coccobacillus

(c) ketabton.com: The Digital Library

Neisseria



Gram ⊖ diplococci. Metabolize glucose and produce IgA proteases. Contain lipooligosaccharides (LOS) with strong endotoxin activity. *N gonorrhoeae* is often intracellular (within neutrophils) A.

Acid production: MeninGococci—Maltose and Glucose; Gonococci—Glucose.

Gonococci	Meningococci
No polysaccharide capsule	Polysaccharide capsule
No maltose acid detection	Maltose acid detection
No vaccine due to antigenic variation of pilus proteins	Vaccine (type B vaccine available for at-risk individuals)
Sexually or perinatally transmitted	Transmitted via respiratory and oral secretions
Causes gonorrhea, septic arthritis, neonatal conjunctivitis (2–5 days after birth), pelvic inflammatory disease (PID), and Fitz-Hugh–Curtis syndrome	Causes meningococcemia with petechial hemorrhages and gangrene of toes B, meningitis, Waterhouse-Friderichsen syndrome (adrenal insufficiency, fever, DIC, shock)
Diagnosed with NAT	Diagnosed via culture-based tests or PCR
Condoms ↓ sexual transmission, erythromycin eye ointment prevents neonatal blindness	Rifampin, ciprofloxacin, or ceftriaxone prophylaxis in close contacts
Treatment: ceftriaxone (+ azithromycin or doxycycline, for possible chlamydial coinfection)	Treatment: ceftriaxone or penicillin G

Haemophilus influenzae





Small gram ⊖ (coccobacillary) rod. Aerosol transmission. Nontypeable (unencapsulated) strains are the most common cause of mucosal infections (otitis media, conjunctivitis, bronchitis) as well as invasive infections since the vaccine for capsular type b was introduced. Produces IgA protease.

Culture on chocolate agar, which contains factors V (NAD^+) and X (hematin) for growth; can also be grown with *S aureus*, which provides factor V via RBC hemolysis.

HaEMOPhilus causes Epiglottitis (endoscopic appearance in A, can be "cherry red" in children; "thumb sign" on lateral neck x-ray B), Meningitis, Otitis media, and Pneumonia.

Vaccine contains type b capsular polysaccharide (polyribosylribitol phosphate) conjugated to diphtheria toxoid or other protein. Given between 2 and 18 months of age.

Does not cause the flu (influenza virus does).

Treatment: amoxicillin +/– clavulanate for mucosal infections; ceftriaxone for meningitis; rifampin prophylaxis for close contacts.

Acinetobacter baumannii

Gram ⊖, strictly aerobic, oxidase ⊖ coccobacillus. Commensal opportunist but increasingly associated with resistant hospital-acquired infections, especially in ICU. Can cause ventilator-associated pneumonia and septicemia in immunocompromised patients.

(c) ketabton.com: The Digital Library

Bordetella pertussis

Gram ⊖, aerobic coccobacillus. Virulence factors include pertussis toxin (disables G_i), adenylate cyclase toxin (↑ cAMP), and tracheal cytotoxin. Three clinical stages:

- **C**atarrhal—low-grade fevers, **C**oryza.
- **P**aroxysmal—paroxysms of intense cough followed by inspiratory "whoo**P**" ("whooping cough"), posttussive vomiting.
- Convalescent—gradual recovery of chronic cough.

Prevented by Tdap, DTaP vaccines. May be mistaken as viral infection due to lymphocytic infiltrate resulting from immune response.

Treatment: macrolides; if allergic use TMP-SMX.

Brucella

Gram ⊖, aerobic coccobacillus. Transmitted via ingestion of contaminated animal products (eg, **un**pasteurized milk). Survives in macrophages in the reticuloendothelial system. Can form non-caseating granulomas. Typically presents with **un**dulant fever, night sweats, and arthralgia.

Treatment: doxycycline + rifampin or streptomycin.

Legionella pneumophila



Gram ⊖ rod. Gram stains poorly—use **silver** stain. Grow on **charcoal** yeast extract medium with **iron** and **cysteine**. Detected by presence of antigen in urine. Labs may show hyponatremia.

Aerosol transmission from environmental water source habitat (eg, air conditioning systems, hot water tanks). No person-to-person transmission.

Treatment: macrolide or quinolone.

Think of a French **legionnaire** (soldier) with his **silver** helmet, sitting around a campfire (**charcoal**) with his **iron** dagger—he is no **sissy** (cysteine).

Legionnaires' disease—severe pneumonia (often unilateral and lobar A), fever, GI and CNS symptoms. Common in smokers and in chronic lung disease.

Pontiac fever—mild flu-like syndrome.

Pseudomonas aeruginosa



Aeruginosa—**aer**obic; motile, catalase ⊕, gram ⊖ rod. Non-lactose fermenting. Oxidase ⊕. Frequently found in water. Has a grape-like odor.

PSEUDOMONAS is associated with: **P**neumonia, **S**epsis, **E**cthyma gangrenosum, **U**TIs, **D**iabetes, **O**steomyelitis, **M**ucoid polysaccharide capsule, **O**titis externa (swimmer's ear), **N**osocomial infections (eg, catheters, equipment), **A**ddicts (drug abusers), **S**kin infections (eg, hot tub folliculitis, wound infection in burn victims).

Mucoid polysaccharide capsule may contribute to chronic pneumonia in cystic fibrosis patients due to biofilm formation.

Produces **PEEP**: **P**hospholipase C (degrades cell membranes); **E**ndotoxin (fever, shock); **E**xotoxin A (inactivates EF-2); **P**igments: pyoverdine and pyocyanin (blue-green pigment A; also generates reactive oxygen species).

Corneal ulcers/keratitis in contact lens wearers/minor eye trauma.

Ecthyma gangrenosum—rapidly progressive, necrotic cutaneous lesion B caused by *Pseudomonas* bacteremia. Typically seen in immunocompromised patients.

Treatments include "**CAMPFIRE**" drugs:

- **C**arbapenems
- **A**minoglycosides
- **M**onobactams
- **P**olymyxins (eg, polymyxin B, colistin)
- **F**luoroquinolones (eg, ciprofloxacin, levofloxacin)
- Th**IR**d- and fourth-generation cephalosporins (eg, ceftazidime, cefepime)
- **E**xtended-spectrum penicillins (eg, piperacillin, ticarcillin)

(c) ketabton.com: The Digital Library

Salmonella vs *Shigella*

Both *Salmonella* and *Shigella* are gram ⊖ rods, non-lactose fermenters, oxidase ⊖, and can invade the GI tract via M cells of Peyer patches.

	Salmonella typhi (ty-Vi)	*Salmonella* spp. (except *S typhi*)	*Shigella*
RESERVOIRS	Humans only	Humans and animals	Humans only
SPREAD	Can disseminate hematogenously	Can disseminate hematogenously	Cell to cell; no hematogenous spread
H_2S PRODUCTION	Yes	Yes	No
FLAGELLA	Yes (**salmon swim**)	Yes (**salmon swim**)	No
VIRULENCE FACTORS	Endotoxin; **Vi** capsule	Endotoxin	Endotoxin; Shiga toxin (enterotoxin)
INFECTIOUS DOSE (ID_{50})	High—large inoculum required; acid-labile (inactivated by gastric acids)	High	Low—very small inoculum required; acid stable (resistant to gastric acids)
EFFECT OF ANTIBIOTICS ON FECAL EXCRETION	Prolongs duration	Prolongs duration	Shortens duration
IMMUNE RESPONSE	Primarily monocytes	PMNs in disseminated disease	Primarily PMN infiltration
GI MANIFESTATIONS	Constipation, followed by diarrhea	Diarrhea (possibly bloody)	Crampy abdominal pain → tenesmus, bloody mucoid stools (bacillary dysentery)
VACCINE	Oral vaccine contains live attenuated *S typhi* IM vaccine contains Vi capsular polysaccharide	No vaccine	No vaccine
UNIQUE PROPERTIES	▪ Causes typhoid fever (rose spots on abdomen, constipation, abdominal pain, fever; later GI ulceration and hemorrhage); treat with ceftriaxone or fluoroquinolone ▪ Carrier state with gallbladder colonization	▪ Poultry, eggs, pets, and turtles are common sources ▪ Antibiotics not indicated ▪ Gastroenteritis is usually caused by non-typhoidal *Salmonella*	▪ **4 F**'s: **F**ingers, **F**lies, **F**ood, **F**eces ▪ In order of decreasing severity (less toxin produced): *S dysenteriae*, *S flexneri*, *S boydii*, *S sonnei* ▪ Invasion of M cells is key to pathogenicity: organisms that produce little toxin can cause disease

Yersinia enterocolitica

Gram ⊖ pleomorphic rod/coccobacillus. Usually transmitted from pet feces (eg, puppies), contaminated milk, or pork. Can cause acute bloody diarrhea, pseudoappendicitis (right lower abdominal pain due to mesenteric adenitis and/or terminal ileitis), reactive arthritis in adults.

Lactose-fermenting enteric bacteria

Fermentation of **lactose** → pink colonies on MacCon**key** agar. Examples include *E coli*, *Enterobacter*, *Klebsiella*. *E coli* produces β-galactosidase, which breaks down lactose into glucose and galactose.

Lactose is **key**.

EMB agar—lactose fermenters grow as purple/black colonies. *E coli* grows colonies with a green sheen.

(c) ketabton.com: The Digital Library

Escherichia coli

Gram ⊖, indole ⊕ rod. *E coli* virulence factors: fimbriae—cystitis and pyelonephritis (P pili); K capsule—pneumonia, neonatal meningitis; LPS endotoxin—septic shock.

STRAIN	TOXIN AND MECHANISM	PRESENTATION
Enteroinvasive *E coli*	Microbe invades intestinal mucosa and causes necrosis and inflammation.	EIEC is Invasive; dysentery. Clinical manifestations similar to *Shigella*.
Enterotoxigenic *E coli*	Produces heat-labile and heat-stable enteroToxins. No inflammation or invasion.	ETEC; Traveler's diarrhea (watery).
Enteropathogenic *E coli*	No toxin produced. Adheres to apical surface, flattens villi, prevents absorption.	Diarrhea, usually in children (think EPEC and Pediatrics).
Enterohemorrhagic *E coli*	O157:H7 is most common serotype in US. Often transmitted via undercooked meat, raw leafy vegetables. Shiga-like toxin causes hemolytic-uremic syndrome: triad of anemia, thrombocytopenia, and acute kidney injury due to microthrombi forming on damaged endothelium → mechanical hemolysis (with schistocytes on peripheral blood smear), platelet consumption, and ↓ renal blood flow.	Dysentery (toxin alone causes necrosis and inflammation). Does not ferment sorbitol (vs other *E coli*). Hemorrhagic, Hamburgers, Hemolytic-uremic syndrome.

Klebsiella



Gram ⊖ rod; intestinal flora that causes lobar pneumonia in alcoholics and diabetics when aspirated. Very mucoid colonies A caused by abundant polysaccharide capsules. Dark red "currant jelly" sputum (blood/mucus).
Also cause of nosocomial UTIs. Associated with evolution of multidrug resistance (MDR).

ABCDE's of Klebsiella:
Aspiration pneumonia
aBscess in lungs and liver
"Currant jelly" sputum
Diabetes
EtOH abuse

Campylobacter jejuni



Gram ⊖, comma or S shaped (with polar flagella) A, oxidase ⊕, grows at 42°C ("*Camp*ylobacter likes the hot campfire").
Major cause of bloody diarrhea, especially in children. Fecal-oral transmission through person-to-person contact or via ingestion of undercooked contaminated poultry or meat, unpasteurized milk. Contact with infected animals (dogs, cats, pigs) is also a risk factor.
Common antecedent to Guillain-Barré syndrome and reactive arthritis.

(c) ketabton.com: The Digital Library

Vibrio cholerae

Gram ⊖, flagellated, comma shaped A, oxidase ⊕, grows in alkaline media. Endemic to developing countries. Produces profuse rice-water diarrhea via enterotoxin that permanently activates G_s, ↑ cAMP. Sensitive to stomach acid (acid labile); requires large inoculum (high ID_{50}) unless host has ↓ gastric acidity. Transmitted via ingestion of contaminated water or uncooked food (eg, raw shellfish). Treat promptly with oral rehydration solution.

Helicobacter pylori



Curved, flagellated (motile), gram ⊖ rod A that is **triple** ⊕: catalase ⊕, oxidase ⊕, and urease ⊕ (can use urea breath test or fecal antigen test for diagnosis). Urease produces ammonia, creating an alkaline environment, which helps *H pylori* survive in acidic mucosa. Colonizes mainly antrum of stomach; causes gastritis and peptic ulcers (especially duodenal). Risk factor for peptic ulcer disease, gastric adenocarcinoma, and MALT lymphoma.

Most common initial treatment is **triple** therapy: **A**moxicillin (metronidazole if penicillin allergy) + **C**larithromycin + **P**roton pump inhibitor; **A**ntibiotics **C**ure *Pylori*. Bismuth-based quadruple therapy if concerned about macrolide resistance.

Spirochetes



Spiral-shaped bacteria A with axial filaments. Includes ***B**orrelia* (big size), ***L**eptospira*, and ***T**reponema*. Only *Borrelia* can be visualized using aniline dyes (Wright or Giemsa stain) in light microscopy due to size. *Treponema* is visualized by dark-field microscopy or direct fluorescent antibody (DFA) microscopy.

BLT.
Borrelia is **B**ig.

Lyme disease



Caused by *Borrelia burgdorferi*, which is transmitted by the *Ixodes* deer tick A (also vector for *Anaplasma* spp. and protozoa *Babesia*). Natural reservoir is the mouse; deer are essential to tick life cycle but do not harbor *Borrelia*.

Common in northeastern United States.

Stage 1—early localized: erythema migrans (typical "bulls-eye" configuration B is pathognomonic but not always present), flu-like symptoms.

Stage 2—early disseminated: secondary lesions, carditis, AV block, facial nerve (Bell) palsy, migratory myalgias/transient arthritis.

Stage 3—late disseminated: encephalopathy, chronic arthritis.

A Key **Lyme** pie to the **FACE**:
Facial nerve palsy (typically bilateral)
Arthritis
Cardiac block
Erythema migrans

Treatment: doxycycline (1st line); amoxicillin and, if severe illness, CNS signs, or heart block, ceftriaxone

(c) ketabton.com: The Digital Library

Leptospira interrogans Spirochete with hook-shaped ends found in water contaminated with animal urine.

Leptospirosis—flu-like symptoms, myalgias (classically of calves), jaundice, photophobia with conjunctival suffusion (erythema without exudate). Prevalent among surfers and in tropics (eg, Hawaii).

Weil disease (icterohemorrhagic leptospirosis)—severe form with jaundice and azotemia from liver and kidney dysfunction, fever, hemorrhage, and anemia.

Syphilis	Caused by spirochete *Treponema pallidum*. Treatment: penicillin G.
Primary syphilis	Localized disease presenting with **painless** chancre A. Use fluorescent or dark-field microscopy to visualize treponemes in fluid from chancre B. VDRL ⊕ in ~ 80%.
Secondary syphilis	Disseminated disease with constitutional symptoms, maculopapular rash C (including palms D and soles), condylomata lata E (smooth, painless, wart-like white lesions on genitals), lymphadenopathy, patchy hair loss; also confirmable with dark-field microscopy. Serologic testing: VDRL/RPR (nonspecific), confirm diagnosis with specific test (eg, FTA-ABS). Secondary syphilis = Systemic. Latent syphilis (⊕ serology without symptoms) may follow.
Tertiary syphilis	Gummas F (chronic granulomas), aortitis (vasa vasorum destruction), neurosyphilis (tabes dorsalis, "general paresis"), Argyll Robertson pupil (constricts with accommodation but is not reactive to light; also called "prostitute's pupil" since it accommodates but does not react). Signs: broad-based ataxia, ⊕ Romberg, Charcot joint, stroke without hypertension. For neurosyphilis: test spinal fluid with VDRL, FTA-ABS, and PCR.
Congenital syphilis	Presents with facial abnormalities such as rhagades (linear scars at angle of mouth, black arrow in G), snuffles (nasal discharge, red arrow in G), saddle nose, notched (Hutchinson) teeth H, mulberry molars, and short maxilla; saber shins; CN VIII deafness. To prevent, treat mother early in pregnancy, as placental transmission typically occurs after first trimester.

(c) ketabton.com: The Digital Library

VDRL false positives

VDRL detects nonspecific antibody that reacts with beef cardiolipin. Quantitative, inexpensive, and widely available test for syphilis (sensitive but not specific).

False-**P**ositive results on **VDRL** with:
Pregnancy
Viral infection (eg, EBV, hepatitis)
Drugs (eg, chlorpromazine, procainamide)
Rheumatic fever (rare)
Lupus and leprosy



Jarisch-Herxheimer reaction

Flu-like syndrome (fever, chills, headache, myalgia) after antibiotics are started; due to killed bacteria (usually spirochetes) releasing toxins.

Gardnerella vaginalis



A pleomorphic, gram-variable rod involved in bacterial vaginosis. Presents as a gray vaginal discharge with a **fishy** smell; nonpainful (vs vaginitis). Associated with sexual activity, but not sexually transmitted. Bacterial **vagin**osis is also characterized by overgrowth of certain anaerobic bacteria in vagina (due to ↓ lactobacilli). **Clue** cells (vaginal epithelial cells covered with ***Gardner**ella*) have stippled appearance along outer margin (arrow in **A**).

I don't have a **clue** why I smell **fish** in the **vagin**a **garden**!
Amine whiff test—mixing discharge with 10% KOH enhances fishy odor.
Treatment: metronidazole or clindamycin.

Chlamydiae



Chlamydiae cannot make their own ATP. They are obligate intracellular organisms that cause mucosal infections. 2 forms:
- **E**lementary body (small, dense) is "**E**nfectious" and **E**nters cell via **E**ndocytosis; transforms into reticulate body.
- **R**eticulate body **R**eplicates in cell by fission; **R**eorganizes into elementary bodies.

Chlamydia trachomatis causes neonatal and follicular adult conjunctivitis **A**, nongonococcal urethritis, PID, and reactive arthritis.
Chlamydophila pneumoniae and *Chlamydophila psittaci* cause atypical pneumonia; transmitted by aerosol.

Chlamydial cell wall lacks classic peptidoglycan (due to reduced muramic acid), rendering β-lactam antibiotics ineffective.
Chlamys = cloak (intracellular).
C psittaci—has an avian reservoir (**p**arrots), causes atypical **p**neumonia.
Lab diagnosis: PCR, nucleic acid amplification test. Cytoplasmic inclusions (reticulate bodies) seen on Giemsa or fluorescent antibody–stained smear.
Treatment: azithromycin (favored because one-time treatment) or doxycycline. Add ceftriaxone for possible concomitant gonorrhea.

(c) ketabton.com: The Digital Library

Chlamydia trachomatis serotypes

Types A, B, and C	Chronic infection, cause blindness due to follicular conjunctivitis in Africa.	**A**BC = **A**frica, **B**lindness, **C**hronic infection.
Types D–K	Urethritis/PID, ectopic pregnancy, neonatal pneumonia (staccato cough) with eosinophilia, neonatal conjunctivitis (1–2 weeks after birth).	D–K = everything else. Neonatal disease can be acquired during passage through infected birth canal.
Types L1, L2, and L3	Lymphogranuloma venereum—small, painless ulcers on genitals → swollen, painful inguinal lymph nodes that ulcerate (buboes). Treat with doxycycline.	

Zoonotic bacteria

Zoonosis—infectious disease transmitted between animals and humans.

SPECIES	DISEASE	TRANSMISSION AND SOURCE
***Anaplasma* spp**	Anaplasmosis	*Ixodes* ticks (live on deer and mice)
***Bartonella* spp**	Cat scratch disease, bacillary angiomatosis	Cat scratch
Borrelia burgdorferi	Lyme disease	*Ixodes* ticks (live on deer and mice)
Borrelia recurrentis	**R**elapsing fever	Louse (recurrent due to variable surface antigens)
***Brucella* spp**	Brucellosis/**un**dulant fever	**Un**pasteurized dairy
Campylobacter	Bloody diarrhea	Feces from infected pets/animals; contaminated meats/foods/hands
Chlamydophila psittaci	Psittacosis	Parrots, other birds
Coxiella burnetii	Q fever	Aerosols of cattle/sheep amniotic fluid
Ehrlichia chaffeensis	Ehrlichiosis	*Amblyomma* (Lone Star tick)
Francisella tularensis	Tularemia	Ticks, rabbits, deer flies
***Leptospira* spp**	Leptospirosis	Animal urine in water; recreational water use
Mycobacterium leprae	Leprosy	Humans with lepromatous leprosy; armadillo (rare)
Pasteurella multocida	Cellulitis, osteomyelitis	Animal bite, cats, dogs
Rickettsia prowazekii	Epidemic typhus	Human to human via human body louse
Rickettsia rickettsii	Rocky Mountain spotted fever	*Dermacentor* (dog tick)
Rickettsia typhi	Endemic typhus	Fleas
***Salmonella* spp (except *S typhi*)**	Diarrhea (which may be bloody), vomiting, fever, abdominal cramps	Reptiles and poultry
Yersinia pestis	Plague	Fleas (rats and prairie dogs are reservoirs)

(c) ketabton.com: The Digital Library

Rickettsial diseases and vector-borne illnesses

Treatment: doxycycline.

RASH COMMON		
Rocky Mountain spotted fever	*Rickettsia rickettsii*, vector is tick. Despite its name, disease occurs primarily in the South Atlantic states, especially North Carolina. Rash typically starts at wrists A and ankles and then spreads to trunk, palms, and soles.	Classic triad—headache, fever, rash (vasculitis). **P**alms and **s**oles rash is seen in **C**oxsackievirus **A** infection (hand, foot, and mouth disease), **R**ocky Mountain spotted fever, and 2° **S**yphilis (you drive **CARS** using your **p**alms and **s**oles).
Typhus	Endemic (fleas)—*R typhi*. Epidemic (human body louse)—*R prowazekii*. Rash starts centrally and spreads out, sparing palms and soles.	*R*ickettsii on the w**R**ists, **T**yphus on the **T**runk.
RASH RARE		
Ehrlichiosis	*E*hrlichia, vector is tick. **M**onocytes with morulae B (mul**b**erry-like inclusions) in cytoplasm.	**M**EGA **b**erry— **M**onocytes = **E**hrlichiosis **G**ranulocytes = **A**naplasmosis
Anaplasmosis	*A*naplasma, vector is tick. **G**ranulocytes with morulae C in cytoplasm.	
Q fever	*C*oxiella *b*urnetii, no arthropod vector. Spores inhaled as aerosols from cattle/sheep amniotic fluid. Presents with headache, cough, influenza-like symptoms, pneumonia, possibly in combination with hepatitis. Common cause of culture ⊖ endocarditis.	**Q** fever is caused by a **Q**uite **C**omplicated **B**ug because it has no rash or vector and its causative organism can survive outside in its endospore form. Not in the *Rickettsia* genus, but closely related.



Mycoplasma pneumoniae



Classic cause of atypical "walking pneumonia" (insidious onset, headache, nonproductive cough, patchy or diffuse interstitial infiltrate).
Occurs frequently in those <30 years old; outbreaks in military recruits, prisons, colleges.
X-ray looks worse than patient. High titer of **c**old agglutinins (IgM), which can agglutinate RBCs.
Treatment: macrolides, doxycycline, or fluoroquinolone (penicillin ineffective since *Mycoplasma* has no cell wall).

Not seen on Gram stain. Pleomorphic A. Bacterial membrane contains sterols for stability. Grown on Eaton agar.

Mycoplasma gets **c**old without a **c**oat (no cell wall).
Can cause atypical variant of Stevens-Johnson syndrome, typically in children and adolescents.

(c) ketabton.com: The Digital Library

▸ MICROBIOLOGY—MYCOLOGY

Systemic mycoses

All of the following can cause pneumonia and can disseminate.
All are caused by dimorphic fungi: **cold** (20°C) = **mold**; **heat** (37°C) = **yeast**. Only exception is *Coccidioides*, which is a spherule (not yeast) in tissue.
Systemic mycoses can form granulomas (like TB); cannot be transmitted person-to-person (unlike TB).
Treatment: fluconazole or itraconazole for **local** infection; amphotericin B for **systemic** infection.

DISEASE	ENDEMIC LOCATION	PATHOLOGIC FEATURES	UNIQUE SIGNS/SYMPTOMS	NOTES
Histoplasmosis 	Mississippi and Ohio River Valleys	Macrophage filled with *Histoplasma* (smaller than RBC) A	Palatal/tongue ulcers, splenomegaly, pancytopenia	**H**isto **h**ides (within macrophages) Associated with bird or bat droppings (eg, spelunking) Diagnosis via urine/serum antigen
Blastomycosis	Eastern and Central US, Great Lakes	**Broad**-based budding of *Blastomyces* (same size as RBC) B	Inflammatory lung disease Disseminates to bone/skin (may mimic SCC) Forms granulomatous nodules	**B**lasto **b**uds **broad**ly
Coccidioidomycosis 	Southwestern US, California	Spherule (much larger than RBC) filled with endospores of *Coccidioides* C	Disseminates to skin/bone Erythema nodosum (desert bumps) or multiforme Arthralgias (desert rheumatism) Can cause meningitis	Associated with dust exposure in endemic areas (eg, archeological excavations, earthquakes)
Para-coccidioidomycosis 	**Latin America**	Budding yeast of *Paracoccidioides* with "**captain's wheel**" formation (much larger than RBC) D	Similar to blastomycosis, males > females	**Para**coccidio **para**sails with the **captain's wheel** all the way to **Latin America**



(c) ketabton.com: The Digital Library

Cutaneous mycoses

Tinea (dermatophytes)	Clinical name for dermatophyte (cutaneous fungal) infections. Dermatophytes include *Microsporum*, *Trichophyton*, and *Epidermophyton*. Branching septate hyphae visible on KOH preparation with blue fungal stain A. Associated with pruritus.
Tinea capitis	Occurs on head, scalp. Associated with lymphadenopathy, alopecia, scaling B.
Tinea corporis	Occurs on body (usually torso). Characterized by enlarging erythematous, scaly rings ("ringworm") with central clearing C. Can be acquired from contact with infected pets or farm animals.
Tinea cruris	Occurs in inguinal area D. Often does not show the central clearing seen in tinea corporis.
Tinea pedis	Three varieties: ▪ Interdigital E; most common ▪ Moccasin distribution F ▪ Vesicular type
Tinea unguium	Onychomycosis; occurs on nails.
Tinea (pityriasis) versicolor	Caused by *Malassezia* spp. (*Pityrosporum* spp.), a yeast-like fungus (not a dermatophyte despite being called tinea). Degradation of lipids produces acids that inhibit tyrosinase (involved in melanin synthesis) → hypopigmentation G; hyperpigmentation and/or pink patches can also occur due to inflammatory response. Less pruritic than dermatophytes. Can occur any time of year, but more common in summer (hot, humid weather). "Spaghetti and meatballs" appearance on microscopy H. Treatment: selenium sulfide, topical and/or oral antifungal medications.

(c) ketabton.com: The Digital Library

Opportunistic fungal infections

Candida albicans	*alba* = white. Dimorphic; forms pseudohyphae and budding yeasts at 20°C A, germ tubes at 37°C B. Systemic or superficial fungal infection. Causes oral C and esophageal thrush in immunocompromised (neonates, steroids, diabetes, AIDS), vulvovaginitis (diabetes, use of antibiotics), diaper rash, endocarditis (IV drug users), disseminated candidiasis (especially in neutropenic patients), chronic mucocutaneous candidiasis. Treatment: oral fluconazole/topical azoles for vaginal; nystatin, azoles, or, rarely, echinocandins for oral; fluconazole, echinocandins, or amphotericin B for esophageal or systemic disease.
Aspergillus fumigatus	Septate hyphae that branch at 45° Acute Angle D E. Causes invasive aspergillosis in immunocompromised patients, neutrophil dysfunction (eg, chronic granulomatous disease). Can cause aspergillomas F in pre-existing lung cavities, especially after TB infection. Some species of *Aspergillus* produce Aflatoxins (associated with hepatocellular carcinoma). Treatment: voriconazole or echinocandins (2nd-line). Allergic bronchopulmonary aspergillosis (ABPA)—hypersensitivity response to *Aspergillus* growing in lung mucus. Associated with asthma and cystic fibrosis; may cause bronchiectasis and eosinophilia.
Cryptococcus neoformans	5–10 μm with narrow budding. Heavily encapsulated yeast. Not dimorphic. Found in soil, pigeon droppings. Acquired through inhalation with hematogenous dissemination to meninges. Highlighted with India ink (clear halo G) and mucicarmine (red inner capsule H). Latex agglutination test detects polysaccharide capsular antigen and is more sensitive and specific. Causes cryptococcosis, cryptococcal meningitis, cryptococcal encephalitis ("soap bubble" lesions in brain), primarily in immunocompromised. Treatment: amphotericin B + flucytosine followed by fluconazole for cryptococcal meningitis.
***Mucor* and *Rhizopus* spp**	Irregular, broad, nonseptate hyphae branching at wide angles I. Causes mucormycosis, mostly in ketoacidotic diabetic and/or neutropenic patients (eg, leukemia). Inhalation of spores → fungi proliferate in blood vessel walls, penetrate cribriform plate, and enter brain. Rhinocerebral, frontal lobe abscess; cavernous sinus thrombosis. Headache, facial pain, black necrotic eschar on face J; may have cranial nerve involvement. Treatment: surgical debridement, amphotericin B or isavuconazole.



FAS1_2019_03-Microbiology.indd 153 11/14/19 12:21 PM

(c) ketabton.com: The Digital Library

Pneumocystis jirovecii

Causes *Pneumocystis* pneumonia (PCP), a diffuse interstitial pneumonia A. Yeast-like fungus (originally classified as protozoan). Most infections are asymptomatic. Immunosuppression (eg, AIDS) predisposes to disease. Diffuse, bilateral ground-glass opacities on chest imaging, with pneumatoceles B. Diagnosed by bronchoalveolar lavage or lung biopsy. Disc-shaped yeast seen on methenamine silver stain of lung tissue C or with fluorescent antibody.

Treatment/prophylaxis: TMP-SMX, pentamidine, dapsone (prophylaxis as single agent, or treatment in combination with TMP), atovaquone. Start prophylaxis when CD4+ count drops to < 200 cells/mm^3 in HIV patients.



Sporothrix schenckii

Causes sporotrichosis. Dimorphic fungus. Exists as a **cigar**-shaped yeast at 37 °C in the human body and as hyphae with spores in soil (conidia). Lives on vegetation. When spores are traumatically introduced into the skin, typically by a thorn ("**rose gardener**'s disease"), causes local pustule or ulcer with nodules along draining lymphatics (ascending lymphangitis A). Disseminated disease possible in immunocompromised host.

Treatment: itraconazole or **pot**assium iodide (only for cutaneous/lymphocutaneous).

Think of a **rose gardener** who smokes a **cigar** and **pot**.

(c) ketabton.com: The Digital Library

▸ MICROBIOLOGY—PARASITOLOGY

Protozoa—gastrointestinal infections

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
Giardia lamblia	**Giardiasis**—bloating, flatulence, foul-smelling, fatty diarrhea (often seen in campers/hikers)—think fat-rich Ghirardelli chocolates for fatty stools of *Giardia*	Cysts in water	Multinucleated trophozoites A or cysts B in stool, antigen detection	Metronidazole
Entamoeba histolytica	**Amebiasis**—bloody diarrhea (dysentery), liver abscess ("anchovy paste" exudate), RUQ pain; histology of colon biopsy shows flask-shaped ulcers	Cysts in water	Serology, antigen testing, and/or trophozoites (with engulfed RBCs C in the cytoplasm) or cysts with up to 4 nuclei in stool D; Entamoeba Eats Erythrocytes	Metronidazole; paromomycin or iodoquinol for asymptomatic cyst passers
Cryptosporidium	Severe diarrhea in AIDS Mild disease (watery diarrhea) in immunocompetent hosts	Oocysts in water	Oocysts on acid-fast stain E, antigen detection	Prevention (by filtering city water supplies); nitazoxanide in immunocompetent hosts









(c) ketabton.com: The Digital Library

Protozoa—CNS infections

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
Toxoplasma gondii	Immunocompetent: mononucleosis-like symptoms, ⊖ heterophile antibody test Reactivation in AIDS → brain abscesses usually seen as multiple ring-enhancing lesions on MRI A Congenital toxoplasmosis: classic triad of chorioretinitis, hydrocephalus, and intracranial calcifications	Cysts in meat (most common); oocysts in cat feces; crosses placenta (pregnant women should avoid cats)	Serology, biopsy (tachyzoite) B	Sulfadiazine + pyrimethamine
Naegleria fowleri	Rapidly fatal meningoencephalitis	Swimming in warm freshwater; enters via cribriform plate	Amoebas in CSF C	Amphotericin B has been effective for a few survivors
Trypanosoma brucei	**African sleeping sickness**—enlarged lymph nodes, recurring fever (due to antigenic variation), somnolence, coma	Tsetse fly, a painful bite	Trypomastigote in blood smear D	**Sur**amin for blood-borne disease or **mel**arsoprol for CNS penetration ("I **sur**e am **mel**low when I'm **sleeping**")



FAS1_2019_03-Microbiology.indd 156 11/14/19 12:21 PM

(c) ketabton.com: The Digital Library

Protozoa—hematologic infections

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
Plasmodium ***P vivax/ovale*** ***P falciparum*** ***P malariae***  	**Malaria**—fever, headache, anemia, splenomegaly *P vivax/ovale*—48-hr cycle (tertian; includes fever on first day and third day, thus fevers are actually 48 hr apart); dormant form (hypnozoite) in liver *P falciparum*—severe; irregular fever patterns; parasitized RBCs occlude capillaries in brain (cerebral malaria), kidneys, lungs *P malariae*—72-hr cycle (quartan)	*Anopheles* mosquito	Blood smear: trophozoite ring form within RBC A, schizont containing merozoites; red granules (Schüffner stippling) B throughout RBC cytoplasm seen with *P vivax/ovale*	Chloroquine (for sensitive species); if resistant, use mefloquine or atovaquone/proguanil If life-threatening, use intravenous quinidine or artesunate (test for G6PD deficiency) For *P vivax/ovale*, add primaquine for hypnozoite (test for G6PD deficiency)
Babesia 	**Babesiosis**—fever and hemolytic anemia; predominantly in northeastern United States; asplenia ↑ risk of severe disease	*Ixodes* tick (also vector for *Borrelia burgdorferi* and *Anaplasma* spp)	Blood smear: ring form C1, "Maltese cross" C2; PCR	Atovaquone + azithromycin

(c) ketabton.com: The Digital Library

Protozoa—others

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
Visceral infections				
Trypanosoma cruzi	**Chagas disease**—dilated cardiomyopathy with apical atrophy, megacolon, megaesophagus; predominantly in South America Unilateral periorbital swelling (Romaña sign) characteristic of acute stage	Triatomine insect (kissing bug) bites and defecates around the mouth or eyes; fecal transmission into bite site or mucosa	Trypomastigote in blood smear A	**Benz**nidazole or ni**fur**timox; **cruzi**ng in my **Benz**, with a **fur** coat on
***Leishmania* spp**	**Visceral leishmaniasis (kala-azar)**—spiking fevers, hepatosplenomegaly, pancytopenia **Cutaneous leishmaniasis**—skin ulcers B	Sandfly	Macrophages containing amastigotes C	Amphotericin B, sodium stibogluconate
Sexually transmitted infections				
Trichomonas vaginalis	**Vaginitis**—foul-smelling, greenish discharge; itching and burning; do not confuse with *Gardnerella vaginalis*, a gram-variable bacterium associated with bacterial vaginosis	Sexual (cannot exist outside human because it cannot form cysts)	Trophozoites (motile) D on wet mount; punctate cervical hemorrhages ("strawberry cervix")	Metronidazole for patient and partner (prophylaxis; check for STI)



Nematode routes of infection

Ingested—*Enterobius*, *Ascaris*, *Toxocara*, *Trichinella*, *Trichuris*	You'll get sick if you **EATTT** these!
Cutaneous—*Strongyloides*, *Ancylostoma*, *Necator*	These get into your feet from the **SAN**d
Bites—*Loa loa*, *Onchocerca volvulus*, *Wuchereria bancrofti*	Lay **LOW** to avoid getting bitten

(c) ketabton.com: The Digital Library

Nematodes (roundworms)

ORGANISM	DISEASE	TRANSMISSION	TREATMENT
Intestinal			
***Enterobius vermicularis* (pinworm)**	Causes anal pruritus (diagnosed by seeing egg A via the tape test).	Fecal-oral.	Bendazoles (bendy worms), pyrantel pamoate.
***Ascaris lumbricoides* (giant roundworm)**	May cause obstruction at ileocecal valve, biliary obstruction, intestinal perforation, migrates from nose/mouth.	Fecal-oral; knobby-coated, oval eggs seen in feces under microscope B.	Bendazoles.
***Strongyloides stercoralis* (threadworm)**	GI (eg, duodenitis), pulmonary (eg, dry cough, hemoptysis), and cutaneous (eg, pruritus) symptoms. Hyperinfection syndrome caused by autoinfection (larvae enter bloodstream).	Larvae in soil penetrate skin; rhabditiform larvae seen in feces under microscope.	Ivermectin or bendazoles.
***Ancylostoma* spp, *Necator americanus* (hookworms)**	Cause microcytic anemia by sucking blood from intestinal wall. Cutaneous larva migrans—pruritic, serpiginous rash C from walking barefoot on contaminated beach.	Larvae penetrate skin.	Bendazoles or pyrantel pamoate.
Trichinella spiralis	Larvae enter bloodstream, encyst in striated muscle D → myositis. Trichinosis—fever, vomiting, nausea, periorbital edema, myalgia.	Undercooked meat (especially pork); fecal-oral (less likely).	Bendazoles.
***Trichuris trichiura* (whipworm)**	Often asymptomatic; loose stools, anemia, rectal prolapse in children.	Fecal-oral.	Bendazoles.
Tissue			
Toxocara canis	Visceral larva migrans—nematodes migrate to blood through intestinal wall → inflammation affecting liver, eyes (visual impairment, blindness), CNS (seizures, coma), heart (myocarditis).	Fecal-oral.	Bendazoles.
Onchocerca volvulus	Skin changes, loss of elastic fibers, river blindness (black skin nodules, "black sight"); allergic reaction possible.	Female black fly.	Ivermectin (ivermectin for river blindness).
Loa loa	Swelling in skin, worm in conjunctiva.	Deer fly, horse fly, mango fly.	Diethylcarbamazine.
Wuchereria bancrofti	Lymphatic filariasis (elephantiasis)—worms invade lymph nodes. → inflammation → lymphedema E; symptom onset after 9 mo–1 yr.	Female mosquito.	Diethylcarbamazine.







(c) ketabton.com: The Digital Library

Cestodes (tapeworms)

ORGANISM	DISEASE	TRANSMISSION	TREATMENT
Taenia solium A	Intestinal tapeworm	Ingestion of larvae encysted in undercooked pork	Praziquantel
	Cysticercosis, neurocysticercosis (cystic CNS lesions, seizures) B	Ingestion of eggs in food contaminated with human feces	Praziquantel; albendazole for neurocysticercosis
Diphyllobothrium latum	Vitamin B_{12} deficiency (tapeworm competes for B_{12} in intestine) → megaloblastic anemia	Ingestion of larvae in raw freshwater fish	Praziquantel
Echinococcus granulosus C	Hydatid cysts D ("eggshell calcification") in liver E; cyst rupture can cause anaphylaxis	Ingestion of eggs in food contaminated with dog feces Sheep are an intermediate host	Albendazole

Trematodes (flukes)

ORGANISM	DISEASE	TRANSMISSION	TREATMENT
Schistosoma  	Liver and spleen enlargement (*S mansoni*, egg with lateral spine A), fibrosis, inflammation, portal hypertension Chronic infection with *S haematobium* (egg with terminal spine B) can lead to squamous cell carcinoma of the bladder (painless hematuria) and pulmonary hypertension	Snails are intermediate host; cercariae penetrate skin of humans in contact with contaminated fresh water (eg, swimming or bathing)	Praziquantel
Clonorchis sinensis	Biliary tract inflammation → pigmented gallstones Associated with cholangiocarcinoma	Undercooked fish	Praziquantel

(c) ketabton.com: The Digital Library

Ectoparasites

Sarcoptes scabiei 	Mites burrow into stratum corneum and cause **scabies**—pruritus (worse at night) and serpiginous burrows (lines) often between fingers and toes A.	Common in children, crowded populations (jails, nursing homes); transmission through skin-to-skin contact (most common) or via fomites. Treatment: permethrin cream, washing/drying all clothing/bedding, treat close contacts.
Pediculus humanus/ Phthirus pubis 	Blood-sucking lice that cause intense pruritus with associated excoriations, commonly on scalp and neck (head lice), waistband and axilla (body lice), or pubic and perianal regions (pubic lice).	Body lice can transmit *Rickettsia prowazekii* (epidemic typhus), *Borrelia recurrentis* (relapsing fever), *Bartonella quintana* (trench fever). Treatment: pyrethroids, malathion, or ivermectin lotion, and nit B combing. Children with head lice can be treated at home without interrupting school attendance.

Parasite hints

ASSOCIATIONS	ORGANISM
Biliary tract disease, cholangiocarcinoma	*Clonorchis sinensis*
Brain cysts, seizures	*Taenia solium* (neurocysticercosis)
Hematuria, squamous cell bladder cancer	*Schistosoma haematobium*
Liver (hydatid) cysts	*Echinococcus granulosus*
Microcytic anemia	*Ancylostoma*, *Necator*
Myalgias, periorbital edema	*Trichinella spiralis*
Perianal pruritus	*Enterobius*
Portal hypertension	*Schistosoma mansoni*, *Schistosoma japonicum*
Vitamin B_{12} deficiency	*Diphyllobothrium latum*

(c) ketabton.com: The Digital Library

▸ MICROBIOLOGY—VIROLOGY

Viral structure—general features



Naked virus with icosahedral capsid | Enveloped virus with icosahedral capsid | Enveloped virus with helical capsid | Bacteriophage

Viral genetics

Recombination	Exchange of genes between 2 chromosomes by crossing over within regions of significant base sequence homology.	
Reassortment	When viruses with segmented genomes (eg, influenza virus) exchange genetic material. For example, the 2009 novel H1N1 influenza A pandemic emerged via complex viral reassortment of genes from human, swine, and avian viruses. Has potential to cause antigenic shift.	
Complementation	When 1 of 2 viruses that infect the cell has a mutation that results in a nonfunctional protein, the nonmutated virus "complements" the mutated one by making a functional protein that serves both viruses. For example, hepatitis D virus requires the presence of replicating hepatitis B virus to supply HBsAg, the envelope protein for HDV.	 Functional Nonfunctional Functional
Phenotypic mixing	Occurs with simultaneous infection of a cell with 2 viruses. For progeny 1, genome of virus A can be partially or completely coated (forming pseudovirion) with the surface proteins of virus B. Type B protein coat determines the tropism (infectivity) of the hybrid virus. Progeny from subsequent infection of a cell by progeny 1 will have a type A coat that is encoded by its type A genetic material.	 Virus A Virus B Progeny 1 Progeny 2

(c) ketabton.com: The Digital Library

DNA viral genomes

All DNA viruses have dsDNA genomes except Parvoviridae (ssDNA).
All are linear except papilloma-, polyoma-, and hepadnaviruses (circular).

All are dsDNA (like our cells), except "**part-of-a-virus**" (**parvovirus**) is ssDNA.
Parvus = small.

RNA viral genomes

All RNA viruses have ssRNA genomes except Reoviridae (dsRNA).
⊕ stranded RNA viruses: I went to a **retro** (**retro**virus) **toga** (**toga**virus) party, where I drank **flav**ored (**flav**ivirus) **Corona** (**corona**virus) and ate **hippie** (**hepe**virus) **Cali**fornia (**cali**civirus) **pic**kles (**pic**ornavirus).

All are ssRNA, except "**repeat**o-virus" (**reo**virus) is dsRNA.

Naked viral genome infectivity

Purified nucleic acids of most dsDNA viruses (except poxviruses and HBV) and ⊕ strand ssRNA (≈ mRNA) viruses are infectious. Naked nucleic acids of ⊖ strand ssRNA and dsRNA viruses are not infectious. They require polymerases contained in the complete virion.

Viral envelopes

Generally, enveloped viruses acquire their envelopes from plasma membrane when they exit from cell. Exceptions include herpesviruses, which acquire envelopes from nuclear membrane.
Naked (nonenveloped) viruses include **P**apillomavirus, **A**denovirus, **P**arvovirus, **P**olyomavirus, **C**alicivirus, **P**icornavirus, **R**eovirus, and **Hep**evirus.

DNA = **PAPP**; RNA = **CPR** and **hep**evirus.
Give **PAPP** smears and **CPR** to a **naked hip**pie (**hepe**virus).
Enveloped DNA viruses **H**ave **H**elpful **P**rotection (**H**erpesvirus, **H**epadnavirus, **P**oxvirus).

DNA virus characteristics

Some general rules—all DNA viruses:

GENERAL RULE	COMMENTS
Are **HHAPPPP**y viruses	**H**epadna, **H**erpes, **A**deno, **P**ox, **P**arvo, **P**apilloma, **P**olyoma.
Are double stranded	Except parvo (single stranded).
Have linear genomes	Except papilloma and polyoma (circular, supercoiled) and hepadna (circular, incomplete).
Are icosahedral	Except pox (complex).
Replicate in the nucleus	Except pox (carries own DNA-dependent RNA polymerase).

(c) ketabton.com: The Digital Library

DNA viruses

All replicate in the nucleus (except poxvirus). "Pox is out of the box (nucleus)."

VIRAL FAMILY	ENVELOPE	DNA STRUCTURE	MEDICAL IMPORTANCE
Herpesviruses	Yes	DS and linear	See Herpesviruses entry
Poxvirus	Yes	DS and linear (largest DNA virus)	Smallpox eradicated world wide by use of the live-attenuated vaccine Cowpox ("milkmaid blisters") **Molluscum contagiosum**—flesh-colored papule with central umbilication
Hepadnavirus	Yes	Partially DS and circular	HBV: ▪ Acute or chronic hepatitis ▪ Not a retrovirus but has reverse transcriptase
Adenovirus  A	No	DS and linear	Febrile pharyngitis A—sore throat Acute hemorrhagic cystitis Pneumonia Conjunctivitis—"pink eye" Gastroenteritis Myocarditis
Papillomavirus	No	DS and circular	HPV–warts (serotypes 1, 2, 6, 11), CIN, cervical cancer (most commonly 16, 18)
Polyomavirus	No	DS and circular	JC virus—progressive multifocal leukoencephalopathy (PML) in HIV BK virus—transplant patients, commonly targets kidney JC: Junky Cerebrum; BK: Bad Kidney
Parvovirus	No	SS and linear (smallest DNA virus)	B19 virus—aplastic crises in sickle cell disease, "slapped cheek" rash in children (erythema infectiosum, or fifth disease); infects RBC precursors and endothelial cells → RBC destruction → hydrops fetalis and death in fetus, pure RBC aplasia and rheumatoid arthritis–like symptoms in adults

Herpesviruses

Enveloped, DS, and linear viruses

VIRUS	ROUTE OF TRANSMISSION	CLINICAL SIGNIFICANCE	NOTES
Herpes simplex virus-1	Respiratory secretions, saliva	Gingivostomatitis, keratoconjunctivitis A, herpes labialis (cold sores) B, herpetic whitlow on finger, temporal lobe encephalitis, esophagitis, erythema multiforme	Most commonly latent in trigeminal ganglia Most common cause of sporadic encephalitis, can present as altered mental status, seizures, and/or aphasia
Herpes simplex virus-2	Sexual contact, perinatal	Herpes genitalis C, neonatal herpes	Most commonly latent in sacral ganglia Viral meningitis more common with HSV-2 than with HSV-1

(c) ketabton.com: The Digital Library

Herpesviruses *(continued)*

VIRUS	ROUTE OF TRANSMISSION	CLINICAL SIGNIFICANCE	NOTES
Varicella-Zoster virus (HHV-3)	Respiratory secretions, contact with fluid from vesicles	Varicella-zoster (chickenpox D, shingles E), encephalitis, pneumonia Most common complication of shingles is post-herpetic neuralgia	Latent in dorsal root or trigeminal ganglia; CN V_1 branch involvement can cause herpes zoster ophthalmicus
Epstein-Barr virus (HHV-4)	Respiratory secretions, saliva; aka "kissing disease," (common in teens, young adults)	**Mononucleosis**—fever, hepatosplenomegaly F, pharyngitis, and lymphadenopathy (especially posterior cervical nodes); avoid contact sports until resolution due to risk of splenic rupture Associated with lymphomas (eg, endemic Burkitt lymphoma), nasopharyngeal carcinoma (especially Asian adults), lymphoproliferative disease in transplant patients	Infects B cells through CD21, "Must be 21 to drink Beer in a Barr" Atypical lymphocytes on peripheral blood smear G—not infected B cells but reactive cytotoxic T cells ⊕ Monospot test—heterophile antibodies detected by agglutination of sheep or horse RBCs Use of amoxicillin in mononucleosis can cause characteristic maculopapular rash
Cytomegalo-virus (HHV-5)	Congenital, transfusion, sexual contact, saliva, urine, transplant	Mononucleosis (⊖ Monospot) in immunocompetent patients; infection in immunocompromised, especially pneumonia in transplant patients; esophagitis; AIDS **retinitis** ("**sight**omegalovirus"): hemorrhage, cotton-wool exudates, vision loss Congenital CMV	Infected cells have characteristic "owl eye" intranuclear inclusions H Latent in mononuclear cells
Human herpes-viruses 6 and 7	Saliva	Roseola infantum (exanthem subitum): high fevers for several days that can cause seizures, followed by diffuse macular rash (starts on trunk then spreads to extremities) I	**Ros**eo**la**: fever first, **Ros**y (rash) **la**ter. HHV-7—less common cause of roseola
Human herpesvirus 8	Sexual contact	Kaposi sarcoma (neoplasm of endothelial cells). Seen in HIV/AIDS and transplant patients. Dark/violaceous plaques or nodules J representing vascular proliferations	Can also affect GI tract and lungs



(c) ketabton.com: The Digital Library

HSV identification



Viral culture for skin/genitalia.
CSF PCR for herpes encephalitis.
Tzanck test—a smear of an opened skin vesicle to detect multinucleated giant cells A commonly seen in HSV-1, HSV-2, and VZV infection. PCR of skin lesions is test of choice.
Tzanck heavens I do not have herpes.
Intranuclear eosinophilic Cowdry A inclusions also seen with HSV-1, HSV-2, VZV.

Receptors used by viruses

VIRUS	RECEPTORS
CMV	Integrins (heparan sulfate)
EBV	CD21
HIV	CD4, CXCR4, CCR5
Parvovirus B19	**P** antigen on RBCs
Rabies	Nicotinic AChR
Rhinovirus	I**CAM**-1 (**I cam**e to see the **rhino**)

(c) ketabton.com: The Digital Library

RNA viruses All replicate in the cytoplasm (except retrovirus and influenza virus). "Retro flu is outta cyt (sight)."

VIRAL FAMILY	ENVELOPE	RNA STRUCTURE	CAPSID SYMMETRY	MEDICAL IMPORTANCE
Reoviruses	No	DS linear Multisegmented	Icosahedral (double)	Coltivirus[a]—Colorado tick fever Rotavirus—cause of fatal diarrhea in children
Picornaviruses	No	SS ⊕ linear	Icosahedral	Poliovirus—polio-Salk/Sabin vaccines—IPV/OPV Echovirus—aseptic meningitis Rhinovirus—"common cold" Coxsackievirus—aseptic meningitis; herpangina (mouth blisters, fever); hand, foot, and mouth disease; myocarditis; pericarditis HAV—acute viral hepatitis PERCH
Hepevirus	No	SS ⊕ linear	Icosahedral	HEV
Caliciviruses	No	SS ⊕ linear	Icosahedral	Norovirus—viral gastroenteritis
Flaviviruses	Yes	SS ⊕ linear	Icosahedral	HCV Yellow fever[a] Dengue[a] St. Louis encephalitis[a] West Nile virus[a]—meningoencephalitis, flaccid paralysis Zika virus[a]
Togaviruses	Yes	SS ⊕ linear	Icosahedral	Toga CREW—Chikungunya virus[a] (co-infection with dengue virus can occur), Rubella, Eastern and Western equine encephalitis
Retroviruses	Yes	SS ⊕ linear 2 copies	Icosahedral (HTLV), complex and conical (HIV)	Have reverse transcriptase HTLV—T-cell leukemia HIV—AIDS
Coronaviruses	Yes	SS ⊕ linear	Helical	"Common cold," SARS, MERS
Orthomyxoviruses	Yes	SS ⊖ linear 8 segments	Helical	Influenza virus
Paramyxoviruses	Yes	SS ⊖ linear Nonsegmented	Helical	PaRaMyxovirus: Parainfluenza—croup RSV—bronchiolitis in babies Measles, Mumps
Rhabdoviruses	Yes	SS ⊖ linear	Helical	Rabies
Filoviruses	Yes	SS ⊖ linear	Helical	Ebola/Marburg hemorrhagic fever—often fatal.
Arenaviruses	Yes	SS ⊕ and ⊖ circular 2 segments	Helical	LCMV—lymphocytic choriomeningitis virus Lassa fever encephalitis—spread by rodents
Bunyaviruses	Yes	SS ⊖ circular 3 segments	Helical	California encephalitis[a] Sandfly/Rift Valley fevers[a] Crimean-Congo hemorrhagic fever[a] Hantavirus—hemorrhagic fever, pneumonia
Delta virus	Yes	SS ⊖ circular	Uncertain	HDV is a "defective" virus that requires the presence of HBV to replicate

SS, single-stranded; DS, double-stranded; ⊕, positive sense; ⊖, negative sense; [a]= arbovirus, arthropod borne (mosquitoes, ticks).

(c) ketabton.com: The Digital Library

Negative-stranded viruses	Must transcribe ⊖ strand to ⊕. Virion brings its own RNA-dependent RNA polymerase. They include **A**renaviruses, **B**unyaviruses, **P**aramyxoviruses, **O**rthomyxoviruses, **F**iloviruses, and **R**habdoviruses.	**A**lways **B**ring **P**olymerase **O**r **F**ail **R**eplication.
Segmented viruses	All are RNA viruses. They include **B**unyaviruses (**3** segments), **O**rthomyxoviruses (influenza viruses) (**8** segments), **A**renaviruses (**2** segments), and **R**eoviruses (**10-12** segments).	**BOAR**ding flight **382** in **10-12** minutes.
Picornavirus	Includes **P**oliovirus, **E**chovirus, **R**hinovirus, **C**oxsackievirus, and **H**AV. RNA is translated into 1 large polypeptide that is cleaved by virus-encoded proteases into functional viral proteins. Can cause aseptic (viral) meningitis (except rhinovirus and HAV). All are enteroviruses except rhinovirus and HAV.	Pico**RNA**virus = small **RNA** virus. **PERCH** on a "**peak**" (**pic**o).
Rhinovirus	A picornavirus. Nonenveloped RNA virus. Cause of common cold; > 100 serologic types. Acid labile—destroyed by stomach acid; therefore, does not infect the GI tract (unlike the other picornaviruses).	**Rhino** has a runny **nose**.
Yellow fever virus	A flavivirus (also an arbovirus) transmitted by *Aedes* mosquitoes. Virus has a monkey or human reservoir. Symptoms: high fever, black vomitus, and jaundice. May see Councilman bodies (eosinophilic apoptotic globules) on liver biopsy.	*Flavi* = yellow, jaundice.
Rotavirus 	Segmented dsRNA virus (a reovirus) **A**. Most important global cause of infantile gastroenteritis. Major cause of acute diarrhea in the United States during winter, especially in day care centers, kindergartens. Villous destruction with atrophy leads to ↓ absorption of Na^+ and loss of K^+.	**ROTA**virus = **R**ight **O**ut **T**he **A**nus. CDC recommends routine vaccination of all infants except those with a history of intussusception or SCID.

(c) ketabton.com: The Digital Library

Influenza viruses

Orthomyxoviruses. Enveloped, ⊖ ssRNA viruses with 8-segment genome. Contain hemagglutinin (binds sialic acid and promotes viral entry) and neuraminidase (promotes progeny virion release) antigens. Patients at risk for fatal bacterial superinfection, most commonly *S aureus*, *S pneumoniae*, and *H influenzae*.

Reformulated vaccine ("the flu shot") contains viral strains most likely to appear during the flu season, due to the virus' rapid genetic change.
Killed viral vaccine is most frequently used.
Live attenuated vaccine contains temperature-sensitive mutant that replicates in the nose but not in the lung; administered intranasally.
Treatment: supportive +/– neuraminidase inhibitor (eg, oseltamivir, zanamivir).

Genetic/antigenic shift

Infection of 1 cell by 2 different segmented viruses (eg, swine influenza and human influenza viruses) → RNA segment reassortment → dramatically different virus (genetic shift) → major global outbreaks (pandemics).

Sudden shift is more deadly than gradual drift.

Genetic/antigenic drift

Random mutation in hemagglutinin or neuraminidase genes → minor changes (antigenic drift) → local outbreaks (epidemics).



Rubella virus

A togavirus. Causes rubella, once known as German (3-day) measles. Fever, postauricular and other lymphadenopathy, arthralgias, and fine, maculopapular rash that starts on face and spreads centrifugally to involve trunk and extremities A.
Causes mild disease in children but serious congenital disease (a TORCH infection). Congenital rubella findings include "blueberry muffin" appearance due to dermal extramedullary hematopoiesis.

Paramyxoviruses

Paramyxoviruses cause disease in children. They include those that cause parainfluenza (croup), mumps, measles, RSV, and human metapneumovirus, which causes respiratory tract infection (bronchiolitis, pneumonia) in infants. All contain surface F (fusion) protein, which causes respiratory epithelial cells to fuse and form multinucleated cells. Palivizumab (monoclonal antibody against F protein) prevents pneumonia caused by RSV infection in premature infants.
Palivizumab for Paramyxovirus (RSV) Prophylaxis in Preemies.

(c) ketabton.com: The Digital Library

Acute laryngotracheobronchitis



Also called croup. Caused by parainfluenza viruses. Virus membrane contains hemagglutinin (binds sialic acid and promotes viral entry) and neuraminidase (promotes progeny virion release) antigens. Results in a "seal-like" barking cough and inspiratory stridor. Narrowing of upper trachea and subglottis leads to characteristic steeple sign on x-ray A. Severe croup can result in pulsus paradoxus 2° to upper airway obstruction.

Measles (rubeola) virus



Usual presentation involves prodromal fever with cough, coryza, and conjunctivitis, then eventually Koplik spots (bright red spots with blue-white center on buccal mucosa A), followed 1–2 days later by a maculopapular rash B that starts at the head/neck and spreads downward.

Lymphadenitis with Warthin-Finkeldey giant cells (fused lymphocytes) in a background of paracortical hyperplasia. Possible sequelae:

- Subacute sclerosing panencephalitis (SSPE): personality changes, dementia, autonomic dysfunction, death (occurs years later)
- Encephalitis (1:1000): symptoms appear within few days of rash
- Giant cell pneumonia (rare except in immunosuppressed)

4 C's of measles:
Cough
Coryza
Conjunctivitis
"C"oplik spots

Vitamin A supplementation can reduce morbidity and mortality from measles, particularly in malnourished children.

Pneumonia is the most common cause of measles-associated death in children.

Mumps virus

Uncommon due to effectiveness of MMR vaccine.

Symptoms: Parotitis A, Orchitis (inflammation of testes), aseptic Meningitis, and Pancreatitis. Can cause sterility (especially after puberty).

Mumps makes your parotid glands and testes as big as POM-Poms.

(c) ketabton.com: The Digital Library

Rabies virus





Bullet-shaped virus A. Negri bodies (cytoplasmic inclusions B) commonly found in Purkinje cells of cerebellum and in hippocampal neurons. Rabies has long incubation period (weeks to months) before symptom onset. Postexposure prophylaxis is wound cleaning plus immunization with killed vaccine and rabies immunoglobulin. Example of passive-active immunity.

Travels to the CNS by migrating in a retrograde fashion (via dynein motors) up nerve axons after binding to ACh receptors.

Progression of disease: fever, malaise → agitation, photophobia, hydrophobia, hypersalivation → paralysis, coma → death.

Infection more commonly from bat, raccoon, and skunk bites than from dog bites in the United States; aerosol transmission (eg, bat caves) also possible.

Ebola virus



A filovirus A that targets endothelial cells, phagocytes, hepatocytes. Following an incubation period of up to 21 days, presents with abrupt onset of flu-like symptoms, diarrhea/vomiting, high fever, myalgia. Can progress to DIC, diffuse hemorrhage, shock.

Diagnosed with RT-PCR within 48 hr of symptom onset. High mortality rate.

Transmission requires direct contact with bodily fluids, fomites (including dead bodies), infected bats or primates (apes/monkeys); high incidence of nosocomial infection.

Supportive care, no definitive treatment. Strict isolation of infected individuals and barrier practices for health care workers are key to preventing transmission.

Zika virus

A flavivirus most commonly transmitted by *Aedes* mosquito bites.

Causes conjunctivitis, low-grade pyrexia, and itchy rash in 20% of cases. Can lead to congenital microcephaly or miscarriage if transmitted in utero.

Diagnose with RT-PCR or serology.

Sexual and vertical transmission possible. Outbreaks more common in tropical and subtropical climates. Supportive care, no definitive treatment.

(c) ketabton.com: The Digital Library

Hepatitis viruses

Signs and symptoms of all hepatitis viruses: episodes of fever, jaundice, ↑ ALT and AST. Naked viruses (HAV and HEV) lack an envelope and are not destroyed by the gut: the **vowels** hit your **bowels**.

HBV DNA polymerase has DNA- and RNA-dependent activities. Upon entry into nucleus, the polymerase completes the partial dsDNA. Host RNA polymerase transcribes mRNA from viral DNA to make viral proteins. The DNA polymerase then reverse transcribes viral RNA to DNA, which is the genome of the progeny virus.

HCV lacks 3′-5′ exonuclease activity → no proofreading ability → antigenic variation of HCV envelope proteins. Host antibody production lags behind production of new mutant strains of HCV.

Virus	HAV	HBV	HCV	HDV	HEV
FAMILY	RNA picornavirus	DNA hepadnavirus	RNA flavivirus	RNA deltavirus	RNA hepevirus
TRANSMISSION	Fecal-oral (shellfish, travelers, day care)	Parenteral (Blood), sexual (Baby-making), perinatal (Birthing)	Primarily blood (IVDU, post-transfusion)	Parenteral, sexual, perinatal	Fecal-oral, especially waterborne
INCUBATION	Short (weeks)	Long (months)	Long	Superinfection (HDV after HBV) = short Coinfection (HDV with HBV) = long	Short
CLINICAL COURSE	Acute and self limiting (adults), Asymptomatic (children)	Initially like serum sickness (fever, arthralgias, rash); may progress to carcinoma	May progress to Cirrhosis or Carcinoma	Similar to HBV	Fulminant hepatitis in Expectant (pregnant) women
PROGNOSIS	Good	Adults → mostly full resolution; neonates → worse prognosis	Majority develop stable, Chronic hepatitis C	Superinfection → worse prognosis	High mortality in pregnant women
HCC RISK	No	Yes	Yes	Yes	No
LIVER BIOPSY	Hepatocyte swelling, monocyte infiltration, Councilman bodies	Granular eosinophilic "ground glass" appearance; cytotoxic T cells mediate damage	Lymphoid aggregates with focal areas of macrovesicular steatosis	Similar to HBV	Patchy necrosis
NOTES	No carrier state	Carrier state common	Carrier state very common	Defective virus, Depends on HBV HBsAg coat for entry into hepatocytes	Enteric, Epidemic (eg, in parts of Asia, Africa, Middle East), no carrier state

(c) ketabton.com: The Digital Library

Extrahepatic manifestations of hepatitis B and C

	Hepatitis B	Hepatitis C
HEMATOLOGIC	Aplastic anemia	Essential mixed cryoglobulinemia, ↑ risk B-cell NHL, ITP, autoimmune hemolytic anemia
RENAL	Membranous GN > membranoproliferative GN	Membranoproliferative GN > membranous GN
VASCULAR	Polyarteritis nodosa	Leukocytoclastic vasculitis
DERMATOLOGIC		Sporadic porphyria cutanea tarda, lichen planus
ENDOCRINE		↑ risk of diabetes mellitus, autoimmune hypothyroidism

(c) ketabton.com: The Digital Library

Hepatitis serologic markers

Anti-HAV (IgM)	IgM antibody to HAV; best test to detect acute hepatitis A.
Anti-HAV (IgG)	IgG antibody indicates prior HAV infection and/or prior vaccination; protects against reinfection.
HBsAg	Antigen found on surface of HBV; indicates hepatitis B infection.
Anti-HBs	Antibody to HBsAg; indicates immunity to hepatitis B due to vaccination or recovery from infection.
HBcAg	Antigen associated with core of HBV.
Anti-HBc	Antibody to HBcAg; IgM = acute/recent infection; IgG = prior exposure or chronic infection. IgM anti-HBc may be the sole ⊕ marker of infection during window period.
HBeAg	Secreted by infected hepatocyte into circulation. Not part of mature HBV virion. Indicates active viral replication and therefore high transmissibility and poorer prognosis.
Anti-HBe	Antibody to HBeAg; indicates low transmissibility.



	HBsAg	Anti-HBs	HBeAg	Anti-HBe	Anti-HBc
Acute HBV	✓		✓		IgM
Window				✓	IgM
Chronic HBV (high infectivity)	✓		✓		IgG
Chronic HBV (low infectivity)	✓			✓	IgG
Recovery		✓		✓	IgG
Immunized		✓			

(c) ketabton.com: The Digital Library

HIV



Diploid genome (2 molecules of RNA).

The 3 structural genes (protein coded for):

- *env* (gp120 and gp41):
  - Formed from cleavage of gp160 to form envelope glycoproteins.
  - gp120—attachment to host CD4+ T cell.
  - gp41—fusion and entry.
- *gag* (p24 and p17)—capsid and matrix proteins, respectively.
- ***pol***—**R**everse transcriptase, **I**ntegrase, **P**rotease; **RIP** "**Pol**" (Paul)

Reverse transcriptase synthesizes dsDNA from genomic RNA; dsDNA integrates into host genome.

Virus binds CD4 as well as a coreceptor, either CCR5 on macrophages (early infection) or CXCR4 on T cells (late infection).

Homozygous CCR5 mutation = immunity.

Heterozygous CCR5 mutation = slower course.

HIV diagnosis

Presumptive diagnosis made with HIV-1/2 Ag/Ab immunoassays. These immunoassays detect viral p24 Ag capsid protein and IgG Abs to HIV-1/2. Very high sensitivity/specificity.

Viral load tests determine the amount of viral RNA in the plasma. High viral load associated with poor prognosis. Also use viral load to monitor effect of drug therapy. Use HIV genotyping to determine appropriate therapy.

AIDS diagnosis: ≤ 200 CD4+ cells/mm^3 (normal: 500–1500 cells/mm^3) or HIV ⊕ with AIDS-defining condition (eg, *Pneumocystis* pneumonia).

Western blot tests are no longer recommended by the CDC for confirmatory testing.

HIV-1/2 Ag/Ab testing is not recommended in babies with suspected HIV due to maternally transferred antibody. Use HIV viral load instead.



(c) ketabton.com: The Digital Library

Time course of untreated HIV infection



Dashed lines on CD4+ count axis indicate moderate immunocompromise (< 400 CD4+ cells/mm^3) and when AIDS-defining illnesses emerge (< 200 CD4+ cells/mm^3).

Most patients who do not receive treatment eventually die of complications of HIV infection.

Four stages of untreated infection:

1. Flu-like (acute)
2. Feeling fine (latent)
3. Falling count
4. Final crisis

During clinical latency phase, virus replicates in lymph nodes

FAS1_2019_03-Microbiology.indd 176 11/14/19 12:22 PM

(c) ketabton.com: The Digital Library

Common diseases of HIV-positive adults

↓ CD4+ cell count → reactivation of past infections (eg, TB, HSV, shingles), dissemination of bacterial infections and fungal infections (eg, coccidioidomycosis), and non-Hodgkin lymphomas.

PATHOGEN	PRESENTATION	FINDINGS
CD4+ cell count < 500/mm^3		
Candida albicans	Oral thrush	Scrapable white plaque, pseudohyphae on microscopy
EBV	Oral hairy leukoplakia	Unscrapable white plaque on lateral tongue
HHV-8	Kaposi sarcoma	Biopsy with lymphocytic inflammation
HPV	Squamous cell carcinoma, commonly of anus (men who have sex with men) or cervix	
CD4+ cell count < 200/mm^3		
Histoplasma capsulatum	Fever, weight loss, fatigue, cough, dyspnea, nausea, vomiting, diarrhea	Oval yeast cells within macrophages
HIV	Dementia	Cerebral atrophy on neuroimaging
JC virus (reactivation)	Progressive multifocal leukoencephalopathy	Nonenhancing areas of demyelination on MRI
Pneumocystis jirovecii	*Pneumocystis* pneumonia	"Ground-glass" opacities on chest imaging
CD4+ cell count < 100/mm^3		
Aspergillus fumigatus	Hemoptysis, pleuritic pain	Cavitation or infiltrates on chest imaging
Bartonella spp	Bacillary angiomatosis	Biopsy with neutrophilic inflammation
Candida albicans	Esophagitis	White plaques on endoscopy; yeast and pseudohyphae on biopsy
CMV	**C**olitis, **R**etinitis, **E**sophagitis, **E**ncephalitis, **P**neumonitis (**CREEP**)	Linear ulcers on endoscopy, cotton-wool spots on fundoscopy Biopsy reveals cells with intranuclear (owl eye) inclusion bodies
Cryptococcus neoformans	Meningitis	Encapsulated yeast on India ink stain or capsular antigen ⊕
Cryptosporidium spp	Chronic, watery diarrhea	Acid-fast oocysts in stool
EBV	B-cell lymphoma (eg, non-Hodgkin lymphoma, CNS lymphoma)	CNS lymphoma—ring enhancing, may be solitary (vs *Toxoplasma*)
Mycobacterium avium–intracellulare, Mycobacterium avium complex	Nonspecific systemic symptoms (fever, night sweats, weight loss) or focal lymphadenitis	Most common if CD4+ cell count < 50/mm^3
Toxoplasma gondii	Brain abscesses	Multiple ring-enhancing lesions on MRI

(c) ketabton.com: The Digital Library

Prions

Prion diseases are caused by the conversion of a normal (predominantly α-helical) protein termed prion protein (PrP^{c}) to a β-pleated form (PrP^{sc}), which is transmissible via CNS-related tissue (iatrogenic CJD) or food contaminated by BSE-infected animal products (variant CJD). PrP^{sc} resists protease degradation and facilitates the conversion of still more PrP^{c} to PrP^{sc}. Resistant to standard sterilizing procedures, including standard autoclaving. Accumulation of PrP^{sc} results in spongiform encephalopathy and dementia, ataxia, and death.

Creutzfeldt-Jakob disease—rapidly progressive dementia, typically sporadic (some familial forms).

Bovine spongiform encephalopathy—also called "mad cow disease."

Kuru—acquired prion disease noted in tribal populations practicing human cannibalism.

▸ MICROBIOLOGY—SYSTEMS

Normal flora: dominant

Neonates delivered by C-section have no flora but are rapidly colonized after birth.

LOCATION	MICROORGANISM
Skin	*S epidermidis*
Nose	*S epidermidis*; colonized by *S aureus*
Oropharynx	Viridans group streptococci
Dental plaque	*S mutans*
Colon	*B fragilis* > *E coli*
Vagina	*Lactobacillus*; colonized by *E coli* and group B strep

Bugs causing food-borne illness

S aureus and *B cereus* food poisoning starts quickly and ends quickly.

MICROORGANISM	SOURCE OF INFECTION
B cereus	Reheated rice. "Food poisoning from reheated rice? **Be serious**!" (***B cereus***)
C botulinum	Improperly canned foods (toxins), raw honey (spores)
C perfringens	Reheated meat
E coli O157:H7	Undercooked meat
L monocytogenes	Deli meats, soft cheeses
Salmonella	Poultry, meat, and eggs
S aureus	Meats, mayonnaise, custard; preformed toxin
V parahaemolyticus and *V vulnificus*[a]	Raw/undercooked seafood

[a] *V vulnificus* can also cause wound infections from contact with contaminated water or shellfish.

(c) ketabton.com: The Digital Library

Bugs causing diarrhea

Bloody diarrhea	
Campylobacter	Comma- or S-shaped organisms; growth at 42°C
E histolytica	Protozoan; amebic dysentery; liver abscess
Enterohemorrhagic *E coli*	O157:H7; can cause HUS; makes Shiga-like toxin
Enteroinvasive *E coli*	Invades colonic mucosa
Salmonella (non-typhoidal)	Lactose ⊖; flagellar motility; has animal reservoir, especially poultry and eggs
Shigella	Lactose ⊖; very low ID_{50}; produces Shiga toxin; human reservoir only; bacillary dysentery
Y enterocolitica	Day care outbreaks; pseudoappendicitis
Watery diarrhea	
C difficile	Pseudomembranous colitis; associated with antibiotics and PPIs; occasionally bloody diarrhea
C perfringens	Also causes gas gangrene
Enterotoxigenic *E coli*	Travelers' diarrhea; produces heat-labile (LT) and heat-stable (ST) toxins
Protozoa	*Giardia*, *Cryptosporidium*
V cholerae	Comma-shaped organisms; rice-water diarrhea; often from infected seafood
Viruses	Rotavirus, norovirus, enteric adenovirus

Common causes of pneumonia

NEONATES (< 4 WK)	CHILDREN (4 WK–18 YR)	ADULTS (18–40 YR)	ADULTS (40–65 YR)	ELDERLY
Group B streptococci *E coli*	Viruses (RSV) *Mycoplasma* *C trachomatis* (infants–3 yr) *C pneumoniae* (school-aged children) *S pneumoniae* Runts May Cough Chunky Sputum	*Mycoplasma* *C pneumoniae* *S pneumoniae* Viruses (eg, influenza)	*S pneumoniae* *H influenzae* Anaerobes Viruses *Mycoplasma*	*S pneumoniae* Influenza virus Anaerobes *H influenzae* Gram ⊖ rods

Special groups	
Alcoholic	*Klebsiella*, anaerobes usually due to aspiration (eg, *Peptostreptococcus*, *Fusobacterium*, *Prevotella*, *Bacteroides*)
IV drug users	*S pneumoniae*, *S aureus*
Aspiration	Anaerobes
Atypical	*Mycoplasma*, *Chlamydophila*, *Legionella*, viruses (RSV, CMV, influenza, adenovirus)
Cystic fibrosis	*Pseudomonas*, *S aureus*, *S pneumoniae*, *Burkholderia cepacia*
Immunocompromised	*S aureus*, enteric gram ⊖ rods, fungi, viruses, *P jirovecii* (with HIV)
Nosocomial (hospital acquired)	*S aureus*, *Pseudomonas*, other enteric gram ⊖ rods
Postviral	*S pneumoniae*, *S aureus*, *H influenzae*

(c) ketabton.com: The Digital Library

Common causes of meningitis

NEWBORN (0–6 MO)	CHILDREN (6 MO–6 YR)	6–60 YR	60 YR +
Group B *Streptococcus*	*S pneumoniae*	*N meningitidis*	*S pneumoniae*
E coli	*N meningitidis*	*S pneumoniae*	*N meningitidis*
Listeria	*H influenzae* type b	Enteroviruses	*H influenzae* type b
	Group B *Streptococcus*	HSV	Group B *Streptococcus*
	Enteroviruses		*Listeria*

Give ceftriaxone and vancomycin empirically (add ampicillin if *Listeria* is suspected).

Viral causes of meningitis: enteroviruses (especially coxsackievirus), HSV-2 (HSV-1 = encephalitis), HIV, West Nile virus (also causes encephalitis), VZV.

In HIV: *Cryptococcus* spp.

Note: Incidence of Group B streptococcal meningitis in neonates has ↓ greatly due to screening and antibiotic prophylaxis in pregnancy. Incidence of *H influenzae* meningitis has ↓ greatly due to conjugate *H influenzae* vaccinations. Today, cases are usually seen in unimmunized children.

Cerebrospinal fluid findings in meningitis

	OPENING PRESSURE	CELL TYPE	PROTEIN	GLUCOSE
Bacterial	↑	↑ PMNs	↑	↓
Fungal/TB	↑	↑ lymphocytes	↑	↓
Viral	Normal/↑	↑ lymphocytes	Normal/↑	Normal

Infections causing brain abscess

Most commonly viridans streptococci and *Staphylococcus aureus*. If dental infection or extraction precedes abscess, oral anaerobes commonly involved.

Multiple abscesses are usually from bacteremia; single lesions from contiguous sites: otitis media and mastoiditis → temporal lobe and cerebellum; sinusitis or dental infection → frontal lobe.

Toxoplasma reactivation in AIDS.

Osteomyelitis

RISK FACTOR	ASSOCIATED INFECTION
Assume if no other information is available	*S aureus* (most common overall)
Sexually active	*Neisseria gonorrhoeae* (rare), septic arthritis more common
Sickle cell disease	*Salmonella* and *S aureus*
Prosthetic joint replacement	*S aureus* and *S epidermidis*
Vertebral involvement	*S aureus*, *M tuberculosis* (Pott disease)
Cat and dog bites	*Pasteurella multocida*
IV drug abuse	*S aureus*; also *Pseudomonas*, *Candida*

Elevated ESR and CRP sensitive but not specific.

Radiographs are insensitive early but can be useful in chronic osteomyelitis (A, left). MRI is best for detecting acute infection and detailing anatomic involvement (A, right).

(c) ketabton.com: The Digital Library

Urinary tract infections

Cystitis presents with dysuria, frequency, urgency, suprapubic pain, and WBCs (but not WBC casts) in urine. Primarily caused by ascension of microbes from urethra to bladder. Ascension to kidney results in pyelonephritis, which presents with fever, chills, flank pain, costovertebral angle tenderness, hematuria, and WBC casts.

Ten times more common in women (shorter urethras colonized by fecal flora).

Risk factors: obstruction (eg, kidney stones, enlarged prostate), kidney surgery, catheterization, congenital GU malformation (eg, vesicoureteral reflux), diabetes, pregnancy.

SPECIES	FEATURES	COMMENTS
Escherichia coli	Leading cause of UTI. Colonies show strong pink lactose-fermentation on MacConkey agar.	Diagnostic markers: ⊕ Leukocyte esterase = evidence of WBC activity. ⊕ Nitrite test = reduction of urinary nitrates by gram ⊖ bacterial species (eg, *E coli*). ⊕ Urease test = urease-producing bugs (eg, *S saprophyticus*, *Proteus*, *Klebsiella*).
Staphylococcus saprophyticus	2nd leading cause of UTI in sexually active women.	
Klebsiella pneumoniae	3rd leading cause of UTI. Large mucoid capsule and viscous colonies.	
Serratia marcescens	Some strains produce a red pigment; often nosocomial and drug resistant.	
Enterococcus	Often nosocomial and drug resistant.	
Proteus mirabilis	Motility causes "swarming" on agar; associated with struvite stones.	
Pseudomonas aeruginosa	Blue-green pigment and fruity odor; usually nosocomial and drug resistant.	

Common vaginal infections

	Bacterial vaginosis	*Trichomonas* vaginitis	*Candida* vulvovaginitis
SIGNS AND SYMPTOMS	No inflammation Thin, white discharge A with fishy odor	Inflammation ("strawberry cervix") Frothy, yellow-green, foul-smelling discharge	Inflammation Thick, white, "cottage cheese" discharge C
LAB FINDINGS	Clue cells pH > 4.5 ⊕ KOH whiff test	Motile pear-shaped trichomonads B pH > 4.5	Pseudohyphae pH normal (4.0–4.5)
TREATMENT	Metronidazole or clindamycin	Metronidazole Treat sexual partner(s)	Azoles







(c) ketabton.com: The Digital Library

TORCH infections

Microbes that may pass from mother to fetus. Transmission is transplacental in most cases, or via delivery (especially HSV-2). Nonspecific signs common to many ToRCHHeS infections include hepatosplenomegaly, jaundice, thrombocytopenia, and growth retardation.

Other important infectious agents include *Streptococcus agalactiae* (group B streptococci), *E coli*, and *Listeria monocytogenes*—all causes of meningitis in neonates. Parvovirus B19 causes hydrops fetalis.

AGENT	MODES OF MATERNAL TRANSMISSION	MATERNAL MANIFESTATIONS	NEONATAL MANIFESTATIONS
Toxoplasma gondii	Cat feces or ingestion of undercooked meat	Usually asymptomatic; lymphadenopathy (rarely)	Classic triad: chorioretinitis, hydrocephalus, and intracranial calcifications, +/– "blueberry muffin" rash A
Rubella	Respiratory droplets	Rash, lymphadenopathy, polyarthritis, polyarthralgia	Classic triad: abnormalities of **eye** (cataracts B) and **ear** (deafness) and congenital **heart** disease (PDA); +/– "blueberry muffin" rash. "**I** (eye) ♥ **ru**by (**rub**ella) **ear**rings"
Cytomegalovirus	Sexual contact, organ transplants	Usually asymptomatic; mononucleosis-like illness	Hearing loss, seizures, petechial rash, "blueberry muffin" rash, chorioretinitis, periventricular calcifications C
HIV	Sexual contact, needlestick	Variable presentation depending on CD4+ cell count	Recurrent infections, chronic diarrhea
Herpes simplex virus-2	Skin or mucous membrane contact	Usually asymptomatic; herpetic (vesicular) lesions	Meningoencephalitis, herpetic (vesicular) lesions
Syphilis	Sexual contact	Chancre (1°) and disseminated rash (2°) are the two stages likely to result in fetal infection	Often results in stillbirth, hydrops fetalis; if child survives, presents with facial abnormalities (eg, notched teeth, saddle nose, short maxilla), saber shins, CN VIII deafness

FAS1_2019_03-Microbiology.indd 182 11/14/19 12:22 PM

(c) ketabton.com: The Digital Library

Red rashes of childhood

AGENT	ASSOCIATED SYNDROME/DISEASE	CLINICAL PRESENTATION
Coxsackievirus type A	Hand-foot-mouth disease	Oval-shaped vesicles on palms and soles A; vesicles and ulcers in oral mucosa (herpangina)
Human herpesvirus 6	Roseola (exanthem subitum)	Asymptomatic rose-colored macules appear on body after several days of high fever; can present with febrile seizures; usually affects infants
Measles virus	Measles (rubeola)	Confluent rash beginning at head and moving down; preceded by cough, coryza, conjunctivitis, and blue-white (Koplik) spots on buccal mucosa
Parvovirus B19	Erythema infectiosum (fifth disease)	"Slapped cheek" rash on face B (can cause hydrops fetalis in pregnant women)
Rubella virus	Rubella	Pink macules and papules begin at head and move down, remain discrete → fine desquamating truncal rash; postauricular lymphadenopathy
Streptococcus pyogenes	Scarlet fever	Flushed cheeks and circumoral pallor C on face; erythematous, sandpaper-like rash from neck to trunk and extremities; fever, sore throat, strawberry tongue
Varicella-Zoster virus	Chickenpox	Vesicular rash begins on trunk; spreads to face D and extremities with lesions of different stages

(c) ketabton.com: The Digital Library

Sexually transmitted infections

DISEASE	CLINICAL FEATURES	ORGANISM
AIDS	Opportunistic infections, Kaposi sarcoma, lymphoma	HIV
Chancroid	Painful genital ulcer with exudate, inguinal adenopathy A	*Haemophilus ducreyi* (it's so painful, you "do cry")
Chlamydia	Urethritis, cervicitis, epididymitis, conjunctivitis, reactive arthritis, PID	*Chlamydia trachomatis* (D–K)
Condylomata acuminata	Genital warts, koilocytes	HPV-6 and -11
Genital herpes	Painful penile, vulvar, or cervical vesicles and ulcers; can cause systemic symptoms such as fever, headache, myalgia	HSV-2, less commonly HSV-1
Gonorrhea	Urethritis, cervicitis, PID, prostatitis, epididymitis, arthritis, creamy purulent discharge	*Neisseria gonorrhoeae*
Granuloma inguinale (Donovanosis)	Painless, beefy red ulcer that bleeds readily on contact B Uncommon in US	*Klebsiella (Calymmatobacterium) granulomatis*; cytoplasmic Donovan bodies (bipolar staining) seen on microscopy
Hepatitis B	Jaundice	HBV
Lymphogranuloma venereum	Infection of lymphatics; painless genital ulcers, painful lymphadenopathy (ie, buboes)	*C trachomatis* (L1–L3)
Primary syphilis	Painless chancre	*Treponema pallidum*
Secondary syphilis	Fever, lymphadenopathy, skin rashes, condylomata lata	
Tertiary syphilis	Gummas, tabes dorsalis, general paresis, aortitis, Argyll Robertson pupil	
Trichomoniasis	Vaginitis, strawberry cervix, motile in wet prep	*Trichomonas vaginalis*

(c) ketabton.com: The Digital Library

Pelvic inflammatory disease





Top bugs—*Chlamydia trachomatis* (subacute, often undiagnosed), *Neisseria gonorrhoeae* (acute).

C trachomatis—most common bacterial STI in the United States.

Signs include cervical motion tenderness, adnexal tenderness, purulent cervical discharge A.

PID may include salpingitis, endometritis, hydrosalpinx, and tubo-ovarian abscess.

Salpingitis is a risk factor for ectopic pregnancy, infertility, chronic pelvic pain, and adhesions.

Can lead to perihepatitis (**Fitz-Hugh–Curtis syndrome**)—infection and inflammation of liver capsule and "violin string" adhesions of peritoneum to liver B.

Nosocomial infections

E coli (UTI) and *S aureus* (wound infection) are the two most common causes.

RISK FACTOR	PATHOGEN	UNIQUE SIGNS/SYMPTOMS
Antibiotic use	*Clostridium difficile*	Watery diarrhea, leukocytosis
Aspiration (2° to altered mental status, old age)	Polymicrobial, gram ⊖ bacteria, often anaerobes	Right lower lobe infiltrate or right upper/middle lobe (patient recumbent); purulent malodorous sputum
Decubitus ulcers, surgical wounds, drains	*S aureus* (including MRSA), gram ⊖ anaerobes *(Bacteroides, Prevotella, Fusobacterium)*	Erythema, tenderness, induration, drainage from surgical wound sites
Intravascular catheters	*S aureus* (including MRSA), *S epidermidis* (long term), *Enterobacter*	Erythema, induration, tenderness, drainage from access sites
Mechanical ventilation, endotracheal intubation	Late onset: *P aeruginosa, Klebsiella, Acinetobacter, S aureus*	New infiltrate on CXR, ↑ sputum production; sweet odor *(Pseudomonas)*
Renal dialysis unit, needlestick	HBV, HCV	
Urinary catheterization	***P**roteus* spp, ***E** **c**oli*, ***K**lebsiella* (infections in your **PEcK**er)	Dysuria, leukocytosis, flank pain or costovertebral angle tenderness
Water aerosols	*Legionella*	Signs of pneumonia, GI symptoms (diarrhea, nausea, vomiting), neurologic abnormalities

(c) ketabton.com: The Digital Library

Bugs affecting unvaccinated children

CLINICAL PRESENTATION	FINDINGS/LABS	PATHOGEN
Dermatologic		
Rash	Beginning at head and moving down with postauricular lymphadenopathy	Rubella virus
	Beginning at head and moving down; preceded by cough, coryza, conjunctivitis, and Koplik spots	Measles virus
Neurologic		
Meningitis	Microbe colonizes nasopharynx	*H influenzae* type b
	Can also lead to myalgia and paralysis	Poliovirus
Tetanus	Muscle spasms and spastic paralysis (eg, lockjaw, opisthotonus)	*Clostridium tetani*
Respiratory		
Epiglottitis	Fever with dysphagia, drooling, and difficulty breathing due to edema	*H influenzae* type b (also capable of causing epiglottitis in fully immunized children)
Pertussis	Low-grade fevers, coryza → whooping cough, post-tussive vomiting → gradual recovery	*Bordetella pertussis*
Pharyngitis	Grayish pseudomembranes (may obstruct airways)	*Corynebacterium diphtheriae*

Bug hints

CHARACTERISTIC	ORGANISM
Asplenic patients	Encapsulated microbes, especially **SHiN** (***S*** *pneumoniae* >> ***H*** *influenzae* type b > ***N*** *meningitidis*)
Branching rods in oral infection, sulfur granules	*Actinomyces israelii*
Chronic granulomatous disease	Catalase ⊕ microbes, especially *S aureus*
"Currant jelly" sputum	*Klebsiella*
Dog or cat bite	*Pasteurella multocida*
Facial nerve palsy (typically bilateral)	*Borrelia burgdorferi* (Lyme disease)
Human bite	Human oral flora (eg, *Eikenella*, *Fusobacterium*)
Neutropenic patients	*Candida albicans* (systemic), *Aspergillus*
Organ transplant recipient	CMV
PAS ⊕	*Tropheryma whipplei* (Whipple disease)
Pediatric infection	*Haemophilus influenzae* (including epiglottitis)
Pneumonia in cystic fibrosis, burn infection	*Pseudomonas aeruginosa*
Puncture wound, lockjaw	*Clostridium tetani*
Pus, empyema, abscess	*S aureus*
Rash on hands and feet	**C**oxsackie **A**, ***R*** *rickettsii*, **S**yphilis (**CARS**)
Sepsis/meningitis in newborn	Group B strep
Sinus/CNS infection in diabetics	*Mucor* or *Rhizopus* spp.
Surgical wound	*S aureus*
Traumatic open wound	*Clostridium perfringens*

(c) ketabton.com: The Digital Library

▸ MICROBIOLOGY—ANTIMICROBIALS

Antimicrobial therapy



Penicillin G, V	Penicillin G (IV and IM form), penicillin V (oral). Prototype β-lactam antibiotics.
MECHANISM	D-Ala-D-Ala structural analog. Bind penicillin-binding proteins (transpeptidases). Block transpeptidase cross-linking of peptidoglycan in cell wall. Activate autolytic enzymes.
CLINICAL USE	Mostly used for gram ⊕ organisms (*S pneumoniae*, *S pyogenes*, *Actinomyces*). Also used for gram ⊖ cocci (mainly *N meningitidis*) and spirochetes (mainly *T pallidum*). Bactericidal for gram ⊕ cocci, gram ⊕ rods, gram ⊖ cocci, and spirochetes. β-lactamase sensitive.
ADVERSE EFFECTS	Hypersensitivity reactions, direct Coombs ⊕ hemolytic anemia, drug-induced interstitial nephritis.
RESISTANCE	β-lactamase cleaves the β-lactam ring. Mutations in PBPs.

FAS1_2019_03-Microbiology.indd 187 11/14/19 12:22 PM

(c) ketabton.com: The Digital Library

Penicillinase-sensitive penicillins	Amoxicillin, ampicillin; aminopenicillins.	
MECHANISM	Same as penicillin. Wider spectrum; penicillinase sensitive. Also combine with clavulanic acid to protect against destruction by β-lactamase.	**AM**ino**P**enicillins are **AMP**ed-up penicillin. Am**O**xicillin has greater **O**ral bioavailability than ampicillin.
CLINICAL USE	Extended-spectrum penicillin—***H*** *influenzae*, ***H*** *pylori*, ***E*** *coli*, ***L****isteria monocytogenes*, ***P****roteus mirabilis*, ***S****almonella*, ***S****higella*, enterococci.	Coverage: ampicillin/amoxicillin **HHELPSS** kill enterococci.
ADVERSE EFFECTS	Hypersensitivity reactions, rash, pseudomembranous colitis.	
MECHANISM OF RESISTANCE	Penicillinase (a type of β-lactamase) cleaves β-lactam ring.	

Penicillinase-resistant penicillins	Dicloxacillin, nafcillin, oxacillin.	
MECHANISM	Same as penicillin. Narrow spectrum; penicillinase resistant because bulky R group blocks access of β-lactamase to β-lactam ring.	
CLINICAL USE	*S aureus* (except MRSA).	"Use **naf** (nafcillin) for **staph**."
ADVERSE EFFECTS	Hypersensitivity reactions, interstitial nephritis.	
MECHANISM OF RESISTANCE	MRSA has altered penicillin-binding protein target site.	

Antipseudomonal penicillins	Piperacillin, ticarcillin.
MECHANISM	Same as penicillin. Extended spectrum. Penicillinase sensitive; use with β-lactamase inhibitors.
CLINICAL USE	*Pseudomonas* spp. and gram ⊖ rods.
ADVERSE EFFECTS	Hypersensitivity reactions.

(c) ketabton.com: The Digital Library

Cephalosporins

MECHANISM	β-lactam drugs that inhibit cell wall synthesis but are less susceptible to penicillinases. Bactericidal.	Organisms typically not covered by 1st–4th generation cephalosporins are **LAME**: ***L****isteria*, **A**typicals (*Chlamydia*, *Mycoplasma*), **M**RSA, and **E**nterococci.
CLINICAL USE	1st generation (cefazolin, cephalexin)—gram ⊕ cocci, ***P****roteus mirabilis*, ***E c****oli*, ***K****lebsiella pneumoniae*. Cefazolin used prior to surgery to prevent *S aureus* wound infections.	1st generation—⊕ **PEcK**.
	2nd generation (ce**fa**clor, ce**fox**itin, ce**fur**oxime, cefo**te**tan)—gram ⊕ cocci, ***H*** *influenzae*, ***E****nterobacter aerogenes*, ***N****eisseria* spp., ***S****erratia marcescens*, ***P****roteus mirabilis*, ***E c****oli*, ***K****lebsiella pneumoniae*.	**2nd** graders wear **fa**ke **fox fur** to **te**a parties. 2nd generation—⊕ **HENS PEcK**.
	3rd generation (ceftriaxone, cefotaxime, cefpodoxime, ceftazidime)—serious gram ⊖ infections resistant to other β-lactams.	Can cross blood-brain barrier. Ceftriaxone—meningitis, gonorrhea, disseminated Lyme disease. Ceftazidime—*Pseudomonas*.
	4th generation (cefepime)—gram ⊖ organisms, with ↑ activity against *Pseudomonas* and gram ⊕ organisms.	
	5th generation (ceftaroline)—broad gram ⊕ and gram ⊖ organism coverage; unlike 1st–4th generation cephalosporins, ceftaroline covers MRSA, and *Enterococcus faecalis*—does not cover *Pseudomonas*.	
ADVERSE EFFECTS	Hypersensitivity reactions, autoimmune hemolytic anemia, disulfiram-like reaction, vitamin K deficiency. Low rate of cross-reactivity even in penicillin-allergic patients. ↑ nephrotoxicity of aminoglycosides.	
MECHANISM OF RESISTANCE	Inactivated by cephalosporinases (a type of β-lactamase). Structural change in penicillin-binding proteins (transpeptidases).	

β-lactamase inhibitors	Include **C**lavulanic acid, **A**vibactam, **S**ulbactam, **T**azobactam. Often added to penicillin antibiotics to protect the antibiotic from destruction by β-lactamase.	**CAST** (eg, amoxicillin-clavulanate, ceftazidime-avibactam, ampicillin-sulbactam, piperacillin-tazobactam).

(c) ketabton.com: The Digital Library

Carbapenems

Doripenem, **I**mipenem, **M**eropenem, **E**rtapenem (**DIME** antibiotics are given when there is a 10/10 [life-threatening] infection).

MECHANISM

Imipenem is a broad-spectrum, β-lactamase–resistant carbapenem. Always administered with cilastatin (inhibitor of renal dehydropeptidase I) to ↓ inactivation of drug in renal tubules.

With imipenem, "the kill is **lastin'** with ci**lastatin**."

Newer carbapenems include ertapenem (limited *Pseudomonas* coverage) and doripenem.

CLINICAL USE

Gram ⊕ cocci, gram ⊖ rods, and anaerobes. Wide spectrum and significant side effects limit use to life-threatening infections or after other drugs have failed. Meropenem has a ↓ risk of seizures and is stable to dehydropeptidase I.

ADVERSE EFFECTS

GI distress, rash, and CNS toxicity (seizures) at high plasma levels.

MECHANISM OF RESISTANCE

Inactivated by carbapenemases produced by, eg, *K pneumoniae*, *E coli*, *E aerogenes*.

Monobactams

Aztreonam

MECHANISM

Less susceptible to β-lactamases. Prevents peptidoglycan cross-linking by binding to penicillin-binding protein 3. Synergistic with aminoglycosides. No cross-allergenicity with penicillins.

CLINICAL USE

Gram ⊖ rods only—no activity against gram ⊕ rods or anaerobes. For penicillin-allergic patients and those with renal insufficiency who cannot tolerate aminoglycosides.

ADVERSE EFFECTS

Usually nontoxic; occasional GI upset.

Vancomycin

MECHANISM

Inhibits cell wall peptidoglycan formation by binding D-Ala-D-Ala portion of cell wall precursors. Bactericidal against most bacteria (bacteriostatic against *C difficile*). Not susceptible to β-lactamases.

CLINICAL USE

Gram ⊕ bugs only—for serious, multidrug-resistant organisms, including MRSA, *S epidermidis*, sensitive *Enterococcus* species, and *Clostridium difficile* (oral dose for pseudomembranous colitis).

ADVERSE EFFECTS

Well tolerated in general but **NOT** trouble **F**ree.

Nephrotoxicity, **O**totoxicity, **T**hrombophlebitis, diffuse **F**lushing (red man syndrome **A** idiopathic reaction largely preventable by pretreatment with antihistamines), DRESS syndrome.



MECHANISM OF RESISTANCE

Occurs in bacteria (eg, *Enterococcus*) via amino acid modification of D-Ala-D-Ala to **D-Ala**-D-**Lac**. "If you **Lac**k a **D-Ala** (dollar), you can't ride the **van** (**van**comycin)."

(c) ketabton.com: The Digital Library

Protein synthesis inhibitors



Specifically target smaller bacterial ribosome (70S, made of 30S and 50S subunits), leaving human ribosome (80S) unaffected.

All are bacteriostatic, except aminoglycosides (bactericidal) and linezolid (variable).

30S inhibitors

Aminoglycosides
Tetracyclines

50S inhibitors

Chloramphenicol, Clindamycin
Erythromycin (macrolides)
Linezolid

"Buy AT 30, CCEL (sell) at 50."

Aminoglycosides	Gentamicin, Neomycin, Amikacin, Tobramycin, Streptomycin.	"Mean" (aminoglycoside) GNATS caNNOT kill anaerobes.
MECHANISM	Bactericidal; irreversible inhibition of initiation complex through binding of the 30S subunit. Can cause misreading of mRNA. Also block translocation. Require O_2 for uptake; therefore ineffective against anaerobes.	
CLINICAL USE	Severe gram ⊖ rod infections. Synergistic with β-lactam antibiotics. Neomycin for bowel surgery.	
ADVERSE EFFECTS	Nephrotoxicity, Neuromuscular blockade (absolute contraindication with myasthenia gravis), Ototoxicity (especially with loop diuretics), Teratogenicity.	
MECHANISM OF RESISTANCE	Bacterial transferase enzymes inactivate the drug by acetylation, phosphorylation, or adenylation.	

(c) ketabton.com: The Digital Library

Tetracyclines	Tetracycline, doxycycline, minocycline.
MECHANISM	Bacteriostatic; bind to 30S and prevent attachment of aminoacyl-tRNA. Limited CNS penetration. Doxycycline is fecally eliminated and can be used in patients with renal failure. Do not take tetracyclines with milk (Ca^{2+}), antacids (eg, Ca^{2+} or Mg^{2+}), or iron-containing preparations because divalent cations inhibit drugs' absorption in the gut.
CLINICAL USE	*Borrelia burgdorferi*, *M pneumoniae*. Drugs' ability to accumulate intracellularly makes them very effective against *Rickettsia* and *Chlamydia*. Also used to treat acne. Doxycycline effective against community-acquired MRSA.
ADVERSE EFFECTS	GI distress, discoloration of teeth and inhibition of bone growth in children, photosensitivity. Contraindicated in pregnancy.
MECHANISM OF RESISTANCE	↓ uptake or ↑ efflux out of bacterial cells by plasmid-encoded transport pumps.

Tigecycline	
MECHANISM	Tetracycline derivative. Binds to 30S, inhibiting protein synthesis. Generally bacteriostatic.
CLINICAL USE	Broad-spectrum anaerobic, gram ⊖, and gram ⊕ coverage. Multidrug-resistant organisms (MRSA, VRE) or infections requiring deep tissue penetration.
ADVERSE EFFECTS	GI symptoms: nausea, vomiting.

Chloramphenicol	
MECHANISM	Blocks peptidyltransferase at 50S ribosomal subunit. Bacteriostatic.
CLINICAL USE	Meningitis (*Haemophilus influenzae*, *Neisseria meningitidis*, *Streptococcus pneumoniae*) and rickettsial diseases (eg, Rocky Mountain spotted fever *[Rickettsia rickettsii]*). Limited use due to toxicity but often still used in developing countries because of low cost.
ADVERSE EFFECTS	Anemia (dose dependent), aplastic anemia (dose independent), gray baby syndrome (in premature infants because they lack liver UDP-glucuronosyltransferase).
MECHANISM OF RESISTANCE	Plasmid-encoded acetyltransferase inactivates the drug.

Clindamycin		
MECHANISM	Blocks peptide transfer (translocation) at 50S ribosomal subunit. Bacteriostatic.	
CLINICAL USE	Anaerobic infections (eg, *Bacteroides* spp., *Clostridium perfringens*) in aspiration pneumonia, lung abscesses, and oral infections. Also effective against invasive group A streptococcal infection.	Treats anaerobic infections **above** the diaphragm vs metronidazole (anaerobic infections **below** diaphragm).
ADVERSE EFFECTS	Pseudomembranous colitis (*C difficile* overgrowth), fever, diarrhea.	

(c) ketabton.com: The Digital Library

Linezolid

MECHANISM	Inhibits protein synthesis by binding to 50S subunit and preventing formation of the initiation complex.
CLINICAL USE	Gram ⊕ species including MRSA and VRE.
ADVERSE EFFECTS	Bone marrow suppression (especially thrombocytopenia), peripheral neuropathy, serotonin syndrome (due to partial MAO inhibition).
MECHANISM OF RESISTANCE	Point mutation of ribosomal RNA.

Macrolides

Azithromycin, clarithromycin, erythromycin.

MECHANISM	Inhibit protein synthesis by blocking translocation ("macro**slides**"); bind to the 23S rRNA of the 50S ribosomal subunit. Bacteriostatic.
CLINICAL USE	Atypical pneumonias (*Mycoplasma*, *Chlamydia*, *Legionella*), STIs (*Chlamydia*), gram ⊕ cocci (streptococcal infections in patients allergic to penicillin), and *B pertussis*.
ADVERSE EFFECTS	**MACRO**: Gastrointestinal **M**otility issues, **A**rrhythmia caused by prolonged QT interval, acute **C**holestatic hepatitis, **R**ash, e**O**sinophilia. Increases serum concentration of theophylline, oral anticoagulants. Clarithromycin and erythromycin inhibit cytochrome P-450.
MECHANISM OF RESISTANCE	Methylation of 23S rRNA-binding site prevents binding of drug.

Polymyxins

Colistin (polymyxin E), polymyxin B.

MECHANISM	Cation polypeptides that bind to phospholipids on cell membrane of gram ⊖ bacteria. Disrupt cell membrane integrity → leakage of cellular components → cell death.
CLINICAL USE	Salvage therapy for multidrug-resistant gram ⊖ bacteria (eg, *P aeruginosa*, *E coli*, *K pneumoniae*). Polymyxin B is a component of a triple antibiotic ointment used for superficial skin infections.
ADVERSE EFFECTS	Nephrotoxicity, neurotoxicity (eg, slurred speech, weakness, paresthesias), respiratory failure.

(c) ketabton.com: The Digital Library

Sulfonamides

Sulfamethoxazole (SMX), sulfisoxazole, sulfadiazine.

MECHANISM	Inhibit dihydropteroate synthase, thus inhibiting folate synthesis. Bacteriostatic (bactericidal when combined with trimethoprim).
CLINICAL USE	Gram ⊕, gram ⊖, *Nocardia*. TMP-SMX for simple UTI.
ADVERSE EFFECTS	Hypersensitivity reactions, hemolysis if G6PD deficient, nephrotoxicity (tubulointerstitial nephritis), photosensitivity, Stevens-Johnson syndrome, kernicterus in infants, displace other drugs from albumin (eg, warfarin).
MECHANISM OF RESISTANCE	Altered enzyme (bacterial dihydropteroate synthase), ↓ uptake, or ↑ PABA synthesis.

Dapsone

MECHANISM	Similar to sulfonamides, but structurally distinct agent.
CLINICAL USE	Leprosy (lepromatous and tuberculoid), *Pneumocystis jirovecii* prophylaxis, or treatment when used in combination with TMP.
ADVERSE EFFECTS	Hemolysis if G6PD deficient, methemoglobinemia, agranulocytosis.

Trimethoprim

MECHANISM	Inhibits bacterial dihydrofolate reductase. Bacteriostatic.
CLINICAL USE	Used in combination with sulfonamides (trimethoprim-sulfamethoxazole [TMP-SMX]), causing sequential block of folate synthesis. Combination used for UTIs, *Shigella*, *Salmonella*, *Pneumocystis jirovecii* pneumonia treatment and prophylaxis, toxoplasmosis prophylaxis.
ADVERSE EFFECTS	Hyperkalemia (high doses), megaloblastic anemia, leukopenia, granulocytopenia, which may be avoided with coadministration of leucovorin (folinic acid). **TMP T**reats **M**arrow **P**oorly.



(c) ketabton.com: The Digital Library

Fluoroquinolones	Ciprofloxacin, enoxacin, norfloxacin, ofloxacin; respiratory fluoroquinolones—gemifloxacin, levofloxacin, moxifloxacin.	
MECHANISM	Inhibit prokaryotic enzymes topoisomerase II (DNA gyrase) and topoisomerase IV. Bactericidal. Must not be taken with antacids.	
CLINICAL USE	Gram ⊖ rods of urinary and GI tracts (including *Pseudomonas*), some gram ⊕ organisms, otitis externa.	
ADVERSE EFFECTS	GI upset, superinfections, skin rashes, headache, dizziness. Less commonly, can cause leg cramps and myalgias. Contraindicated in pregnant women, nursing mothers, and children < 18 years old due to possible damage to cartilage. Some may prolong QT interval. May cause tendonitis or tendon rupture in people > 60 years old and in patients taking prednisone. Ciprofloxacin inhibits cytochrome P-450.	Fluoroquino**lones** hurt attachments to your **bones**.
MECHANISM OF RESISTANCE	Chromosome-encoded mutation in DNA gyrase, plasmid-mediated resistance, efflux pumps.	

Daptomycin		
MECHANISM	Lipopeptide that disrupts cell membranes of gram ⊕ cocci by creating transmembrane channels.	
CLINICAL USE	*S aureus* skin infections (especially MRSA), bacteremia, endocarditis, VRE.	Not used for pneumonia (avidly binds to and is inactivated by surfactant). "Daptomy**skin**" is used for **skin** infections.
ADVERSE EFFECTS	Myopathy, rhabdomyolysis.	

Metronidazole		
MECHANISM	Forms toxic free radical metabolites in the bacterial cell that damage DNA. Bactericidal, antiprotozoal.	
CLINICAL USE	Treats *G*iardia, *E*ntamoeba, *T*richomonas, *G*ardnerella vaginalis, *A*naerobes (*Bacteroides*, *C difficile*). Can be used in place of amoxicillin in *H pylori* "triple therapy" in case of penicillin allergy.	**GET GAP** on the **Metro** with **metro**nidazole! Treats anaerobic infection **below** the diaphragm vs clindamycin (anaerobic infections **above** diaphragm).
ADVERSE EFFECTS	Disulfiram-like reaction (severe flushing, tachycardia, hypotension) with alcohol; headache, metallic taste.	

(c) ketabton.com: The Digital Library

Antimycobacterial therapy

BACTERIUM	PROPHYLAXIS	TREATMENT
M tuberculosis	Isoniazid	**R**ifampin, **I**soniazid, **P**yrazinamide, **E**thambutol (**RIPE** for treatment)
M avium–intracellulare	Azithromycin, rifabutin	Azithromycin or clarithromycin + ethambutol Can add rifabutin or ciprofloxacin
M leprae	N/A	Long-term treatment with dapsone and rifampin for tuberculoid form Add clofazimine for lepromatous form



Rifamycins

Rifampin, rifabutin.

MECHANISM	Inhibit DNA-dependent RNA polymerase.
CLINICAL USE	*Mycobacterium tuberculosis*; delay resistance to dapsone when used for leprosy. Used for meningococcal prophylaxis and chemoprophylaxis in contacts of children with *H influenzae* type b.
ADVERSE EFFECTS	Minor hepatotoxicity and drug interactions (↑ cytochrome P-450); orange body fluids (nonhazardous side effect). Rifabutin favored over rifampin in patients with HIV infection due to less cytochrome P-450 stimulation.
MECHANISM OF RESISTANCE	Mutations reduce drug binding to RNA polymerase. Monotherapy rapidly leads to resistance.

Rifampin's **4 R**'s:
RNA polymerase inhibitor
Ramps up microsomal cytochrome P-450
Red/orange body fluids
Rapid resistance if used alone

Rifam**p**in **ramp**s up cytochrome P-450, **but** rifa**but**in does not.

FAS1_2019_03-Microbiology.indd 196 11/14/19 12:22 PM

(c) ketabton.com: The Digital Library

Isoniazid

MECHANISM	↓ synthesis of mycolic acids. Bacterial catalase-peroxidase (encoded by KatG) needed to convert INH to active metabolite.	
CLINICAL USE	*Mycobacterium tuberculosis*. The only agent used as solo prophylaxis against TB. Also used as monotherapy for latent TB.	Different INH half-lives in fast vs slow acetylators.
ADVERSE EFFECTS	Hepatotoxicity, cytochrome P-450 inhibition, drug-induced SLE, anion gap metabolic acidosis, vitamin B_6 deficiency (peripheral neuropathy, sideroblastic anemia), seizures (in high doses, refractory to benzodiazepines). Administer with pyridoxine (B_6).	**INH** **I**njures **N**eurons and **H**epatocytes.
MECHANISM OF RESISTANCE	Mutations leading to underexpression of KatG.	

Pyrazinamide

MECHANISM	Mechanism uncertain. Pyrazinamide is a prodrug that is converted to the active compound pyrazinoic acid. Works best at acidic pH (eg, in host phagolysosomes).
CLINICAL USE	*Mycobacterium tuberculosis*.
ADVERSE EFFECTS	Hyperuricemia, hepatotoxicity.

Ethambutol

MECHANISM	↓ carbohydrate polymerization of mycobacterium cell wall by blocking arabinosyltransferase.
CLINICAL USE	*Mycobacterium tuberculosis*.
ADVERSE EFFECTS	**Optic** neuropathy (red-green color blindness, usually reversible). Pronounce "**eye**thambutol."

Streptomycin

MECHANISM	Interferes with 30S component of ribosome.
CLINICAL USE	*Mycobacterium tuberculosis* (2nd line).
ADVERSE EFFECTS	Tinnitus, vertigo, ataxia, nephrotoxicity.

(c) ketabton.com: The Digital Library

Antimicrobial prophylaxis

CLINICAL SCENARIO	MEDICATION
Exposure to meningococcal infection	Ceftriaxone, ciprofloxacin, or rifampin
High risk for endocarditis and undergoing surgical or dental procedures	Amoxicillin
History of recurrent UTIs	TMP-SMX
Malaria prophylaxis for travelers	Atovaquone-proguanil, mefloquine, doxycycline, primaquine, or chloroquine (for areas with sensitive species)
Pregnant woman carrying group B strep	Intrapartum penicillin G or ampicillin
Prevention of gonococcal conjunctivitis in newborn	Erythromycin ointment on eyes
Prevention of postsurgical infection due to *S aureus*	Cefazolin
Prophylaxis of strep pharyngitis in child with prior rheumatic fever	Benzathine penicillin G or oral penicillin V

Prophylaxis in HIV/AIDS patients

CELL COUNT	PROPHYLAXIS	INFECTION
CD4 < 200 cells/mm^3	TMP-SMX	*Pneumocystis* pneumonia
CD4 < 100 cells/mm^3	TMP-SMX	*Pneumocystis* pneumonia and toxoplasmosis
CD4 < 50 cells/mm^3	Azithromycin or clarithromycin	*Mycobacterium avium* complex

Treatment of highly resistant bacteria

MRSA: vancomycin, daptomycin, linezolid, tigecycline, ceftaroline, doxycycline.
VRE: linezolid, tigecycline, and streptogramins (quinupristin, dalfopristin).
Multidrug-resistant *P aeruginosa*, multidrug-resistant *Acinetobacter baumannii*: polymyxins B and E (colistin).

Antifungal therapy



(c) ketabton.com: The Digital Library

Amphotericin B

MECHANISM	Binds ergosterol (unique to fungi); forms membrane pores that allow leakage of electrolytes.
CLINICAL USE	Serious, systemic mycoses. *Cryptococcus* (amphotericin B with +/– without flucytosine for cryptococcal meningitis), *Blastomyces*, *Coccidioides*, *Histoplasma*, *Candida*, *Mucor*. Intrathecally for coccidioidal meningitis.
ADVERSE EFFECTS	Fever/chills ("shake and bake"), hypotension, nephrotoxicity, arrhythmias, anemia, IV phlebitis ("**amphoterri**ble"). Hydration ↓ nephrotoxicity. Liposomal amphotericin ↓ toxicity.

Amphoter**ic**in "**tear**s" holes in the fungal membrane by forming pores.

Supplement K^+ and Mg^{2+} because of altered renal tubule permeability.

Nystatin

MECHANISM	Same as amphotericin B. Topical use only as too toxic for systemic use.
CLINICAL USE	"Swish and swallow" for oral candidiasis (thrush); topical for diaper rash or vaginal candidiasis.

Flucytosine

MECHANISM	Inhibits DNA and RNA biosynthesis by conversion to 5-fluorouracil by cytosine deaminase.
CLINICAL USE	Systemic fungal infections (especially meningitis caused by *Cryptococcus*) in combination with amphotericin B.
ADVERSE EFFECTS	Bone marrow suppression.

Azoles

Clotrimazole, fluconazole, isavuconazole, itraconazole, ketoconazole, miconazole, voriconazole.

MECHANISM	Inhibit fungal sterol (ergosterol) synthesis by inhibiting the cytochrome P-450 enzyme that converts lanosterol to ergosterol.
CLINICAL USE	Local and less serious systemic mycoses. Fluconazole for chronic suppression of cryptococcal meningitis in AIDS patients and candidal infections of all types. Itraconazole may be used for *Blastomyces*, *Coccidioides*, *Histoplasma*, *Sporothrix schenckii*. Clotrimazole and miconazole for topical fungal infections. Voriconazole for *Aspergillus* and some *Candida*. Isavuconazole for serious *Aspergillus* and *Mucor* infections.
ADVERSE EFFECTS	Testosterone synthesis inhibition (gynecomastia, especially with ketoconazole), liver dysfunction (inhibits cytochrome P-450).

Terbinafine

MECHANISM	Inhibits the fungal enzyme squalene epoxidase.
CLINICAL USE	Dermatophytoses (especially onychomycosis—fungal infection of finger or toe nails).
ADVERSE EFFECTS	GI upset, headaches, hepatotoxicity, taste disturbance.

(c) ketabton.com: The Digital Library

Echinocandins	Anidulafungin, caspofungin, micafungin.
MECHANISM	Inhibit cell wall synthesis by inhibiting synthesis of β-glucan.
CLINICAL USE	Invasive aspergillosis, *Candida*.
ADVERSE EFFECTS	GI upset, flushing (by histamine release).

Griseofulvin	
MECHANISM	Interferes with microtubule function; disrupts mitosis. Deposits in keratin-containing tissues (eg, nails).
CLINICAL USE	Oral treatment of superficial infections; inhibits growth of dermatophytes (tinea, ringworm).
ADVERSE EFFECTS	Teratogenic, carcinogenic, confusion, headaches, disulfiram-like reaction, ↑ cytochrome P-450 and warfarin metabolism.

Antiprotozoal therapy	Pyrimethamine (toxoplasmosis), suramin and melarsoprol (*Trypanosoma brucei*), nifurtimox (*T cruzi*), sodium stibogluconate (leishmaniasis).

Anti-mite/louse therapy	Permethrin (inhibits Na^+ channel deactivation → neuronal membrane depolarization), malathion (acetylcholinesterase inhibitor), topical +/– oral ivermectin. Used to treat scabies (*Sarcoptes scabiei*) and lice (*Pediculus* and *Pthirus*).

Chloroquine	
MECHANISM	Blocks detoxification of heme into hemozoin. Heme accumulates and is toxic to plasmodia.
CLINICAL USE	Treatment of plasmodial species other than *P falciparum* (frequency of resistance in *P falciparum* is too high). Resistance due to membrane pump that ↓ intracellular concentration of drug. Treat *P falciparum* with artemether/lumefantrine or atovaquone/proguanil. For life-threatening malaria, use quinidine in US (quinine elsewhere) or artesunate.
ADVERSE EFFECTS	Retinopathy; pruritus (especially in dark-skinned individuals).

Antihelminthic therapy	**P**yrantel pamoate, **I**vermectin, **M**ebendazole (microtubule inhibitor), **P**raziquantel (↑ Ca^{2+} permeability, ↑ vacuolization), **D**iethylcarbamazine. Helminths get **PIMP'D**.

(c) ketabton.com: The Digital Library

Antiviral therapy



Oseltamivir, zanamivir

MECHANISM	Inhibit influenza neuraminidase → ↓ release of progeny virus.
CLINICAL USE	Treatment and prevention of influenza A and B. Beginning therapy within 48 hours of symptom onset may shorten duration of illness.

Acyclovir, famciclovir, valacyclovir

MECHANISM	Guanosine analogs. Monophosphorylated by HSV/VZV thymidine kinase and not phosphorylated in uninfected cells → few adverse effects. Triphosphate formed by cellular enzymes. Preferentially inhibit viral DNA polymerase by chain termination.
CLINICAL USE	HSV and VZV. Weak activity against EBV. No activity against CMV. Used for HSV-induced mucocutaneous and genital lesions as well as for encephalitis. Prophylaxis in immunocompromised patients. No effect on latent forms of HSV and VZV. Valacyclovir, a prodrug of acyclovir, has better oral bioavailability. For herpes zoster, use famciclovir.
ADVERSE EFFECTS	Obstructive crystalline nephropathy and acute kidney injury if not adequately hydrated.
MECHANISM OF RESISTANCE	Mutated viral thymidine kinase.

(c) ketabton.com: The Digital Library

Ganciclovir

MECHANISM	5′-monophosphate formed by a CMV viral kinase. Guanosine analog. Triphosphate formed by cellular kinases. Preferentially inhibits viral DNA polymerase.
CLINICAL USE	CMV, especially in immunocompromised patients. Valganciclovir, a prodrug of ganciclovir, has better oral bioavailability.
ADVERSE EFFECTS	Bone marrow suppression (leukopenia, neutropenia, thrombocytopenia), renal toxicity. More toxic to host enzymes than acyclovir.
MECHANISM OF RESISTANCE	Mutated viral kinase.

Foscarnet

MECHANISM	Viral DNA/RNA polymerase inhibitor and HIV reverse transcriptase inhibitor. Binds to pyrophosphate-binding site of enzyme. Does not require any kinase activation.	Foscarnet = pyrofosphate analog.
CLINICAL USE	CMV retinitis in immunocompromised patients when ganciclovir fails; acyclovir-resistant HSV.	
ADVERSE EFFECTS	Nephrotoxicity, electrolyte abnormalities (hypo- or hypercalcemia, hypo- or hyperphosphatemia, hypokalemia, hypomagnesemia) can lead to seizures.	
MECHANISM OF RESISTANCE	Mutated DNA polymerase.	

Cidofovir

MECHANISM	Preferentially inhibits viral DNA polymerase. Does not require phosphorylation by viral kinase.
CLINICAL USE	CMV retinitis in immunocompromised patients; acyclovir-resistant HSV. Long half-life.
ADVERSE EFFECTS	Nephrotoxicity (coadminister with probenecid and IV saline to ↓ toxicity).

(c) ketabton.com: The Digital Library

HIV therapy

Antiretroviral therapy (ART): often initiated at the time of HIV diagnosis.
Strongest indication for use with patients presenting with AIDS-defining illness, low CD4+ cell counts (< 500 cells/mm^3), or high viral load. Regimen consists of 3 drugs to prevent resistance: 2 NRTIs and preferably an integrase inhibitor.
All ARTs are active against HIV-1 and HIV-2 with the exception of NNRTIs and enfuvirtide.

DRUG	MECHANISM	TOXICITY
NRTIs		
Abacavir (ABC) Didanosine (ddI) Emtricitabine (FTC) Lamivudine (3TC) Stavudine (d4T) Tenofovir (TDF) Zidovudine (ZDV, formerly AZT)	Competitively inhibit nucleotide binding to reverse transcriptase and terminate the DNA chain (lack a 3′ OH group). **T**enofovir is a nucleo**T**ide; the others are nucleosides. All need to be phosphorylated to be active. ZDV can be used for general prophylaxis and during pregnancy to ↓ risk of fetal transmission. Ha**ve you dined** (**vudine**) with my **nuclear** (**nucleo**sides) family?	Bone marrow suppression (can be reversed with granulocyte colony-stimulating factor [G-CSF] and erythropoietin), peripheral neuropathy, lactic acidosis (nucleosides), anemia (ZDV), pancreatitis (didanosine). Abacavir contraindicated if patient has HLA-B*5701 mutation due to ↑ risk of hypersensitivity.
NNRTIs		
Delavirdine Efavirenz Nevirapine	Bind to reverse transcriptase at site different from NRTIs. Do not require phosphorylation to be active or compete with nucleotides.	Rash and hepatotoxicity are common to all NNRTIs. Vivid dreams and CNS symptoms are common with efavirenz.
Integrase inhibitors		
Bictegravir Dolutegravir Elvitegravir Raltegravir	Inhibits HIV genome integration into host cell chromosome by reversibly inhibiting HIV integrase.	↑ creatine kinase.
Protease inhibitors		
Atazanavir Darunavir Fosamprenavir Indinavir Lopinavir Ritonavir Saquinavir	Assembly of virions depends on HIV-1 protease (*pol* gene), which cleaves the polypeptide products of HIV mRNA into their functional parts. Thus, protease inhibitors prevent maturation of new viruses. Ritonavir can "boost" other drug concentrations by inhibiting cytochrome P-450. **Navir** (never) **tease** a **protease**.	Hyperglycemia, GI intolerance (nausea, diarrhea), lipodystrophy (Cushing-like syndrome). Nephropathy, hematuria, thrombocytopenia (indinavir). Rifampin (potent CYP/UGT inducer) reduces protease inhibitor concentrations; use rifabutin instead.
Entry inhibitors		
Enfuvirtide	Binds gp41, inhibiting viral entry.	Skin reaction at injection sites. En**fu**virtide inhibits **fu**sion.
Maraviroc	Binds CCR-5 on surface of T cells/monocytes, inhibiting interaction with gp120.	Maravir**oc** inhibits d**oc**king.

(c) ketabton.com: The Digital Library

Hepatitis C therapy

Chronic HCV infection treated with multidrug therapy that targets specific steps within HCV replication cycle (HCV-encoded proteins). Examples of drugs are provided.

DRUG	MECHANISM	TOXICITY
NS5A inhibitors		
Ledipasvir **Ombitasvir** **Velpatasvir**	Inhibits NS5A, a viral phosphoprotein that plays a key role in RNA replication Exact mechanism unknown	Headache, diarrhea
NS5B inhibitors		
Sofosbuvir **Dasabuvir**	Inhibits NS5B, an RNA-dependent RNA polymerase acting as a chain terminator Prevents viral RNA replication	Fatigue, headache
NS3/4A inhibitors		
Grazoprevir **Simeprevir**	Inhibits NS3/4A, a viral protease, preventing viral replication	Grazoprevir: headache, fatigue Simeprevir: photosensitivity reactions, rash
Alternative drugs		
Ribavirin	Inhibits synthesis of guanine nucleotides by competitively inhibiting IMP dehydrogenase Used as adjunct in cases refractory to newer medications	Hemolytic anemia, severe teratogen

Disinfection and sterilization

Goals include the reduction of pathogenic organism counts to safe levels (disinfection) and the inactivation of all microbes including spores (sterilization).

Autoclave	Pressurized steam at > 120°C. Sporicidal. May not reliably inactivate prions.
Alcohols	Denature proteins and disrupt cell membranes. Not sporicidal.
Chlorhexidine	Denatures proteins and disrupts cell membranes. Not sporicidal.
Chlorine	Oxidizes and denatures proteins. Sporicidal.
Ethylene oxide	Alkylating agent. Sporicidal.
Hydrogen peroxide	Free radical oxidation. Sporicidal.
Iodine and iodophors	Halogenation of DNA, RNA, and proteins. May be sporicidal.
Quaternary amines	Impair permeability of cell membranes. Not sporicidal.

Antimicrobials to avoid in pregnancy

ANTIMICROBIAL	ADVERSE EFFECT
Sulfonamides	Kernicterus
Aminoglycosides	Ototoxicity
Fluoroquinolones	Cartilage damage
Clarithromycin	Embryotoxic
Tetracyclines	Discolored teeth, inhibition of bone growth
Ribavirin	Teratogenic
Griseofulvin	Teratogenic
Chloramphenicol	Gray baby syndrome

SAFe **C**hildren **T**ake **R**eally **G**ood **C**are.

(c) ketabton.com: The Digital Library

HIGH-YIELD PRINCIPLES IN

Pathology

"Digressions, objections, delight in mockery, carefree mistrust are signs of health; everything unconditional belongs in pathology."

—Friedrich Nietzsche

"You cannot separate passion from pathology any more than you can separate a person's spirit from his body."

—Richard Selzer

The fundamental principles of pathology are key to understanding diseases in all organ systems. Major topics such as inflammation and neoplasia appear frequently in questions across different organ systems, and such topics are definitely high yield. For example, the concepts of cell injury and inflammation are key to understanding the inflammatory response that follows myocardial infarction, a very common subject of board questions. Similarly, a familiarity with the early cellular changes that culminate in the development of neoplasias—for example, esophageal or colon cancer—is critical. Finally, make sure you recognize the major tumor-associated genes and are comfortable with key cancer concepts such as tumor staging and metastasis.

Cellular Injury 206

Inflammation 213

Neoplasia 219

FAS1_2019_04-Pathol.indd 205 11/7/19 4:02 PM

(c) ketabton.com: The Digital Library

▸ PATHOLOGY—CELLULAR INJURY

Cellular adaptations	Reversible changes that can be physiologic (eg, uterine enlargement during pregnancy) or pathologic (eg, myocardial hypertrophy 2° to systemic HTN). If stress is excessive or persistent, adaptations can progress to cell injury (eg, significant LV hypertrophy → injury to myofibrils → HF).
Hypertrophy	↑ structural proteins and organelles → ↑ in size of cells. Example: cardiac hypertrophy.
Hyperplasia	Controlled proliferation of stem cells and differentiated cells → ↑ in number of cells. Excessive stimulation → pathologic hyperplasia (eg, endometrial hyperplasia), which may progress to dysplasia and cancer. Example: benign prostatic hyperplasia.
Atrophy	↓ in tissue mass due to ↓ in size (↑ cytoskeleton degradation via ubiquitin-proteasome pathway and autophagy; ↓ protein synthesis) and/or number of cells (apoptosis). Causes include disuse, denervation, loss of blood supply, loss of hormonal stimulation, poor nutrition.
Metaplasia	Reprogramming of stem cells → replacement of one cell type by another that can adapt to a new stress. Usually due to exposure to an irritant, such as gastric acid (→ Barrett esophagus) or cigarette smoke (→ respiratory ciliated columnar epithelium replaced by stratified squamous epithelium). May progress to dysplasia → malignant transformation with persistent insult (eg, Barrett esophagus → esophageal adenocarcinoma). Metaplasia of connective tissue can also occur (eg, myositis ossificans, the formation of bone within muscle after trauma).
Dysplasia	Disordered, precancerous epithelial cell growth; not considered a true adaptive response. Characterized by loss of uniformity of cell size and shape (pleomorphism); loss of tissue orientation; nuclear changes (eg, ↑ nuclear:cytoplasmic ratio and clumped chromatin). Mild and moderate dysplasias (ie, do not involve entire thickness of epithelium) may regress with alleviation of inciting cause. Severe dysplasia often becomes irreversible and progresses to carcinoma in situ. Usually preceded by persistent metaplasia or pathologic hyperplasia.



(c) ketabton.com: The Digital Library

Cell injury

Reversible cell injury

- ↓ ATP → ↓ activity of Ca^{2+} and Na^{+}/K^{+} pumps → cellular swelling (earliest morphologic manifestation), mitochondrial swelling
- Ribosomal/polysomal detachment → ↓ protein synthesis
- Plasma membrane changes (eg, blebbing)
- Nuclear changes (eg, chromatin clumping)
- Rapid loss of function (eg, myocardial cells are noncontractile after 1-2 minutes of ischemia)
- Myelin figures (aggregation of peroxidized lipids)

Irreversible cell injury

- Breakdown of plasma membrane → cytosolic enzymes (eg, troponin) leak outside of cell, influx of Ca^{2+} → activation of degradative enzymes
- Mitochondrial damage/dysfunction → loss of electron transport chain → ↓ ATP
- Cytoplasmic vacuolization accompanies programmed cell death (apoptosis)
- Rupture of lysosomes → autolysis
- Nuclear degradation: pyknosis (nuclear condensation) → karyorrhexis (nuclear fragmentation caused by endonuclease-mediated cleavage) → karyolysis (nuclear dissolution)
- Amorphous densities/inclusions in mitochondria



(c) ketabton.com: The Digital Library

Apoptosis

ATP-dependent programmed cell death.
Intrinsic and extrinsic pathways; both pathways activate caspases (cytosolic proteases) → cellular breakdown including cell shrinkage, chromatin condensation, membrane blebbing, and formation of apoptotic bodies, which are then phagocytosed.
Characterized by deeply eosinophilic cytoplasm and basophilic nucleus, pyknosis, and karyorrhexis.
Cell membrane typically remains intact without significant inflammation (unlike necrosis).
DNA laddering (fragments in multiples of 180 bp) is a sensitive indicator of apoptosis.

Intrinsic (mitochondrial) pathway

Involved in tissue remodeling in embryogenesis. Occurs when a regulating factor is withdrawn from a proliferating cell population (eg, ↓ IL-2 after a completed immunologic reaction → apoptosis of proliferating effector cells). Also occurs after exposure to injurious stimuli (eg, radiation, toxins, hypoxia).
Regulated by Bcl-2 family of proteins. BAX and BAK are proapoptotic, while Bcl-2 and Bcl-xL are antiapoptotic.
BAX and BAK form pores in the mitochondrial membrane → release of cytochrome C from inner mitochondrial membrane into the cytoplasm → activation of caspases.
Bcl-2 keeps the mitochondrial membrane impermeable, thereby preventing cytochrome C release. Bcl-2 overexpression (eg, follicular lymphoma t[14;18]) → ↓ caspase activation → tumorigenesis.

Extrinsic (death receptor) pathway

2 pathways:
- Ligand receptor interactions (FasL binding to Fas [CD95] or TNF-α binding to its receptor)
- Immune cell (cytotoxic T-cell release of perforin and granzyme B)

Fas-FasL interaction is necessary in thymic medullary negative selection.
Fas mutations ↑ numbers of circulating self-reacting lymphocytes due to failure of clonal deletion.
Defective Fas-FasL interactions cause autoimmune lymphoproliferative syndrome.



FAS1_2019_04-Pathol.indd 208 11/7/19 4:02 PM

(c) ketabton.com: The Digital Library

Necrosis

Exogenous injury → plasma membrane damage → cell undergoes enzymatic degradation and protein denaturation, intracellular components leak → local inflammatory reaction (unlike apoptosis).

TYPE	SEEN IN	DUE TO	HISTOLOGY
Coagulative	Ischemia/infarcts in most tissues (except brain)	Ischemia or infarction; injury denatures enzymes → proteolysis blocked	Preserved cellular architecture (cell outlines seen), but nuclei disappear; ↑ cytoplasmic binding of eosin stain (→ ↑ eosinophilia; red/pink color) A
Liquefactive	Bacterial abscesses, brain infarcts	Neutrophils release lysosomal enzymes that digest the tissue B	Early: cellular debris and macrophages Late: cystic spaces and cavitation (brain) Neutrophils and cell debris seen with bacterial infection
Caseous	TB, systemic fungi (eg, *Histoplasma capsulatum*), *Nocardia*	Macrophages wall off the infecting microorganism → granular debris C	Fragmented cells and debris surrounded by lymphocytes and macrophages (granuloma)
Fat	Enzymatic: acute pancreatitis (saponification of peripancreatic fat) Nonenzymatic: traumatic (eg, injury to breast tissue)	Damaged pancreatic cells release lipase, which breaks down triglycerides; liberated fatty acids bind calcium → saponification (chalky-white appearance)	Outlines of dead fat cells without peripheral nuclei; saponification of fat (combined with Ca^{2+}) appears dark blue on H&E stain D
Fibrinoid	Immune vascular reactions (eg, PAN) Nonimmune vascular reactions (eg, hypertensive emergency, preeclampsia)	Immune complex deposition (type III hypersensitivity reaction) and/or plasma protein (eg, fibrin) leakage from damaged vessel	Vessel walls are thick and pink E
Gangrenous	Distal extremity and GI tract, after chronic ischemia	Dry: ischemia F	Coagulative
		Wet: superinfection	Liquefactive superimposed on coagulative



(c) ketabton.com: The Digital Library

Ischemia



Inadequate blood supply to meet demand. Mechanisms include ↓ arterial perfusion (eg, atherosclerosis), ↓ venous drainage (eg, testicular torsion, Budd-Chiari syndrome), shock.

Regions most vulnerable to hypoxia/ischemia and subsequent infarction:

ORGAN	REGION
Brain	ACA/MCA/PCA boundary areas[a,b]
Heart	Subendocardium (LV) A
Kidney	Straight segment of proximal tubule (medulla) Thick ascending limb (medulla)
Liver	Area around central vein (zone III)
Colon	Splenic flexure (Griffith point),[a] rectosigmoid junction (Sudeck point)[a]

[a]Watershed areas (border zones) receive blood supply from most distal branches of 2 arteries with limited collateral vascularity. These areas are susceptible to ischemia from hypoperfusion.

[b]Neurons most vulnerable to hypoxic-ischemic insults include Purkinje cells of the cerebellum and pyramidal cells of the hippocampus and neocortex (zones 3, 5, 6).

Types of infarcts

Red infarct

Occurs in venous occlusion and tissues with multiple blood supplies (eg, liver, lung A, intestine, testes), and with reperfusion (eg, after angioplasty). Reperfusion injury is due to damage by free radicals.

Pale infarct

Occurs in solid organs with a single (end-arterial) blood supply (eg, heart, kidney B).



Free radical injury

Free radicals damage cells via membrane lipid peroxidation, protein modification, DNA breakage. Initiated via radiation exposure (eg, cancer therapy), metabolism of drugs (phase I), redox reactions, nitric oxide (eg, inflammation), transition metals, WBC (eg, neutrophils, macrophages) oxidative burst.

Free radicals can be eliminated by scavenging enzymes (eg, catalase, superoxide dismutase, glutathione peroxidase), spontaneous decay, antioxidants (eg, vitamins A, C, E), and certain metal carrier proteins (eg, transferrin, ceruloplasmin).

Examples:

- Oxygen toxicity: retinopathy of prematurity (abnormal vascularization), bronchopulmonary dysplasia, reperfusion injury after thrombolytic therapy
- Drug/chemical toxicity: acetaminophen overdose (hepatotoxicity), carbon tetrachloride (converted by cytochrome P-450 into CCl_3 free radical → fatty liver [cell injury → ↓ apolipoprotein synthesis → fatty change], centrilobular necrosis)
- Metal storage diseases: hemochromatosis (iron) and Wilson disease (copper)

(c) ketabton.com: The Digital Library

Types of calcification

Calcium deposits appear deeply basophilic (arrow in A) on H&E stain.

	Dystrophic calcification	Metastatic calcification
Ca^{2+} DEPOSITION	In abnormal (Diseased) tissues	In normal tissues
EXTENT	Tends to be localized (eg, calcific aortic stenosis)	Widespread (ie, diffuse, metastatic)
ASSOCIATED CONDITIONS 	TB (lung and pericardium) and other granulomatous infections, liquefactive necrosis of chronic abscesses, fat necrosis, infarcts, thrombi, schistosomiasis, congenital CMV, toxoplasmosis, rubella, psammoma bodies, CREST syndrome, atherosclerotic plaques can become calcified	Predominantly in interstitial tissues of kidney, lung, and gastric mucosa (these tissues lose acid quickly; ↑ pH favors Ca^{2+} deposition) Nephrocalcinosis of collecting ducts may lead to nephrogenic diabetes insipidus and renal failure
ETIOLOGY	2° to injury or necrosis	2° to hypercalcemia (eg, 1° hyperparathyroidism, sarcoidosis, hypervitaminosis D) or high calcium-phosphate product levels (eg, chronic kidney disease with 2° hyperparathyroidism, long-term dialysis, calciphylaxis, multiple myeloma)
SERUM Ca^{2+} LEVELS	Normal	Usually abnormal

Lipofuscin



A yellow-brown "wear and tear" pigment A associated with normal aging.

Composed of polymers of lipids and phospholipids complexed with protein. May be derived through lipid peroxidation of polyunsaturated lipids of subcellular membranes.

Autopsy of elderly person will reveal deposits in heart, colon, liver, kidney, eye, and other organs.

(c) ketabton.com: The Digital Library

Amyloidosis

Abnormal aggregation of proteins (or their fragments) into β-pleated linear sheets → insoluble fibrils → cellular damage and apoptosis. Amyloid deposits visualized by Congo red stain (red/orange on nonpolarized light [arrows in A]), (apple-green birefringence on polarized light [arrows in B]), and H&E stain (shows deposits in glomerular mesangial areas). Tubular basement membranes are enlarged on light microscopy.

COMMON TYPES	FIBRIL PROTEIN	DESCRIPTION	
Systemic			
Primary amyloidosis	AL (from Ig Light chains)	Seen in Plasma cell disorders (eg, multiple myeloma)	Manifestations include: ▪ Cardiac (eg, restrictive cardiomyopathy) ▪ GI (eg, macroglossia, hepatomegaly) ▪ Renal (eg, nephrotic syndrome) ▪ Hematologic (eg, easy bruising, splenomegaly) ▪ Neurologic (eg, neuropathy) ▪ Musculoskeletal (eg, carpal tunnel syndrome)
Secondary amyloidosis	Serum Amyloid A (AA)	Seen in chronic inflammatory conditions, (eg, rheumatoid arthritis, IBD, familial Mediterranean fever, protracted infection)	
Dialysis-related amyloidosis	β_2-microglobulin	Seen in patients with ESRD and/or on long-term dialysis	
Localized			
Alzheimer disease	β-amyloid protein	Cleaved from amyloid precursor protein (APP)	
Type 2 diabetes mellitus	Islet amyloid polypeptide (IAPP)	Caused by deposition of amylin in pancreatic islets	
Medullary thyroid cancer	Calcitonin		
Isolated atrial amyloidosis	ANP	Common in normal aging ↑ risk of atrial fibrillation	
Systemic senile (age-related) amyloidosis	Normal (wild-type) transthyretin (TTR)	Seen predominantly in cardiac ventricles	Cardiac dysfunction more insidious than in AL amyloidosis
Hereditary			
Familial amyloid cardiomyopathy	Mutated transthyretin (ATTR)	Ventricular endomyocardium deposition → restrictive cardiomyopathy, arrhythmias	5% of African Americans are carriers of mutant allele
Familial amyloid polyneuropathies	Mutated transthyretin (ATTR)	Due to transthyretin gene mutation	



(c) ketabton.com: The Digital Library

▸ PATHOLOGY—INFLAMMATION

Inflammation — Response to eliminate initial cause of cell injury, to remove necrotic cells resulting from the original insult, and to initiate tissue repair. Divided into acute and chronic. The inflammatory response itself can be harmful to the host if the reaction is excessive (eg, septic shock), prolonged (eg, persistent infections such as TB), or inappropriate (eg, autoimmune diseases such as SLE).

Cardinal signs

SIGN	MECHANISM	MEDIATORS
Rubor (redness), calor (warmth)	Vasodilation (relaxation of arteriolar smooth muscle) → ↑ blood flow	Histamine, prostaglandins, bradykinin, NO
Tumor (swelling)	Endothelial contraction/disruption (eg, from tissue damage) → ↑ vascular permeability → leakage of protein-rich fluid from postcapillary venules into interstitial space (exudate) → ↑ interstitial oncotic pressure	Endothelial contraction: leukotrienes (C_4, D_4, E_4), histamine, serotonin
Dolor (pain)	Sensitization of sensory nerve endings	Bradykinin, PGE_2, histamine
Functio laesa (loss of function)	Cardinal signs above impair function (eg, inability to make fist with hand that has cellulitis)	

Systemic manifestations (acute-phase reaction)

Fever	Pyrogens (eg, LPS) induce macrophages to release IL-1 and TNF → ↑ COX activity in perivascular cells of hypothalamus → ↑ PGE_2 → ↑ temperature set point	
Leukocytosis	Elevation of WBC count; type of cell that is predominantly elevated depends on the inciting agent or injury (eg, bacteria → ↑ neutrophils)	
↑ plasma acute-phase proteins	Factors whose serum concentrations change significantly in response to inflammation Produced by the liver in both acute and chronic inflammatory states	Notably induced by IL-6

Acute phase reactants — More **FFiSH** in the **C** (sea).

POSITIVE (UPREGULATED)	
Ferritin	Binds and sequesters iron to inhibit microbial iron scavenging.
Fibrinogen	Coagulation factor; promotes endothelial repair; correlates with ESR.
Serum amyloid A	Prolonged elevation can lead to amyloidosis.
Hepcidin	↓ iron absorption (by degrading ferroportin) and ↓ iron release (from macrophages) → anemia of chronic disease.
C-reactive protein	Opsonin; fixes complement and facilitates phagocytosis. Measured clinically as a nonspecific sign of ongoing inflammation.
NEGATIVE (DOWNREGULATED)	
Albumin	Reduction conserves amino acids for positive reactants.
Transferrin	Internalized by macrophages to sequester iron.

(c) ketabton.com: The Digital Library

Erythrocyte sedimentation rate

RBCs normally remain separated via ⊖ charges. Products of inflammation (eg, fibrinogen) coat RBCs → ↓ ⊖ charge → ↑ RBC aggregation. Denser RBC aggregates fall at a faster rate within a pipette tube → ↑ ESR. Often co-tested with CRP (more specific marker of inflammation).

↑ ESR	↓ ESR
Most anemias	Sickle cell anemia (altered shape)
Infections	Polycythemia (↑ RBCs "dilute" aggregation factors)
Inflammation (eg, giant cell [temporal] arteritis, polymyalgia rheumatica)	HF
Cancer (eg, metastases, multiple myeloma)	Microcytosis
Renal disease (end-stage or nephrotic syndrome)	Hypofibrinogenemia
Pregnancy	

Acute inflammation



Transient and early response to injury or infection. Characterized by neutrophils in tissue A, often with associated edema. Rapid onset (seconds to minutes) and short duration (minutes to days). Represents a reaction of the innate immune system (ie, less specific response than chronic inflammation).

STIMULI	Infections, trauma, necrosis, foreign bodies.	
MEDIATORS	Toll-like receptors, arachidonic acid metabolites, neutrophils, eosinophils, antibodies (pre-existing), mast cells, basophils, complement, Hageman factor (factor XII).	Inflammasome—Cytoplasmic protein complex that recognizes products of dead cells, microbial products, and crystals (eg, uric acid crystals) → activation of IL-1 and inflammatory response.
COMPONENTS	▪ Vascular: vasodilation (→ ↑ blood flow and stasis) and ↑ endothelial permeability ▪ Cellular: extravasation of leukocytes (mainly neutrophils) from postcapillary venules and accumulation in the focus of injury followed by leukocyte activation	To bring cells and proteins to site of injury or infection. Leukocyte extravasation has 4 steps: margination and rolling, adhesion, transmigration, and migration (chemoattraction).
OUTCOMES	▪ Resolution and healing (IL-10, TGF-β) ▪ Persistent acute inflammation (IL-8) ▪ Abscess (acute inflammation walled off by fibrosis) ▪ Chronic inflammation (antigen presentation by macrophages and other APCs → activation of $CD4^+$ Th cells) ▪ Scarring	Macrophages predominate in the late stages of acute inflammation (peak 2–3 days after onset) and influence outcome by secreting cytokines.

(c) ketabton.com: The Digital Library

Leukocyte extravasation

Extravasation predominantly occurs at postcapillary venules.

STEP	VASCULATURE/STROMA	LEUKOCYTE
❶ Margination and rolling—defective in leukocyte adhesion deficiency type 2 (↓ Sialyl Lewis^X)	E-selectin (upregulated by TNF and IL-1)	Sialyl Lewis^X
	P-selectin (released from Weibel-Palade bodies)	Sialyl Lewis^X
	GlyCAM-1, CD34	L-selectin
❷ Tight binding (adhesion)—defective in leukocyte adhesion deficiency type 1 (↓ CD18 integrin subunit)	ICAM-1 (CD54)	CD11/18 integrins (LFA-1, Mac-1)
	VCAM-1 (CD106)	VLA-4 integrin
❸ Diapedesis (transmigration)—WBC travels between endothelial cells and exits blood vessel	PECAM-1 (CD31)	PECAM-1 (CD31)
❹ Migration—WBC travels through interstitium to site of injury or infection guided by chemotactic signals	Chemotactic factors: C5a, IL-8, LTB_4, kallikrein, platelet-activating factor	Various



(c) ketabton.com: The Digital Library

Chronic inflammation	Prolonged inflammation characterized by mononuclear infiltration (macrophages, lymphocytes, plasma cells), which leads to simultaneous tissue destruction and repair (including angiogenesis and fibrosis). May be preceded by acute inflammation.
STIMULI	Persistent infections (eg, TB, *T pallidum*, certain fungi and viruses) → type IV hypersensitivity, autoimmune diseases, prolonged exposure to toxic agents (eg, silica) and foreign material.
MEDIATORS	Macrophages are the dominant cells. Interaction of macrophages and T lymphocytes → chronic inflammation. ▪ Th1 cells secrete IFN-γ → macrophage classical activation (proinflammatory) ▪ Th2 cells secrete IL-4 and IL-13 → macrophage alternative activation (repair and anti-inflammatory)
OUTCOMES	Scarring, amyloidosis, and neoplastic transformation (eg, chronic HCV infection → chronic inflammation → hepatocellular carcinoma; *Helicobacter pylori* infection → chronic gastritis → gastric adenocarcinoma).

Wound healing

	MEDIATOR	ROLE
Tissue mediators	FGF	Stimulates angiogenesis
	TGF-β	Angiogenesis, fibrosis
	VEGF	Stimulates angiogenesis
	PDGF	Secreted by activated platelets and macrophages Induces vascular remodeling and smooth muscle cell migration Stimulates fibroblast growth for collagen synthesis
	Metalloproteinases	Tissue remodeling
	EGF	Stimulates cell growth via tyrosine kinases (eg, EGFR/*ErbB1*)

PHASE OF WOUND HEALING	EFFECTOR CELLS	CHARACTERISTICS
Inflammatory (up to 3 days after wound)	Platelets, neutrophils, macrophages	Clot formation, ↑ vessel permeability and neutrophil migration into tissue; macrophages clear debris 2 days later
Proliferative (day 3–weeks after wound)	Fibroblasts, myofibroblasts, endothelial cells, keratinocytes, macrophages	Deposition of granulation tissue and type III collagen, angiogenesis, epithelial cell proliferation, dissolution of clot, and wound contraction (mediated by myofibroblasts) Delayed second phase of wound healing in vitamin C and copper deficiency
Remodeling (1 week–6+ months after wound)	Fibroblasts	Type III collagen replaced by type I collagen, ↑ tensile strength of tissue Collagenases (require zinc to function) break down type III collagen Zinc deficiency → delayed wound healing

(c) ketabton.com: The Digital Library

Granulomatous inflammation

A pattern of chronic inflammation. Can be induced by persistent T-cell response to certain infections (eg, TB), immune-mediated diseases, and foreign bodies. Granulomas "wall off" a resistant stimulus without completely eradicating or degrading it → persistent inflammation→ fibrosis, organ damage.

HISTOLOGY



Focus of epithelioid cells (activated macrophages with abundant pink cytoplasm) surrounded by lymphocytes and multinucleated giant cells (formed by fusion of several activated macrophages). Two types:

Caseating: associated with Central necrosis. Seen with infectious etiologies (eg, TB, fungal).

Noncaseating A: no central necrosis. Seen with autoimmune diseases (eg, sarcoidosis, Crohn disease).

MECHANISM

❶ APCs present antigens to CD4+ Th cells and secrete IL-12 → CD4+ Th cells differentiate into Th1 cells

❷ Th1 secretes IFN-γ → macrophage activation

❸ Macrophages ↑ cytokine secretion (eg, TNF) → formation of epithelioid macrophages and giant cells.

Anti-TNF therapy can cause sequestering granulomas to break down → disseminated disease. Always test for latent TB before starting anti-TNF therapy.

Associated with hypercalcemia due to ↑ 1α-hydroxylase activity in activated macrophages, resulting in ↑ vitamin D activity.



ETIOLOGIES

INFECTIOUS	NONINFECTIOUS
Bacterial: *Mycobacteria* (tuberculosis, leprosy), *Bartonella henselae* (cat scratch disease; stellate necrotizing granulomas), *Listeria monocytogenes* (granulomatosis infantiseptica), *Treponema pallidum* (3° syphilis)	Immune-mediated: sarcoidosis, Crohn disease, 1° biliary cholangitis, subacute (de Quervain/ granulomatous) thyroiditis
Fungal: endemic mycoses (eg, histoplasmosis)	Vasculitis: granulomatosis with polyangiitis (Wegener), eosinophilic granulomatosis with polyangiitis (Churg-Strauss), giant cell (temporal) arteritis, Takayasu arteritis
Parasitic: schistosomiasis	Foreign bodies: berylliosis, talcosis, hypersensitivity pneumonitis
	Chronic granulomatous disease

(c) ketabton.com: The Digital Library

Scar formation Occurs when repair cannot be accomplished by cell regeneration alone. Nonregenerated cells (2° to severe acute or chronic injury) are replaced by connective tissue. 70–80% of tensile strength regained at 3 months; little tensile strength regained thereafter. Associated with excess TGF-β.

SCAR TYPE	Hypertrophic A	Keloid B
COLLAGEN SYNTHESIS	↑ (type III collagen)	↑↑↑ (types I and III collagen)
COLLAGEN ORGANIZATION	Parallel	Disorganized
EXTENT OF SCAR	Confined to borders of original wound	Extends beyond borders of original wound with "claw-like" projections typically on earlobes, face, upper extremities
RECURRENCE	Infrequent	Frequent
PREDISPOSITION	None	↑ incidence in ethnic groups with darker skin

(c) ketabton.com: The Digital Library

▸ PATHOLOGY—NEOPLASIA

Neoplasia and neoplastic progression

Uncontrolled, monoclonal proliferation of cells. Can be benign or malignant. Any neoplastic growth has two components: parenchyma (neoplastic cells) and supporting stroma (non-neoplastic; eg, blood vessels, connective tissue).



Normal cells	❶ Normal cells with basal → apical polarity. See cervical example A, which shows normal cells and spectrum of dysplasia, as discussed below.
Dysplasia	❷ Loss of uniformity in cell size and shape (pleomorphism); loss of tissue orientation; nuclear changes (eg, ↑ nuclear:cytoplasmic ratio) A.
Carcinoma in situ/ preinvasive	❸ Irreversible severe dysplasia that involves the entire thickness of epithelium but does not penetrate the intact basement membrane A.
Invasive carcinoma	❹ Cells have invaded basement membrane using collagenases and hydrolases (metalloproteinases). Cell-cell contacts lost by inactivation of E-cadherin.
Metastasis	❺ Spread to distant organ(s) via lymphatics or blood.



(c) ketabton.com: The Digital Library

Tumor nomenclature

Carcinoma implies epithelial origin, whereas **sarcoma** denotes mesenchymal origin. Both terms generally imply malignancy.

Benign tumors are usually well-differentiated and well-demarcated, with low mitotic activity, no metastases, and no necrosis.

Malignant tumors (cancers) may show poor differentiation, erratic growth, local invasion, metastasis, and ↓ apoptosis.

Terms for non-neoplastic malformations include hamartoma (disorganized overgrowth of tissues in their native location, eg, Peutz-Jeghers polyps) and choristoma (normal tissue in a foreign location, eg, gastric tissue located in distal ileum in Meckel diverticulum).

CELL TYPE	BENIGN	MALIGNANT
Epithelium	Adenoma, papilloma	Adenocarcinoma, papillary carcinoma
Mesenchyme		
Blood cells		Leukemia, lymphoma
Blood vessels	Hemangioma	Angiosarcoma
Smooth muscle	Leiomyoma	Leiomyosarcoma
Striated muscle	Rhabdomyoma	Rhabdomyosarcoma
Connective tissue	Fibroma	Fibrosarcoma
Bone	Osteoma	Osteosarcoma
Fat	Lipoma	Liposarcoma
Melanocyte	Nevus/mole	Melanoma

Tumor grade vs stage

Differentiation—degree to which a tumor resembles its tissue of origin. Well-differentiated tumors (often less aggressive) closely resemble their tissue of origin, whereas poorly differentiated tumors (often more aggressive) do not.

Anaplasia—complete lack of differentiation of cells in a malignant neoplasm.

Grade

Degree of cellular differentiation and mitotic activity on histology. Ranges from low grade (well-differentiated) to high grade (poorly differentiated, undifferentiated, or anaplastic).



Stage

Degree of localization/spread based on site and size of 1° lesion, spread to regional lymph nodes, presence of metastases. Based on clinical (c) or pathologic (p) findings. Stage generally has more prognostic value than grade (eg, a high-stage yet low-grade tumor is usually worse than a low-stage yet high-grade tumor). Stage determines Survival.

TNM staging system (Stage = Spread):
T = Tumor size/invasiveness, N = Node involvement, M = Metastases, eg, cT3N1M0. Each TNM factor has independent prognostic value; N and M are often most important.



(c) ketabton.com: The Digital Library

Hallmarks of cancer

Cancer is caused by (mostly acquired) DNA mutations that affect fundamental cellular processes (eg, growth, DNA repair, survival).

HALLMARK	MECHANISM
Growth signal self-sufficiency	Mutations in genes encoding: ▪ Proto-oncogenes → ↑ growth factors → autocrine loop (eg, ↑ PDGF in brain tumors) ▪ Growth factor receptors → constitutive signalling (eg, *HER2/neu* in breast cancer) ▪ Signaling molecules (eg, *RAS*) ▪ Transcription factors (eg, *MYC*) ▪ Cell cycle regulators (eg, cyclins, CDKs)
Anti-growth signal insensitivity	▪ Mutations in tumor suppressor genes (eg, *Rb*) ▪ Loss of E-cadherin function → loss of contact inhibition (eg, *NF2* mutations)
Evasion of apoptosis	Mutations in genes that regulate apoptosis (eg, *TP53*, *BCL2* → follicular B cell lymphoma).
Limitless replicative potential	Reactivation of telomerase → maintenance and lengthening of telomeres → prevention of chromosome shortening and cell aging.
Sustained angiogenesis	↑ pro-angiogenic factors (eg, VEGF) or ↓ inhibitory factors. Factors may be produced by tumor or stromal cells. Vessels can sprout from existing capillaries (neoangiogenesis) or endothelial cells are recruited from bone marrow (vasculogenesis). Vessels may be leaky and/or dilated.
Tissue invasion	Loss of E-cadherin function → loosening of intercellular junctions → metalloproteinases degrade basement membrane and ECM → cells attach to ECM proteins (eg, laminin, fibronectin) → cells migrate through degraded ECM ("locomotion") → vascular dissemination.
Metastasis	Tumor cells or emboli spread via lymphatics or blood → adhesion to endothelium → extravasation and homing. Site of metastasis can be predicted by site of 1° tumor, as the target organ is often the first-encountered capillary bed. Some cancers show organ tropism (eg, lung cancers commonly metastasize to adrenals).
Warburg effect	Shift of glucose metabolism away from mitochondrial oxidative phosphorylation toward glycolysis.
Immune evasion in cancer	Normally, immune cells can recognize and attack tumor cells. For successful tumorigenesis, tumor cells must evade the immune system. Multiple escape mechanisms exist: ▪ ↓ MHC class I expression by tumor cells → cytotoxic T cells are unable to recognize tumor cells. ▪ Tumor cells secrete immunosuppressive factors (eg, TGF-β) and recruit regulatory T cells to down regulate immune response. ▪ Tumor cells up regulate immune checkpoint molecules, which inhibit immune response.

(c) ketabton.com: The Digital Library

Immune checkpoint interactions

Signals that modulate T cell activation and function → ↓ immune response against tumor cells. Targeted by several cancer immunotherapies. Examples:

- Interaction between PD-1 (on T cells) and PD-L1/2 (on tumor cells or immune cells in tumor microenvironment) → T cell dysfunction (exhaustion). Inhibited by antibodies against PD-1 (eg, pembrolizumab, nivolumab) or PD-L1 (eg, atezolizumab, durvalumab, avelumab).
- CTLA-4 on T cells outcompetes CD28 for B7 on APCs → loss of T cell costimulatory signal. Inhibited by ipilimumab (anti-CTLA-4 antibody).



Cancer epidemiology

Skin cancer (basal > squamous >> melanoma) is the most common cancer (not included below).

	MEN	WOMEN	CHILDREN (AGE 0–14)	NOTES
Cancer incidence	1. Prostate 2. Lung 3. Colon/rectum	1. Breast 2. Lung 3. Colon/rectum	1. Leukemia 2. CNS 3. Neuroblastoma	Lung cancer incidence has ↓ in men, but has not changed significantly in women.
Cancer mortality	1. Lung 2. Prostate 3. Colon/rectum	1. Lung 2. Breast 3. Colon/rectum	1. Leukemia 2. CNS 3. Neuroblastoma	Cancer is the 2nd leading cause of death in the United States (heart disease is 1st).

(c) ketabton.com: The Digital Library

Common metastases Most sarcomas spread hematogenously; most carcinomas spread via lymphatics. However, **F**our **C**arcinomas **R**oute **H**ematogenously: **F**ollicular thyroid carcinoma, **C**horiocarcinoma, **R**enal cell carcinoma, and **H**epatocellular carcinoma.

SITE OF METASTASIS	1° TUMOR	NOTES
Brain	Lung > breast > melanoma, colon, kidney	50% of brain tumors are from metastases A B Commonly seen as multiple well-circumscribed tumors at gray/white matter junction
Liver	**C**olon >> **S**tomach > **P**ancreas (**C**ancer **S**ometimes **P**enetrates liver)	Liver C D and lung are the most common sites of metastasis after the regional lymph nodes
Bone	**P**rostate, **B**reast > **K**idney, **T**hyroid, **L**ung (**P**ainful **B**ones **K**ill **T**he **L**ungs)	Bone metastasis E F >> 1° bone tumors (eg, multiple myeloma) Predilection for axial skeleton G Bone metastasis can be: ▪ Lytic (eg, thyroid, kidney, non-small cell lung cancer) ▪ Blastic (eg, prostate, small cell lung cancer) ▪ Mixed (eg, breast cancer)



(c) ketabton.com: The Digital Library

Oncogenes

Gain of function mutation converts proto-oncogene (normal gene) to oncogene → ↑ cancer risk. Requires damage to only one allele of a proto-oncogene.

GENE	GENE PRODUCT	ASSOCIATED NEOPLASM
ALK	Receptor tyrosine Kinase	Lung Adenocarcinoma (Adenocarcinoma of the Lung Kinase)
BCR-ABL	Non-receptor tyrosine kinase	CML, ALL
BCL-2	Antiapoptotic molecule (inhibits apoptosis)	Follicular and diffuse large B Cell Lymphomas
BRAF	Serine/threonine kinase	Melanoma, non-Hodgkin lymphoma, papillary thyroid carcinoma, hairy cell leukemia
c-KIT	CytoKIne receptor	Gastrointestinal stromal tumor (GIST)
c-MYC	Transcription factor	Burkitt lymphoma
HER2/neu (c-erbB2)	Receptor tyrosine kinase	Breast and gastric carcinomas
JAK2	Tyrosine kinase	Chronic myeloproliferative disorders
KRAS	GTPase	Colon cancer, lung cancer, pancreatic cancer
MYCL1	Transcription factor	Lung tumor
N-*myc (MYCN)*	Transcription factor	Neuroblastoma
RET	Receptor tyrosine kinase	MEN 2A and 2B, papillary thyroid carcinoma, pheochromocytoma

Tumor suppressor genes

Loss of function → ↑ cancer risk; both (two) alleles of a tumor suppressor gene must be lost for expression of disease.

GENE	GENE PRODUCT	ASSOCIATED CONDITION
APC	Negative regulator of β-catenin/WNT pathway	Colorectal cancer (associated with FAP)
BRCA1/BRCA2	BRCA1/BRCA2 proteins	Breast, ovarian, and pancreatic cancers
CDKN2A	p16, blocks G_1 → S phase	Melanoma, pancreatic cancer
DCC	DCC—Deleted in Colon Cancer	Colon cancer
SMAD4 (DPC4)	DPC—Deleted in Pancreatic Cancer	Pancreatic cancer
MEN1	Menin	Multiple Endocrine Neoplasia type 1
NF1	Neurofibromin (Ras GTPase activating protein)	Neurofibromatosis type 1
NF2	Merlin (schwannomin) protein	Neurofibromatosis type 2
PTEN	Negative regulator of PI3k/AKT pathway	Prostate, breasT, and ENdometrial cancers
Rb	Inhibits E2F; blocks G_1 → S phase	Retinoblastoma, osteosarcoma (bone cancer)
TP53	p53, activates p21, blocks G_1 → S phase	Most human cancers, Li-Fraumeni syndrome (multiple malignancies at early age, aka, SBLA cancer syndrome: Sarcoma, Breast, Leukemia, Adrenal gland)
TSC1	Hamartin protein	Tuberous sclerosis
TSC2	Tuberin protein	Tuberous sclerosis
VHL	Inhibits hypoxia-inducible factor 1a	von Hippel-Lindau disease
WT1	Urogenital development transcription factor	Wilms tumor (nephroblastoma)

(c) ketabton.com: The Digital Library

Carcinogens

TOXIN	EXPOSURE	ORGAN	IMPACT
Aflatoxins (*Aspergillus*)	Stored grains and nuts	Liver	Hepatocellular carcinoma
Alkylating agents	Oncologic chemotherapy	Blood	Leukemia/lymphoma
Aromatic amines (eg, benzidine, 2-naphthylamine)	Textile industry (dyes), cigarette smoke (2-naphthylamine)	Bladder	Transitional cell carcinoma
Arsenic	Herbicides (vineyard workers), metal smelting	Liver Lung Skin	Angiosarcoma Lung cancer Squamous cell carcinoma
Asbestos	Old roofing material, shipyard workers	Lung	Bronchogenic carcinoma > mesothelioma
Cigarette smoke		Bladder Cervix Esophagus Kidney Larynx Lung Oropharynx Pancreas	Transitional cell carcinoma Squamous cell carcinoma Squamous cell carcinoma/ adenocarcinoma Renal cell carcinoma Squamous cell carcinoma Squamous cell and small cell carcinoma Oropharyngeal cancer Pancreatic adenocarcinoma
Ethanol		Esophagus Liver	Squamous cell carcinoma Hepatocellular carcinoma
Ionizing radiation		Thyroid	Papillary thyroid carcinoma, leukemias
Nickel, chromium, beryllium, silica	Occupational exposure	Lung	Lung cancer
Nitrosamines	Smoked foods	Stomach	Gastric cancer
Radon	Byproduct of uranium decay, accumulates in basements	Lung	Lung cancer (2nd leading cause after cigarette smoke)
Vinyl chloride	Used to make PVC pipes (plumbers)	Liver	Angiosarcoma

(c) ketabton.com: The Digital Library

Oncogenic microbes

Microbe	Associated cancer
EBV	Burkitt lymphoma, Hodgkin lymphoma, nasopharyngeal carcinoma, 1° CNS lymphoma (in immunocompromised patients)
HBV, HCV	Hepatocellular carcinoma
HHV-8	Kaposi sarcoma
HPV	Cervical and penile/anal carcinoma (types 16, 18), head and neck cancer
H pylori	Gastric adenocarcinoma and MALT lymphoma
HTLV-1	Adult T-cell Leukemia/Lymphoma
Liver fluke *(Clonorchis sinensis)*	Cholangiocarcinoma
Schistosoma haematobium	Squamous cell bladder cancer

Serum tumor markers

Tumor markers should not be used as the 1° tool for cancer diagnosis or screening. They may be used to monitor tumor recurrence and response to therapy, but definitive diagnosis is made via biopsy. Some can be associated with non-neoplastic conditions.

MARKER	IMPORTANT ASSOCIATIONS	NOTES
Alkaline phosphatase	Metastases to bone or liver, Paget disease of bone, seminoma (placental ALP).	Exclude hepatic origin by checking LFTs and GGT levels.
α-fetoprotein	Hepatocellular carcinoma, Endodermal sinus (yolk sac) tumor, Mixed germ cell tumor, Ataxia-telangiectasia, Neural tube defects. (HE-MAN is the alpha male!)	Normally made by fetus. Transiently elevated in pregnancy. High levels associated with neural tube and abdominal wall defects, low levels associated with Down syndrome.
hCG	Hydatidiform moles and Choriocarcinomas (Gestational trophoblastic disease), testicular cancer, mixed germ cell tumor.	Produced by syncytiotrophoblasts of the placenta.
CA 15-3/CA 27-29	Breast cancer.	
CA 19-9	Pancreatic adenocarcinoma.	
CA 125	Ovarian cancer.	
Calcitonin	Medullary thyroid carcinoma (alone and in MEN2A, MEN2B).	
CEA	Colorectal and pancreatic cancers. Minor associations: gastric, breast, and medullary thyroid carcinomas.	Carcinoembryonic antigen. Very nonspecific.
Chromogranin	Neuroendocrine tumors.	
LDH	Testicular germ cell tumors, ovarian dysgerminoma, other cancers.	Can be used as an indicator of tumor burden.
Neuron-specific enolase	Neuroendocrine tumors (eg, small cell lung cancer, carcinoid tumor, neuroblastoma)	
PSA	Prostate cancer.	Prostate-specific antigen. Also elevated in BPH and prostatitis. Questionable risk/benefit for screening. Marker for recurrence after treatment.

(c) ketabton.com: The Digital Library

Important immunohistochemical stains

Determine primary site of origin for metastatic tumors and characterize tumors that are difficult to classify. Can have prognostic and predictive value.

STAIN	TARGET	TUMORS IDENTIFIED
Chromogranin and synaptophysin	Neuroendocrine cells	Small cell carcinoma of the lung, carcinoid tumor
Cytokeratin	Epithelial cells	Epithelial tumors (eg, squamous cell carcinoma)
DesMin	Muscle	Muscle tumors (eg, rhabdomyosarcoma)
GFAP	NeuroGlia (eg, astrocytes, Schwann cells, oligodendrocytes)	Astrocytoma, Glioblastoma
Neurofilament	Neurons	Neuronal tumors (eg, neuroblastoma)
PSA	Prostatic epithelium	Prostate cancer
S-100	Neural crest cells	Melanoma, schwannoma, Langerhans cell histiocytosis
TRAP	Tartrate-resistant acid phosphatase	Hairy cell leukemia
Vimentin	Mesenchymal tissue (eg, fibroblasts, endothelial cells, macrophages)	Mesenchymal tumors (eg, sarcoma), but also many other tumors (eg, endometrial carcinoma, renal cell carcinoma, meningioma)

P-glycoprotein

Also known as multidrug resistance protein 1 (MDR1). Classically seen in adrenocortical carcinoma but also expressed by other cancer cells (eg, colon, liver). Used to pump out toxins, including chemotherapeutic agents (one mechanism of ↓ responsiveness or resistance to chemotherapy over time).

Psammoma bodies



Laminated, concentric spherules with dystrophic calcification A, PSaMMOMa bodies are seen in:

- Papillary carcinoma of thyroid
- Somatostatinoma
- Meningioma
- Malignant Mesothelioma
- Ovarian serous papillary cystadenocarcinoma
- Prolactinoma (Milk)

Cachexia

Weight loss, muscle atrophy, and fatigue that occur in chronic disease (eg, cancer, AIDS, heart failure, COPD). Mediated by TNF-α, IFN-γ, IL-1, and IL-6.

(c) ketabton.com: The Digital Library

Paraneoplastic syndromes

MANIFESTATION	DESCRIPTION/MECHANISM	MOST COMMONLY ASSOCIATED TUMOR(S)
Musculoskeletal and cutaneous		
Dermatomyositis	Progressive proximal muscle weakness, Gottron papules, heliotrope rash	Adenocarcinomas, especially ovarian
Acanthosis nigricans	Hyperpigmented velvety plaques in axilla and neck	Gastric adenocarcinoma and other visceral malignancies
Sign of Leser-Trélat	Sudden onset of multiple seborrheic keratoses	GI adenocarcinomas and other visceral malignancies
Hypertrophic osteoarthropathy	Abnormal proliferation of skin and bone at distal extremities → clubbing, arthralgia, joint effusions, periostosis of tubular bones	Adenocarcinoma of the lung
Endocrine		
Hypercalcemia	PTHrP	Squamous cell carcinomas of lung, head, and neck; renal, bladder, breast, and ovarian carcinomas
	↑ 1,25-$(OH)_2$ vitamin D_3 (calcitriol)	Lymphoma
Cushing syndrome	↑ ACTH	Small cell lung cancer
Hyponatremia (SIADH)	↑ ADH	
Hematologic		
Polycythemia	↑ Erythropoietin **P**araneoplastic **r**ise to **h**igh **h**ematocrit **l**evels	**P**heochromocytoma, **r**enal cell carcinoma, **H**CC, **h**emangioblastoma, **l**eiomyoma
Pure red cell aplasia	Anemia with low reticulocytes	Thymoma
Good syndrome	Hypogammaglobulinemia	
Trousseau syndrome	Migratory superficial thrombophlebitis	Adenocarcinomas, especially pancreatic
Nonbacterial thrombotic (marantic) endocarditis	Deposition of sterile platelet thrombi on heart valves	
Neuromuscular		
Anti-NMDA receptor encephalitis	Psychiatric disturbance, memory deficits, seizures, dyskinesias, autonomic instability, language dysfunction	Ovarian teratoma
Opsoclonus-myoclonus ataxia syndrome	"Dancing eyes, dancing feet"	Neuroblastoma (children), small cell lung cancer (adults)
Paraneoplastic cerebellar degeneration	Antibodies against antigens in Purkinje cells	Small cell lung cancer (anti-Hu), gynecologic and breast cancers (anti-Yo), and Hodgkin lymphoma (anti-Tr)
Paraneoplastic encephalomyelitis	Antibodies against Hu antigens in neurons	Small cell lung cancer
Lambert-Eaton myasthenic syndrome	Antibodies against presynaptic (P/Q-type) Ca^{2+} channels at NMJ	
Myasthenia gravis	Antibodies against postsynaptic ACh receptors at NMJ	Thymoma

(c) ketabton.com: The Digital Library

HIGH-YIELD PRINCIPLES IN

Pharmacology

"One pill makes you larger, and one pill makes you small."

—Grace Slick

"I was under medication when I made the decision not to burn the tapes."

—Richard Nixon

"I wondher why ye can always read a doctor's bill an' ye niver can read his purscription."

—Finley Peter Dunne

"One of the first duties of the physician is to educate the masses not to take medicine."

—William Osler

Preparation for pharmacology questions is straightforward. Know all the mechanisms, clinical use, and important adverse effects of key drugs and their major variants. Obscure derivatives are low-yield. Learn their classic and distinguishing toxicities as well as major drug-drug interactions. Reviewing associated biochemistry, physiology, and microbiology concepts can be useful while studying pharmacology. The exam has a strong emphasis on ANS, CNS, antimicrobial, and cardiovascular agents as well as on NSAIDs, which are covered throughout the text. Specific drug dosages or trade names are generally not testable. The exam may use graphs to test various pharmacology content, so make sure you are comfortable interpreting them.

- Pharmacokinetics and Pharmacodynamics 230
- Autonomic Drugs 236
- Toxicities and Side Effects 248
- Miscellaneous 253

FAS1_2019_05-Pharmacology.indd 229 11/7/19 4:08 PM

(c) ketabton.com: The Digital Library

▸ PHARMACOLOGY—PHARMACOKINETICS AND PHARMACODYNAMICS

Enzyme kinetics

Michaelis-Menten kinetics

K_m is inversely related to the affinity of the enzyme for its substrate.

V_{max} is directly proportional to the enzyme concentration.

Most enzymatic reactions follow a hyperbolic curve (ie, Michaelis-Menten kinetics); however, enzymatic reactions that exhibit a sigmoid curve usually indicate cooperative kinetics (eg, hemoglobin).

[S] = concentration of substrate; V = velocity.



Effects of enzyme inhibition



Lineweaver-Burk plot

The closer to 0 on the Y-axis, the higher the V_{max}.

The closer to 0 on the X-axis, the higher the K_m.

The higher the K_m, the lower the affinity.

Competitive inhibitors cross each other, whereas noncompetitive inhibitors do not.

Kompetitive inhibitors increase K_m.



Effects of enzyme inhibition



	Competitive inhibitors, reversible	Competitive inhibitors, irreversible	Noncompetitive inhibitors
Resemble substrate	Yes	Yes	No
Overcome by ↑ [S]	Yes	No	No
Bind active site	Yes	Yes	No
Effect on V_{max}	Unchanged	↓	↓
Effect on K_m	↑	Unchanged	Unchanged
Pharmacodynamics	↓ potency	↓ efficacy	↓ efficacy

(c) ketabton.com: The Digital Library

Pharmacokinetics

Bioavailability (F)

Fraction of administered drug reaching systemic circulation unchanged. For an IV dose, F = 100%.

Orally: F typically < 100% due to incomplete absorption and first-pass metabolism. Can be calculated from the area under the curve in a plot of plasma concentration over time.



Volume of distribution (V_d)

Theoretical volume occupied by the total amount of drug in the body relative to its plasma concentration. Apparent V_d of plasma protein–bound drugs can be altered by liver and kidney disease (↓ protein binding, ↑ V_d). Drugs may distribute in more than one compartment.

$$V_d = \frac{\text{amount of drug in the body}}{\text{plasma drug concentration}}$$

V_d	COMPARTMENT	DRUG TYPES
Low	Intravascular	Large/charged molecules; plasma protein bound
Medium	ECF	Small hydrophilic molecules
High	All tissues including fat	Small lipophilic molecules, especially if bound to tissue protein

Clearance (CL)

The volume of plasma cleared of drug per unit time. Clearance may be impaired with defects in cardiac, hepatic, or renal function.

$$CL = \frac{\text{rate of elimination of drug}}{\text{plasma drug concentration}} = V_d \times K_e \text{ (elimination constant)}$$

Half-life ($t_{1/2}$)

The time required to change the amount of drug in the body by ½ during elimination.

In first-order kinetics, a drug infused at a constant rate takes 4–5 half-lives to reach steady state. It takes 3.3 half-lives to reach 90% of the steady-state level.

$$t_{1/2} = \frac{0.7 \times V_d}{CL} \text{ in first-order elimination}$$

# of half-lives	1	2	3	4
% remaining	50%	25%	12.5%	6.25%

Dosage calculations

$$\text{Loading dose} = \frac{C_p \times V_d}{F}$$

$$\text{Maintenance dose} = \frac{C_p \times CL \times \tau}{F}$$

C_p = target plasma concentration at steady state

τ = dosage interval (time between doses), if not administered continuously

In renal or liver disease, maintenance dose ↓ and loading dose is usually unchanged.

Time to steady state depends primarily on $t_{1/2}$ and is independent of dose and dosing frequency.

(c) ketabton.com: The Digital Library

Drug metabolism

Geriatric patients lose phase I first. Patients who are slow acetylators have ↑ side effects from certain drugs because of ↓ rate of metabolism (eg, isoniazid).



Elimination of drugs

Zero-order elimination

Rate of elimination is constant regardless of C_p (ie, constant **amount** of drug eliminated per unit time). C_p ↓ linearly with time. Examples of drugs—**P**henytoin, **E**thanol, and **A**spirin (at high or toxic concentrations).

Capacity-limited elimination.
PEA (a pea is round, shaped like the "0" in **zero**-order).

First-order elimination

Rate of **F**irst-order elimination is directly proportional to the drug concentration (ie, constant **F**raction of drug eliminated per unit time). C_p ↓ exponentially with time. Applies to most drugs.

Flow-dependent elimination.





(c) ketabton.com: The Digital Library

Urine pH and drug elimination

Ionized species are trapped in urine and cleared quickly. Neutral forms can be reabsorbed.

Weak acids

Examples: phenobarbital, methotrexate, aspirin (salicylates). Trapped in basic environments. Treat overdose with sodium bicarbonate to alkalinize urine.

$$\underset{\text{(lipid soluble)}}{RCOOH} \rightleftharpoons \underset{\text{(trapped)}}{RCOO^- + H^+}$$

Weak bases

Examples: TCAs, amphetamines. Trapped in acidic environments. Treat overdose with ammonium chloride to acidify urine.

$$\underset{\text{(trapped)}}{RNH_3^+} \rightleftharpoons \underset{\text{(lipid soluble)}}{RNH_2 + H^+}$$

TCA toxicity is generally treated with sodium bicarbonate to overcome the sodium channel-blocking activity of TCAs, but not for accelerating drug elimination.

pKa

pH at which drugs (weak acid or base) are 50% ionized and 50% nonionized. The pKa represents the strength of the weak acid or base.

Efficacy vs potency

Efficacy

Maximal effect a drug can produce. Represented by the y-value (V_{max}). ↑ y-value = ↑ V_{max} = ↑ efficacy. Unrelated to potency (ie, efficacious drugs can have high or low potency). Partial agonists have less efficacy than full agonists.



Potency

Amount of drug needed for a given effect. Represented by the x-value (EC_{50}). Left shifting = ↓ EC_{50} = ↑ potency = ↓ drug needed. Unrelated to efficacy (ie, potent drugs can have high or low efficacy).



(c) ketabton.com: The Digital Library

Receptor binding



AGONIST WITH	POTENCY	EFFICACY	REMARKS	EXAMPLE
A Competitive antagonist	↓	No change	**Can** be overcome by ↑ agonist concentration	Diazepam (agonist) + flumazenil (competitive antagonist) on $GABA_A$ receptor.
B Noncompetitive antagonist	No change	↓	**Cannot** be overcome by ↑ agonist concentration	Norepinephrine (agonist) + phenoxybenzamine (noncompetitive antagonist) on α-receptors.
C Partial agonist (alone)	Independent	↓	Acts at same site as full agonist	Morphine (full agonist) vs buprenorphine (partial agonist) at opioid μ-receptors.

Therapeutic index

Measurement of drug safety.

$$\frac{TD_{50}}{ED_{50}} = \frac{\text{median toxic dose}}{\text{median effective dose}}$$

Therapeutic window—dosage range that can safely and effectively treat disease.

TITE: **T**herapeutic **I**ndex = TD_{50} / ED_{50}.

Safer drugs have higher TI values. Drugs with lower TI values frequently require monitoring (eg, **W**arfarin, **T**heophylline, **D**igoxin, **A**ntiepileptic drugs, **L**ithium; **W**arning! **T**hese **D**rugs **A**re **L**ethal!).

LD_{50} (lethal median dose) often replaces TD_{50} in animal studies.



(c) ketabton.com: The Digital Library

Types of drug interactions

TERM	DEFINITION	EXAMPLE
Additive	Effect of substances A and B together is equal to the sum of their individual effects	Aspirin and acetaminophen "2 + 2 = 4"
Permissive	Presence of substance A is required for the full effects of substance B	Cortisol on catecholamine responsiveness
Synergistic	Effect of substances A and B together is greater than the sum of their individual effects	Clopidogrel with aspirin "2 + 2 > 4"
Potentiation	Similar to synergism, but drug B with no therapeutic action enhances the therapeutic action of drug A	Carbidopa only blocks enzyme to prevent peripheral conversion of levodopa "2 + 0 > 2"
Antagonistic	Effect of substances A and B together is less than the sum of their individual effects	Ethanol antidote for methanol toxicity "2 + 2 < 4"
Tachyphylactic	Acute decrease in response to a drug after initial/repeated administration	Nitrates, niacin, phenylephrine, LSD, MDMA

(c) ketabton.com: The Digital Library

▶ PHARMACOLOGY—AUTONOMIC DRUGS

Autonomic receptors



Pelvic splanchnic nerves and CNs III, VII, IX and X are part of the parasympathetic nervous system.
Adrenal medulla is directly innervated by preganglionic sympathetic fibers.
Sweat glands are part of the **sympathetic** pathway but are innervated by **chol**inergic fibers (**sympathetic** nervous system results in a "**chol**d" **sweat**).

Acetylcholine receptors

Nicotinic ACh receptors are ligand-gated Na^+/K^+ channels. Two subtypes: N_N (found in autonomic ganglia, adrenal medulla) and N_M (found in neuromuscular junction of skeletal muscle).
Muscarinic ACh receptors are G-protein–coupled receptors that usually act through 2nd messengers. 5 subtypes: $M_{1–5}$ found in heart, smooth muscle, brain, exocrine glands, and on sweat glands (cholinergic sympathetic).

(c) ketabton.com: The Digital Library

Micturition control

Micturition center in pons regulates involuntary bladder function via coordination of sympathetic and parasympathetic nervous systems.

- ⊕ sympathetic → ↑ urinary retention
- ⊕ parasympathetic → ↑ urine voiding. Some autonomic drugs act on smooth muscle receptors to treat bladder dysfunction.



DRUGS	MECHANISM	USE
1 **Muscarinic antagonists** (eg, oxybutynin)	⊖ M_3 receptor → relaxation of detrusor smooth muscle → ↓ detrusor overactivity	Urgency incontinence
1 **Muscarinic agonists** (eg, bethanechol)	⊕ M_3 receptor → contraction of detrusor smooth muscle → ↑ bladder emptying	Urinary retention
2 **Sympathomimetics** (eg, mirabegron)	⊕ β_3 receptor → relaxation of detrusor smooth muscle → ↑ bladder capacity	Urgency incontinence
3 **α_1-blockers** (eg, tamsulosin)	⊖ α_1-receptor → relaxation of smooth muscle (bladder neck, prostate) → ↓ urinary obstruction	BPH

(c) ketabton.com: The Digital Library

G-protein–linked second messengers

RECEPTOR	G-PROTEIN CLASS	MAJOR FUNCTIONS
Adrenergic		
α_1	q	↑ vascular smooth muscle contraction, ↑ pupillary dilator muscle contraction (mydriasis), ↑ intestinal and bladder sphincter muscle contraction
α_2	i	↓ sympathetic (adrenergic) outflow, ↓ insulin release, ↓ lipolysis, ↑ platelet aggregation, ↓ aqueous humor production
β_1	s	↑ heart rate, ↑ contractility (one heart), ↑ renin release, ↑ lipolysis
β_2	s	Vasodilation, bronchodilation (two lungs), ↑ lipolysis, ↑ insulin release, ↑ glycogenolysis, ↓ uterine tone (tocolysis), ↑ aqueous humor production, ↑ cellular K^+ uptake
β_3	s	↑ lipolysis, ↑ thermogenesis in skeletal muscle, ↑ bladder relaxation
Cholinergic		
M_1	q	Mediates higher cognitive functions, stimulates enteric nervous system
M_2	i	↓ heart rate and contractility of atria
M_3	q	↑ exocrine gland secretions (eg, lacrimal, sweat, salivary, gastric acid), ↑ gut peristalsis, ↑ bladder contraction, bronchoconstriction, ↑ pupillary sphincter muscle contraction (miosis), ciliary muscle contraction (accommodation), ↑ insulin release, endothelium-mediated vasodilation
Dopamine		
D_1	s	Relaxes renal vascular smooth muscle, activates direct pathway of striatum
D_2	i	Modulates transmitter release, especially in brain, inhibits indirect pathway of striatum
Histamine		
H_1	q	↑ nasal and bronchial mucus production, ↑ vascular permeability, bronchoconstriction, pruritus, pain
H_2	s	↑ gastric acid secretion
Vasopressin		
V_1	q	↑ vascular smooth muscle contraction
V_2	s	↑ H_2O permeability and reabsorption via upregulating aquaporin-2 in collecting twobules (tubules) of kidney, ↑ release of vWF

"After qisses (kisses), you get a qiq (kick) out of siq (sick) sqs (super qinky sex)."



(c) ketabton.com: The Digital Library

Autonomic drugs

Release of norepinephrine from a sympathetic nerve ending is modulated by NE itself, acting on presynaptic α_2-autoreceptors → negative feedback.

Amphetamines use the NE transporter (NET) to enter the presynaptic terminal, where they utilize the vesicular monoamine transporter (VMAT) to enter neurosecretory vesicles. This displaces NE from the vesicles. Once NE reaches a concentration threshold within the presynaptic terminal, the action of NET is reversed, and NE is expelled into the synaptic cleft, contributing to the characteristics and effects of ↑ NE observed in patients taking amphetamines.



(c) ketabton.com: The Digital Library

Cholinomimetic agents

Watch for exacerbation of COPD, asthma, and peptic ulcers in susceptible patients.

DRUG	ACTION	APPLICATIONS
Direct agonists		
Bethanechol	Activates bladder smooth muscle; resistant to AChE. No nicotinic activity. "Bethany, call me to activate your bladder."	Urinary retention.
Carbachol	Carbon copy of acetylcholine (but resistant to AChE).	Constricts pupil and relieves intraocular pressure in open-angle glaucoma.
Methacholine	Stimulates muscarinic receptors in airway when inhaled.	Challenge test for diagnosis of asthma.
Pilocarpine	Contracts ciliary muscle of eye (open-angle glaucoma), pupillary sphincter (closed-angle glaucoma); resistant to AChE, can cross blood-brain barrier (tertiary amine). "You cry, drool, and sweat on your 'pilow.'"	Potent stimulator of sweat, tears, and saliva Open-angle and closed-angle glaucoma, xerostomia (Sjögren syndrome).
Indirect agonists (anticholinesterases)		
Donepezil, rivastigmine, galantamine	↑ ACh.	1st line for Alzheimer disease (Dona Riva dances at the gala).
Edrophonium	↑ ACh.	Historically used to diagnose myasthenia gravis; replaced by anti-AChR Ab (anti-acetylcholine receptor antibody) test.
Neostigmine	↑ ACh. Neo CNS = No CNS penetration (quaternary amine).	Postoperative and neurogenic ileus and urinary retention, myasthenia gravis, reversal of neuromuscular junction blockade (postoperative).
Physostigmine	↑ ACh. Phreely (freely) crosses blood-brain barrier → CNS (tertiary amine).	Antidote for anticholinergic toxicity; physostigmine "phyxes" atropine overdose.
Pyridostigmine	↑ ACh; ↑ muscle strength. Used with glycopyrrolate, hyoscyamine, or propantheline to control pyridostigmine side effects. Pyridostigmine gets rid of myasthenia gravis.	Myasthenia gravis (long acting); does not penetrate CNS (quaternary amine).

Anticholinesterase poisoning

Often due to organophosphates (eg, parathion) that irreversibly inhibit AChE. Organophosphates commonly used as insecticides; poisoning usually seen in farmers.

Muscarinic effects	Diarrhea, Urination, Miosis, Bronchospasm, Bradycardia, Emesis, Lacrimation, Sweating, Salivation.	DUMBBELSS. Reversed by atropine, a competitive inhibitor. Atropine can cross BBB to relieve CNS symptoms.
Nicotinic effects	Neuromuscular blockade (mechanism similar to succinylcholine).	Reversed by pralidoxime, regenerates AChE via dephosphorylation if given early. Pralidoxime (quaternary amine) does not readily cross BBB.
CNS effects	Respiratory depression, lethargy, seizures, coma.	

(c) ketabton.com: The Digital Library

Muscarinic antagonists

DRUGS	ORGAN SYSTEMS	APPLICATIONS
Atropine, homatropine, tropicamide	Eye	Produce mydriasis and cycloplegia.
Benztropine, trihexyphenidyl	CNS	Parkinson disease ("park my Benz"). Acute dystonia.
Glycopyrrolate	GI, respiratory	Parenteral: preoperative use to reduce airway secretions. Oral: drooling, peptic ulcer.
Hyoscyamine, dicyclomine	GI	Antispasmodics for irritable bowel syndrome.
Ipratropium, tiotropium	Respiratory	COPD, asthma ("I pray I can breathe soon!").
Oxybutynin, solifenacin, tolterodine	Genitourinary	Reduce bladder spasms and urge urinary incontinence (overactive bladder).
Scopolamine	CNS	Motion sickness.

Atropine

Muscarinic antagonist. Used to treat bradycardia and for ophthalmic applications.

ORGAN SYSTEM	ACTION	NOTES
Eye	↑ pupil dilation, cycloplegia	Blocks muscarinic effects (DUMBBELSS) of anticholinesterases, but not the nicotinic effects.
Airway	Bronchodilation, ↓ secretions	
Stomach	↓ acid secretion	
Gut	↓ motility	
Bladder	↓ urgency in cystitis	
ADVERSE EFFECTS	↑ body temperature (due to ↓ sweating); ↑ HR; dry mouth; dry, flushed skin; cycloplegia; constipation; disorientation Can cause acute angle-closure glaucoma in elderly (due to mydriasis), urinary retention in men with prostatic hyperplasia, and hyperthermia in infants.	Side effects: Hot as a hare Fast as a fiddle Dry as a bone Red as a beet Blind as a bat Mad as a hatter Full as a flask Jimson weed *(Datura)* → gardener's pupil (mydriasis due to plant alkaloids)

(c) ketabton.com: The Digital Library

Sympathomimetics

DRUG	ACTION	HEMODYNAMIC CHANGES	APPLICATIONS
Direct sympathomimetics			
Albuterol, salmeterol, terbutaline	$\beta_2 > \beta_1$	↑ HR (little effect)	Albuterol for Acute asthma/COPD. Salmeterol for Serial (long-term) asthma/COPD. Terbutaline for acute bronchospasm in asthma and tocolysis.
Dobutamine	$\beta_1 > \beta_2, \alpha$	↔/↓ BP, ↑ HR, ↑ CO	Heart failure (HF), cardiogenic shock (inotropic > chronotropic), cardiac stress testing.
Dopamine	$D_1 = D_2 > \beta > \alpha$	↑ BP (high dose), ↑ HR, ↑ CO	Unstable bradycardia, HF, shock; inotropic and chronotropic effects at lower doses due to β effects; vasoconstriction at high doses due to α effects.
Epinephrine	$\beta > \alpha$	↑ BP (high dose), ↑ HR, ↑ CO	Anaphylaxis, asthma, open-angle glaucoma; α effects predominate at high doses. Significantly stronger effect at β_2-receptor than norepinephrine.
Fenoldopam	D_1	↓ BP (vasodilation), ↑ HR, ↑ CO	Postoperative hypertension, hypertensive crisis. Vasodilator (coronary, peripheral, renal, and splanchnic). Promotes natriuresis. Can cause hypotension and tachycardia.
Isoproterenol	$\beta_1 = \beta_2$	↓ BP (vasodilation), ↑ HR, ↑ CO	Electrophysiologic evaluation of tachyarrhythmias. Can worsen ischemia. Has negligible α effect.
Midodrine	α_1	↑ BP (vasoconstriction), ↓ HR, ↔/↓ CO	Autonomic insufficiency and postural hypotension. May exacerbate supine hypertension.
Mirabegron	β_3		Urinary urgency or incontinence or overactive bladder. Think "mirab3gron."
Norepinephrine	$\alpha_1 > \alpha_2 > \beta_1$	↑ BP, ↑ HR, ↔/↑ CO	Hypotension, septic shock.
Phenylephrine	$\alpha_1 > \alpha_2$	↑ BP (vasoconstriction), ↓ HR, ↔/↓ CO	Hypotension (vasoconstrictor), ocular procedures (mydriatic), rhinitis (decongestant), ischemic priapism.
Indirect sympathomimetics			
Amphetamine	Indirect general agonist, reuptake inhibitor, also releases stored catecholamines		Narcolepsy, obesity, ADHD.
Cocaine	Indirect general agonist, reuptake inhibitor		Causes vasoconstriction and local anesthesia. Caution when giving β-blockers if cocaine intoxication is suspected (can lead to unopposed α_1 activation → extreme hypertension, coronary vasospasm).
Ephedrine	Indirect general agonist, releases stored catecholamines		Nasal decongestion (pseudoephedrine), urinary incontinence, hypotension.

(c) ketabton.com: The Digital Library

Norepinephrine vs isoproterenol

NE ↑ systolic and diastolic pressures as a result of α_1-mediated vasoconstriction → ↑ mean arterial pressure → reflex bradycardia. However, isoproterenol (rarely used) has little α effect but causes β_2-mediated vasodilation, resulting in ↓ mean arterial pressure and ↑ heart rate through β_1 and reflex activity.



Sympatholytics (α_2-agonists)

DRUG	APPLICATIONS	ADVERSE EFFECTS
Clonidine, guanfacine	Hypertensive urgency (limited situations), ADHD, Tourette syndrome, symptom control in opioid withdrawal	CNS depression, bradycardia, hypotension, respiratory depression, miosis, rebound hypertension with abrupt cessation
α-methyldopa	Hypertension in pregnancy	Direct Coombs ⊕ hemolysis, drug-induced lupus, hyperprolactinemia
Tizanidine	Relief of spasticity	Hypotension, weakness, xerostomia

(c) ketabton.com: The Digital Library

α-blockers

DRUG	APPLICATIONS	ADVERSE EFFECTS
Nonselective		
Phenoxybenzamine	Irreversible. Pheochromocytoma (used preoperatively) to prevent catecholamine (hypertensive) crisis	Orthostatic hypotension, reflex tachycardia
Phentolamine	Reversible. Given to patients on MAO inhibitors who eat tyramine-containing foods and for severe cocaine-induced hypertension (2nd line)	
α_1 selective (-osin ending)		
Prazosin, terazosin, doxazosin, tamsulosin	Urinary symptoms of BPH; PTSD (prazosin); hypertension (except tamsulosin)	1st-dose orthostatic hypotension, dizziness, headache
α_2 selective		
Mirtazapine	Depression	Sedation, ↑ serum cholesterol, ↑ appetite



Epinephrine response exhibits reversal of mean arterial pressure from a net increase (the α response) to a net decrease (the β_2 response).



Phenylephrine response is suppressed but not reversed because it is a "pure" α-agonist (lacks β-agonist properties).

FAS1_2019_05-Pharmacology.indd 244 11/7/19 4:08 PM

(c) ketabton.com: The Digital Library

β-blockers Acebutolol, atenolol, betaxolol, bisoprolol, carvedilol, esmolol, labetalol, metoprolol, nadolol, nebivolol, pindolol, propranolol, timolol.

APPLICATION	ACTIONS	NOTES/EXAMPLES
Angina pectoris	↓ heart rate and contractility → ↓ O_2 consumption	
Glaucoma	↓ production of aqueous humor	Timolol
Heart failure	↓ mortality	**B**isoprolol, **C**arvedilol, **M**etoprolol (β-blockers **C**urb **M**ortality)
Hypertension	↓ cardiac output, ↓ renin secretion (due to β_1-receptor blockade on JG cells)	
Hyperthyroidism/ thyroid storm	Symptom control (↓ heart rate, ↓ tremor)	Propranolol
Hypertrophic cardiomyopathy	↓ heart rate → ↑ filling time, relieving obstruction	
Myocardial infarction	↓ O_2 demand (short-term), ↓ mortality (long-term)	
Supraventricular tachycardia	↓ AV conduction velocity (class II antiarrhythmic)	Metoprolol, esmolol
Variceal bleeding	↓ hepatic venous pressure gradient and portal hypertension (prophylactic use)	Nadolol, propranolol, carvedilol
ADVERSE EFFECTS	Erectile dysfunction, cardiovascular (bradycardia, AV block, HF), CNS (seizures, sleep alterations), dyslipidemia (metoprolol), and asthma/COPD exacerbations	Use of β-blockers for acute cocaine-associated chest pain remains controversial due to unsubstantiated concern for unopposed α-adrenergic stimulation
SELECTIVITY	β_1-selective antagonists ($\beta_1 > \beta_2$)—**a**cebutolol (partial agonist), **a**tenolol, **b**etaxolol, **b**isoprolol, **e**smolol, **m**etoprolol	Selective antagonists mostly go from **A** to **M** (β_1 with **1**st half of alphabet)
	Nonselective antagonists ($\beta_1 = \beta_2$)—**n**adolol, **p**indolol (partial agonist), **p**ropranolol, **t**imolol	**N**on**Z**elective antagonists mostly go from **N** to **Z** (β_2 with **2**nd half of alphabet)
	Nonselective α- and β-antagonists—carved**ilol**, labet**alol**	Nonselective α- and β-antagonists have **modified suffixes** (instead of "-olol")
	Nebiv**o**lol combines cardiac-selective β_1-adrenergic blockade with stimulation of β_3-receptors (activate **n**itric **o**xide synthase in the vasculature and ↓ SVR)	**N**ebiv**o**lol increases **NO**

(c) ketabton.com: The Digital Library

Phosphodiesterase inhibitors

Phosphodiesterase (PDE) inhibitors inhibit PDE, which catalyzes the hydrolysis of cAMP and/or cGMP, and thereby increase cAMP and/or cGMP. These inhibitors have varying specificity for PDE isoforms and thus have different clinical uses.

TYPE OF INHIBITOR	MECHANISM OF ACTION	CLINICAL USES	ADVERSE EFFECTS
Nonspecific PDE inhibitor Theophylline	↓ cAMP hydrolysis → ↑ cAMP → bronchial smooth muscle relaxation → bronchodilation	COPD/asthma (rarely used)	Cardiotoxicity (eg, tachycardia, arrhythmia), neurotoxicity (eg, headache), abdominal pain
PDE-5 inhibitors Sildena**fil**, varden**afil**, tadala**fil**, avana**fil**	↓ hydrolysis of cGMP → ↑ cGMP → ↑ smooth muscle relaxation by enhancing NO activity → pulmonary vasodilation and ↑ blood flow in corpus cavernosum **fil**ls the penis	Erectile dysfunction Pulmonary hypertension BPH (tadalafil only)	Facial flushing, headache, dyspepsia, hypotension in patients taking nitrates; "**H**ot and sweaty," then **H**eadache, **H**eartburn, **H**ypotension Sildenafil only: cyanopia (blue-tinted vision) via inhibition of PDE-6 in retina
PDE-4 inhibitor Roflumilast	↑ cAMP in neutrophils, granulocytes, and bronchial epithelium	Severe COPD	Abdominal pain, weight loss, mental disorders (eg, depression)
PDE-3 inhibitor Milrinone	In cardiomyocytes: ↑ cAMP → ↑ Ca^{2+} influx → ↑ ionotropy and chronotropy In vascular smooth muscle: ↑ cAMP → MLCK inhibition → vasodilation → ↓ preload and afterload	Acute decompensated HF with cardiogenic shock	Tachycardia, ventricular arrhythmias (thus not for chronic use), hypotension
"Platelet inhibitors" Cilostazol[a] Dipyridamole[b]	In platelets: ↑ cAMP → inhibition of platelet aggregation	Intermittent claudication Stroke or TIA prevention (with aspirin) Cardiac stress testing (dipyridamole only, due to coronary vasodilation) Prevention of coronary stent restenosis	Nausea, headache, facial flushing, hypotension, abdominal pain

[a]Cilostazol is a PDE-3 inhibitor, but due to its indications is categorized as a platelet inhibitor together with dipyridamole.
[b]Dipyridamole is a nonspecific PDE inhibitor, leading to inhibition of platelet aggregation. It also prevents adenosine reuptake by platelets → ↑ extracellular adenosine → ↑ vasodilation.

(c) ketabton.com: The Digital Library

Ingested seafood toxins

Toxin actions include **H**istamine release, **T**otal block of Na^+ channels, or opening of Na^+ channels to **C**ause depolarization.

TOXIN	SOURCE	ACTION	SYMPTOMS	TREATMENT
Histamine (scombroid poisoning)	Spoiled dark-meat fish such as tuna, mahi-mahi, mackerel, and bonito	Bacterial histidine decarboxylase converts histidine to histamine Frequently misdiagnosed as fish allergy	Mimics anaphylaxis: acute burning sensation of mouth, flushing of face, erythema, urticaria, itching May progress to bronchospasm, angioedema, hypotension	Antihistamines Albuterol and epinephrine if needed
Tetrodotoxin	Pufferfish	Highly potent toxin; binds fast voltage-gated Na^+ channels in nerve tissue, preventing depolarization	Nausea, diarrhea, paresthesias, weakness, dizziness, loss of reflexes	Supportive
Ciguatoxin	Reef fish such as barracuda, snapper, and moray eel	Opens Na^+ channels, causing depolarization	Nausea, vomiting, diarrhea; perioral numbness; reversal of hot and cold sensations; bradycardia, heart block, hypotension	Supportive

Beers criteria

Widely used criteria developed to reduce potentially inappropriate prescribing and harmful polypharmacy in the geriatric population. Includes > 50 medications that should be avoided in elderly patients due to ↓ efficacy and/or ↑ risk of adverse events. Examples:

- α-blockers (↑ risk of hypotension)
- Anticholinergics, antidepressants, antihistamines, opioids (↑ risk of delirium, sedation, falls, constipation, urinary retention)
- Benzodiazepines (↑ risk of delirium, sedation, falls)
- NSAIDs (↑ risk of GI bleeding, especially with concomitant anticoagulation)
- PPIs (↑ risk of *C difficile* infection)

(c) ketabton.com: The Digital Library

▸ PHARMACOLOGY—TOXICITIES AND SIDE EFFECTS

Specific toxicity treatments

TOXIN	TREATMENT
Acetaminophen	*N*-acetylcysteine (replenishes glutathione)
AChE inhibitors, organophosphates	Atropine > pralidoxime
Antimuscarinic, anticholinergic agents	Physostigmine (crosses BBB), control hyperthermia
Arsenic	Dimercaprol, succimer
Benzodiazepines	Flumazenil
β-blockers	Atropine, glucagon, saline
Carbon monoxide	100% O_2, hyperbaric O_2
Copper	"**Penny**"cillamine (penicillamine), **tri**entine (**copper penny** × **3**)
Cyanide	Hydroxocobalamin, nitrites + sodium thiosulfate
Digitalis (digoxin)	Digoxin-specific antibody fragments
Heparin	Protamine sulfate
Iron (**Fe**)	**Fe**roxamine, **fe**rasirox, **fe**riprone
Lead	Calcium disodium EDTA, dimercaprol, succimer, penicillamine
Mercury	Di**mer**caprol, succi**mer**
Methanol, ethylene glycol (antifreeze)	Fomepizole > ethanol, dialysis
Methemoglobin	**Meth**ylene blue, vitamin C (reducing agent)
Opi**O**ids	Nal**O**x**O**ne
Salicylates	$NaHCO_3$ (alkalinize urine), dialysis
TCAs	$NaHCO_3$ (stabilizes cardiac cell membrane)
Warfarin	Vitamin K (delayed effect), PCC (prothrombin complex concentrate)/FFP (immediate effect)

Drug reactions—cardiovascular

DRUG REACTION	CAUSAL AGENTS
Coronary vasospasm	**C**ocaine, **A**mphetamines, **S**umatriptan, **E**rgot alkaloids (**CASE**)
Cutaneous flushing	**V**ancomycin, **A**denosine, **N**iacin, **C**a^{2+} channel blockers, **E**chinocandins, **N**itrates (**flush**ed from **VANCEN** [dancing]) Red man syndrome—rate-dependent infusion reaction to vancomycin causing widespread pruritic erythema due to histamine release. Manage with diphenhydramine, slower infusion rate.
Dilated cardiomyopathy	Anthracyclines (eg, **D**oxorubicin, **D**aunorubicin); prevent with **D**exrazoxane
Torsades de pointes	Agents that prolong QT interval: anti**A**rrhythmics (class IA, III), anti**B**iotics (eg, macrolides), anti"**C**"ychotics (eg, ziprasidone), anti**D**epressants (eg, TCAs), anti**E**metics (eg, ondansetron) (**ABCDE**)

(c) ketabton.com: The Digital Library

Drug reactions—endocrine/reproductive

DRUG REACTION	CAUSAL AGENTS	NOTES
Adrenocortical insufficiency	HPA suppression 2° to glucocorticoid withdrawal	
Diabetes insipidus	Lithium, demeclocycline	
Hot flashes	SERMs (eg, tamoxifen, clomiphene, raloxifene)	
Hyperglycemia	**T**acrolimus, **P**rotease inhibitors, **N**iacin, **H**CTZ, **C**orticosteroids	**T**he **P**eople **N**eed **H**ard **C**andies
Hyperprolactinemia	Typical antipsychotics (eg, haloperidol), atypical antipsychotics (eg, risperidone), metoclopramide, methyldopa, reserpine	Presents with hypogonadism (eg, infertility, amenorrhea, erectile dysfunction) and galactorrhea
Hyperthyroidism	Amiodarone, iodine	
Hypothyroidism	**AM**iodarone, **SU**lfonamides, **L**ithium	I **AM SU**ddenly **L**ethargic
SIADH	**C**arbamazepine, **C**yclophosphamide, **SS**RIs	**C**an't **C**oncentrate **S**erum **S**odium

Drug reactions—gastrointestinal

DRUG REACTION	CAUSAL AGENTS	NOTES
Acute cholestatic hepatitis, jaundice	Macrolides (eg, erythromycin)	
Diarrhea	Acamprosate, antidiabetic agents (acarbose, metformin, pramlintide), colchicine, cholinesterase inhibitors, lipid-lowering agents (eg, ezetimibe, orlistat), macrolides (eg, erythromycin), SSRIs, chemotherapy (eg, irinotecan)	
Focal to massive hepatic necrosis	**H**alothane, **A***manita phalloides* (death cap mushroom), **V**alproic acid, **Ac**etaminophen	Liver "**HAVAc**"
Hepatitis	Rifampin, isoniazid, pyrazinamide, statins, fibrates	
Pancreatitis	**D**idanosine, **C**orticosteroids, **A**lcohol, **V**alproic acid, **A**zathioprine, **D**iuretics (eg, furosemide, HCTZ)	**D**rugs **C**ausing **A** **V**iolent **A**bdominal **D**istress
Pill-induced esophagitis	Bisphosphonates, ferrous sulfate, NSAIDs, potassium chloride, tetracyclines	Caustic effect minimized with upright posture and adequate water ingestion
Pseudomembranous colitis	Ampicillin, cephalosporins, clindamycin, fluoroquinolones, PPIs	Antibiotics predispose to superinfection by resistant *C difficile*

(c) ketabton.com: The Digital Library

Drug reactions—hematologic

DRUG REACTION	CAUSAL AGENTS	NOTES
Agranulocytosis	Dapsone, Clozapine, Carbamazepine, Propylthiouracil, Methimazole, Colchicine, Ganciclovir	Drugs Can Cause Pretty Major Collapse of Granulocytes
Aplastic anemia	Carbamazepine, Methimazole, NSAIDs, Benzene, Chloramphenicol, Propylthiouracil	Can't Make New Blood Cells Properly
Direct Coombs ⊕ hemolytic anemia	Penicillin, methylDopa, Cephalosporins	P Diddy Coombs
Drug reaction with eosinophilia and systemic symptoms (DRESS)	Allopurinol, anticonvulsants, antibiotics, sulfa drugs	Potentially fatal delayed hypersensitivity reaction. Latency period (2- 8 weeks), then fever, morbilliform skin rash, frequent multiorgan involvement. Treatment: withdrawal of offending drug, corticosteroids
Gray baby syndrome	Chloramphenicol	
Hemolysis in G6PD deficiency	Isoniazid, Sulfonamides, Dapsone, Primaquine, Aspirin, Ibuprofen, Nitrofurantoin	Hemolysis IS D PAIN
Megaloblastic anemia	Hydroxyurea, Phenytoin, Methotrexate, Sulfa drugs	You're having a mega blast with PMS
Thrombocytopenia	Heparin, vancomycin, linezolid, quinidine, indinavir, ganciclovir, abciximab	
Thrombotic complications	Combined oral contraceptives, hormone replacement therapy, SERMs (eg, tamoxifen)	Estrogen-mediated side effect

Drug reactions—musculoskeletal/skin/connective tissue

DRUG REACTION	CAUSAL AGENTS	NOTES
Drug-induced lupus	Methyldopa, Minocycline, Hydralazine, Isoniazid, Phenytoin, Sulfa drugs, Etanercept, Procainamide	Lupus Makes My HIPS Extremely Painful
Fat redistribution	Protease inhibitors, Glucocorticoids	Fat PiG
Gingival hyperplasia	Cyclosporine, Ca^{2+} channel blockers, Phenytoin	Can Cause Puffy gums
Hyperuricemia (gout)	Pyrazinamide, Thiazides, Furosemide, Niacin, Cyclosporine	Painful Tophi and Feet Need Care
Myopathy	Statins, fibrates, niacin, colchicine, daptomycin, hydroxychloroquine, interferon-α, penicillamine, glucocorticoids	
Osteoporosis	Corticosteroids, depot medroxyprogesterone acetate, GnRH agonists, aromatase inhibitors, anticonvulsants, heparin, PPIs	
Photosensitivity	Sulfonamides, Amiodarone, Tetracyclines, 5-FU	SAT For Photo
Rash (Stevens-Johnson syndrome)	Anti-epileptic drugs (especially lamotrigine), allopurinol, sulfa drugs, penicillin	Steven Johnson has epileptic allergy to sulfa drugs and penicillin
Teeth discoloration	Tetracyclines	Teethracyclines
Tendon/cartilage damage	Fluoroquinolones	

(c) ketabton.com: The Digital Library

Drug reactions—neurologic

DRUG REACTION	CAUSAL AGENTS	NOTES
Cinchonism	Quinidine, quinine	Can present with tinnitus, hearing/vision loss, psychosis, and cognitive impairment
Parkinson-like syndrome	Antipsychotics, Reserpine, Metoclopramide	Cogwheel rigidity of ARM
Peripheral neuropathy	Isoniazid, phenytoin, platinum agents (eg, cisplatin), vincristine	
Idiopathic intracranial hypertension	Growth hormones, tetracyclines, vitamin A	
Seizures	Isoniazid, Bupropion, Imipenem/cilastatin, Tramadol, Enflurane	With seizures, I BITE my tongue
Tardive dyskinesia	Antipsychotics, metoclopramide	
Visual disturbance	Topiramate (blurred vision/diplopia, haloes), Digoxin (yellow-tinged vision), Isoniazid (optic neuritis), Vigabatrin (bilateral visual field defects), PDE-5 inhibitors (blue-tinged vision), Ethambutol (color vision changes)	These Drugs Irritate Very Precious Eyes

Drug reactions—renal/genitourinary

DRUG REACTION	CAUSAL AGENTS	NOTES
Fanconi syndrome	Cisplatin, ifosfamide, expired tetracyclines, tenofovir	
Hemorrhagic cystitis	Cyclophosphamide, ifosfamide	Prevent by coadministering with mesna
Interstitial nephritis	Diuretics (Pee), NSAIDs (Pain-free), Penicillins and cephalosporins, PPIs, rifamPin, and sulfa drugs	Remember the 5 P's

Drug reactions—respiratory

DRUG REACTION	CAUSAL AGENTS	NOTES
Dry cough	ACE inhibitors	
Pulmonary fibrosis	Methotrexate, Nitrofurantoin, Carmustine, Bleomycin, Busulfan, Amiodarone	My Nose Cannot Breathe Bad Air

Drug reactions—multiorgan

DRUG REACTION	CAUSAL AGENTS	NOTES
Antimuscarinic	Atropine, TCAs, H_1-blockers, antipsychotics	
Disulfiram-like reaction	1st-generation Sulfonylureas, Procarbazine, certain Cephalosporins, Griseofulvin, Metronidazole	Sorry Pals, Can't Go Mingle
Nephrotoxicity/ ototoxicity	Loop diuretics, Aminoglycosides, cisPlatin, Vancomycin, amphoTERicin B	Listen And Pee Very TERriBly Cisplatin toxicity may respond to amifostine

(c) ketabton.com: The Digital Library

Drugs affecting pupil size

↑ pupil size	↓ pupil size
Anticholinergics (eg, atropine, TCAs, tropicamide, scopolamine, antihistamines)	Sympatholytics (eg, α_2-agonists)
Drugs of abuse (eg, amphetamines, cocaine, LSD), meperidine	Drugs of abuse (eg, heroin/opioids)
Sympathomimetics	Parasympathomimetics (eg, pilocarpine), organophosphates

Cytochrome P-450 interactions (selected)

Inducers (+)	Substrates	Inhibitors (–)
Modafinil	Warfarin	Sodium valproate
Chronic alcohol use	Anti-epileptics	Isoniazid
St. John's wort	Theophylline	Cimetidine
Phenytoin	OCPs	Ketoconazole
Phenobarbital		Fluconazole
Nevirapine		Acute alcohol abuse
Rifampin		Chloramphenicol
Griseofulvin		Erythromycin/clarithromycin
Carbamazepine		Sulfonamides
		Ciprofloxacin
		Omeprazole
		Metronidazole
		Amiodarone
		Ritonavir
		Grapefruit juice
Most chronic alcoholics Steal Phen-Phen and Never Refuse Greasy Carbs	War Against The OCPs	SICKFACES.COM (when I Am Really drinking Grapefruit juice)

Sulfa drugs

Sulfonamide antibiotics, Sulfasalazine, Probenecid, Furosemide, Acetazolamide, Celecoxib, Thiazides, Sulfonylureas.

Patients with sulfa allergies may develop fever, urinary tract infection, Stevens-Johnson syndrome, hemolytic anemia, thrombocytopenia, agranulocytosis, acute interstitial nephritis, and urticaria (hives).

Scary Sulfa Pharm FACTS

(c) ketabton.com: The Digital Library

▸ PHARMACOLOGY—MISCELLANEOUS

Drug names

ENDING	CATEGORY	EXAMPLE
Antimicrobial		
-bendazole	Antiparasitic/antihelminthic	Mebendazole
-cillin	Transpeptidase inhibitor	Ampicillin
-conazole	Ergosterol synthesis inhibitor	Ketoconazole
-cycline	Protein synthesis inhibitor	Tetracycline
-ivir	Neuraminidase inhibitor	Oseltamivir
-navir	Protease inhibitor	Ritonavir
-ovir	Viral DNA polymerase inhibitor	Acyclovir
-tegravir	Integrase inhibitor	Elvitegravir, raltegravir
-thromycin	Macrolide antibiotic	Azithromycin
CNS		
-apine, -idone	Atypical antipsychotic	Quetiapine, risperidone
-azine	Typical antipsychotic	Thioridazine
-barbital	Barbiturate	Phenobarbital
-ipramine, -triptyline	TCA	Imipramine, amitriptyline
-triptan	5-$HT_{1B/1D}$ agonist	Sumatriptan
-zepam, -zolam	Benzodiazepine	Diazepam, alprazolam
Autonomic		
-chol	Cholinergic agonist	Bethanechol, carbachol
-olol	β-blocker	Propranolol
-stigmine	AChE inhibitor	Neostigmine
-terol	β_2-agonist	Albuterol
-zosin	α_1-blocker	Prazosin
Cardiovascular		
-afil	PDE-5 inhibitor	Sildenafil
-dipine	Dihydropyridine Ca^{2+} channel blocker	Amlodipine
-pril	ACE inhibitor	Captopril
-sartan	Angiotensin-II receptor blocker	Losartan
-xaban	Direct factor **Xa** inhibitor	Api**xa**ban, edo**xa**ban, rivaro**xa**ban
Metabolic		
-gliflozin	SGLT-2 inhibitor	Dapagliflozin, canagliflozin
-glinide	Meglitinide	Repaglinide, nateglinide
-gliptin	DPP-4 inhibitor	Sitagliptin
-glitazone	PPAR-γ activator	Rosiglitazone
-glutide	GLP-1 analog	Liraglutide, albiglutide

(c) ketabton.com: The Digital Library

Drug names *(continued)*

ENDING	CATEGORY	EXAMPLE
Other		
-dronate	Bisphosphonate	Alendronate
-prazole	Proton pump inhibitor	Omeprazole
-prost	Prostaglandin analog	Latanoprost
-sentan	Endothelin receptor antagonist	Bosentan
-tidine	H_2-antagonist	Cimetidine
-vaptan	ADH antagonist	Tolvaptan

Biologic agents

ENDING	CATEGORY	EXAMPLE
Monoclonal antibodies (-mab)—target overexpressed cell surface receptors		
-ximab	Chimeric human-mouse monoclonal antibody	Rituximab
-zumab	Humanized mouse monoclonal antibody	Bevacizumab
-umab	Human monoclonal antibody	Denosumab
Small molecule inhibitors (-ib)—target intracellular molecules		
-tinib	Tyrosine kinase inhibitor	Imatinib
-zomib	Proteasome inhibitor	Bortezomib
-ciclib	Cyclin-dependent kinase inhibitor	Palbociclib
Receptor fusion proteins (-cept)		
-cept	TNF-α antagonist	Etanercept
Interleukin receptor modulators (-kin)—agonists and antagonists of interleukin receptors		
-leukin	IL-2 agonist/analog	Aldesleukin
-kinra	Interleukin receptor antagonist	Anakinra

(c) ketabton.com: The Digital Library

HIGH-YIELD PRINCIPLES IN

Public Health Sciences

"Medicine is a science of uncertainty and an art of probability."
—William Osler

"There are two kinds of statistics: the kind you look up and the kind you make up."
—Rex Stout

"On a long enough timeline, the survival rate for everyone drops to zero."
—Chuck Palahniuk

"There are three kinds of lies: lies, damned lies, and statistics."
—Mark Twain

A heterogenous mix of epidemiology, biostatistics, ethics, law, healthcare delivery, patient safety, quality improvement, and more falls under the heading of public health sciences. Biostatistics and epidemiology are the foundations of evidence-based medicine and are very high yield. Make sure you can quickly apply biostatistical equations such as sensitivity, specificity, and predictive values in a problem-solving format. Also, know how to set up your own 2 × 2 tables. Quality improvement and patient safety topics were introduced a few years ago on the exam and represent trends in health system science. Medical ethics questions often require application of principles. Typically, you are presented with a patient scenario and then asked how you would respond.

- Epidemiology and Biostatistics 256
- Ethics 265
- The Well Patient 270
- Healthcare Delivery 270
- Quality and Safety 273

FAS1_2019_06-PubHealth.indd 255 11/7/19 4:16 PM

(c) ketabton.com: The Digital Library

▸ PUBLIC HEALTH SCIENCES—EPIDEMIOLOGY AND BIOSTATISTICS

Observational studies

STUDY TYPE	DESIGN	MEASURES/EXAMPLE
Cross-sectional study	Frequency of disease and frequency of risk-related factors are assessed in the present. Asks, "What is happening?"	Disease prevalence. Can show risk factor association with disease, but does not establish causality.
Case-control study	Compares a group of people with disease to a group without disease. Looks to see if odds of prior exposure or risk factor differ by disease state. Asks, "What happened?"	Odds ratio (OR). Patients with COPD had higher odds of a smoking history than those without COPD.
Cohort study	Compares a group with a given exposure or risk factor to a group without such exposure. Looks to see if exposure or risk factor is associated with later development of disease. Can be prospective or retrospective.	Relative risk (RR). Smokers had a higher risk of developing COPD than nonsmokers. Cohort = relative risk.
Crossover study	Compares the effect of a series of ≥2 treatments on a participant. Order in which participants receive treatments is randomized. Washout period occurs between each treatment.	Allows participants to serve as their own controls.
Twin concordance study	Compares the frequency with which both monozygotic twins vs both dizygotic twins develop the same disease.	Measures heritability and influence of environmental factors ("nature vs nurture").
Adoption study	Compares siblings raised by biological vs adoptive parents.	Measures heritability and influence of environmental factors.

Clinical trial

Experimental study involving humans. Compares therapeutic benefits of ≥2 treatments, or of treatment and placebo. Study quality improves when study is randomized, controlled, and double-blinded (ie, neither patient nor doctor knows whether the patient is in the treatment or control group). Triple-blind refers to the additional blinding of the researchers analyzing the data.
Four phases ("Does the drug **SWIM**?").

DRUG TRIALS	TYPICAL STUDY SAMPLE	PURPOSE
Phase I	Small number of either healthy volunteers or patients with disease of interest.	"Is it **S**afe?" Assesses safety, toxicity, pharmacokinetics, and pharmacodynamics.
Phase II	Moderate number of patients with disease of interest.	"Does it **W**ork?" Assesses treatment efficacy, optimal dosing, and adverse effects.
Phase III	Large number of patients randomly assigned either to the treatment under investigation or to the standard of care (or placebo).	"Is it as good or better?" Compares the new treatment to the current standard of care (any **I**mprovement?).
Phase IV	Postmarketing surveillance of patients after treatment is approved.	"Can it stay?" Detects rare or long-term adverse effects (eg, black box warnings). Can result in treatment being withdrawn from **M**arket.

(c) ketabton.com: The Digital Library

Evaluation of diagnostic tests

Sensitivity and specificity are fixed properties of a test. PPV and NPV vary depending on disease prevalence in population being tested.



Sensitivity (true-positive rate)

Proportion of all people with disease who test positive, or the probability that when the disease is present, the test is positive.
Value approaching 100% is desirable for **ruling out** disease and indicates a **low false-negative rate**.

= TP / (TP + FN)
= 1 – FN rate
SN-N-OUT = highly SeNsitive test, when Negative, rules OUT disease
High sensitivity test used for screening

Specificity (true-negative rate)

Proportion of all people without disease who test negative, or the probability that when the disease is absent, the test is negative.
Value approaching 100% is desirable for **ruling in** disease and indicates a **low false-positive rate**.

= TN / (TN + FP)
= 1 – FP rate
SP-P-IN = highly SPecific test, when Positive, rules IN disease
High specificity test used for confirmation after a positive screening test

Positive predictive value

Probability that a person who has a positive test result actually has the disease.

PPV = TP / (TP + FP)
PPV varies directly with pretest probability (baseline risk, such as prevalence of disease): high pretest probability → high PPV

Negative predictive value

Probability that a person with a negative test result actually does not have the disease.

NPV = TN / (TN + FN)
NPV varies inversely with prevalence or pretest probability



Possible cutoff values for (+) vs (-) test result
A = 100% sensitivity cutoff value
B = practical compromise between specificity and sensitivity
C = 100% specificity cutoff value

Lowering the cutoff value: **B** → **A** (↑ FP ↓ FN)	↑ Sensitivity ↑ NPV ↓ Specificity ↓ PPV
Raising the cutoff value: **B** → **C** (↑ FN ↓ FP)	↑ Specificity ↑ PPV ↓ Sensitivity ↓ NPV

Likelihood ratio

Likelihood that a given test result would be expected in a patient with the target disorder compared to the likelihood that the same result would be expected in a patient without the target disorder.
$LR^+ > 10$ indicates a highly specific test, while $LR^- < 0.1$ indicates a highly sensitive test.
LRs can be multiplied with pretest odds of disease to estimate posttest odds.

$$LR^+ = \frac{\text{sensitivity}}{1 - \text{specificity}} = \frac{\text{TP rate}}{\text{FP rate}}$$

$$LR^- = \frac{1 - \text{sensitivity}}{\text{specificity}} = \frac{\text{FN rate}}{\text{TN rate}}$$

(c) ketabton.com: The Digital Library

Quantifying risk

Definitions and formulas are based on the classic 2 × 2 or contingency table.

Exposure or intervention	Disease or outcome ⊕	Disease or outcome ⊖
⊕	a	b
⊖	c	d

TERM	DEFINITION	EXAMPLE	FORMULA
Odds ratio	Typically used in case-control studies. Represents the odds of exposure among cases (a/c) vs odds of exposure among controls (b/d).	If in a case-control study, 20/30 lung cancer patients and 5/25 healthy individuals report smoking, the OR is 8; so the lung cancer patients are 8 times more likely to have a history of smoking.	$OR = \frac{a/c}{b/d} = \frac{ad}{bc}$ a 20 | b 5 c 10 | d 20
Relative risk	Typically used in cohort studies. Risk of developing disease in the exposed group divided by risk in the unexposed group. RR = 1 → no association between exposure and disease. RR > 1 → exposure associated with ↑ disease occurrence. RR < 1 → exposure associated with ↓ disease occurrence.	If 5/10 people exposed to radiation are diagnosed with cancer, and 1/10 people not exposed to radiation are diagnosed with cancer, the RR is 5; so people exposed to radiation have a 5 times greater risk of developing cancer. For rare diseases (low prevalence), OR approximates RR.	$RR = \frac{a/(a+b)}{c/(c+d)}$ a 5 | b 5 c 1 | d 9
Relative risk reduction	The proportion of risk reduction attributable to the intervention as compared to a control.	If 2% of patients who receive a flu shot develop the flu, while 8% of unvaccinated patients develop the flu, then RR = 2/8 = 0.25, and RRR = 0.75.	$RRR = 1 - RR$
Attributable risk	The difference in risk between exposed and unexposed groups.	If risk of lung cancer in smokers is 21% and risk in nonsmokers is 1%, then the attributable risk is 20%.	$AR = \frac{a}{a+b} - \frac{c}{c+d}$ $AR\% = \frac{RR-1}{RR} \times 100$
Absolute risk reduction	The difference in risk (not the proportion) attributable to the intervention as compared to a control.	If 8% of people who receive a placebo vaccine develop the flu vs 2% of people who receive a flu vaccine, then ARR = 8%–2% = 6% = 0.06.	$ARR = \frac{c}{c+d} - \frac{a}{a+b}$
Number needed to treat	Number of patients who need to be treated for 1 patient to benefit. Lower number = better treatment.		$NNT = 1/ARR$
Number needed to harm	Number of patients who need to be exposed to a risk factor for 1 patient to be harmed. Higher number = safer exposure.		$NNH = 1/AR$
Case fatality rate	Percentage of deaths occurring among those with disease.	If 4 patients die among 10 cases of meningitis, case fatality rate is 40%.	$CFR\% = \frac{deaths}{cases} \times 100$

(c) ketabton.com: The Digital Library

Incidence vs prevalence



$$\text{Incidence} = \frac{\#\text{ of new cases}}{\#\text{ of people at risk}} \quad \text{(per unit of time)}$$

$$\text{Prevalence} = \frac{\#\text{ of existing cases}}{\text{Total \# of people in a population}} \quad \text{(at a point in time)}$$

$$\frac{\text{Prevalence}}{1 - \text{prevalence}} = \text{Incidence rate} \times \text{average duration of disease}$$

Incidence looks at new cases (**incidents**).

Prevalence looks at **all** current cases.

Prevalence ≈ incidence for short duration disease (eg, common cold).
Prevalence > incidence for chronic diseases, due to large # of existing cases (eg, diabetes).

Prevalence ~ pretest probability.
↑ prevalence → ↑ PPV and ↓ NPV.

SITUATION	INCIDENCE	PREVALENCE
↑ survival time	—	↑
↑ mortality	—	↓
Faster recovery time	—	↓
Extensive vaccine administration	↓	↓
↓ risk factors	↓	↓

Precision vs accuracy

Precision (reliability)	The consistency and reproducibility of a test. The absence of random variation in a test.	Random error ↓ precision in a test. ↑ precision → ↓ standard deviation. ↑ precision → ↑ statistical power (1 – β).
Accuracy (validity)	The closeness of test results to the true values. The absence of systematic error or bias in a test.	Systematic error ↓ accuracy in a test.



(c) ketabton.com: The Digital Library

Receiving operating characteristic curve

ROC curve demonstrates how well a diagnostic test can distinguish between 2 groups (eg, disease vs healthy). Plots the true-positive rate (sensitivity) against the false-positive rate (1 – specificity).

The better performing test will have a higher area under the curve (AUC), with the curve closer to the upper left corner.



Bias and study errors

TYPE	DEFINITION	EXAMPLES	STRATEGIES TO REDUCE BIAS
Recruiting participants			
Selection bias	Nonrandom sampling or treatment allocation of subjects such that study population is not representative of target population. Most commonly a sampling bias.	Berkson bias—cases and/or controls selected from hospitals are less healthy and have different exposures than general population Attrition bias—participants lost to follow up have a different prognosis than those who complete the study	Randomization Ensure the choice of the right comparison/reference group
Performing study			
Recall bias	Awareness of disorder alters recall by subjects; common in retrospective studies	Patients with disease recall exposure after learning of similar cases	Decrease time from exposure to follow-up
Measurement bias	Information is gathered in a systemically distorted manner	Using a faulty automatic sphygmomanometer to measure BP Hawthorne effect—participants change behavior upon awareness of being observed	Use objective, standardized, and previously tested methods of data collection that are planned ahead of time Use placebo group
Procedure bias	Subjects in different groups are not treated the same	Patients in treatment group spend more time in highly specialized hospital units	Blinding (masking) and use of placebo reduce influence of participants and researchers on procedures and interpretation of outcomes as neither are aware of group assignments
Observer-expectancy bias	Researcher's belief in the efficacy of a treatment changes the outcome of that treatment (aka, Pygmalion effect)	An observer expecting treatment group to show signs of recovery is more likely to document positive outcomes	

(c) ketabton.com: The Digital Library

Bias and study errors *(continued)*

TYPE	DEFINITION	EXAMPLES	STRATEGY TO REDUCE BIAS
Interpreting results			
Confounding bias	Factor related to both exposure and outcome (but not on causal path) distorts effect of exposure on outcome (vs effect modification, in which the exposure leads to different outcomes in subgroups stratified by the factor)	An uncontrolled study shows an association between drinking coffee and lung cancer. However, coffee drinkers also smoke more, which can account for the association	Multiple/repeated studies Crossover studies (subjects act as their own controls) Matching (patients with similar characteristics in both treatment and control groups)
Lead-time bias	Early detection is confused with ↑ survival	Early detection makes it seem like survival has increased, but the disease's natural history has not changed	Measure "back-end" survival (adjust survival according to the severity of disease at the time of diagnosis)
Length-time bias	Screening test detects diseases with long latency period, while those with shorter latency period become symptomatic earlier	A slowly progressive cancer is more likely detected by a screening test than a rapidly progressive cancer	A randomized controlled trial assigning subjects to the screening program or to no screening

(c) ketabton.com: The Digital Library

Statistical distribution

Measures of central tendency	Mean = (sum of values)/(total number of values).	Most affected by outliers (extreme values).
	Median = middle value of a list of data sorted from least to greatest.	If there is an even number of values, the median will be the average of the middle two values.
	Mode = most common value.	Least affected by outliers.
Measures of dispersion	Standard deviation = how much variability exists in a set of values, around the mean of these values. Standard error = an estimate of how much variability exists in a (theoretical) set of sample means around the true population mean.	σ = SD; n = sample size. Variance = $(SD)^2$. SE = $\sigma/\sqrt{n}$. SE ↓ as n ↑.
Normal distribution	Gaussian, also called bell-shaped. Mean = median = mode.	

Nonnormal distributions

Bimodal	Suggests two different populations (eg, metabolic polymorphism such as fast vs slow acetylators; age at onset of Hodgkin lymphoma; suicide rate by age).	
Positive skew	Typically, mean > median > mode. Asymmetry with longer tail on right.	
Negative skew	Typically, mean < median < mode. Asymmetry with longer tail on left.	

Statistical hypotheses

Null (H_0)	Hypothesis of no difference or relationship (eg, there is no association between the disease and the risk factor in the population).
Alternative (H_1)	Hypothesis of some difference or relationship (eg, there is some association between the disease and the risk factor in the population).

(c) ketabton.com: The Digital Library

Outcomes of statistical hypothesis testing

Correct result

Stating that there is an effect or difference when one exists (null hypothesis rejected in favor of alternative hypothesis).
Stating that there is no effect or difference when none exists (null hypothesis not rejected).



Blue shading = correct result.

Incorrect result

Type I error (α)

Stating that there is an effect or difference when none exists (null hypothesis incorrectly rejected in favor of alternative hypothesis).
α is the probability of making a type I error. p is judged against a preset α level of significance (usually 0.05). If $p < 0.05$ for a study outcome, the probability of obtaining that result purely by chance is < 5%.
Statistical significance ≠ clinical significance.

Also called false-positive error.

α = you **a**ccused an innocent man.
You can never "prove" the alternate hypothesis, but you can reject the null hypothesis as being very unlikely.

Type II error (β)

Stating that there is not an effect or difference when one exists (null hypothesis is not rejected when it is in fact false).
β is the probability of making a type II error. β is related to statistical power (1 – β), which is the probability of rejecting the null hypothesis when it is false.
↑ power and ↓ β by:
- ↑ sample size
- ↑ expected effect size
- ↑ precision of measurement

Also called false-negative error.

β = you **b**lindly let the guilty man go free.
If you ↑ sample size, you ↑ power. There is **power in numbers**.

Confidence interval

Range of values within which the true mean of the population is expected to fall, with a specified probability.
CI for sample mean = $\bar{x} \pm Z(SE)$
The 95% CI (corresponding to α = .05) is often used. As sample size increases, CI narrows.
For the 95% CI, Z = 1.96.
For the 99% CI, Z = 2.58.

If the 95% CI for a mean difference between 2 variables includes 0, then there is no significant difference and H_0 is not rejected.
If the 95% CI for odds ratio or relative risk includes 1, H_0 is not rejected.
If the CIs between 2 groups do not overlap → statistically significant difference exists.
If the CIs between 2 groups overlap → usually no significant difference exists.

(c) ketabton.com: The Digital Library

Meta-analysis

A method of statistical analysis that pools summary data (eg, means, RRs) from multiple studies for a more precise estimate of the size of an effect. Also estimates heterogeneity of effect sizes between studies.

Improves power, strength of evidence, and generalizability of study findings. Limited by quality of individual studies and bias in study selection.

Common statistical tests

t-test	Checks differences between **mean**s of **2** groups.	**T**ea is **mean**t for **2**. Example: comparing the mean blood pressure between men and women.
ANOVA	Checks differences between means of **3** or more groups.	**3** words: **AN**alysis **O**f **VA**riance. Example: comparing the mean blood pressure between members of 3 different ethnic groups.
Chi-square (χ^2)	Checks differences between 2 or more percentages or proportions of **categorical** outcomes (not mean values).	Pronounce **Chi-tegorical**. Example: comparing the percentage of members of 3 different ethnic groups who have essential hypertension.
Fisher's exact test	Checks differences between 2 percentages or proportions of categorical, nominal outcomes. Use instead of chi-square test with small populations.	Example: comparing the percentage of 20 men and 20 women with hypertension.



Pearson correlation coefficient

r is always between –1 and +1. The closer the absolute value of *r* is to 1, the stronger the linear correlation between the 2 variables. Variance is how much the measured values differ from the average value in a data set.

Positive *r* value → positive correlation (as one variable ↑, the other variable ↑).

Negative *r* value → negative correlation (as one variable ↑, the other variable ↓).

Coefficient of determination = r^2 (amount of variance in one variable that can be explained by variance in another variable).



(c) ketabton.com: The Digital Library

▸ PUBLIC HEALTH SCIENCES—ETHICS

Core ethical principles

Autonomy	Obligation to respect patients as individuals (truth-telling, confidentiality), to create conditions necessary for autonomous choice (informed consent), and to honor their preference in accepting or not accepting medical care.
Beneficence	Physicians have a special ethical (fiduciary) duty to act in the patient's best interest. May conflict with autonomy (an informed patient has the right to decide) or what is best for society (eg, mandatory TB treatment). Traditionally, patient interest supersedes.
Nonmaleficence	"Do no harm." Must be balanced against beneficence; if the benefits outweigh the risks, a patient may make an informed decision to proceed (most surgeries and medications fall into this category).
Justice	To treat persons fairly and equitably. This does not always imply equally (eg, triage).

Informed consent

A process (not just a document/signature) that requires:

- Disclosure: discussion of pertinent information (using medical interpreter, if needed)
- Understanding: ability to comprehend
- Capacity: ability to reason and make one's own decisions (distinct from competence, a legal determination)
- Voluntariness: freedom from coercion and manipulation

Patients must have an intelligent understanding of their diagnosis and the risks/benefits of proposed treatment and alternative options, including no treatment.

Patient must be informed that he or she can revoke written consent at any time, even orally.

Exceptions to informed consent (**WIPE** it away):

- **W**aiver—patient explicitly waives the right of informed consent
- Legally **I**ncompetent—patient lacks decision-making capacity (obtain consent from legal surrogate)
- Therapeutic **P**rivilege—withholding information when disclosure would severely harm the patient or undermine informed decision-making capacity
- **E**mergency situation—implied consent may apply

Consent for minors

A minor is generally any person < 18 years old. Parental consent laws in relation to healthcare vary by state. In general, parental consent should be obtained, but exceptions exist for emergency treatment (eg, blood transfusions) or if minor is legally emancipated (eg, married, self-supporting, or in the military).

Situations in which parental consent is usually not required:

- **Sex** (contraception, STIs, pregnancy)
- **Drugs** (substance abuse)
- **Rock and roll** (emergency/trauma)

Physicians should always encourage healthy minor-guardian communication.

Physician should seek a minor's assent even if their consent is not required.

(c) ketabton.com: The Digital Library

Decision-making capacity

Physician must determine whether the patient is psychologically and legally capable of making a particular healthcare decision. Note that decisions made with capacity cannot be revoked simply if the patient later loses capacity. Intellectual disability alone (eg, Down syndrome, autism) is not an exclusion criterion for informed decision-making.

Capacity is determined by a physician for a specific healthcare-related decision (eg, to refuse medical care). Competency is determined by a judge and usually refers to more global categories of decision making (eg, legally unable to make any healthcare-related decision).

Components (think **GIEMSA**):

- Decision is consistent with patient's values and **G**oals
- Patient is **I**nformed (knows and understands)
- Patient **E**xpresses a choice
- Decision is not a result of altered **M**ental status (eg, delirium, psychosis, intoxication), **M**ood disorder
- Decision remains **S**table over time
- Patient is ≥ 18 years of **A**ge or otherwise legally emancipated

Advance directives

Instructions given by a patient in anticipation of the need for a medical decision. Details vary per state law.

Oral advance directive

Incapacitated patient's prior oral statements commonly used as guide. Problems arise from variance in interpretation. If patient was informed, directive was specific, patient made a choice, and decision was repeated over time to multiple people, then the oral directive is more valid.

Written advance directive

Specifies specific healthcare interventions that a patient anticipates he or she would accept or reject during treatment for a critical or life-threatening illness. A living will is an example.

Medical power of attorney

Patient designates an agent to make medical decisions in the event that he/she loses decision-making capacity. Patient may also specify decisions in clinical situations. Can be revoked by patient if decision-making capacity is intact. More flexible than a living will.

Do not resuscitate order

DNR order prohibits cardiopulmonary resuscitation (CPR). Other resuscitative measures that may follow (eg, feeding tube) are also typically avoided.

Surrogate decision-maker

If a patient loses decision-making capacity and has not prepared an advance directive, individuals (surrogates) who know the patient must determine what the patient would have done. Priority of surrogates: **spouse** → adult **Chi**ldren → **P**arents → **S**iblings → other relatives (the **spouse ChiPS** in).

(c) ketabton.com: The Digital Library

Confidentiality

Confidentiality respects patient privacy and autonomy. If the patient is incapacitated or the situation is emergent, disclosing information to family and friends should be guided by professional judgment of patient's best interest. The patient may voluntarily waive the right to confidentiality (eg, insurance company request).

General principles for exceptions to confidentiality:

- Potential physical harm to others is serious and imminent
- Alternative means to warn or protect those at risk is not possible
- Self-harm is likely
- Steps can be taken to prevent harm

Examples of exceptions to patient confidentiality (many are state specific) include the following ("The physician's good judgment **SAVED** the day"):

- **S**uicidal/homicidal patients.
- **A**buse (children, elderly, and/or prisoners).
- Duty to protect—state-specific laws that sometimes allow physician to inform or somehow protect potential **V**ictim from harm.
- **E**pileptic patients and other impaired automobile drivers.
- Reportable **D**iseases (eg, STIs, hepatitis, food poisoning); physicians may have a duty to warn public officials, who will then notify people at risk. Dangerous communicable diseases, such as TB or Ebola, may require involuntary treatment.

(c) ketabton.com: The Digital Library

Ethical situations

SITUATION	APPROPRIATE RESPONSE
Patient is not adherent.	Attempt to identify the reason for nonadherence and determine his/her willingness to change; do not coerce the patient into adhering and do not refer him/her to another physician.
Patient desires an unnecessary procedure.	Attempt to understand why the patient wants the procedure and address underlying concerns. Do not refuse to see the patient and do not refer him/her to another physician. Avoid performing unnecessary procedures.
Patient has difficulty taking medications.	Provide written instructions; attempt to simplify treatment regimens; use teach-back method (ask patient to repeat regimen back to physician) to ensure comprehension.
Family members ask for information about patient's prognosis.	Avoid discussing issues with relatives without the patient's permission.
A patient's family member asks you not to disclose the results of a test if the prognosis is poor because the patient will be "unable to handle it."	Attempt to identify why the family member believes such information would be detrimental to the patient's condition. Explain that as long as the patient has decision-making capacity and does not indicate otherwise, communication of information concerning his/her care will not be withheld. However, if you believe the patient might seriously harm himself/herself or others if informed, then you may invoke therapeutic privilege and withhold the information.
A 17-year-old girl is pregnant and requests an abortion.	Many states require parental notification or consent for minors for an abortion. Unless there are specific medical risks associated with pregnancy, a physician should not sway the patient's decision for, or against, an elective abortion (regardless of maternal age or fetal condition).
A 15-year-old girl is pregnant and wants to keep the child. Her parents want you to tell her to give the child up for adoption.	The patient retains the right to make decisions regarding her child, even if her parents disagree. Provide information to the teenager about the practical issues of caring for a baby. Discuss the options, if requested. Encourage discussion between the teenager and her parents to reach the best decision.
A terminally ill patient requests physician assistance in ending his/her own life.	Overwhelming majority of states refuse involvement in any form of physician-assisted death. Physicians may, however, prescribe medically appropriate analgesics even if they shorten the patient's life.
Patient is suicidal.	Assess the seriousness of the threat. If it is serious, suggest that the patient remain in the hospital voluntarily; patient can be hospitalized involuntarily if he/she refuses.
Patient states that he/she finds you attractive.	Ask direct, closed-ended questions and use a chaperone if necessary. Romantic relationships with patients are never appropriate. It may be necessary to transition care to another physician.
A woman who had a mastectomy says she now feels "ugly."	Find out why the patient feels this way. Do not offer falsely reassuring statements (eg, "You still look good").
Patient is angry about the long time he/she spent in the waiting room.	Acknowledge the patient's anger, but do not take a patient's anger personally. Apologize for any inconvenience. Stay away from efforts to explain the delay.
Patient is upset with the way he/she was treated by another doctor.	Suggest that the patient speak directly to that physician regarding his/her concerns. If the problem is with a member of the office staff, tell the patient you will speak to that person.
An invasive test is performed on the wrong patient.	Regardless of the outcome, a physician is ethically obligated to inform a patient that a mistake has been made.

(c) ketabton.com: The Digital Library

Ethical situations *(continued)*

SITUATION	APPROPRIATE RESPONSE
A patient requires a treatment not covered by his/her insurance.	Never limit or deny care because of the expense in time or money. Discuss all treatment options with patients, even if some are not covered by their insurance companies.
A 7-year-old boy loses a sister to cancer and now feels responsible.	At ages 5–7, children begin to understand that death is permanent, that all life functions end completely at death, and that everything that is alive eventually dies. Provide a direct, concrete description of his sister's death. Avoid clichés and euphemisms. Reassure the boy that he is not responsible. Identify and normalize fears and feelings. Encourage play and healthy coping behaviors (eg, remembering her in his own way).
Patient is victim of intimate partner violence.	Ask if patient is safe and has an emergency plan. Do not necessarily pressure patient to leave his or her partner, or disclose the incident to the authorities (unless required by state law).
Patient wants to try alternative or holistic medicine.	Explore any underlying reasons with the patient in a supportive, nonjudgmental manner. Advise the patient of known benefits and risks of treatment, including adverse effects, contraindications, and medication interactions.
Physician colleague presents to work impaired.	If impaired or incompetent, colleague is a threat to patient safety. Report the situation to local supervisory personnel. Should the organization fail to take action, alert the state licensing board.
Patient is officially determined to suffer brain death. Patient's family insists on maintaining life support indefinitely because patient is still moving when touched.	Gently explain to family that there is no chance of recovery, and that brain death is equivalent to death. Movement is due to spinal arc reflex and is not voluntary. Bring case to appropriate ethics board regarding futility of care and withdrawal of life support.
A pharmaceutical company offers you a sponsorship in exchange for advertising its new drug.	Reject this offer. Generally, decline gifts and sponsorships to avoid any appearance of conflict of interest. The AMA Code of Ethics does make exceptions for gifts directly benefitting patients; gifts of minimal value; special funding for medical education of students, residents, fellows; grants whose recipients are chosen by independent institutional criteria; and funds that are distributed without attribution to sponsors.
Patient requests a nonemergent procedure that is against your personal or religious beliefs.	Provide accurate and unbiased information so patients can make an informed decision. Explain to the patient that you do not perform the procedure but offer to refer him/her to another physician.
Mother and 15-year-old daughter are unresponsive following a car accident and are bleeding internally. Father says do not transfuse because they are Jehovah's Witnesses.	Transfuse daughter, but do not transfuse mother. Emergent care can be refused by the healthcare proxy for an adult, particularly when patient preferences are known or reasonably inferred, but not for a minor based solely on faith.
A child presents with injuries inconsistent with parental story.	Contact child protective services and ensure child is in a safe location. Physicians are required by law to report any reasonable suspicion of child abuse or endangerment.

(c) ketabton.com: The Digital Library

▸ PUBLIC HEALTH SCIENCES—THE WELL PATIENT

Changes in the elderly

Sexual changes:
- Men—slower erection/ejaculation, longer refractory period, but unchanged libido.
- Women—vaginal shortening, thinning, and dryness

Sleep patterns: ↓ REM and slow-wave sleep, ↑ sleep latency, ↑ early awakenings
↑ suicide rate
↓ vision and hearing
↓ immune response
↓ renal, pulmonary, and GI function
↓ muscle mass, ↑ fat
Intelligence does not decrease

▸ PUBLIC HEALTH SCIENCES—HEALTHCARE DELIVERY

Disease prevention

Primary disease prevention	Prevent disease before it occurs (eg, HPV vaccination)
Secondary disease prevention	Screen early for and manage existing but asymptomatic disease (eg, Pap smear for cervical cancer)
Tertiary disease prevention	Treatment to reduce complications from disease that is ongoing or has long-term effects (eg, chemotherapy)
Quaternary disease prevention	Quit (avoid) unnecessary medical interventions to minimize incidental harm (eg, imaging studies, optimizing medications to reduce polypharmacy).

(c) ketabton.com: The Digital Library

Major medical insurance plans

PLAN	PROVIDERS	PAYMENTS	SPECIALIST CARE
Exclusive provider organization	Restricted to limited panel (except emergencies)		No referral required
Health maintenance organization	Restricted to limited panel (except emergencies)	Denied for any service that does not meet established, evidence-based guidelines	Requires referral from primary care provider
Point of service	Patient can see providers outside network	Higher copays and deductibles for out-of-network services	Requires referral from primary care provider
Preferred provider organization	Patient can see providers outside network	Higher copays and deductibles for all services	No referral required
Accountable care organization	Providers voluntarily enroll	Medicare	Specialists voluntarily enroll

Healthcare payment models

Bundled payment	Healthcare organization receives a set amount per service, regardless of ultimate cost, to be divided among all providers and facilities involved.
Capitation	Physicians receive a set amount per patient assigned to them per period of time, regardless of how much the patient uses the healthcare system. Used by some HMOs.
Discounted fee-for-service	Patient pays for each individual service at a discounted rate predetermined by providers and payers (eg, PPOs).
Fee-for-service	Patient pays for each individual service.
Global payment	Patient pays for all expenses associated with a single incident of care with a single payment. Most commonly used during elective surgeries, as it covers the cost of surgery as well as the necessary pre- and postoperative visits.

(c) ketabton.com: The Digital Library

Medicare and Medicaid

Medicare and Medicaid—federal social healthcare programs that originated from amendments to the Social Security Act.

Medicare is available to patients ≥ 65 years old, < 65 with certain disabilities, and those with end-stage renal disease.

Medicaid is joint federal and state health assistance for people with limited income and/or resources.

MedicarE is for Elderly.
MedicaiD is for Destitute.

The 4 parts of Medicare:

- Part A: HospitAl insurance, home hospice care
- Part B: Basic medical bills (eg, doctor's fees, diagnostic testing)
- Part C: (parts A + B = Combo) delivered by approved private companies
- Part D: Prescription Drugs

Hospice care

Medical care focused on providing comfort and palliation instead of definitive cure. Available to patients on Medicare or Medicaid and in most private insurance plans whose life expectancy is < 6 months.

During end-of-life care, priority is given to improving the patient's comfort and relieving pain (often includes opioid, sedative, or anxiolytic medications). Facilitating comfort is prioritized over potential side effects (eg, respiratory depression). This prioritization of positive effects over negative effects is called the principle of double effect.

Common causes of death (US) by age

	< 1 YR	1–14 YR	15–34 YR	35–44 YR	45–64 YR	65+ YR
#1	Congenital malformations	Unintentional injury	Unintentional injury	Unintentional injury	Cancer	Heart disease
#2	Preterm birth	Cancer	Suicide	Cancer	Heart disease	Cancer
#3	Maternal pregnancy complications	Congenital malformations	Homicide	Heart disease	Unintentional injury	Chronic respiratory disease

Conditions with frequent hospital readmissions

Readmissions may be reduced by discharge planning and outpatient follow-up appointments. The table below is based on readmission for any reason within 30 days of discharge.

	MEDICARE	MEDICAID	PRIVATE INSURANCE	UNINSURED
#1	Congestive HF	Mood disorders	Maintenance of chemotherapy or radiotherapy	Mood disorders
#2	Septicemia	Schizophrenia/psychotic disorders	Mood disorders	Alcohol-related disorders
#3	Pneumonia	Diabetes mellitus with complications	Complications of surgical procedures or medical care	Diabetes mellitus with complications

(c) ketabton.com: The Digital Library

▸ PUBLIC HEALTH SCIENCES—QUALITY AND SAFETY

Safety culture

Organizational environment in which everyone can freely bring up safety concerns without fear of censure. Facilitates error identification.

Event reporting systems collect data on errors for internal and external monitoring.

Human factors design

Forcing functions (those that prevent undesirable actions [eg, connecting feeding syringe to IV tubing]) are the most effective.
Standardization improves process reliability (eg, clinical pathways, guidelines, checklists).
Simplification reduces wasteful activities (eg, consolidating electronic medical records).

Deficient designs hinder workflow and lead to staff workarounds that bypass safety features (eg, patient ID barcodes affixed to computers due to unreadable wristbands).

PDSA cycle

Process improvement model to test changes in real clinical setting. Impact on patients:
- Plan—define problem and solution
- Do—test new process
- Study—measure and analyze data
- Act—integrate new process into workflow



Quality measurements

	MEASURE	EXAMPLE
Structural	Physical equipment, resources, facilities	Number of diabetes educators
Process	Performance of system as planned	Percentage of diabetic patients whose HbA_{1c} was measured in the past 6 months
Outcome	Impact on patients	Average HbA_{1c} of patients with diabetes
Balancing	Impact on other systems/outcomes	Incidence of hypoglycemia among patients who tried an intervention to lower HbA_{1c}

Swiss cheese model

Focuses on systems and conditions rather than an individual's error. The risk of a threat becoming a reality is mitigated by differing layers and types of defenses. Patient harm can occur despite multiple safeguards when "the holes in the cheese line up."



(c) ketabton.com: The Digital Library

Types of medical errors	May involve patient identification, diagnosis, monitoring, nosocomial infection, medications, procedures, devices, documentation, handoffs. Medical errors should be disclosed to patients, independent of immediate outcome (harmful or not).	
Active error	Occurs at level of frontline operator (eg, wrong IV pump dose programmed).	Immediate impact.
Latent error	Occurs in processes indirect from operator but impacts patient care (eg, different types of IV pumps used within same hospital).	Accident waiting to happen.
Never event	Adverse event that is identifiable, serious, and usually preventable (eg, scalpel retained in a surgical patient's abdomen).	Major error that should never occur.

Burnout vs fatigue

Burnout	Prolonged, excessive stress → cynicism, detachment, ↓ motivation and interest, sense of failure and helplessness, ↓ immunity. Medical errors due to lack of concern.
Fatigue	Sleep deprivation → ↓ energy and motivation, cognitive impairment. Medical errors due to compromised intellectual function.

Medical error analysis

	DESIGN	METHODS
Root cause analysis	Retrospective approach. Applied after failure event to prevent recurrence.	Uses records and participant interviews to identify all the underlying problems (eg, process, people, environment, equipment, materials, management) that led to an error.
Failure mode and effects analysis	Forward-looking approach. Applied before process implementation to prevent failure occurrence.	Uses inductive reasoning to identify all the ways a process might fail and prioritizes them by their probability of occurrence and impact on patients.

FAS1_2019_06-PubHealth.indd 274 11/7/19 4:16 PM

(c) ketabton.com: The Digital Library

SECTION III

High-Yield Organ Systems

"Symptoms, then, are in reality nothing but the cry from suffering organs."
—Jean-Martin Charcot

"Man is an intelligence in servitude to his organs."
—Aldous Huxley

"When every part of the machine is correctly adjusted and in perfect harmony, health will hold dominion over the human organism by laws as natural and immutable as the laws of gravity."
—Andrew T. Still

- Approaching the Organ Systems 276
- Cardiovascular 279
- Endocrine 325
- Gastrointestinal 357
- Hematology and Oncology 403
- Musculoskeletal, Skin, and Connective Tissue 445
- Neurology and Special Senses 489
- Psychiatry 553
- Renal 577
- Reproductive 611
- Respiratory 659

FAS1_2019_07-Cardio.indd 275 11/7/19 4:24 PM

(c) ketabton.com: The Digital Library

APPROACHING THE ORGAN SYSTEMS

In this section, we have divided the High-Yield Facts into the major **Organ Systems.** Within each Organ System are several subsections, including **Embryology, Anatomy, Physiology, Pathology,** and **Pharmacology.** As you progress through each Organ System, refer back to information in the previous subsections to organize these basic science subsections into a "vertically integrated" framework for learning. Below is some general advice for studying the organ systems by these subsections.

Embryology

Relevant embryology is included in each organ system subsection. Embryology tends to correspond well with the relevant anatomy, especially with regard to congenital malformations.

Anatomy

Several topics fall under this heading, including gross anatomy, histology, and neuroanatomy. Do not memorize all the small details; however, do not ignore anatomy altogether. Review what you have already learned and what you wish you had learned. Many questions require two or more steps. The first step is to identify a structure on anatomic cross section, electron micrograph, or photomicrograph. The second step may require an understanding of the clinical significance of the structure.

When studying, stress clinically important material. For example, be familiar with gross anatomy and radiologic anatomy related to specific diseases (eg, Pancoast tumor, Horner syndrome), traumatic injuries (eg, fractures, sensory and motor nerve deficits), procedures (eg, lumbar puncture), and common surgeries (eg, cholecystectomy). There are also many questions on the exam involving x-rays, CT scans, and neuro MRI scans. Many students suggest browsing through a general radiology atlas, pathology atlas, and histology atlas. Focus on learning basic anatomy at key levels in the body (eg, sagittal brain MRI; axial CT of the midthorax, abdomen, and pelvis). Basic neuroanatomy (especially pathways, blood supply, and functional anatomy), associated neuropathology, and neurophysiology have good yield. Please note that many of the photographic images in this book are for illustrative purposes and are not necessarily reflective of Step 1 emphasis.

Physiology

The portion of the examination dealing with physiology is broad and concept oriented and thus does not lend itself as well to fact-based review. Diagrams are often the best study aids, especially given the increasing number of questions requiring the interpretation of diagrams. Learn to apply basic physiologic relationships in a variety of ways (eg, the Fick equation, clearance equations). You are seldom asked to perform complex

(c) ketabton.com: The Digital Library

calculations. Hormones are the focus of many questions, so learn their sites of production and action as well as their regulatory mechanisms.

A large portion of the physiology tested on the USMLE Step 1 is clinically relevant and involves understanding physiologic changes associated with pathologic processes (eg, changes in pulmonary function with COPD). Thus, it is worthwhile to review the physiologic changes that are found with common pathologies of the major organ systems (eg, heart, lungs, kidneys, GI tract) and endocrine glands.

Pathology

Questions dealing with this discipline are difficult to prepare for because of the sheer volume of material involved. Review the basic principles and hallmark characteristics of the key diseases. Given the clinical orientation of Step 1, it is no longer sufficient to know only the "buzzword" associations of certain diseases (eg, café-au-lait macules and neurofibromatosis); you must also know the clinical descriptions of these findings.

Given the clinical slant of the USMLE Step 1, it is also important to review the classic presenting signs and symptoms of diseases as well as their associated laboratory findings. Delve into the signs, symptoms, and pathophysiology of major diseases that have a high prevalence in the United States (eg, alcoholism, diabetes, hypertension, heart failure, ischemic heart disease, infectious disease). Be prepared to think one step beyond the simple diagnosis to treatment or complications.

The examination includes a number of color photomicrographs and photographs of gross specimens that are presented in the setting of a brief clinical history. However, read the question and the choices carefully before looking at the illustration, because the history will help you identify the pathologic process. Flip through an illustrated pathology textbook, color atlases, and appropriate Web sites in order to look at the pictures in the days before the exam. Pay attention to potential clues such as age, sex, ethnicity, occupation, recent activities and exposures, and specialized lab tests.

Pharmacology

Preparation for questions on pharmacology is straightforward. Learning all the key drugs and their characteristics (eg, mechanisms, clinical use, and important side effects) is high yield. Focus on understanding the prototype drugs in each class. Avoid memorizing obscure derivatives. Learn the "classic" and distinguishing toxicities of the major drugs. Do not bother with drug dosages or trade names. Reviewing associated biochemistry, physiology, and microbiology can be useful while studying pharmacology. There is a strong emphasis on ANS, CNS, antimicrobial, and cardiovascular agents as well as NSAIDs. Much of the material is clinically relevant. Newer drugs on the market are also fair game.

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

Cardiovascular

"As for me, except for an occasional heart attack, I feel as young as I ever did."

—Robert Benchley

"Hearts will never be practical until they are made unbreakable."

—The Wizard of Oz

"As the arteries grow hard, the heart grows soft."

—H. L. Mencken

"Nobody has ever measured, not even poets, how much the heart can hold."

—Zelda Fitzgerald

"Only from the heart can you touch the sky."

—Rumi

"It is not the size of the man but the size of his heart that matters."

—Evander Holyfield

The cardiovascular system is one of the highest yield areas for the boards and, for some students, may be the most challenging. Focusing on understanding the mechanisms instead of memorizing the details can make a big difference, especially for this topic. Pathophysiology of atherosclerosis and heart failure, MOA of drugs (particular physiology interactions) and their adverse effects, ECGs of heart blocks, the cardiac cycle, and the Starling curve are some of the more high-yield topics. Differentiating between systolic and diastolic dysfunction is also very important. Heart murmurs and maneuvers that affect these murmurs have also been high yield and may be asked in a multimedia format.

- Embryology 280
- Anatomy 283
- Physiology 284
- Pathology 298
- Pharmacology 316

FAS1_2019_07-Cardio.indd 279 11/7/19 4:24 PM

(c) ketabton.com: The Digital Library

▸ CARDIOVASCULAR—EMBRYOLOGY

Heart morphogenesis

First functional organ in vertebrate embryos; beats spontaneously by week 4 of development.

Cardiac looping

Primary heart tube loops to establish left-right polarity; begins in week 4 of development.

Defect in left-right Dynein (involved in L/R asymmetry) can lead to Dextrocardia, as seen in Kartagener syndrome (1° ciliary Dyskinesia).



Septation of the chambers

Atria

1. Septum primum grows toward endocardial cushions, narrowing foramen primum.
2. Foramen secundum forms in septum primum (foramen primum regresses).
3. Septum secundum develops on the right side of septum primum, as foramen secundum maintains right-to-left shunt.
4. Septum secundum expands and covers most of foramen secundum. The residual foramen is the foramen ovale.
5. Remaining portion of septum primum forms the one-way valve of the foramen ovale.
6. Septum primum closes against septum secundum, sealing the foramen ovale soon after birth because of ↑ LA pressure and ↓ RA pressure.
7. Septum secundum and septum primum fuse during infancy/early childhood, forming the atrial septum.

Patent foramen ovale—caused by failure of septum primum and septum secundum to fuse after birth; most are left untreated. Can lead to paradoxical emboli (venous thromboemboli entering the systemic arterial circulation) as can occur in ASD.



(c) ketabton.com: The Digital Library

Heart morphogenesis *(continued)*

Ventricles

❶ Muscular interventricular septum forms. Opening is called interventricular foramen.
❷ Aorticopulmonary septum rotates and fuses with muscular ventricular septum to form membranous interventricular septum, closing interventricular foramen.
❸ Growth of endocardial cushions separates atria from ventricles and contributes to both atrial septation and membranous portion of the interventricular septum.

Ventricular septal defect—most common congenital cardiac anomaly, usually occurs in membranous septum.



Outflow tract formation

Neural crest and endocardial cell migrations → truncal and bulbar ridges that spiral and fuse to form aorticopulmonary septum → ascending aorta and pulmonary trunk.

Conotruncal abnormalities associated with failure of neural crest cells to migrate:
- Transposition of great vessels.
- Tetralogy of Fallot.
- Persistent truncus arteriosus.

Valve development

Aortic/pulmonary: derived from endocardial cushions of outflow tract.
Mitral/tricuspid: derived from fused endocardial cushions of the AV canal.

Valvular anomalies may be stenotic, regurgitant, atretic (eg, tricuspid atresia), or displaced (eg, Ebstein anomaly).

Heart embryology

EMBRYONIC STRUCTURE	GIVES RISE TO
Truncus arteriosus	Ascending aorta and pulmonary trunk
Bulbus cordis	Smooth parts (outflow tract) of left and right ventricles
Primitive ventricle	Trabeculated part of left and right ventricles
Primitive atrium	Trabeculated part of left and right atria
Left horn of sinus venosus	Coronary sinus
Right horn of sinus venosus	Smooth part of right atrium (sinus venarum)
Endocardial cushion	Atrial septum, membranous interventricular septum; AV and semilunar valves
Right common cardinal vein and right anterior cardinal vein	Superior vena cava (SVC)
Posterior, subcardinal, and supracardinal veins	Inferior vena cava (IVC)
Primitive pulmonary vein	Smooth part of left atrium

(c) ketabton.com: The Digital Library

Fetal circulation



Blood in umbilical vein has a Po_2 of ≈ 30 mm Hg and is ≈ 80% saturated with O_2. Umbilical arteries have low O_2 saturation.

3 important shunts:

1. Blood entering fetus through the umbilical vein is conducted via the **ductus venosus** into the IVC, bypassing hepatic circulation.
2. Most of the highly **O**xygenated blood reaching the heart via the IVC is directed through the **foramen O**vale into the left atrium.
3. **D**eoxygenated blood from the SVC passes through the RA → RV → main pulmonary artery → **Ductus arteriosus** → **D**escending aorta; shunt is due to high fetal pulmonary artery resistance (due partly to low O_2 tension).

At birth, infant takes a breath → ↓ resistance in pulmonary vasculature → ↑ left atrial pressure vs right atrial pressure→ foramen ovale closes (now called fossa ovalis); ↑ in O_2 (from respiration) and ↓ in prostaglandins (from placental separation) → closure of ductus arteriosus.

Indomethacin helps **close** the patent **D**uctus arteriosus → ligamentum arteriosum (remnant of ductus arteriosus). Come **In** and **close** the **D**oor.

Prostaglandins E_1 and E_2 k**EE**p PDA open.

Fetal-postnatal derivatives

FETAL STRUCTURE	POSTNATAL DERIVATIVE	NOTES
Ductus arteriosus	Ligamentum arteriosum	Near the left recurrent laryngeal nerve
Ductus venosus	Ligamentum venosum	
Foramen ovale	**F**ossa **o**valis	
Allantois → urachus	Medi**an** umbilical ligament	Urachus is part of allantoic duct between bladder and umbilicus
Umbilical arteries	Medi**al** umbilical ligaments	
Umbilical vein	Ligamentum teres hepatis (round ligament)	Contained in falciform ligament
Notochord	**N**ucleus pulposus	

(c) ketabton.com: The Digital Library

▸ CARDIOVASCULAR—ANATOMY

Anatomy of the heart



LA is the most posterior part of the heart A ; enlargement of the LA (eg, in mitral stenosis) can lead to compression of the esophagus (dysphagia) and/or the left recurrent laryngeal nerve, a branch of the vagus nerve, causing hoarseness (Ortner syndrome).

RV is the most anterior part of the heart and most commonly injured in trauma.

Pericardium

Consists of 3 layers (from outer to inner):
- Fibrous pericardium
- Parietal layer of serous pericardium
- Visceral layer of serous pericardium

Pericardial cavity lies between parietal and visceral layers.

Pericardium innervated by phrenic nerve.

Pericarditis can cause referred pain to the neck, arms, or one or both shoulders (often left).

Coronary blood supply

LAD and its branches supply anterior 2/3 of interventricular septum, anterolateral papillary muscle, and anterior surface of LV. Most commonly occluded.

PDA supplies AV node (dependent on dominance), posterior 1/3 of interventricular septum, posterior 2/3 walls of ventricles, and posteromedial papillary muscle.

RCA supplies SA node (blood supply independent of dominance). Infarct may cause nodal dysfunction (bradycardia or heart block). Right (acute) marginal artery supplies RV.

Dominance:
- Right-dominant circulation (85%) = PDA arises from RCA.
- Left-dominant circulation (8%) = PDA arises from LCX.
- Codominant circulation (7%) = PDA arises from both LCX and RCA.

Coronary blood flow peaks in early diastole.

Key:
AMA = Acute marginal artery
LAD = Left anterior descending artery
LCA = Left coronary artery
LCX = Left circumflex artery
OMA = Obtuse marginal artery
PDA = Posterior descending artery
PT = Pulmonary trunk
PV = Pulmonary vein
RCA = Right coronary artery



Anterior view Posterior view

(c) ketabton.com: The Digital Library

▸ CARDIOVASCULAR—PHYSIOLOGY

Cardiac output variables

Stroke volume	Stroke Volume affected by Contractility, Afterload, and Preload. ↑ SV with: ▪ ↑ Contractility (eg, anxiety, exercise) ▪ ↑ Preload (eg, early pregnancy) ▪ ↓ Afterload	SV CAP. A failing heart has ↓ SV (systolic and/or diastolic dysfunction).
Contractility	Contractility (and SV) ↑ with: ▪ Catecholamine stimulation via β_1 receptor: ▪ Activated protein kinase A → phospholamban phosphorylation → active Ca^{2+} ATPase → ↑ Ca^{2+} storage in sarcoplasmic reticulum ▪ Activated protein kinase A → Ca^{2+} channel phosphorylation → ↑ Ca^{2+} entry → ↑ Ca^{2+}-induced Ca^{2+} release ▪ ↑ intracellular Ca^{2+} ▪ ↓ extracellular Na^+ (↓ activity of Na^+/Ca^{2+} exchanger) ▪ Digitalis (blocks Na^+/K^+ pump → ↑ intracellular Na^+ → ↓ Na^+/Ca^{2+} exchanger activity → ↑ intracellular Ca^{2+})	Contractility (and SV) ↓ with: ▪ β_1-blockade (↓ cAMP) ▪ HF with systolic dysfunction ▪ Acidosis ▪ Hypoxia/hypercapnia (↓ P_{O_2}/↑ P_{CO_2}) ▪ Non-dihydropyridine Ca^{2+} channel blockers
Preload	Preload approximated by ventricular EDV; depends on venous tone and circulating blood volume.	Vasodilators (eg, nitroglycerin) ↓ preload.
Afterload	Afterload approximated by MAP. ↑ wall tension per Laplace's law → ↑ pressure → ↑ afterload. LV compensates for ↑ afterload by thickening (hypertrophy) in order to ↓ wall stress.	Arterial vasodilators (eg, hydralazine) ↓ Afterload. ACE inhibitors and ARBs ↓ both preload and afterload. Chronic hypertension (↑ MAP) → LV hypertrophy.
Myocardial oxygen demand	Myocardial O_2 demand is ↑ by: ▪ ↑ Contractility ▪ ↑ Afterload (proportional to arterial pressure) ▪ ↑ heart Rate ▪ ↑ Diameter of ventricle (↑ wall tension)	Wall tension follows Laplace's law: Wall tension = pressure × radius $\text{Wall stress} = \frac{\text{pressure} \times \text{radius}}{2 \times \text{wall thickness}}$

(c) ketabton.com: The Digital Library

Cardiac output equations

	EQUATION	NOTES
Stroke volume	$SV = EDV - ESV$	
Ejection fraction	$EF = \frac{SV}{EDV} = \frac{EDV - ESV}{EDV}$	EF is an index of ventricular contractility (↓ in systolic HF; usually normal in diastolic HF).
Cardiac output	$CO = SV \times HR$ Fick principle: $CO = \frac{\text{rate of } O_2 \text{ consumption}}{(\text{arterial } O_2 \text{ content} - \text{venous } O_2 \text{ content})}$	In early stages of exercise, CO maintained by ↑ HR and ↑ SV. In later stages, CO maintained by ↑ HR only (SV plateaus). Diastole is shortened with ↑↑ HR (eg, ventricular tachycardia) → ↓ diastolic filling time → ↓ SV → ↓ CO.
Pulse pressure	$PP = SBP - DBP$	PP directly proportional to SV and inversely proportional to arterial compliance. ↑ PP in hyperthyroidism, aortic regurgitation, aortic stiffening (isolated systolic hypertension in elderly), obstructive sleep apnea (↑ sympathetic tone), anemia, exercise (transient). ↓ PP in aortic stenosis, cardiogenic shock, cardiac tamponade, advanced HF.
Mean arterial pressure	$MAP = CO \times TPR$	MAP (at resting HR) = 2/3 DBP + 1/3 SBP = DBP + 1/3 PP.

Starling curves



Force of contraction is proportional to end-diastolic length of cardiac muscle fiber (preload).
↑ contractility with catecholamines, positive inotropes (eg, digoxin).
↓ contractility with loss of functional myocardium (eg, MI), β-blockers (acutely), non-dihydropyridine Ca^{2+} channel blockers, dilated cardiomyopathy.

(c) ketabton.com: The Digital Library

Resistance, pressure, flow

$\Delta P = Q \times R$

Similar to Ohm's law: $\Delta V = I \times R$

Volumetric flow rate (Q) = flow velocity (v) × cross-sectional area (A)

Resistance $= \frac{\text{driving pressure } (\Delta P)}{Q} = \frac{8\eta \text{ (viscosity)} \times \text{length}}{\pi r^4}$

Total resistance of vessels in series: $R_T = R_1 + R_2 + R_3 \ldots$

Total resistance of vessels in parallel: $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} \ldots$

Capillaries have highest total cross-sectional area and lowest flow velocity.

Pressure gradient drives flow from high pressure to low pressure.

Arterioles account for most of TPR. Veins provide most of blood storage capacity.

Viscosity depends mostly on hematocrit.

Viscosity ↑ in hyperproteinemic states (eg, multiple myeloma), polycythemia.

Viscosity ↓ in anemia.

Cardiac and vascular function curves



Intersection of curves = operating point of heart (ie, venous return and CO are equal, as circulatory system is a closed system).

GRAPH	EFFECT	EXAMPLES
A Inotropy	Changes in contractility → altered SV → altered CO/VR and RA pressure (RAP)	1 Catecholamines, digoxin, exercise ⊕ 2 HF with reduced EF, narcotic overdose, sympathetic inhibition ⊖
B Venous return	Changes in circulating volume → altered RAP → altered SV → change in CO	3 Fluid infusion, sympathetic activity ⊕ 4 Acute hemorrhage, spinal anesthesia ⊖
C Total peripheral resistance	Changes in TPR → altered CO Change in RAP unpredictable.	5 Vasopressors ⊕ 6 Exercise, AV shunt ⊖

Changes often occur in tandem, and may be reinforcing (eg, exercise ↑ inotropy and ↓ TPR to maximize CO) or compensatory (eg, HF ↓ inotropy → fluid retention to ↑ preload to maintain CO).

(c) ketabton.com: The Digital Library

Pressure-volume loops and cardiac cycle



The black loop represents normal cardiac physiology.

Phases—left ventricle:

1. Isovolumetric contraction—period between mitral valve closing and aortic valve opening; period of highest O_2 consumption
2. Systolic ejection—period between aortic valve opening and closing
3. Isovolumetric relaxation—period between aortic valve closing and mitral valve opening
4. Rapid filling—period just after mitral valve opening
5. Reduced filling—period just before mitral valve closing



Heart sounds:

S1—mitral and tricuspid valve closure. Loudest at mitral area.

S2—aortic and pulmonary valve closure. Loudest at left upper sternal border.

S3—in early diastole during rapid ventricular filling phase. Best heard at apex with patient in left lateral decubitus position. Associated with ↑ filling pressures (eg, MR, AR, HF, thyrotoxicosis) and more common in dilated ventricles (but can be normal in children, young adults, athletes, and pregnancy).

S4—in late diastole ("atrial kick"). Best heard at apex with patient in left lateral decubitus position. High atrial pressure. Associated with ventricular noncompliance (eg, hypertrophy). Left atrium must push against stiff LV wall. Considered abnormal if palpable.

Jugular venous pulse (JVP):

a wave—**a**trial contraction. Absent in atrial fibrillation (AF).

c wave—RV **c**ontraction (closed tricuspid valve bulging into atrium).

x descent—downward displacement of closed tricuspid valve during rapid ventricular ejection phase. Reduced or absent in tricuspid regurgitation and right HF because pressure gradients are reduced.

v wave—↑ right atrial pressure due to filling ("**v**illing") against closed tricuspid **v**alve.

y descent—RA empt**y**ing into RV. Prominent in constrictive pericarditis, absent in cardiac tamponade.

(c) ketabton.com: The Digital Library

Pressure-volume loops and valvular disease

Aortic stenosis



↑ LV pressure
↑ ESV
No change in EDV
↓ SV

Ventricular hypertrophy → ↓ ventricular compliance → ↑ EDP for given EDV

Mitral regurgitation



No true isovolumetric phase
↓ ESV due to ↓ resistance and ↑ regurgitation into LA during systole
↑ EDV due to ↑ LA volume/pressure from regurgitation → ↑ ventricular filling
↑ SV

Aortic regurgitation



No true isovolumetric phase
↑ EDV
↑ SV

Mitral stenosis



↑ LA pressure
↓ EDV because of impaired ventricular filling
↓ ESV
↓ SV

(c) ketabton.com: The Digital Library

Splitting of S2

Physiologic splitting	Inspiration → drop in intrathoracic pressure → ↑ venous return → ↑ RV filling → ↑ RV stroke volume → ↑ RV ejection time → delayed closure of pulmonic valve. ↓ pulmonary impedance (↑ capacity of the pulmonary circulation) also occurs during inspiration, which contributes to delayed closure of pulmonic valve.	
Wide splitting	Seen in conditions that delay RV emptying (eg, pulmonic stenosis, right bundle branch block). Causes delayed pulmonic sound (especially on inspiration). An exaggeration of normal splitting.	
Fixed splitting	Heard in ASD. ASD → left-to-right shunt → ↑ RA and RV volumes → ↑ flow through pulmonic valve → delayed pulmonic valve closure (independent of respiration).	
Paradoxical splitting	Heard in conditions that delay aortic valve closure (eg, aortic stenosis, left bundle branch block). Normal order of semilunar valve closure is reversed so that P2 sound occurs before delayed A2 sound. On inspiration, P2 closes later and moves closer to A2, "paradoxically" eliminating the split. On expiration, the split can be heard (opposite to physiologic splitting).	

(c) ketabton.com: The Digital Library

Auscultation of the heart



MANEUVER	CARDIOVASCULAR CHANGES	MURMURS THAT INCREASE WITH MANEUVER	MURMURS THAT DECREASE WITH MANEUVER
Standing Valsalva (strain phase)	↓ preload (↓ LV volume)	MVP (↓ LV volume) HCM (↓ LV volume)	Most murmurs (↓ flow through stenotic or regurgitant valve)
Passive leg raise	↑ preload (↑ LV volume)	Most murmurs (↑ flow through stenotic or regurgitant valve)	MVP (↑ LV volume) HCM (↑ LV volume)
Squatting	↑ preload, ↑ afterload (↑ LV volume)		
Hand grip	↑↑ afterload → ↑ reverse flow across aortic valve (↑ LV volume)	Most other left-sided murmurs (AR, MR, VSD)	AS (↓ transaortic valve pressure gradient) HCM (↑ LV volume)
Inspiration	↑ venous return to right heart, ↓ venous return to left heart	Most right-sided murmurs	Most left-sided murmurs

(c) ketabton.com: The Digital Library

Heart murmurs

Systolic

Aortic stenosis



Crescendo-decrescendo systolic ejection murmur and soft S2 (ejection click may be present). LV >> aortic pressure during systole. Loudest at heart base; radiates to carotids. "Pulsus parvus et tardus"—pulses are weak with a delayed peak. Can lead to **S**yncope, **A**ngina, and **D**yspnea on exertion (**SAD**). Most commonly due to age-related calcification in older patients (> 60 years old) or in younger patients with early-onset calcification of bicuspid aortic valve.

Mitral/tricuspid regurgitation



Holosystolic, high-pitched "blowing murmur."
Mitral—loudest at apex and radiates toward axilla. MR is often due to ischemic heart disease (post-MI), MVP, LV dilatation.
Tricuspid—loudest at tricuspid area. TR commonly caused by RV dilatation.
Rheumatic fever and infective endocarditis can cause either MR or TR.

Mitral valve prolapse



Late systolic crescendo murmur with midsystolic click (MC) due to sudden tensing of chordae tendineae as mitral leaflets prolapse into the LA (**C**hordae cause **C**rescendo with **C**lick). Most frequent valvular lesion. Best heard over apex. Loudest just before S2. Usually benign. Can predispose to infective endocarditis. Can be caused by myxomatous degeneration (1° or 2° to connective tissue disease such as Marfan or Ehlers-Danlos syndrome), rheumatic fever (particularly in developing countries), chordae rupture.

Ventricular septal defect



Holosystolic, harsh-sounding murmur. Loudest at tricuspid area. Larger VSDs have a lower intensity murmur than smaller VSDs.

Diastolic

Aortic regurgitation



High-pitched "blowing" early diastolic decrescendo murmur. Best heard at base (aortic root dilation) or left sternal border (valvular disease). Long diastolic murmur, hyperdynamic pulse, and head bobbing when severe and chronic. Wide pulse pressure. Causes include **B**icuspid aortic valve, **E**ndocarditis, **A**ortic root dilation, **R**heumatic fever (**BEAR**). Progresses to left HF.

Mitral stenosis



Follows opening snap (OS; due to abrupt halt in leaflet motion in diastole, after rapid opening due to fusion at leaflet tips). Delayed rumbling mid-to-late diastolic murmur (↓ interval between S2 and OS correlates with ↑ severity). LA >> LV pressure during diastole.
Often a late (and highly specific) sequela of rheumatic fever. Chronic MS can result in pulmonary congestion/hypertension and LA dilation → atrial fibrillation and Ortner syndrome.

Continuous

Patent ductus arteriosus



Continuous machine-like murmur. Best heard at left infraclavicular area. Loudest at S2. Often due to congenital rubella or prematurity.
"**PDA**s (**P**ublic **D**isplays of **A**ffection) are **continuous**ly annoying."

FAS1_2019_07-Cardio.indd 291 11/7/19 4:24 PM

(c) ketabton.com: The Digital Library

Myocardial action potential

Phase 0 = rapid upstroke and depolarization—voltage-gated Na^+ channels open.
Phase 1 = initial repolarization—inactivation of voltage-gated Na^+ channels. Voltage-gated K^+ channels begin to open.
Phase 2 = plateau—Ca^{2+} influx through voltage-gated Ca^{2+} channels balances K^+ efflux. Ca^{2+} influx triggers Ca^{2+} release from sarcoplasmic reticulum and myocyte contraction.
Phase 3 = rapid repolarization—massive K^+ efflux due to opening of voltage-gated slow delayed-rectifier K^+ channels and closure of voltage-gated Ca^{2+} channels.
Phase 4 = resting potential—high K^+ permeability through K^+ channels.

In contrast to skeletal muscle:

- Cardiac muscle action potential has a plateau due to Ca^{2+} influx and K^+ efflux.
- Cardiac muscle contraction requires Ca^{2+} influx from ECF to induce Ca^{2+} release from sarcoplasmic reticulum (Ca^{2+}-induced Ca^{2+} release).
- Cardiac myocytes are electrically coupled to each other by gap junctions.



Occurs in all cardiac myocytes except for those in the SA and AV nodes.

Pacemaker action potential

Occurs in the SA and AV nodes. Key differences from the ventricular action potential include:
Phase 0 = upstroke—opening of voltage-gated Ca^{2+} channels. Fast voltage-gated Na^+ channels are permanently inactivated because of the less negative resting potential of these cells. Results in a slow conduction velocity that is used by the AV node to prolong transmission from the atria to ventricles.
Phases 1 and 2 are absent.
Phase 3 = repolarization—inactivation of the Ca^{2+} channels and ↑ activation of K^+ channels → ↑ K^+ efflux.
Phase 4 = slow spontaneous diastolic depolarization due to I_f ("funny current"). I_f channels responsible for a slow, mixed Na^+/K^+ inward current; different from I_{Na} in phase 0 of ventricular action potential. Accounts for automaticity of SA and AV nodes. The slope of phase 4 in the SA node determines HR. ACh/adenosine ↓ the rate of diastolic depolarization and ↓ HR, while catecholamines ↑ depolarization and ↑ HR. Sympathetic stimulation ↑ the chance that I_f channels are open and thus ↑ HR.



(c) ketabton.com: The Digital Library

Electrocardiogram

Conduction pathway: SA node → atria → AV node → bundle of His → right and left bundle branches → Purkinje fibers → ventricles; left bundle branch divides into left anterior and posterior fascicles.

SA node—located at junction of RA and SVC; "pacemaker" inherent dominance with slow phase of upstroke.

AV node—located in posteroinferior part of interatrial septum. Blood supply usually from RCA. 100-msec delay allows time for ventricular filling.

Pacemaker rates: SA > AV > bundle of His/Purkinje/ventricles.

Speed of conduction: **H**is-**P**urkinje > **At**ria > **Vent**ricles > **AV** node. **H**e **P**arks **At** **Vent**ura **Av**enue.

P wave—atrial depolarization.

PR interval—time from start of atrial depolarization to start of ventricular depolarization (normally 120-200 msec).

QRS complex—ventricular depolarization (normally < 100 msec).

QT interval—ventricular depolarization, mechanical contraction of the ventricles, ventricular repolarization.

T wave—ventricular repolarization. T-wave inversion may indicate ischemia or recent MI.

J point—junction between end of QRS complex and start of ST segment.

ST segment—isoelectric, ventricles depolarized.

U wave—prominent in hypokalemia (think hyp"**U**"kalemia), bradycardia.





(c) ketabton.com: The Digital Library

Torsades de pointes



Polymorphic ventricular tachycardia, characterized by shifting sinusoidal waveforms on ECG; can progress to ventricular fibrillation (VF). Long QT interval predisposes to torsades de pointes. Caused by drugs, ↓ K^+, ↓ Mg^{2+}, ↓ Ca^{2+}, congenital abnormalities. Treatment includes magnesium sulfate.

Drug-induced long QT (**ABCDE**):
AntiArrhythmics (class IA, III)
AntiBiotics (eg, macrolides)
Anti"C"ychotics (eg, haloperidol)
AntiDepressants (eg, TCAs)
AntiEmetics (eg, ondansetron)
Torsades de pointes = twisting of the points

Congenital long QT syndrome

Inherited disorder of myocardial repolarization, typically due to ion channel defects (most commonly loss-of-function mutations affecting K^+ channels); ↑ risk of sudden cardiac death (SCD) due to torsades de pointes. Includes:

- Romano-Ward syndrome—autosomal dominant, pure cardiac phenotype (**no** deafness).
- Jervell and Lange-Nielsen syndrome—autosomal recessive, sensorineural deafness.

Brugada syndrome

Autosomal dominant disorder most common in Asian males. ECG pattern of pseudo-right bundle branch block and ST elevations in V_1-V_3. ↑ risk of ventricular tachyarrhythmias and SCD. Prevent SCD with implantable cardioverter-defibrillator (ICD).

Wolff-Parkinson-White syndrome

Most common type of ventricular pre-excitation syndrome. Abnormal fast accessory conduction pathway from atria to ventricle (bundle of Kent) bypasses the rate-slowing AV node → ventricles begin to partially depolarize earlier → characteristic delta wave with widened QRS complex and shortened PR interval on ECG. May result in reentry circuit → supraventricular tachycardia.



(c) ketabton.com: The Digital Library

ECG tracings

RHYTHM	DESCRIPTION	EXAMPLE
Atrial fibrillation	Chaotic and erratic baseline with no discrete P waves in between irregularly spaced QRS complexes. Irregularly irregular heartbeat. Most common risk factors include hypertension and coronary artery disease (CAD). Occasionally seen after binge drinking ("holiday heart syndrome"). Can lead to thromboembolic events, particularly stroke. Treatment: anticoagulation, rate and rhythm control and/or cardioversion.	
Atrial flutter	A rapid succession of identical, back-to-back atrial depolarization waves. The identical appearance accounts for the "sawtooth" appearance of the flutter waves. Treat like atrial fibrillation +/– catheter ablation.	
Ventricular fibrillation	A completely erratic rhythm with no identifiable waves. Fatal arrhythmia without immediate CPR and defibrillation.	
AV block		
First-degree AV block	The PR interval is prolonged (> 200 msec). Benign and asymptomatic. No treatment required.	
Second-degree AV block		
Mobitz type I (Wenckebach)	Progressive lengthening of PR interval until a beat is "dropped" (a P wave not followed by a QRS complex). Usually asymptomatic. Variable RR interval with a pattern (regularly irregular).	
Mobitz type II	Dropped beats that are not preceded by a change in the length of the PR interval (as in type I). May progress to 3rd-degree block. Often treated with pacemaker.	
Third-degree (complete) AV block	The atria and ventricles beat independently of each other. P waves and QRS complexes not rhythmically associated. Atrial rate > ventricular rate. Usually treated with pacemaker. Can be caused by Lym3 disease.	

(c) ketabton.com: The Digital Library

Atrial natriuretic peptide

Released from **atrial myocytes** in response to ↑ blood volume and atrial pressure. Acts via cGMP. Causes vasodilation and ↓ Na^+ reabsorption at the renal collecting tubule. Dilates afferent renal arterioles and constricts efferent arterioles, promoting diuresis and contributing to "aldosterone escape" mechanism.

B-type (brain) natriuretic peptide

Released from **ventricular myocytes** in response to ↑ tension. Similar physiologic action to ANP, with longer half-life. BNP blood test used for diagnosing HF (very good negative predictive value). Available in recombinant form (nesiritide) for treatment of HF.

Baroreceptors and chemoreceptors



Receptors:

- Aortic arch transmits via vagus nerve to solitary nucleus of medulla (responds to changes in BP).
- Carotid sinus (dilated region at carotid bifurcation) transmits via glossopharyngeal nerve to solitary nucleus of medulla (responds to changes in BP).

Baroreceptors:

- Hypotension—↓ arterial pressure → ↓ stretch → ↓ afferent baroreceptor firing → ↑ efferent sympathetic firing and ↓ efferent parasympathetic stimulation → vasoconstriction, ↑ HR, ↑ contractility, ↑ BP. Important in the response to severe hemorrhage.
- Carotid massage—↑ pressure on carotid sinus → ↑ stretch → ↑ afferent baroreceptor firing → ↑ AV node refractory period → ↓ HR.
- Component of Cushing reflex (triad of hypertension, bradycardia, and respiratory depression)—↑ intracranial pressure constricts arterioles → cerebral ischemia → ↑ pCO_2 and ↓ pH → central reflex sympathetic ↑ in perfusion pressure (hypertension) → ↑ stretch → peripheral reflex baroreceptor–induced bradycardia.

Chemoreceptors:

- Peripheral—carotid and aortic bodies are stimulated by ↑ P_{CO_2}, ↓ pH of blood, and ↓ P_{O_2} (< 60 mm Hg).
- Central—are stimulated by changes in pH and P_{CO_2} of brain interstitial fluid, which in turn are influenced by arterial CO_2 as H^+ cannot cross the blood-brain barrier. Do not directly respond to P_{O_2}. Central chemoreceptors become less responsive with chronically ↑ P_{CO_2} (eg, COPD) → ↑ dependence on peripheral chemoreceptors to detect ↓ O_2 to drive respiration.

FAS1_2019_07-Cardio.indd 296 11/7/19 4:24 PM

(c) ketabton.com: The Digital Library

Normal cardiac pressures

Pulmonary capillary wedge pressure (PCWP; in mm Hg) is a good approximation of left atrial pressure. In mitral stenosis, PCWP > LV end diastolic pressure. PCWP is measured with pulmonary artery catheter (Swan-Ganz catheter).



Autoregulation

How blood flow to an organ remains constant over a wide range of perfusion pressures.

ORGAN	FACTORS DETERMINING AUTOREGULATION
Heart	Local metabolites (vasodilatory): adenosine, NO, CO_2, ↓ O_2
Brain	Local metabolites (vasodilatory): CO_2 (pH)
Kidneys	Myogenic and tubuloglomerular feedback
Lungs	Hypoxia causes vasoconstriction
Skeletal muscle	Local metabolites during exercise (vasodilatory): CO_2, H^+, Adenosine, Lactate, K^+ At rest: sympathetic tone in arteries
Skin	Sympathetic vasoconstriction most important mechanism for temperature control

The pulmonary vasculature is unique in that alveolar hypoxia causes vasoconstriction so that only well-ventilated areas are perfused. In other organs, hypoxia causes vasodilation.

CHALK

Capillary fluid exchange

Starling forces determine fluid movement through capillary membranes:

- P_c = capillary hydrostatic pressure—pushes fluid out of capillary
- P_i = interstitial hydrostatic pressure—pushes fluid into capillary
- π_c = plasma colloid osmotic (oncotic) pressure—pulls fluid into capillary
- π_i = interstitial fluid colloid osmotic pressure—pulls fluid out of capillary

J_v = net fluid flow = $K_f [(P_c - P_i) - \sigma(\pi_c - \pi_i)]$

K_f = capillary permeability to fluid

σ = reflection coefficient (measure of capillary permeability to protein)

Edema—excess fluid outflow into interstitium commonly caused by:

- ↑ capillary pressure (↑ P_c; eg, HF)
- ↑ capillary permeability (↑ K_f; eg, toxins, infections, burns)
- ↑ interstitial fluid colloid osmotic pressure (↑ π_i; eg, lymphatic blockage)
- ↓ plasma proteins (↓ π_c; eg, nephrotic syndrome, liver failure, protein malnutrition)



(c) ketabton.com: The Digital Library

▸ CARDIOVASCULAR—PATHOLOGY

Congenital heart diseases

RIGHT-TO-LEFT SHUNTS	Early cyanosis—"blue babies." Often diagnosed prenatally or become evident immediately after birth. Usually require urgent surgical treatment and/or maintenance of a PDA.	The 5 T's: 1. Truncus arteriosus (1 vessel) 2. Transposition (2 switched vessels) 3. Tricuspid atresia (3 = Tri) 4. Tetralogy of Fallot (4 = Tetra) 5. TAPVR (5 letters in the name)
Persistent truncus arteriosus	Truncus arteriosus fails to divide into pulmonary trunk and aorta due to failure of aorticopulmonary septum formation; most patients have accompanying VSD.	
D-transposition of great vessels	Aorta leaves RV (anterior) and pulmonary trunk leaves LV (posterior) → separation of systemic and pulmonary circulations. Not compatible with life unless a shunt is present to allow mixing of blood (eg, VSD, PDA, or patent foramen ovale). Due to failure of the aorticopulmonary septum to spiral ("egg on a string" appearance on CXR). Without surgical intervention, most infants die within the first few months of life.	
Tricuspid atresia	Absence of tricuspid valve and hypoplastic RV; requires both ASD and VSD for viability.	
Tetralogy of Fallot 	Caused by anterosuperior displacement of the infundibular septum. Most common cause of early childhood cyanosis. ❶ Pulmonary infundibular stenosis (most important determinant for prognosis) ❷ Right ventricular hypertrophy (RVH)—boot-shaped heart on CXR A ❸ Overriding aorta ❹ VSD Pulmonary stenosis forces right-to-left flow across VSD → RVH, "tet spells" (often caused by crying, fever, and exercise due to exacerbation of RV outflow obstruction).	PROVe. Squatting: ↑ SVR, ↓ right-to-left shunt, improves cyanosis. Associated with 22q11 syndromes. 
Total anomalous pulmonary venous return	Pulmonary veins drain into right heart circulation (SVC, coronary sinus, etc); associated with ASD and sometimes PDA to allow for right-to-left shunting to maintain CO.	
Ebstein anomaly	Displacement of tricuspid valve leaflets downward into RV, artificially "atrializing" the ventricle. Associated with tricuspid regurgitation, accessory conduction pathways, right-sided HF.	Can be caused by lithium exposure in utero.

(c) ketabton.com: The Digital Library

Congenital heart diseases *(continued)*

LEFT-TO-RIGHT SHUNTS	Acyanotic at presentation; cyanosis may occur years later. Frequency: VSD > ASD > PDA.	**R**ight-to-**L**eft shunts: ea**RL**y cyanosis. **L**eft-to-**R**ight shunts: "**L**ate**R**" cyanosis.
Ventricular septal defect 	Asymptomatic at birth, may manifest weeks later or remain asymptomatic throughout life. Most self resolve; larger lesions B may lead to LV overload and HF.	O_2 saturation ↑ in RV and pulmonary artery.
Atrial septal defect 	Defect in interatrial septum C; wide, fixed split S2. Ostium secundum defects most common and usually an isolated finding; ostium primum defects rarer and usually occur with other cardiac anomalies. Symptoms range from none to HF. Distinct from patent foramen ovale in that septa are missing tissue rather than unfused.	O_2 saturation ↑ in RA, RV, and pulmonary artery. May lead to paradoxical emboli (systemic venous emboli use ASD to bypass lungs and become systemic arterial emboli). Associated with Down syndrome.
Patent ductus arteriosus 	In fetal period, shunt is right to left (normal). In neonatal period, ↓ pulmonary vascular resistance → shunt becomes left to right → progressive RVH and/or LVH and HF. Associated with a continuous, "machine-like" murmur. Patency is maintained by PGE synthesis and low O_2 tension. Uncorrected PDA D can eventually result in late cyanosis in the lower extremities (differential cyanosis).	PDA is normal in utero and normally closes only after birth.
Eisenmenger syndrome 	Uncorrected left-to-right shunt (VSD, ASD, PDA) → ↑ pulmonary blood flow → pathologic remodeling of vasculature → pulmonary arterial hypertension. RVH occurs to compensate → shunt becomes right to left. Causes late cyanosis, clubbing E, and polycythemia. Age of onset varies.	
OTHER ANOMALIES		
Coarctation of the aorta 	Aortic narrowing F near insertion of ductus arteriosus ("juxtaductal"). Associated with bicuspid aortic valve, other heart defects, and Turner syndrome. Hypertension in upper extremities and weak, delayed pulse in lower extremities (brachial-femoral delay). With age, intercostal arteries enlarge due to collateral circulation; arteries erode ribs → notched appearance on CXR. Complications include HF, ↑ risk of cerebral hemorrhage (berry aneurysms), aortic rupture, and possible endocarditis.	

(c) ketabton.com: The Digital Library

Congenital cardiac defect associations

DISORDER	DEFECT
Alcohol exposure in utero (fetal alcohol syndrome)	VSD, PDA, ASD, tetralogy of Fallot
Congenital rubella	PDA, pulmonary artery stenosis, septal defects
Down syndrome	AV septal defect (endocardial cushion defect), VSD, ASD
Infant of diabetic mother	Transposition of great vessels, VSD
Marfan syndrome	MVP, thoracic aortic aneurysm and dissection, aortic regurgitation
Prenatal lithium exposure	Ebstein anomaly
Turner syndrome	Bicuspid aortic valve, coarctation of aorta
Williams syndrome	Supravalvular aortic stenosis
22q11 syndromes	Truncus arteriosus, tetralogy of Fallot

Hypertension

Persistent systolic BP ≥ 130 mm Hg and/or diastolic BP ≥ 80 mm Hg.

RISK FACTORS

↑ age, obesity, diabetes, physical inactivity, excess salt intake, excess alcohol intake, cigarette smoking, family history; African American > Caucasian > Asian.

FEATURES



90% of hypertension is 1° (essential) and related to ↑ CO or ↑ TPR. Remaining 10% mostly 2° to renal/renovascular diseases such as fibromuscular dysplasia (characteristic "string of beads" appearance of renal artery A, usually seen in women of child-bearing age) and atherosclerotic renal artery stenosis or to 1° hyperaldosteronism.

Hypertensive urgency—severe (≥ 180/≥ 120 mm Hg) hypertension without acute end-organ damage.

Hypertensive emergency—severe hypertension with evidence of acute end-organ damage (eg, encephalopathy, stroke, retinal hemorrhages and exudates, papilledema, MI, HF, aortic dissection, kidney injury, microangiopathic hemolytic anemia, eclampsia).

PREDISPOSES TO

CAD, LVH, HF, atrial fibrillation; aortic dissection, aortic aneurysm; stroke; CKD (hypertensive nephropathy); retinopathy.

(c) ketabton.com: The Digital Library

Hyperlipidemia signs

Xanthomas	Plaques or nodules composed of lipid-laden histiocytes in skin A, especially the eyelids (xanthelasma B).
Tendinous xanthoma	Lipid deposit in tendon C, especially Achilles.
Corneal arcus	Lipid deposit in cornea. Common in elderly (arcus senilis D), but appears earlier in life with hypercholesterolemia.



Arteriosclerosis	Hardening of arteries, with arterial wall thickening and loss of elasticity.
Arteriolosclerosis	Common. Affects small arteries and arterioles. Two types: hyaline (thickening of vessel walls 2° to plasma protein leak into endothelium in essential hypertension or diabetes mellitus A) and hyperplastic ("onion skinning" in severe hypertension B with proliferation of smooth muscle cells).
Mönckeberg sclerosis (Medial calcific sclerosis)	Uncommon. Affects Medium-sized arteries. Calcification of internal elastic lamina and media of arteries → vascular stiffening without obstruction. "Pipestem" appearance on x-ray C. Does not obstruct blood flow; intima not involved.



(c) ketabton.com: The Digital Library

Atherosclerosis

Very common. Disease of elastic arteries and large- and medium-sized muscular arteries; a form of arteriosclerosis caused by buildup of cholesterol plaques in intima.

LOCATION

Abdominal aorta > Coronary artery > Popliteal artery > Carotid artery > circle of Willis.
A CoPy Cat named Willis.

RISK FACTORS

Modifiable: smoking, hypertension, dyslipidemia (↑ LDL, ↓ HDL), diabetes.
Non-modifiable: age, sex (↑ in men and postmenopausal women), family history.

SYMPTOMS

Angina, claudication, but can be asymptomatic.

PROGRESSION



Inflammation important in pathogenesis: endothelial cell dysfunction → macrophage and LDL accumulation → foam cell formation → fatty streaks → smooth muscle cell migration (involves PDGF and FGF), proliferation, and extracellular matrix deposition → fibrous plaque → complex atheromas A → calcification (calcium content correlates with risk of complications).

COMPLICATIONS

Aneurysms, ischemia, infarcts, peripheral vascular disease, thrombus, emboli.



Aortic aneurysm

Localized pathologic dilation of the aorta. May cause abdominal and/or back pain, which is a sign of leaking, dissection, or imminent rupture.

Abdominal aortic aneurysm



Usually associated with atherosclerosis. Risk factors include history of tobacco use, ↑ age, male sex, family history. May present as palpable pulsatile abdominal mass (arrows in A point to outer dilated calcified aortic wall, with partial crescent-shaped non-opacification of aorta due to flap/clot). Most often infrarenal (distal to origin of renal arteries).

Thoracic aortic aneurysm

Associated with cystic medial degeneration. Risk factors include hypertension, bicuspid aortic valve, connective tissue disease (eg, Marfan syndrome). Also associated with 3° syphilis (obliterative endarteritis of the vasa vasorum). Aortic root dilatation may lead to aortic valve regurgitation.

(c) ketabton.com: The Digital Library

Traumatic aortic rupture

Due to trauma and/or deceleration injury, most commonly at aortic isthmus (proximal descending aorta just distal to origin of left subclavian artery). X-ray may reveal widened mediastinum.

Aortic dissection



Longitudinal intimal tear forming a false lumen. Associated with hypertension, bicuspid aortic valve, inherited connective tissue disorders (eg, Marfan syndrome). Can present with tearing, sudden-onset chest pain radiating to the back +/– markedly unequal BP in arms. CXR can show mediastinal widening. Can result in organ ischemia, aortic rupture, death. Two types:

- Stanford type **A** (proximal): involves **A**scending aorta (red arrow in A). May extend to aortic arch or descending aorta (blue arrow in A). May result in acute aortic regurgitation or cardiac tamponade. Treatment: surgery.
- Stanford type **B** (distal): involves only descending aorta (**B**elow left subclavian artery). Treatment: β-blockers, then vasodilators.



(c) ketabton.com: The Digital Library

Ischemic heart disease manifestations

Angina

Chest pain due to ischemic myocardium 2° to coronary artery narrowing or spasm; no myocyte necrosis.

- Stable—usually 2° to atherosclerosis (≥ 70% occlusion); exertional chest pain in classic distribution (usually with ST depression on ECG), resolving with rest or nitroglycerin.
- Vasospastic (also called Prinzmetal or Variant)—occurs at rest 2° to coronary artery spasm; transient ST elevation on ECG. Smoking is a risk factor; hypertension and hypercholesterolemia are not. Triggers include cocaine, alcohol, and triptans. Treat with Ca^{2+} channel blockers, nitrates, and smoking cessation (if applicable).
- Unstable—thrombosis with incomplete coronary artery occlusion; +/– ST depression and/or T-wave inversion on ECG but no cardiac biomarker elevation (unlike NSTEMI); ↑ in frequency or intensity of chest pain or any chest pain at rest.

Coronary steal syndrome

Distal to coronary stenosis, vessels are maximally dilated at baseline. Administration of vasodilators (eg, dipyridamole, regadenoson) dilates normal vessels → blood is shunted toward well-perfused areas → ischemia in myocardium perfused by stenosed vessels. Principle behind pharmacologic stress tests with coronary vasodilators.

Sudden cardiac death

Death from cardiac causes within 1 hour of onset of symptoms, most commonly due to a lethal arrhythmia (eg, VF). Associated with CAD (up to 70% of cases), cardiomyopathy (hypertrophic, dilated), and hereditary ion channelopathies (eg, long QT syndrome, Brugada syndrome). Prevent with ICD.

Chronic ischemic heart disease

Progressive onset of HF over many years due to chronic ischemic myocardial damage.

Myocardial infarction

Most often due to rupture of coronary artery atherosclerotic plaque → acute thrombosis. ↑ cardiac biomarkers (CK-MB, troponins) are diagnostic.

Non–ST-segment elevation MI (NSTEMI)	ST-segment elevation MI (STEMI)
Subendocardial infarcts	Transmural infarcts
Subendocardium (inner 1/3) especially vulnerable to ischemia	Full thickness of myocardial wall involved
ST depression on ECG	ST elevation, pathologic Q waves on ECG





(c) ketabton.com: The Digital Library

Evolution of myocardial infarction

Commonly occluded coronary arteries: LAD > RCA > circumflex.
Symptoms: diaphoresis, nausea, vomiting, severe retrosternal pain, pain in left arm and/or jaw, shortness of breath, fatigue.

TIME	GROSS	LIGHT MICROSCOPE	COMPLICATIONS
0–24 hr	Dark mottling 	Early coagulative necrosis → cell content released into blood; edema, hemorrhage, wavy fibers Reperfusion injury → free radicals and ↑ Ca^{2+} influx → hypercontraction of myofibrils (dark eosinophilic stripes) 	Ventricular arrhythmia, HF, cardiogenic shock
1–3 days		Extensive coagulative necrosis Tissue surrounding infarct shows acute inflammation with neutrophils 	Postinfarction fibrinous pericarditis
3–14 days		Macrophages, then granulation tissue at margins 	Free wall rupture → tamponade; papillary muscle rupture → mitral regurgitation; interventricular septal rupture due to macrophage-mediated structural degradation → left-to-right shunt LV pseudoaneurysm (risk of rupture)
2 weeks to several months		Contracted scar complete 	Dressler syndrome, HF, arrhythmias, true ventricular aneurysm (risk of mural thrombus)

(c) ketabton.com: The Digital Library

Diagnosis of myocardial infarction

In the first 6 hours, ECG is the gold standard.

Cardiac troponin I rises after 4 hours (peaks at 24 hr) and is ↑ for 7–10 days; more specific than other protein markers.

CK-MB rises after 6–12 hours (peaks at 16–24 hr) and is predominantly found in myocardium but can also be released from skeletal muscle. Useful in diagnosing reinfarction following acute MI because levels return to normal after 48 hours.

Large MIs lead to greater elevations in troponin I and CK-MB. Exact curves vary with testing procedure.

ECG changes can include ST elevation (STEMI, transmural infarct), ST depression (NSTEMI, subendocardial infarct), hyperacute (peaked) T waves, T-wave inversion, new left bundle branch block, and pathologic Q waves or poor R wave progression (evolving or old transmural infarct).



ECG localization of STEMI

INFARCT LOCATION	LEADS WITH ST-SEGMENT ELEVATIONS OR Q WAVES
Anteroseptal (LAD)	V_1–V_2
Anteroapical (distal LAD)	V_3–V_4
Anterolateral (LAD or LCX)	V_5–V_6
Lateral (LCX)	I, aVL
InFerior (RCA)	II, III, aVF
Posterior (PDA)	V_7–V_9, ST depression in V_1–V_3 with tall R waves





(c) ketabton.com: The Digital Library

Myocardial infarction complications

Cardiac arrhythmia	Occurs within the first few days after MI. Important cause of death before reaching the hospital and within the first 24 hours post-MI.
Postinfarction fibrinous pericarditis	1–3 days: friction rub.
Papillary muscle rupture	2–7 days: posteromedial papillary muscle rupture A ↑ risk due to single blood supply from posterior descending artery. Can result in severe mitral regurgitation.
Interventricular septal rupture	3–5 days: macrophage-mediated degradation → VSD → ↑ O_2 saturation and pressure in RV.
Ventricular pseudoaneurysm formation	3–14 days: free wall rupture contained by adherent pericardium or scar tissue B; ↓ CO, risk of arrhythmia, embolus from mural thrombus.
Ventricular free wall rupture	5–14 days: free wall rupture C → cardiac tamponade. LV hypertrophy and previous MI protect against free wall rupture. Acute form usually leads to sudden death.
True ventricular aneurysm	2 weeks to several months: outward bulge with contraction ("dyskinesia"), associated with fibrosis.
Dressler syndrome	Several weeks: autoimmune phenomenon resulting in fibrinous pericarditis.
LV failure and pulmonary edema	Can occur 2° to LV infarction, VSD, free wall rupture, papillary muscle rupture with mitral regurgitation.



Acute coronary syndrome treatments

Unstable angina/NSTEMI—Anticoagulation (eg, heparin), antiplatelet therapy (eg, aspirin) + ADP receptor inhibitors (eg, clopidogrel), β-blockers, ACE inhibitors, statins. Symptom control with nitroglycerin and morphine.

STEMI—In addition to above, reperfusion therapy most important (percutaneous coronary intervention preferred over fibrinolysis).

(c) ketabton.com: The Digital Library

Cardiomyopathies

Dilated cardiomyopathy  	Most common cardiomyopathy (90% of cases). Often idiopathic or familial (eg, due to mutation of *TTN* gene encoding the sarcomeric protein titin). Other etiologies include drugs (eg, alcohol, cocaine, doxorubicin), infection (eg, coxsackie B virus, Chagas disease), ischemia (eg, CAD), systemic conditions (eg, hemochromatosis, sarcoidosis, thyrotoxicosis, wet beriberi), peripartum cardiomyopathy. Findings: HF, S3, systolic regurgitant murmur, dilated heart on echocardiogram, balloon appearance of heart on CXR. Treatment: Na^+ restriction, ACE inhibitors, β-blockers, diuretics, mineralocorticoid receptor blockers (eg, spironolactone), digoxin, ICD, heart transplant.	Leads to systolic dysfunction. Dilated cardiomyopathy **A** displays eccentric hypertrophy (sarcomeres added in series). Takotsubo cardiomyopathy: broken heart syndrome—ventricular apical ballooning likely due to increased sympathetic stimulation (eg, stressful situations).
Hypertrophic obstructive cardiomyopathy  	60–70% of cases are familial, autosomal dominant (most commonly due to mutations in genes encoding sarcomeric proteins, such as myosin binding protein C and β-myosin heavy chain). Causes syncope during exercise and may lead to sudden death (eg, in young athletes) due to ventricular arrhythmia. Findings: S4, systolic murmur. May see mitral regurgitation due to impaired mitral valve closure. Treatment: cessation of high-intensity athletics, use of β-blocker or non-dihydropyridine Ca^{2+} channel blockers (eg, verapamil). ICD if syncope occurs.	Diastolic dysfunction ensues. Marked ventricular concentric hypertrophy (sarcomeres added in parallel) **B**, often septal predominance. Myofibrillar disarray and fibrosis. Physiology of HOCM—asymmetric septal hypertrophy and systolic anterior motion of mitral valve → outflow obstruction → dyspnea, possible syncope. Other causes of concentric LV hypertrophy: chronic HTN, Friedreich ataxia.
Restrictive/infiltrative cardiomyopathy	**P**ostradiation fibrosis, **L**öffler endocarditis, **E**ndocardial fibroelastosis (thick fibroelastic tissue in endocardium of young children), **A**myloidosis, **S**arcoidosis, **H**emochromatosis (although dilated cardiomyopathy is more common) (**P**uppy **LEASH**).	Diastolic dysfunction ensues. Can have low-voltage ECG despite thick myocardium (especially in amyloidosis). Löffler endocarditis—associated with hypereosinophilic syndrome; histology shows eosinophilic infiltrates in myocardium.

(c) ketabton.com: The Digital Library

Heart failure

Clinical syndrome of cardiac pump dysfunction → congestion and low perfusion. Symptoms include dyspnea, orthopnea, fatigue; signs include S3 heart sound, rales, jugular venous distention (JVD), pitting edema A.

Systolic dysfunction—reduced EF, ↑ EDV; ↓ contractility often 2° to ischemia/MI or dilated cardiomyopathy.

Diastolic dysfunction—preserved EF, normal EDV; ↓ compliance (↑ EDP) often 2° to myocardial hypertrophy.

Right HF most often results from left HF. Cor pulmonale refers to isolated right HF due to pulmonary cause.

ACE inhibitors or angiotensin II receptor blockers, β-blockers (except in acute decompensated HF), and spironolactone ↓ mortality. Loop and thiazide diuretics are used mainly for symptomatic relief. Hydralazine with nitrate therapy improves both symptoms and mortality in select patients.

Left heart failure

Orthopnea — Shortness of breath when supine: ↑ venous return from redistribution of blood (immediate gravity effect) exacerbates pulmonary vascular congestion.

Paroxysmal nocturnal dyspnea — Breathless awakening from sleep: ↑ venous return from redistribution of blood, reabsorption of peripheral edema, etc.

Pulmonary edema — ↑ pulmonary venous pressure → pulmonary venous distention and transudation of fluid. Presence of hemosiderin-laden macrophages ("HF" cells) in lungs.

Right heart failure

Hepatomegaly (nutmeg liver) — ↑ central venous pressure → ↑ resistance to portal flow. Rarely, leads to "cardiac cirrhosis."

Jugular venous distention — ↑ venous pressure.

Peripheral edema — ↑ venous pressure → fluid transudation.







(c) ketabton.com: The Digital Library

Shock

Inadequate organ perfusion and delivery of nutrients necessary for normal tissue and cellular function. Initially may be reversible but life threatening if not treated promptly.

	CAUSED BY	SKIN	PCWP (PRELOAD)	CO	SVR (AFTERLOAD)	TREATMENT
Hypovolemic shock	Hemorrhage, dehydration, burns	Cold, clammy	↓↓	↓	↑	IV fluids
Cardiogenic shock	Acute MI, HF, valvular dysfunction, arrhythmia	Cold, clammy	↑ or ↓	↓↓	↑	Inotropes, diuresis
Obstructive shock	Cardiac tamponade, pulmonary embolism, tension pneumothorax					Relieve obstruction
Distributive shock	Sepsis, anaphylaxis CNS injury	Warm Dry	↓ ↓	↑ ↓	↓↓ ↓↓	IV fluids, pressors, epinephrine (anaphylaxis)

Cardiac tamponade





Compression of the heart by fluid (eg, blood, effusions [arrows in **A**] in pericardial space) → ↓ CO.
Equilibration of diastolic pressures in all 4 chambers.
Findings: Beck triad (hypotension, distended neck veins, distant heart sounds), ↑ HR, pulsus paradoxus. ECG shows low-voltage QRS and electrical alternans **B** (due to "swinging" movement of heart in large effusion).

Pulsus paradoxus—↓ in amplitude of systolic BP by > 10 mm Hg during inspiration. Seen in constrictive **P**ericarditis, obstructive pulmonary disease (eg, **C**roup, **O**SA, **A**sthma, COPD), cardiac **T**amponade (**P**ea **COAT**).

(c) ketabton.com: The Digital Library

Bacterial endocarditis

Acute—*S aureus* (high virulence). Large vegetations on previously normal valves **A**. Rapid onset.

Subacute—viridans streptococci (low virulence). Smaller vegetations on congenitally abnormal or diseased valves. Sequela of dental procedures. Gradual onset.

Symptoms: fever (most common), new murmur, Roth spots (round white spots on retina surrounded by hemorrhage **B**), Osler nodes (Ouchy raised lesions on finger or toe pads **C** due to immune complex deposition), Janeway lesions (small, painless, erythematous lesions on palm or sole) **D**, splinter hemorrhages **E** on nail bed.

Associated with glomerulonephritis, septic arterial or pulmonary emboli.

May be nonbacterial (marantic/thrombotic) 2° to malignancy, hypercoagulable state, or lupus.

FROM JANE with ♥:
- **F**ever
- **R**oth spots
- **O**sler nodes
- **M**urmur
- **J**aneway lesions
- **A**nemia
- **N**ail-bed hemorrhage
- **E**mboli

Requires multiple blood cultures for diagnosis.

If culture ⊖, most likely *Coxiella burnetii*, *Bartonella* spp.

Mitral valve is most frequently involved.

Tricuspid valve endocarditis is associated with IV **drug** abuse (don't "**tri**" **drugs**). Associated with *S aureus*, *Pseudomonas*, and *Candida*.

S bovis (gallolyticus) is present in colon cancer, *S epidermidis* on prosthetic valves.

Native valve endocarditis may be due to **HACEK** organisms (***H**aemophilus*, ***A**ggregatibacter* [formerly *Actinobacillus*], ***C**ardiobacterium*, ***E**ikenella*, ***K**ingella*).

(c) ketabton.com: The Digital Library

Rheumatic fever



A consequence of pharyngeal infection with group A β-hemolytic streptococci. Late sequelae include rheumatic heart disease, which affects heart valves—**m**itral > **a**ortic >> **t**ricuspid (high-pressure valves affected most). Early lesion is mitral valve regurgitation; late lesion is mitral stenosis.

Associated with Aschoff bodies (granuloma with giant cells [blue arrows in A]), Anitschkow cells (enlarged macrophages with ovoid, wavy, rod-like nucleus [red arrow in A]), ↑ anti-streptolysin O (ASO) and ↑ anti-DNase B titers.

Immune mediated (type II hypersensitivity); not a direct effect of bacteria. Antibodies to **M** protein cross-react with self antigens, often **m**yosin (**m**olecular **m**imicry).

Treatment/prophylaxis: penicillin.

J♥NES (major criteria):
- **J**oint (migratory polyarthritis)
- **♥** (carditis)
- **N**odules in skin (subcutaneous)
- **E**rythema marginatum (evanescent rash with ring margin)
- **S**ydenham chorea

Syphilitic heart disease

3° syphilis disrupts the vasa vasorum of the aorta with consequent atrophy of vessel wall and dilation of aorta and valve ring.

May see calcification of aortic root, ascending aortic arch, and thoracic aorta. Leads to "tree bark" appearance of aorta.

Can result in aneurysm of ascending aorta or aortic arch, aortic insufficiency.

(c) ketabton.com: The Digital Library

Acute pericarditis



Inflammation of the pericardium [A, red arrows]. Commonly presents with sharp pain, aggravated by inspiration, and relieved by sitting up and leaning forward. Often complicated by pericardial effusion [between yellow arrows in A]. Presents with friction rub. ECG changes include widespread ST-segment elevation and/or PR depression.

Causes include idiopathic (most common; presumed viral), confirmed infection (eg, coxsackievirus B), neoplasia, autoimmune (eg, SLE, rheumatoid arthritis), uremia, cardiovascular (acute STEMI or Dressler syndrome), radiation therapy.

Treatment: NSAIDs, colchicine, glucocorticoids, dialysis (uremia).

Myocarditis

Inflammation of myocardium → global enlargement of heart and dilation of all chambers. Major cause of SCD in adults < 40 years old.

Presentation highly variable, can include dyspnea, chest pain, fever, arrhythmias (persistent tachycardia out of proportion to fever is characteristic).

Multiple causes:

- Viral (eg, adenovirus, coxsackie B, parvovirus B19, HIV, HHV-6); lymphocytic infiltrate with focal necrosis highly indicative of viral myocarditis.
- Parasitic (eg, *Trypanosoma cruzi*, *Toxoplasma gondii*)
- Bacterial (eg, *Borrelia burgdorferi*, *Mycoplasma pneumoniae*, *Corynebacterium diphtheriae*)
- Toxins (eg, carbon monoxide, black widow venom)
- Rheumatic fever
- Drugs (eg, doxorubicin, cocaine)
- Autoimmune (eg, Kawasaki disease, sarcoidosis, SLE, polymyositis/dermatomyositis)

Complications include sudden death, arrhythmias, heart block, dilated cardiomyopathy, HF, mural thrombus with systemic emboli.

(c) ketabton.com: The Digital Library

Vasculitides

	EPIDEMIOLOGY/PRESENTATION	NOTES
Large-vessel vasculitis		
Giant cell (temporal) arteritis	Usually elderly females. Unilateral headache, possible temporal artery tenderness, jaw claudication. May lead to irreversible blindness due to ophthalmic artery occlusion. Associated with polymyalgia rheumatica.	Most commonly affects branches of carotid artery. Focal granulomatous inflammation A. ↑ ESR. Treat with high-dose corticosteroids prior to temporal artery biopsy to prevent blindness.
Takayasu arteritis	Usually Asian females < 40 years old. "Pulseless disease" (weak upper extremity pulses), fever, night sweats, arthritis, myalgias, skin nodules, ocular disturbances.	Granulomatous thickening and narrowing of aortic arch and proximal great vessels B. ↑ ESR. Treatment: corticosteroids.
Medium-vessel vasculitis		
Buerger disease (thromboangiitis obliterans)	Heavy smokers, males < 40 years old. Intermittent claudication. May lead to gangrene C, autoamputation of digits, superficial nodular phlebitis. Raynaud phenomenon is often present.	Segmental thrombosing vasculitis with vein and nerve involvement. Treatment: smoking cessation.
Kawasaki disease (mucocutaneous lymph node syndrome)	Asian children < 4 years old. Conjunctival injection, Rash (polymorphous → desquamating), Adenopathy (cervical), Strawberry tongue (oral mucositis) D, Hand-foot changes (edema, erythema), fever.	CRASH and burn on a Kawasaki. May develop coronary artery aneurysms E; thrombosis or rupture can cause death. Treatment: IV immunoglobulin and aspirin.
Polyarteritis nodosa	Usually middle-aged men. Hepatitis B seropositivity in 30% of patients. Fever, weight loss, malaise, headache. GI: abdominal pain, melena. Hypertension, neurologic dysfunction, cutaneous eruptions, renal damage.	Typically involves renal and visceral vessels, not pulmonary arteries. Different stages of transmural inflammation with fibrinoid necrosis. Innumerable renal microaneurysms F and spasms on arteriogram (string of pearls appearance). Treatment: corticosteroids, cyclophosphamide.
Small-vessel vasculitis		
Behçet syndrome	High incidence in people of Turkish and eastern Mediterranean descent. Recurrent aphthous ulcers, genital ulcerations, uveitis, erythema nodosum. Can be precipitated by HSV or parvovirus. Flares last 1–4 weeks.	Immune complex vasculitis. Associated with HLA-B51.
Cutaneous small-vessel vasculitis	Occurs 7-10 days after certain medications (penicillin, cephalosporins, phenytoin, allopurinol) or infections (eg, HCV, HIV). Palpable purpura, no visceral involvement.	Immune complex–mediated leukocytoclastic vasculitis; late involvement indicates systemic vasculitis.

(c) ketabton.com: The Digital Library

Vasculitides *(continued)*

	EPIDEMIOLOGY/PRESENTATION	NOTES
Small-vessel vasculitis *(continued)*		
Eosinophilic granulomatosis with polyangiitis (Churg-Strauss)	Asthma, sinusitis, skin nodules or purpura, peripheral neuropathy (eg, wrist/foot drop). Can also involve heart, GI, kidneys (pauci-immune glomerulonephritis).	Granulomatous, necrotizing vasculitis with eosinophilia G. MPO-ANCA/p-ANCA, ↑ IgE level.
Granulomatosis with polyangiitis (Wegener)	Upper respiratory tract: perforation of nasal septum, chronic sinusitis, otitis media, mastoiditis. Lower respiratory tract: hemoptysis, cough, dyspnea. Renal: hematuria, red cell casts.	Triad: ▪ Focal necrotizing vasculitis ▪ Necrotizing granulomas in lung and upper airway ▪ Necrotizing glomerulonephritis PR3-ANCA/c-ANCA H (anti-proteinase 3). CXR: large nodular densities. Treatment: cyclophosphamide, corticosteroids.
Immunoglobulin A vasculitis	Also called Henoch-Schönlein purpura. Most common childhood systemic vasculitis. Often follows URI. Classic triad: ▪ Skin: palpable purpura on buttocks/legs I ▪ Arthralgias ▪ GI: abdominal pain (associated with intussusception)	Vasculitis 2° to IgA immune complex deposition. Associated with IgA nephropathy (Berger disease). Treatment: supportive care, possibly corticosteroids.
Microscopic polyangiitis	Necrotizing vasculitis commonly involving lung, kidneys, and skin with pauci-immune glomerulonephritis and palpable purpura. Presentation similar to granulomatosis with polyangiitis but without nasopharyngeal involvement.	No granulomas. MPO-ANCA/p-ANCA J (anti-myeloperoxidase). Treatment: cyclophosphamide, corticosteroids.
Mixed cryoglobulinemia	Often due to viral infections, especially HCV. Triad of palpable purpura, weakness, arthralgias. May also have peripheral neuropathy and renal disease (eg, glomerulonephritis).	Cryoglobulins are immunoglobulins that precipitate in the Cold. Vasculitis due to mixed IgG and IgM immune complex deposition.



(c) ketabton.com: The Digital Library

Cardiac tumors	Most common heart tumor is a metastasis (eg, melanoma).
Myxomas 	Most common 1° cardiac tumor in **adults** (arrows in A). 90% occur in the atria (mostly left atrium). Myxomas are usually described as a "ball valve" obstruction in the left atrium (associated with multiple syncopal episodes). IL-6 production by tumor → constitutional symptoms (eg, fever, weight loss). May auscultate early diastolic "tumor plop" sound. Histology: gelatinous material, myxoma cells immersed in glycosaminoglycans. **Adults** make **myx**ed drinks.
Rhabdomyomas	Most frequent 1° cardiac tumor in children (associated with tuberous sclerosis). Histology: hamartomatous growths.
Kussmaul sign	↑ in JVP on inspiration instead of a normal ↓. Inspiration → negative intrathoracic pressure not transmitted to heart → impaired filling of right ventricle → blood backs up into vena cava → JVD. May be seen with constrictive pericarditis, restrictive cardiomyopathies, right heart failure, massive pulmonary embolism, right atrial or ventricular tumors.
Hereditary hemorrhagic telangiectasia	Also called Osler-Weber-Rendu syndrome. Autosomal dominant disorder of blood vessels. Findings: blanching lesions (telangiectasias) on skin and mucous membranes, recurrent epistaxis, skin discolorations, arteriovenous malformations (AVMs), GI bleeding, hematuria.

▸ CARDIOVASCULAR—PHARMACOLOGY

Hypertension treatment

Primary (essential) hypertension	Thiazide diuretics, ACE inhibitors, angiotensin II receptor blockers (ARBs), dihydropyridine Ca^{2+} channel blockers.	
Hypertension with heart failure	Diuretics, ACE inhibitors/ARBs, β-blockers (compensated HF), aldosterone antagonists.	β-blockers must be used cautiously in decompensated HF and are contraindicated in cardiogenic shock. In HF, ARBs may be combined with the neprilysin inhibitor sacubitril.
Hypertension with diabetes mellitus	ACE inhibitors/ARBs, Ca^{2+} channel blockers, thiazide diuretics, β-blockers.	ACE inhibitors/ARBs are protective against diabetic nephropathy. β-blockers can mask hypoglycemia symptoms.
Hypertension in asthma	ARBs, Ca^{2+} channel blockers, thiazide diuretics, cardioselective β-blockers.	Avoid nonselective β-blockers to prevent β_2-receptor–induced bronchoconstriction. Avoid ACE inhibitors to prevent confusion between drug or asthma-related cough.
Hypertension in pregnancy	**H**ydralazine, **l**abetalol, **m**ethyldopa, **n**ifedipine.	"**H**e **l**ikes **m**y **n**eonate."

(c) ketabton.com: The Digital Library

Cardiac therapy



(c) ketabton.com: The Digital Library

Calcium channel blockers	Amlodipine, clevidipine, nicardipine, nifedipine, nimodipine (dihydropyridines, act on vascular smooth muscle); diltiazem, verapamil (non-dihydropyridines, act on heart).
MECHANISM	Block voltage-dependent L-type calcium channels of cardiac and smooth muscle → ↓ muscle contractility. Vascular smooth muscle—amlodipine = nifedipine > diltiazem > verapamil. Heart—verapamil > diltiazem > amlodipine = nifedipine (**v**erapamil = **v**entricle).
CLINICAL USE	Dihydropyridines (except nimodipine): hypertension, angina (including vasospastic type), Raynaud phenomenon. Nimodipine: subarachnoid hemorrhage (prevents cerebral vasospasm). Nicardipine, clevidipine: hypertensive urgency or emergency. Non-dihydropyridines: hypertension, angina, atrial fibrillation/flutter.
ADVERSE EFFECTS	Gingival hyperplasia. Dihydropyridine: peripheral edema, flushing, dizziness. Non-dihydropyridine: cardiac depression, AV block, hyperprolactinemia (verapamil), constipation.

Hydralazine	
MECHANISM	↑ cGMP → smooth muscle relaxation. Vasodilates arterioles > veins; afterload reduction.
CLINICAL USE	Severe hypertension (particularly acute), HF (with organic nitrate). Safe to use during pregnancy. Frequently coadministered with a β-blocker to prevent reflex tachycardia.
ADVERSE EFFECTS	Compensatory tachycardia (contraindicated in angina/CAD), fluid retention, headache, angina, drug-induced lupus.

Hypertensive emergency	Treat with labetalol, clevidipine, fenoldopam, nicardipine, nitroprusside.
Nitroprusside	Short acting vasodilator (arteries = veins); ↑ cGMP via direct release of NO. Can cause cyanide toxicity (releases cyanide).
Fenoldopam	**Dopam**ine D_1 receptor agonist—coronary, peripheral, renal, and splanchnic vasodilation. ↓ BP, ↑ natriuresis. Also used postoperatively as an antihypertensive. Can cause hypotension and tachycardia.

Nitrates	Nitroglycerin, isosorbide dinitrate, isosorbide mononitrate.
MECHANISM	Vasodilate by ↑ NO in vascular smooth muscle → ↑ in cGMP and smooth muscle relaxation. Dilate veins >> arteries. ↓ preload.
CLINICAL USE	Angina, acute coronary syndrome, pulmonary edema.
ADVERSE EFFECTS	Reflex tachycardia (treat with β-blockers), hypotension, flushing, headache, "Monday disease" in industrial exposure: development of tolerance for the vasodilating action during the work week and loss of tolerance over the weekend → tachycardia, dizziness, headache upon reexposure. Contraindicated in right ventricular infarction, hypertrophic cardiomyopathy, and with concurrent PDE-5 inhibitor use.

(c) ketabton.com: The Digital Library

Antianginal therapy

Goal is reduction of myocardial O_2 consumption (MVO_2) by ↓ 1 or more of the determinants of MVO_2: end-diastolic volume, BP, HR, contractility.

COMPONENT	NITRATES	β-BLOCKERS	NITRATES + β-BLOCKERS
End-diastolic volume	↓	No effect or ↑	No effect or ↓
Blood pressure	↓	↓	↓
Contractility	↑ (reflex response)	↓	Little/no effect
Heart rate	↑ (reflex response)	↓	No effect or ↓
Ejection time	↓	↑	Little/no effect
MVO_2	↓	↓	↓↓

Verapamil is similar to β-blockers in effect.
Pindolol and acebutolol are partial β-agonists that should be used with caution in angina.

Ranolazine

MECHANISM	Inhibits the late phase of inward sodium current thereby reducing diastolic wall tension and oxygen consumption. Does not affect heart rate or blood pressure.
CLINICAL USE	Angina refractory to other medical therapies.
ADVERSE EFFECTS	Constipation, dizziness, headache, nausea.

Sacubitril

MECHANISM	A neprilysin inhibitor; prevents degradation of natriuretic peptides, angiotensin II, and substance P → ↑ vasodilation, ↓ ECF volume.
CLINICAL USE	Used in combination with valsartan (an ARB) to treat HFrEF.
ADVERSE EFFECTS	Hypotension, hyperkalemia, cough, dizziness; contraindicated with ACE inhibitors due to angioedema.

(c) ketabton.com: The Digital Library

Lipid-lowering agents

DRUG	LDL	HDL	TRIGLYCERIDES	MECHANISMS OF ACTION	ADVERSE EFFECTS/PROBLEMS
HMG-CoA reductase inhibitors (eg, atorvastatin, simvastatin)	↓↓↓	↑	↓	Inhibit conversion of HMG-CoA to mevalonate, a cholesterol precursor; ↑ LDL recycling; ↓ mortality in CAD patients	Hepatotoxicity (↑ LFTs), myopathy (esp when used with fibrates or niacin)
Bile acid resins Cholestyramine, colestipol, colesevelam	↓↓	↑ slightly	↑ slightly	Prevent intestinal reabsorption of bile acids; liver must use cholesterol to make more	GI upset, ↓ absorption of other drugs and fat-soluble vitamins
Ezetimibe	↓↓	↑/—	↓/—	Prevents cholesterol absorption at small intestine brush border	Rare ↑ LFTs, diarrhea
Fibrates Gemfibrozil, bezafibrate, fenofibrate	↓	↑	↓↓↓	Upregulate LPL → ↑ TG clearance Activates PPAR-α to induce HDL synthesis	Myopathy (↑ risk with statins), cholesterol gallstones (via inhibition of cholesterol 7α-hydroxylase)
Niacin	↓↓	↑↑	↓	Inhibits lipolysis (hormone-sensitive lipase) in adipose tissue; reduces hepatic VLDL synthesis	Flushed face (↓ by NSAIDs or long-term use) Hyperglycemia Hyperuricemia
PCSK9 inhibitors Alirocumab, evolocumab	↓↓↓	↑	↓	Inactivation of LDL-receptor degradation → ↑ removal of LDL from bloodstream	Myalgias, delirium, dementia, other neurocognitive effects
Fish oil and marine omega-3 fatty acids	↑ slightly	↑ slightly	↓ at high doses	Believed to decrease FFA delivery to liver and decrease activity of TG-synthesizing enzymes	Nausea, fish-like taste

(c) ketabton.com: The Digital Library

Lipid-lowering agents *(continued)*



Cardiac glycosides

Digoxin.

MECHANISM

Direct inhibition of Na^+/K^+ ATPase → indirect inhibition of Na^+/Ca^{2+} exchanger. ↑ $[Ca^{2+}]_i$ → positive inotropy. Stimulates vagus nerve → ↓ HR.



CLINICAL USE

HF (↑ contractility); atrial fibrillation (↓ conduction at AV node and depression of SA node).

ADVERSE EFFECTS

Cholinergic effects (nausea, vomiting, diarrhea), blurry yellow vision (think van Glow), arrhythmias, AV block.

Can lead to hyperkalemia, which indicates poor prognosis.

Factors predisposing to toxicity: renal failure (↓ excretion), hypokalemia (permissive for digoxin binding at K^+-binding site on Na^+/K^+ ATPase), drugs that displace digoxin from tissue-binding sites, and ↓ clearance (eg, verapamil, amiodarone, quinidine).

ANTIDOTE

Slowly normalize K^+, cardiac pacer, anti-digoxin Fab fragments, Mg^{2+}.

(c) ketabton.com: The Digital Library

Antiarrhythmics—sodium channel blockers (class I)

Slow or block (↓) conduction (especially in depolarized cells). ↓ slope of phase 0 depolarization. Are state dependent (selectively depress tissue that is frequently depolarized [eg, tachycardia]).

Class IA

Quinidine, **Proc**ainamide, **Disopyramid**e.
"The **Q**ueen **Proc**laims **Diso**'s **pyramid**."



MECHANISM
Moderate Na^+ channel blockade.
↑ AP duration, ↑ effective refractory period (ERP) in ventricular action potential, ↑ QT interval, some potassium channel blocking effects.

CLINICAL USE
Both atrial and ventricular arrhythmias, especially re-entrant and ectopic SVT and VT.

ADVERSE EFFECTS
Cinchonism (headache, tinnitus with quinidine), reversible SLE-like syndrome (procainamide), HF (disopyramide), thrombocytopenia, torsades de pointes due to ↑ QT interval.

Class IB

Lidocaine, **Mexi**le**T**ine.
"**I**'d **B**uy **Lid**dy's **Mexi**can **T**acos."



MECHANISM
Weak Na^+ channel blockade.
↓ AP duration. Preferentially affect ischemic or depolarized Purkinje and ventricular tissue. Phenytoin can also fall into the IB category.

CLINICAL USE
Acute ventricular arrhythmias (especially post-MI), digitalis-induced arrhythmias.
IB is **B**est post-MI.

ADVERSE EFFECTS
CNS stimulation/depression, cardiovascular depression.

Class IC

Flecainide, **P**ropafenone.
"**C**an I have **F**ries, **P**lease."



MECHANISM
Strong Na^+ channel blockade.
Significantly prolongs ERP in AV node and accessory bypass tracts. No effect on ERP in Purkinje and ventricular tissue.
Minimal effect on AP duration.

CLINICAL USE
SVTs, including atrial fibrillation. Only as a last resort in refractory VT.

ADVERSE EFFECTS
Proarrhythmic, especially post-MI (contraindicated). **IC** is **C**ontraindicated in structural and ischemic heart disease.

FAS1_2019_07-Cardio.indd 322 11/7/19 4:24 PM

(c) ketabton.com: The Digital Library

Antiarrhythmics—β-blockers (class II)

Metoprolol, propranolol, esmolol, atenolol, timolol, carvedilol.

MECHANISM

Decrease SA and AV nodal activity by ↓ cAMP, ↓ Ca^{2+} currents. Suppress abnormal pacemakers by ↓ slope of phase 4.

AV node particularly sensitive—↑ PR interval. Esmolol very short acting.

CLINICAL USE

SVT, ventricular rate control for atrial fibrillation and atrial flutter.

ADVERSE EFFECTS

Impotence, exacerbation of COPD and asthma, cardiovascular effects (bradycardia, AV block, HF), CNS effects (sedation, sleep alterations). May mask the signs of hypoglycemia.

Metoprolol can cause dyslipidemia. Propranolol can exacerbate vasospasm in vasospastic angina. β-blockers (except the nonselective α- and β-antagonists carvedilol and labetalol) cause unopposed α_1-agonism if given alone for pheochromocytoma or for cocaine toxicity (unsubstantiated). Treat β-blocker overdose with saline, atropine, glucagon.



Pacemaker cell action potential

Antiarrhythmics—potassium channel blockers (class III)

Amiodarone, Ibutilide, Dofetilide, Sotalol. AIDS.

MECHANISM

↑ AP duration, ↑ ERP, ↑ QT interval.

CLINICAL USE

Atrial fibrillation, atrial flutter; ventricular tachycardia (amiodarone, sotalol).

ADVERSE EFFECTS

Sotalol—torsades de pointes, excessive β blockade.

Ibutilide—torsades de pointes.

Amiodarone—pulmonary fibrosis, hepatotoxicity, hypothyroidism or hyperthyroidism (amiodarone is 40% iodine by weight), acts as hapten (corneal deposits, blue/gray skin deposits resulting in photodermatitis), neurologic effects, constipation, cardiovascular effects (bradycardia, heart block, HF).

Remember to check PFTs, LFTs, and TFTs when using amiodarone.

Amiodarone is lipophilic and has class I, II, III, and IV effects.



Cell action potential

(c) ketabton.com: The Digital Library

Antiarrhythmics—calcium channel blockers (class IV)

Diltiazem, Verapamil

MECHANISM	Decrease conduction Velocity, ↑ ERP, ↑ PR interval.
CLINICAL USE	Prevention of nodal arrhythmias (eg, SVT), rate control in atrial fibrillation.
ADVERSE EFFECTS	Constipation, flushing, edema, cardiovascular effects (HF, AV block, sinus node depression).



Other antiarrhythmics

Adenosine	↑ K^+ out of cells → hyperpolarizing the cell and ↓ I_{Ca}, decreasing AV node conduction. Drug of choice in diagnosing/terminating certain forms of SVT. Very short acting (~ 15 sec). Effects blunted by theophylline and caffeine (both are adenosine receptor antagonists). Adverse effects include flushing, hypotension, chest pain, sense of impending doom, bronchospasm.
Magnesium	Effective in torsades de pointes and digoxin toxicity.

Ivabradine

MECHANISM	IVabradine prolongs slow depolarization (phase "IV") by selectively inhibiting "funny" sodium channels (I_f).
CLINICAL USE	Chronic stable angina in patients who cannot take β-blockers. Chronic HFrEF.
ADVERSE EFFECTS	Luminous phenomena/visual brightness, hypertension, bradycardia.

(c) ketabton.com: The Digital Library

Endocrine

"If you skew the endocrine system, you lose the pathways to self."
—Hilary Mantel

"We have learned that there is an endocrinology of elation and despair, a chemistry of mystical insight, and, in relation to the autonomic nervous system, a meteorology and even . . . an astro-physics of changing moods."
—Aldous Huxley

"Chocolate causes certain endocrine glands to secrete hormones that affect your feelings and behavior by making you happy."
—Elaine Sherman, *Book of Divine Indulgences*

The endocrine system comprises widely distributed organs that work in a highly integrated manner to orchestrate a state of hormonal equilibrium within the body. Generally speaking, endocrine diseases can be classified either as diseases of underproduction or overproduction, or as conditions involving the development of mass lesions—which themselves may be associated with underproduction or overproduction of hormones. Therefore, study the endocrine system first by learning the glands, their hormones, and their regulation, and then by integrating disease manifestations with diagnosis and management. Take time to learn the multisystem connections.

Embryology 326
Anatomy 327
Physiology 328
Pathology 338
Pharmacology 352

FAS1_2019_08-Endocrine.indd 325 11/7/19 4:30 PM

(c) ketabton.com: The Digital Library

▸ ENDOCRINE—EMBRYOLOGY

Thyroid development

Thyroid diverticulum arises from floor of primitive pharynx and descends into neck. Connected to tongue by thyroglossal duct, which normally disappears but may persist as cysts or the pyramidal lobe of thyroid. Foramen cecum is normal remnant of thyroglossal duct.

Most common ectopic thyroid tissue site is the tongue (lingual thyroid). Removal may result in hypothyroidism if it is the only thyroid tissue present.

Thyroglossal duct cyst **A** presents as an anterior midline neck mass that moves with swallowing or protrusion of the tongue (vs persistent cervical sinus leading to pharyngeal cleft cyst in lateral neck).

Thyroid follicular cells derived from endoderm.



(c) ketabton.com: The Digital Library

▸ ENDOCRINE—ANATOMY

Pituitary gland

Anterior pituitary (adenohypophysis)	Secretes FSH, LH, ACTH, TSH, prolactin, GH, and β-endorphin. Melanotropin (MSH) secreted from intermediate lobe of pituitary. Derived from oral ectoderm (Rathke pouch). ▪ α subunit—hormone subunit common to TSH, LH, FSH, and hCG. ▪ β subunit—determines hormone specificity.	Proopiomelanocortin derivatives—β-endorphin, ACTH, and MSH. Go pro with a BAM! FLAT PiG: FSH, LH, ACTH, TSH, PRL, GH. B-FLAT: Basophils—FSH, LH, ACTH, TSH. Acid PiG: Acidophils — PRL, GH.
Posterior pituitary (neurohypophysis)	Stores and releases vasopressin (antidiuretic hormone, or ADH) and oxytocin, both made in the hypothalamus (supraoptic and paraventricular nuclei) and transported to posterior pituitary via neurophysins (carrier proteins). Derived from neuroectoderm.	

Adrenal cortex and medulla

Adrenal cortex (derived from mesoderm) and medulla (derived from neural crest).



GFR corresponds with Salt (mineralocorticoids), Sugar (glucocorticoids), and Sex (androgens). "The deeper you go, the sweeter it gets."

(c) ketabton.com: The Digital Library

▸ ENDOCRINE—PHYSIOLOGY

Hypothalamic-pituitary hormones

HORMONE	FUNCTION	CLINICAL NOTES
ADH	↑ water permeability of distal convoluted tubule and collecting duct cells in kidney to ↑ water reabsorption	Stimulus for secretion is ↑ plasma osmolality, except in SIADH, in which ADH is elevated despite ↓ plasma osmolality
CRH	↑ ACTH, MSH, β-endorphin	↓ in chronic exogenous steroid use
Dopamine	↓ prolactin, TSH	Also called prolactin-inhibiting factor Dopamine antagonists (eg, antipsychotics) can cause galactorrhea due to hyperprolactinemia
GHRH	↑ GH	Analog (tesamorelin) used to treat HIV-associated lipodystrophy
GnRH	↑ FSH, LH	Suppressed by hyperprolactinemia Tonic GnRH analog (eg, leuprolide) suppresses hypothalamic–pituitary–gonadal axis. Pulsatile GnRH leads to puberty, fertility
MSH	↑ melanogenesis by melanocytes	Causes hyperpigmentation in Cushing disease, as MSH and ACTH share the same precursor molecule, proopiomelanocortin
Oxytocin	Causes uterine contractions during labor. Responsible for milk letdown reflex in response to suckling.	Modulates fear, anxiety, social bonding, mood, and depression
Prolactin	↓ GnRH Stimulates lactogenesis.	Pituitary prolactinoma → amenorrhea, osteoporosis, hypogonadism, galactorrhea Breastfeeding → ↑ prolactin → ↓ GnRH → delayed postpartum ovulation (natural contraception)
Somatostatin	↓ GH, TSH	Also called growth hormone inhibiting hormone (GHIH) Analogs used to treat acromegaly
TRH	↑ TSH, prolactin	↑ TRH (eg, in 1°/2° hypothyroidism) may increase prolactin secretion → galactorrhea



(c) ketabton.com: The Digital Library

Growth hormone



Also called somatotropin. Secreted by anterior pituitary.

Stimulates linear growth and muscle mass through IGF-1 (somatomedin C) secretion by liver. ↑ insulin resistance (diabetogenic).

Released in pulses in response to growth hormone–releasing hormone (GHRH).

Secretion ↑ during exercise, deep sleep, puberty, hypoglycemia, CKD.

Secretion ↓ by glucose, somatostatin, somatomedin (regulatory molecule secreted by liver in response to GH acting on target tissues).

Excess secretion of GH (eg, pituitary adenoma) may cause acromegaly (adults) or gigantism (children). Treatment: somatostatin analogs (eg, octreotide) or surgery.

Antidiuretic hormone

Also called vasopressin.

SOURCE	Synthesized in hypothalamus (supraoptic and paraventricular nuclei), stored and secreted by posterior pituitary.	
FUNCTION	Regulates blood pressure (V_1-receptors) and serum osmolality (V_2-receptors). Primary function is serum osmolality regulation (ADH ↓ serum osmolality, ↑ urine osmolality) via regulation of aquaporin channel insertion in principal cells of renal collecting duct.	ADH level is ↓ in central diabetes insipidus (DI), normal or ↑ in nephrogenic DI. Nephrogenic DI can be caused by mutation in V_2-receptor. Desmopressin (ADH analog) is a treatment for central DI and nocturnal enuresis.
REGULATION	Plasma osmolality (1°); hypovolemia.	

(c) ketabton.com: The Digital Library

Prolactin

SOURCE	Secreted mainly by anterior pituitary.	Structurally homologous to growth hormone.
FUNCTION	Stimulates milk production in breast; inhibits ovulation in females and spermatogenesis in males by inhibiting GnRH synthesis and release.	Excessive amounts of prolactin associated with ↓ libido.
REGULATION	Prolactin secretion from anterior pituitary is tonically inhibited by dopamine from tuberoinfundibular pathway of hypothalamus. Prolactin in turn inhibits its own secretion by ↑ dopamine synthesis and secretion from hypothalamus. TRH ↑ prolactin secretion (eg, in 1° or 2° hypothyroidism).	Dopamine agonists (eg, bromocriptine) inhibit prolactin secretion and can be used in treatment of prolactinoma. Dopamine antagonists (eg, most antipsychotics, metoclopramide) and estrogens (eg, OCPs, pregnancy) stimulate prolactin secretion.



(c) ketabton.com: The Digital Library

Thyroid hormones

Thyroid produces triiodothyronine (T_3) and thyroxine (T_4), iodine-containing hormones that control the body's metabolic rate.

SOURCE

Follicles of thyroid. 5′-deiodinase converts T_4 (the major thyroid product) to T_3 in peripheral tissue (5, 4, 3). Peripheral conversion is inhibited by glucocorticoids, β-blockers, and propylthiouracil (PTU). Reverse T_3 (rT_3) is a metabolically inactive byproduct of the peripheral conversion of T_4 and its production is increased by growth hormone and glucocorticoids. Functions of thyroid peroxidase include oxidation, organification of iodine, and coupling of monoiodotyrosine (MIT) and diiodotyrosine (DIT). Inhibited by PTU and methimazole. DIT + DIT = T_4. DIT + MIT = T_3. Wolff-Chaikoff effect—excess iodine temporarily turns off thyroid peroxidase → ↓ T_3/T_4 production (protective autoregulatory effect).

FUNCTION

Only free hormone is active. T_3 binds nuclear receptor with greater affinity than T_4. T_3 functions —7 B's:

- Brain maturation
- Bone growth (synergism with GH)
- β-adrenergic effects. ↑ β_1 receptors in heart → ↑ CO, HR, SV, contractility; β-blockers alleviate adrenergic symptoms in thyrotoxicosis
- Basal metabolic rate ↑ (via Na^+/K^+-ATPase activity → ↑ O_2 consumption, RR, body temperature)
- Blood sugar (↑ glycogenolysis, gluconeogenesis)
- Break down lipids (↑ lipolysis)
- Stimulates surfactant synthesis in Babies

REGULATION

TRH ⊕ TSH release → ⊕ follicular cells. Thyroid-stimulating immunoglobulin (TSI) may ⊕ follicular cells in Graves disease.

Negative feedback primarily by free T_3/T_4:

- Anterior pituitary → ↓ sensitivity to TRH
- Hypothalamus → ↓ TRH secretion

Thyroxine-binding globulin (TBG) binds most T_3/T_4 in blood. Bound T_3/T_4 = inactive.

- ↑ TBG in pregnancy, OCP use (estrogen → ↑ TBG) → ↑ total T_3/T_4
- ↓ TBG in steroid use, nephrotic syndrome





(c) ketabton.com: The Digital Library

Parathyroid hormone

SOURCE

Chief cells of parathyroid

FUNCTION

↑ free Ca^{2+} in the blood (1° function)
↑ Ca^{2+} and PO_4^{3-} absorption in GI system
↑ Ca^{2+} and PO_4^{3-} from bone resorption
↑ Ca^{2+} reabsorption from DCT
↓ PO_4^{3-} reabsorption in PCT
↑ 1,25-$(OH)_2D_3$ (calcitriol) production by activating 1α-hydroxylase in **PCT**
Tri to make D_3 in the **PCT**

PTH ↑ serum Ca^{2+}, ↓ serum PO_4^{3-}, ↑ urine PO_4^{3-}, ↑ urine cAMP
↑ RANK-L (receptor activator of NF-κB ligand) secreted by osteoblasts and osteocytes; binds RANK (receptor) on osteoclasts and their precursors to stimulate osteoclasts and ↑ Ca^{2+} → bone resorption (intermittent PTH release can also stimulate bone formation)
PTH = **P**hosphate-**T**rashing **H**ormone
PTH-related peptide (PTHrP) functions like PTH and is commonly increased in malignancies (eg, squamous cell carcinoma of the lung, renal cell carcinoma)

REGULATION

↓ serum Ca^{2+} → ↑ PTH secretion
↑ serum PO_4^{3-} → ↑ PTH secretion
↓ serum Mg^{2+} → ↑ PTH secretion
↓↓ serum Mg^{2+} → ↓ PTH secretion
Common causes of ↓ Mg^{2+} include diarrhea, aminoglycosides, diuretics, alcohol abuse



(c) ketabton.com: The Digital Library

Calcium homeostasis

Plasma Ca^{2+} exists in three forms:
- Ionized/free (~ 45%, active form)
- Bound to albumin (~ 40%)
- Bound to anions (~ 15%)

↑ **pH** (less H^+) → albumin binds more Ca^{2+} → ↓ ionized Ca^{2+} (eg, cramps, pain, paresthesias, carpopedal spasm) → ↑ **PTH**

↓ **pH** (more H^+) → albumin binds less Ca^{2+} → ↑ ionized Ca^{2+} → ↓ **PTH**

Ionized/free Ca^{2+} is 1° regulator of PTH; changes in pH alter PTH secretion, whereas changes in albumin concentration do not

Calcitonin

SOURCE	Parafollicular cells (C cells) of thyroid.
FUNCTION	↓ bone resorption of Ca^{2+}.
REGULATION	↑ serum Ca^{2+} → ↑ calcitonin secretion.

Calcitonin opposes actions of PTH. Not important in normal Ca^{2+} homeostasis

Calcitonin tones down serum Ca^{2+} levels and keeps it in bones

Glucagon

SOURCE	Made by α cells of pancreas.
FUNCTION	Promotes glycogenolysis, gluconeogenesis, lipolysis, ketogenesis. Elevates blood sugar levels to maintain homeostasis when bloodstream glucose levels fall too low (ie, fasting state).
REGULATION	Secreted in response to hypoglycemia. Inhibited by insulin, hyperglycemia, somatostatin.

(c) ketabton.com: The Digital Library

Insulin

SYNTHESIS



Preproinsulin (synthesized in RER of pancreatic β cells) → cleavage of "presignal" → proinsulin (stored in secretory granules) → cleavage of proinsulin → exocytosis of insulin and C-peptide equally. Insulin and C-peptide are ↑ in insulinoma and sulfonylurea use, whereas exogenous insulin lacks C-peptide.

FUNCTION

Binds **in**sulin receptors (tyrosine kinase activity ❶), **in**ducing glucose uptake (carrier-mediated transport) **in**to insulin-dependent tissue ❷ and gene transcription.

Anabolic effects of insulin:

- ↑ glucose transport in skeletal muscle and adipose tissue
- ↑ glycogen synthesis and storage
- ↑ triglyceride synthesis
- ↑ Na^+ retention (kidneys)
- ↑ protein synthesis (muscles)
- ↑ cellular uptake of K^+ and amino acids
- ↓ glucagon release
- ↓ lipolysis in adipose tissue

Unlike glucose, insulin does not cross placenta.

Insulin-dependent glucose transporters:

- GLUT4: adipose tissue, striated muscle (exercise can also ↑ GLUT4 expression)

Insulin-independent transporters:

- GLUT1: RBCs, brain, cornea, placenta
- GLUT**2** (**bi**directional): β islet cells, liver, kidney, GI tract (think **2**-way street)
- GLUT3: brain, placenta
- GLUT**5** (**F**ructose): spermatocytes, GI tract
- SGLT1/SGLT2 (Na^+-glucose cotransporters): kidney, small intestine

Brain prefers glucose, but may use ketone bodies during starvation. RBCs utilize glucose, as they lack mitochondria for aerobic metabolism.

BRICK LIPS (insulin-independent glucose uptake): **B**rain, **R**BCs, **I**ntestine, **C**ornea, **K**idney, **L**iver, **I**slet (β) cells, **P**lacenta, **S**permatocytes.

REGULATION

Glucose is the major regulator of insulin release. ↑ insulin response with oral vs IV glucose due to incretins (eg, glucagon-like peptide 1 [GLP-1], glucose-dependent insulinotropic polypeptide [GIP]), which are released after meals and ↑ β cell sensitivity to glucose. Release ↓ by α_2, ↑ by β_2 stimulation (**2** = regulates **in**sul**in**)

Glucose enters β cells ❸ → ↑ ATP generated from glucose metabolism ❹ closes K^+ channels (target of sulfonylureas) ❺ and depolarizes β cell membrane ❻. Voltage-gated Ca^{2+} channels open → Ca^{2+} influx ❼ and stimulation of insulin exocytosis ❽.



Insulin-dependent glucose uptake



Insulin secretion by pancreatic β cells

(c) ketabton.com: The Digital Library

Adrenal steroids and congenital adrenal hyperplasias



[a]Rate-limiting step.

ENZYME DEFICIENCY	MINERALOCORTICOIDS	[K^+]	BP	CORTISOL	SEX HORMONES	LABS	PRESENTATION
A 17α-hydroxylase[a]	↑	↓	↑	↓	↓	↓ androstenedione	XY: ambiguous genitalia, undescended testes XX: lacks 2° sexual development
B 21-hydroxylase[a]	↓	↑	↓	↓	↑	↑ renin activity ↑ 17-hydroxy-progesterone	Most common Presents in infancy (salt wasting) or childhood (precocious puberty) XX: virilization
C 11β-hydroxylase[a]	↓ aldosterone ↑ 11-deoxycorti-costerone (results in ↑ BP)	↓	↑	↓	↑	↓ renin activity	Presents in infancy (severe hypertension) or childhood (precocious puberty) XX: virilization

[a]All congenital adrenal enzyme deficiencies are autosomal recessive disorders and most are characterized by skin hyperpigmentation (due to ↑ MSH production, which is coproduced and secreted with ACTH) and bilateral adrenal gland enlargement (due to ↑ ACTH stimulation).

If deficient enzyme starts with 1, it causes hypertension; if deficient enzyme ends with 1, it causes virilization in females.

(c) ketabton.com: The Digital Library

Cortisol

SOURCE	Adrenal zona fasciculata.	Bound to corticosteroid-binding globulin.
FUNCTION	↑ **A**ppetite ↑ **B**lood pressure: ▪ Upregulates α_1-receptors on arterioles → ↑ sensitivity to norepinephrine and epinephrine (permissive action) ▪ At high concentrations, can bind to mineralocorticoid (aldosterone) receptors ↑ **I**nsulin resistance (diabetogenic) ↑ **G**luconeogenesis, lipolysis, and proteolysis (↓ glucose utilization) ↓ **F**ibroblast activity (poor wound healing, ↓ collagen synthesis, ↑ striae) ↓ **I**nflammatory and **I**mmune responses: ▪ Inhibits production of leukotrienes and prostaglandins ▪ Inhibits WBC adhesion → neutrophilia ▪ Blocks histamine release from mast cells ▪ Eosinopenia, lymphopenia ▪ Blocks IL-2 production ↓ **B**one formation (↓ osteoblast activity)	Cortisol is **A BIG FIB**. Exogenous corticosteroids can cause reactivation of TB and candidiasis (blocks IL-2 production). 
REGULATION	CRH (hypothalamus) stimulates ACTH release (pituitary) → cortisol production in adrenal zona fasciculata. Excess cortisol ↓ CRH, ACTH, and cortisol secretion.	Chronic stress may induce prolonged cortisol secretion, cortisol resistance, impaired immunocompetency, and dysregulation of HPA axis.

Appetite regulation

Ghrelin	Stimulates hunger (orexigenic effect) and GH release (via GH secretagog receptor). Produced by stomach. Sleep deprivation, fasting, or Prader-Willi syndrome → ↑ ghrelin production. **Ghr**elin makes you hung**hr**e and **ghr**ow. Acts on lateral area of hypothalamus (hunger center) to ↑ appetite.
Leptin	Satiety hormone. Produced by adipose tissue. Mutation of leptin gene → central obesity. (Obese people have ↑ leptin due to ↑ adipose tissue but also appear resistant to leptin's anorexigenic effect.) Sleep deprivation or starvation → ↓ leptin production. Lept**in** keeps you th**in**. Acts on ventromedial area of hypothalamus (satiety center) to ↓ appetite.
Endocannabinoids	Act at cannabinoid receptors in hypothalamus and nucleus accumbens, two key brain areas for the homeostatic and hedonic control of food intake → ↑ appetite. Exogenous cannabinoids cause "the munchies."

(c) ketabton.com: The Digital Library

Signaling pathways of endocrine hormones

cAMP	**F**SH, **L**H, **A**CTH, **T**SH, **C**RH, **h**CG, **A**DH (V_2-receptor), **M**SH, **P**TH, **C**alcitonin, **H**istamine (H_2-receptor), **G**lucagon, **G**HRH	**F**LAT **Ch**AMPs **CH**u**GG**
cGMP	**B**NP, **A**NP, E**D**RF (NO)	**BAD** Gra**MP**a Think vasodilation and diuresis
IP_3	**G**nRH, **O**xytocin, **A**DH (V_1-receptor), **T**RH, **H**istamine (H_1-receptor), **A**ngiotensin II, **G**astrin	**GOAT HAG**
Intracellular receptor	**P**rogesterone, **E**strogen, **T**estosterone, **C**ortisol, **A**ldosterone, **T_3**/**T_4**, **V**itamin D	**PET CAT** on **TV**
Receptor tyrosine kinase	I**GF**-1, F**GF**, PD**GF**, E**GF**, T**GF**-β, **In**sulin	**MAP** kinase pathway **G**et **F**ound **In** the **MAP**
Nonreceptor tyrosine kinase	**P**rolactin, **I**mmunomodulators (eg, cytokines IL-2, IL-6, IFN), **GH**, **G**-CSF, **E**rythropoietin, **T**hrombopoietin	JAK/STAT pathway Think acidophils and cytokines **PIGGLET**

Signaling pathways of steroid hormones



Steroid hormones are lipophilic and therefore must circulate bound to specific binding globulins, which ↑ their solubility.

In men, ↑ sex hormone–binding globulin (SHBG) lowers free testosterone → gynecomastia.

In women, ↓ SHBG raises free testosterone → hirsutism.

↑ estrogen (eg, OCPs, pregnancy) → ↑ SHBG.

(c) ketabton.com: The Digital Library

▸ ENDOCRINE—PATHOLOGY

Syndrome of inappropriate antidiuretic hormone secretion

Characterized by:
- Excessive free water retention
- Euvolemic hyponatremia with continued urinary Na^+ excretion
- Urine osmolality > serum osmolality

Body responds to water retention with ↓ aldosterone and ↑ ANP and BNP → ↑ urinary Na^+ secretion → normalization of extracellular fluid volume → euvolemic hyponatremia. Very low serum Na^+ levels can lead to cerebral edema, seizures. Correct slowly to prevent osmotic demyelination syndrome (formerly called central pontine myelinolysis).

SIADH causes include:
- Ectopic ADH (eg, small cell lung cancer)
- CNS disorders/head trauma
- Pulmonary disease
- Drugs (eg, SSRIs, carbamazepine, cyclophosphamide)

Treatment: fluid restriction (first line), salt tablets, IV hypertonic saline, diuretics, ADH antagonists (eg, conivaptan, tolvaptan, demeclocycline).

Diabetes insipidus

Characterized by intense thirst and polyuria with inability to concentrate urine due to lack of ADH (central) or failure of response to circulating ADH (nephrogenic).

	Central DI	Nephrogenic DI
ETIOLOGY	Pituitary tumor, autoimmune, trauma, surgery, ischemic encephalopathy, idiopathic	Hereditary (ADH receptor mutation), 2° to hypercalcemia, hypokalemia, lithium, demeclocycline (ADH antagonist)
FINDINGS	↓ ADH	Normal or ↑ ADH levels
	Urine specific gravity < 1.006 Urine osmolality < 300 mOsm/kg Serum osmolality > 290 mOsm/kg Hyperosmotic volume contraction	
WATER DEPRIVATION TEST[a]	> 50% ↑ in urine osmolality only after administration of ADH analog	Minimal change in urine osmolality, even after administration of ADH analog
TREATMENT	Desmopressin Hydration	HCTZ, indomethacin, amiloride Hydration, dietary salt restriction, avoidance of offending agent

[a]No water intake for 2–3 hr followed by hourly measurements of urine volume and osmolality as well as plasma Na^+ concentration and osmolality. ADH analog (desmopressin) is administered if serum osmolality > 295–300 mOsm/kg, plasma $Na^+ \geq 145$ mEq/L, or urine osmolality does not rise despite a rising plasma osmolality.

(c) ketabton.com: The Digital Library

Hypopituitarism

Undersecretion of pituitary hormones due to:

- Nonsecreting pituitary adenoma, craniopharyngioma
- Sheehan syndrome—ischemic infarct of pituitary following postpartum bleeding; pregnancy-induced pituitary growth → ↑ susceptibility to hypoperfusion. Usually presents with failure to lactate, absent menstruation, cold intolerance
- Empty sella syndrome—atrophy or compression of pituitary (which lies in the sella turcica), often idiopathic, common in obese women; associated with idiopathic intracranial hypertension
- Pituitary apoplexy—sudden hemorrhage of pituitary gland, often in the presence of an existing pituitary adenoma. Usually presents with sudden onset severe headache, visual impairment (eg, bitemporal hemianopia, diplopia due to CN III palsy), and features of hypopituitarism
- Brain injury
- Radiation

Treatment: hormone replacement therapy (corticosteroids, thyroxine, sex steroids, human growth hormone)

Acromegaly

Excess GH in adults. Typically caused by pituitary adenoma.

FINDINGS

Large tongue with deep furrows, deep voice, large hands and feet, coarsening of facial features with aging A, frontal bossing, diaphoresis (excessive sweating), impaired glucose tolerance (insulin resistance), hypertension. ↑ risk of colorectal polyps and cancer.

↑ GH in children → gigantism (↑ linear bone growth). HF most common cause of death.

DIAGNOSIS

↑ serum IGF-1; failure to suppress serum GH following oral glucose tolerance test; pituitary mass seen on brain MRI.

TREATMENT

Pituitary adenoma resection. If not cured, treat with octreotide (somatostatin analog), pegvisomant (GH receptor antagonist), or dopamine agonists (eg, cabergoline).

(c) ketabton.com: The Digital Library

Hypothyroidism vs hyperthyroidism

FINDINGS	Hypothyroidism	Hyperthyroidism
METABOLIC	Cold intolerance, ↓ sweating, weight gain (↓ basal metabolic rate → ↓ calorigenesis), hyponatremia (↓ free water clearance)	Heat intolerance, ↑ sweating, weight loss (↑ synthesis of Na^+-K^+ ATPase → ↑ basal metabolic rate → ↑ calorigenesis)
SKIN/HAIR	Dry, cool skin (due to ↓ blood flow); coarse, brittle hair; diffuse alopecia; brittle nails; puffy facies and generalized nonpitting edema (myxedema) due to ↑ GAGs in interstitial spaces → ↑ osmotic pressure → water retention	Warm, moist skin (due to vasodilation); fine hair; onycholysis (A); pretibial myxedema in Graves disease
OCULAR	Periorbital edema	Ophthalmopathy in Graves disease (including periorbital edema, exophthalmos), lid lag/retraction (↑ sympathetic stimulation of levator palpebrae superioris and superior tarsal muscle)
GASTROINTESTINAL	Constipation (↓ GI motility), ↓ appetite	Hyperdefecation/diarrhea (↑ GI motility), ↑ appetite
MUSCULOSKELETAL	Hypothyroid myopathy (proximal weakness, ↑ CK), carpal tunnel syndrome, myoedema (small lump rising on the surface of a muscle when struck with a hammer)	Thyrotoxic myopathy (proximal weakness, normal CK), osteoporosis/↑ fracture rate (T_3 directly stimulates bone resorption)
REPRODUCTIVE	Abnormal uterine bleeding, ↓ libido, infertility	Abnormal uterine bleeding, gynecomastia, ↓ libido, infertility
NEUROPSYCHIATRIC	Hypoactivity, lethargy, fatigue, weakness, depressed mood, ↓ reflexes (delayed/slow relaxing)	Hyperactivity, restlessness, anxiety, insomnia, fine tremors (due to ↑ β-adrenergic activity), ↑ reflexes (brisk)
CARDIOVASCULAR	Bradycardia, dyspnea on exertion (↓ cardiac output)	Tachycardia, palpitations, dyspnea, arrhythmias (eg, atrial fibrillation), chest pain and systolic HTN due to ↑ number and sensitivity of β-adrenergic receptors, ↑ expression of cardiac sarcolemmal ATPase and ↓ expression of phospholamban
LABS	↑ TSH (if 1°) ↓ free T_3 and T_4 Hypercholesterolemia (due to ↓ LDL receptor expression)	↓ TSH (if 1°) ↑ free T_3 and T_4 ↓ LDL, HDL, and total cholesterol

(c) ketabton.com: The Digital Library

Hypothyroidism

Hashimoto thyroiditis	Most common cause of hypothyroidism in iodine-sufficient regions; an autoimmune disorder with antithyroid peroxidase (antimicrosomal) and antithyroglobulin antibodies. Associated with HLA-DR3, HLA-DR5, ↑ risk of non-Hodgkin lymphoma (typically of B-cell origin). May be hyperthyroid early in course due to thyrotoxicosis during follicular rupture. Histology: Hürthle cells A, lymphoid aggregates with germinal centers. Findings: moderately enlarged, **nontender** thyroid.
Postpartum thyroiditis	Self-limited thyroiditis arising up to 1 year after delivery. Presents as transient hyperthyroidism, hypothyroidism, or hyperthyroidism followed by hypothyroidism. Majority of women are euthyroid following resolution. Thyroid usually painless and normal in size. Histology: lymphocytic infiltrate with occasional germinal center formation.
Congenital hypothyroidism (cretinism)	Severe fetal hypothyroidism due to antibody-mediated maternal hypothyroidism, thyroid dysgenesis (most common cause in US; eg, agenesis, ectopy, hypoplasia), iodine deficiency, dyshormonogenetic goiter (commonly due to mutations in thyroid peroxidase). Findings (6 P's): Pot-bellied, Pale, Puffy-faced child B with Protruding umbilicus, Protuberant tongue C, and Poor brain development.
Subacute granulomatous thyroiditis (de Quervain)	Self-limited disease often following a flu-like illness (eg, viral infection). May be hyperthyroid early in course, followed by hypothyroidism (permanent in ~15% of cases). Histology: granulomatous inflammation. Findings: ↑ ESR, jaw pain, very **tender** thyroid. (de Quervain is associated with pain.)
Riedel thyroiditis	Thyroid replaced by fibrous tissue and inflammatory infiltrate D. Fibrosis may extend to local structures (eg, trachea, esophagus), mimicking anaplastic carcinoma. ⅓ of patients are hypothyroid. Considered a manifestation of IgG_4-related systemic disease (eg, autoimmune pancreatitis, retroperitoneal fibrosis, noninfectious aortitis). Findings: fixed, hard (rock-like), **painless** goiter.
Other causes	Iodine deficiency (with goiter E), goitrogens (eg, amiodarone, lithium), Wolff-Chaikoff effect (thyroid gland downregulation in response to ↑ iodide).

(c) ketabton.com: The Digital Library

Hyperthyroidism

Graves disease



Most common cause of hyperthyroidism. Thyroid-stimulating immunoglobulin (IgG, can cause transient neonatal hyperthyroidism; type II hypersensitivity) stimulates TSH receptors on thyroid (hyperthyroidism, diffuse goiter), dermal fibroblasts (pretibial myxedema), and orbital fibroblasts (Graves orbitopathy). Activation of T-cells → lymphocytic infiltration of retroorbital space → ↑ cytokines (eg, TNF-α, IFN-γ) → ↑ fibroblast secretion of hydrophilic GAGs → ↑ osmotic muscle swelling, muscle inflammation, and adipocyte count → exophthalmos A. Often presents during stress (eg, pregnancy). Associated with HLA-DR3 and HLA-B8.
Histology: tall, crowded follicular epithelial cells; scalloped colloid.

Toxic multinodular goiter

Focal patches of hyperfunctioning follicular cells distended with colloid working independently of TSH (due to TSH receptor mutations in 60% of cases). ↑ release of T_3 and T_4. Hot nodules are rarely malignant.

Thyroid storm

Uncommon but serious complication that occurs when hyperthyroidism is incompletely treated/untreated and then significantly worsens in the setting of acute stress such as infection, trauma, surgery. Presents with agitation, delirium, fever, diarrhea, coma, and tachyarrhythmia (cause of death). May see ↑ LFTs. Treat with the 4 P's: β-blockers (eg, Propranolol), Propylthiouracil, corticosteroids (eg, Prednisolone), Potassium iodide (Lugol iodine). Iodide load → ↓ T_4 synthesis → Wolff-Chaikoff effect.

Jod-Basedow phenomenon

Iodine-induced hyperthyroidism. Occurs when a patient with iodine deficiency and partially autonomous thyroid tissue (eg, autonomous nodule) is made iodine replete. Can happen after iodine IV contrast or amiodarone use. Opposite to Wolff-Chaikoff effect.

Causes of goiter

Smooth/diffuse: Graves disease, Hashimoto thyroiditis, iodine deficiency, TSH-secreting pituitary adenoma.
Nodular: toxic multinodular goiter, thyroid adenoma, thyroid cancer, thyroid cyst.

Thyroid adenoma



Benign solitary growth of the thyroid. Most are nonfunctional ("cold"), can rarely cause hyperthyroidism via autonomous thyroid hormone production ("hot" or "toxic"). Most common histology is follicular (arrows in A); absence of capsular or vascular invasion (unlike follicular carcinoma).

(c) ketabton.com: The Digital Library

Thyroid cancer

Typically diagnosed with fine needle aspiration; treated with thyroidectomy. Complications of surgery include hypocalcemia (due to removal of parathyroid glands), transection of recurrent laryngeal nerve during ligation of inferior thyroid artery (leads to dysphagia and dysphonia [hoarseness]), and injury to the external branch of the superior laryngeal nerve during ligation of superior thyroid vascular pedicle (may lead to loss of tenor usually noticeable in professional voice users).

Papillary carcinoma



Most common, excellent prognosis. Empty-appearing nuclei with central clearing ("**Orphan Annie**" eyes) A, psam**Moma** bodies, nuclear grooves (**Papi** and **Moma** adopted **Orphan Annie**). ↑ risk with *RET/PTC* rearrangements and *BRAF* mutations, childhood irradiation.

Papillary carcinoma: most **P**revalent, **P**alpable lymph nodes. Good prognosis.

Follicular carcinoma

Good prognosis. Invades thyroid capsule and vasculature (unlike follicular adenoma), uniform follicles; hematogenous spread is common. Associated with *RAS* mutation and *PAX8-PPAR-γ* translocations.

Medullary carcinoma



From parafollicular "**C** cells"; produces **c**alcitonin, sheets of polygonal cells in an amyloid stroma B (stains with **C**ongo red). Associated with MEN 2A and 2B (*RET* mutations).

Undifferentiated/ anaplastic carcinoma

Older patients; presents with rapidly enlarging neck mass → compressive symptoms (eg, dyspnea, dysphagia, hoarseness); very poor prognosis. Associated with *TP53* mutation.

Diagnosing parathyroid disease



(c) ketabton.com: The Digital Library

Hypoparathyroidism

Due to injury to parathyroid glands or their blood supply (usually during surgery), autoimmune destruction, or DiGeorge syndrome. Findings: tetany, hypocalcemia, hyperphosphatemia.

Chvostek sign—tapping of facial nerve (tap the Cheek) → contraction of facial muscles.

Trousseau sign—occlusion of brachial artery with BP cuff (cuff the Triceps) → carpal spasm.

Pseudohypoparathyroidism type 1A—autosomal dominant, maternally transmitted mutations (imprinted *GNAS* gene). GNAS1-inactivating mutation (coupled to PTH receptor) that encodes the G_s protein α subunit → inactivation of adenylate cyclase when PTH binds to its receptor → end-organ resistance (kidney and bone) to PTH.

Physical findings: Albright hereditary osteodystrophy (shortened 4th/5th digits A, short stature, round face, subcutaneous calcifications, developmental delay).

Labs: ↑ PTH, ↓ Ca^{2+}, ↑ PO_4^{3-}.

Pseudopseudohypoparathyroidism—autosomal dominant, paternally transmitted mutations (imprinted *GNAS* gene) but without end-organ resistance to PTH due to normal maternal allele maintaining renal responsiveness to PTH.

Physical findings: same as Albright hereditary osteodystrophy.

Labs: normal PTH, Ca^{2+}, PO_4^{3-}.

Lab values in hypocalcemia

DISORDER	Ca^{2+}	PO_4^{3-}	PTH
Vitamin D deficiency	↓	↓	↑
Hypoparathyroidism	↓	↑	↓
2° hyperparathyroidism (CKD)	↓	↑	↑
Pseudohypoparathyroidism	↓	↑	↑
Hyperphosphatemia	↓	↑	↑

(c) ketabton.com: The Digital Library

Hyperparathyroidism

Primary hyperparathyroidism 	Usually due to parathyroid adenoma or hyperplasia. **Hypercalcemia**, hypercalciuria (renal **stones**), polyuria (**thrones**), hypophosphatemia, ↑ PTH, ↑ ALP, ↑ urinary cAMP. Most often asymptomatic. May present with **bone** pain, weakness, constipation ("**groans**"), abdominal/flank pain (kidney stones, acute pancreatitis), neuropsychiatric disturbances ("**psychiatric overtones**").	**Osteitis fibrosa cystica**—cystic **bone** spaces filled with brown fibrous tissue A ("brown tumor" consisting of osteoclasts and deposited hemosiderin from hemorrhages; causes bone pain). Due to ↑ PTH, classically associated with 1° (but also seen with 2°) hyperparathyroidism. "**Stones**, **thrones**, **bones**, **groans**, and **psychiatric overtones**."
Secondary hyperparathyroidism	2° hyperplasia due to ↓ Ca^{2+} absorption and/or ↑ PO_4^{3-}, most often in chronic kidney disease (causes hypovitaminosis D and hyperphosphatemia → ↓ Ca^{2+}). **Hypocalcemia**, hyperphosphatemia in chronic kidney disease (vs hypophosphatemia with most other causes), ↑ ALP, ↑ PTH.	**Renal osteodystrophy**—renal disease → 2° and 3° hyperparathyroidism → bone lesions.
Tertiary hyperparathyroidism	Refractory (autonomous) hyperparathyroidism resulting from chronic kidney disease. ↑↑ PTH, ↑ Ca^{2+}.	

Familial hypocalciuric hypercalcemia	Defective G-coupled Ca^{2+}-sensing receptors in multiple tissues (eg, parathyroids, kidneys). Higher than normal Ca^{2+} levels required to suppress PTH. Excessive renal Ca^{2+} reabsorption → mild hypercalcemia and hypocalciuria with normal to ↑ PTH levels.

(c) ketabton.com: The Digital Library

Diabetes mellitus

ACUTE MANIFESTATIONS

Polydipsia, polyuria, polyphagia, weight loss, DKA (type 1), hyperosmolar hyperglycemic state (type 2).

Rarely, can be caused by unopposed secretion of GH and epinephrine. Also seen in patients on glucocorticoid therapy (steroid diabetes).

CHRONIC COMPLICATIONS

Nonenzymatic glycation:

- Small vessel disease (diffuse thickening of basement membrane) → retinopathy (hemorrhage, exudates, microaneurysms, vessel proliferation), glaucoma, nephropathy. Nodular glomerulosclerosis → progressive proteinuria (initially microalbuminuria; ACE inhibitors and ARBs are renoprotective) and arteriolosclerosis (causing hypertension) → chronic kidney disease.
- Large vessel atherosclerosis, CAD, peripheral vascular occlusive disease, gangrene → limb loss, cerebrovascular disease. MI most common cause of death.

Osmotic damage (sorbitol accumulation in organs with aldose reductase and ↓ or absent sorbitol dehydrogenase):

- Neuropathy (motor, sensory [glove and stocking distribution], and autonomic degeneration).
- Cataracts.

DIAGNOSIS

TEST	DIAGNOSTIC CUTOFF	NOTES
HbA_{1c}	≥ 6.5%	Reflects average blood glucose over prior 3 months
Fasting plasma glucose	≥ 126 mg/dL	Fasting for > 8 hours
2-hour oral glucose tolerance test	≥ 200 mg/dL	2 hours after consumption of 75 g of glucose in water

Insulin deficiency or severe insulin insensitivity



(c) ketabton.com: The Digital Library

Type 1 vs type 2 diabetes mellitus

	Type 1	Type 2
1° DEFECT	Autoimmune T-cell–mediated destruction of β cells (eg, due to presence of glutamic acid decarboxylase antibodies)	↑ resistance to insulin, progressive pancreatic β-cell failure
INSULIN NECESSARY IN TREATMENT	Always	Sometimes
AGE (EXCEPTIONS COMMON)	< 30 yr	> 40 yr
ASSOCIATION WITH OBESITY	No	Yes
GENETIC PREDISPOSITION	Relatively weak (50% concordance in identical twins), polygenic	Relatively strong (90% concordance in identical twins), polygenic
ASSOCIATION WITH HLA SYSTEM	Yes, HLA-DR4 and -DR3 (4 – 3 = type 1)	No
GLUCOSE INTOLERANCE	Severe	Mild to moderate
INSULIN SENSITIVITY	High	Low
KETOACIDOSIS	Common	Rare
β-CELL NUMBERS IN THE ISLETS	↓	Variable (with amyloid deposits)
SERUM INSULIN LEVEL	↓	↑ initially, but ↓ in advanced disease
CLASSIC SYMPTOMS OF POLYURIA, POLYDIPSIA, POLYPHAGIA, WEIGHT LOSS	Common	Sometimes
HISTOLOGY	Islet leukocytic infiltrate	Islet amyloid polypeptide (IAPP) deposits

Diabetic ketoacidosis

Insulin absent, ketones present (→ complications).
Insulin noncompliance or ↑ requirements from ↑ stress (eg, infection) → excess fat breakdown and ↑ ketogenesis from ↑ free fatty acids → ketone bodies (β-hydroxybutyrate > acetoacetate).

SIGNS/SYMPTOMS	DKA is Deadly: Delirium/psychosis, Kussmaul respirations (rapid, deep breathing), Abdominal pain/nausea/vomiting, Dehydration. Fruity breath odor (due to exhaled acetone).
LABS	Hyperglycemia, ↑ H^+, ↓ HCO_3^- (↑ anion gap metabolic acidosis), ↑ urine and blood ketone levels, leukocytosis. Normal/↑ serum K^+, but depleted intracellular K^+ due to transcellular shift from ↓ insulin and acidosis. Osmotic diuresis → ↑ K^+ loss in urine → total body K^+ depletion.
COMPLICATIONS	Life-threatening mucormycosis, cerebral edema, cardiac arrhythmias, HF.
TREATMENT	IV fluids, IV insulin, K^+ (to replete intracellular stores) +/– glucose to prevent hypoglycemia.

Hyperosmolar hyperglycemic state

Insulin present, ketones absent.
Profound hyperglycemia → excessive osmotic diuresis → dehydration and ↑ serum osmolality → HHS. Classically seen in elderly type 2 diabetics with limited ability to drink.

SIGNS/SYMPTOMS	Thirst, polyuria, lethargy, focal neurologic deficits, seizures.
LABS	Hyperglycemia (often >600 mg/dL), ↑ serum osmolality (> 320 mOsm/kg), normal pH (no acidosis), no ketones. Normal/↑ serum K^+, ↓ intracellular K^+.
COMPLICATIONS	Can progress to coma and death if untreated.
TREATMENT	IV fluids, IV insulin, and K^+ (to replete intracellular stores).

(c) ketabton.com: The Digital Library

Cushing syndrome

ETIOLOGY

↑ cortisol due to a variety of causes:

- Exogenous corticosteroids → ↓ ACTH → bilateral adrenal atrophy. Most common cause.
- Primary adrenal adenoma, hyperplasia, or carcinoma → ↓ ACTH → atrophy of uninvolved adrenal gland.
- ACTH-secreting pituitary adenoma (Cushing disease); paraneoplastic ACTH secretion (eg, small cell lung cancer, bronchial carcinoids)→ bilateral adrenal hyperplasia. Cushing disease is responsible for the majority of endogenous cases of Cushing syndrome.

FINDINGS

CUSHING Syndrome: ↑ Cholesterol, ↑ Urinary free cortisol, Skin changes (thinning, striae A), Hypertension, Immunosuppression, Neoplasm (a cause, not a finding), Growth retardation (in children), ↑ Sugar (hyperglycemia, insulin resistance). Also, amenorrhea, moon facies B, buffalo hump, osteoporosis, ↑ weight (truncal obesity), hirsutism.

DIAGNOSIS

Screening tests include: ↑ free cortisol on 24-hr urinalysis, ↑ late night salivary cortisol, and no suppression with overnight low-dose dexamethasone test.





Nelson syndrome

Enlargement of pre-existing ACTH–secreting pituitary adenoma after bilateral adrenalectomy for refractory Cushing disease → ↑ ACTH (hyperpigmentation), mass effect (headaches, bitemporal hemianopia).

Treatment: transsphenoidal resection, postoperative pituitary irradiation for residual tumor.

(c) ketabton.com: The Digital Library

Adrenal insufficiency

Inability of adrenal glands to generate enough glucocorticoids +/– mineralocorticoids for the body's needs. Symptoms include weakness, fatigue, orthostatic hypotension, muscle aches, weight loss, GI disturbances, sugar and/or salt cravings. Treatment: glucocorticoid/mineralocorticoid replacement.



Primary adrenal insufficiency



↓ gland function → ↓ cortisol, ↓ aldosterone → hypotension (hyponatremic volume contraction), hyperkalemia, metabolic acidosis, skin/mucosal hyperpigmentation A (↑ melanin synthesis due to ↑ MSH, a byproduct of ACTH production from POMC).

- **Acute**—sudden onset (eg, due to massive hemorrhage). May present with shock in acute adrenal crisis.
- **Chronic**—**Addison disease**. Due to adrenal atrophy or destruction by disease (autoimmune destruction most common in the Western world; TB most common in the developing world).

Primary Pigments the skin/mucosa.
Associated with autoimmune polyglandular syndromes.
Waterhouse-Friderichsen syndrome—acute 1° adrenal insufficiency due to adrenal hemorrhage associated with septicemia (usually *Neisseria meningitidis*), DIC, endotoxic shock.

Secondary adrenal insufficiency

Seen with ↓ pituitary ACTH production. No skin/mucosal hyperpigmentation (ACTH is not elevated), no hyperkalemia (aldosterone synthesis preserved due to functioning adrenal gland, intact RAAS).

Secondary Spares the skin/mucosa.

Tertiary adrenal insufficiency

Seen in patients with chronic exogenous steroid use, precipitated by abrupt withdrawal. Aldosterone synthesis unaffected.

Tertiary from Treatment.

Hyperaldosteronism

Increased secretion of aldosterone from adrenal gland. Clinical features include hypertension, ↓ or normal K^+, metabolic alkalosis. 1° hyperaldosteronism does not directly cause edema due to aldosterone escape mechanism. However, certain 2° causes of hyperaldosteronism (eg, heart failure) impair the aldosterone escape mechanism, leading to worsening of edema.

Primary hyperaldosteronism

Seen with adrenal adenoma (Conn syndrome) or bilateral adrenal hyperplasia. ↑ aldosterone, ↓ renin. Leads to treatment-resistant hypertension.

Secondary hyperaldosteronism

Seen in patients with renovascular hypertension, juxtaglomerular cell tumors (renin-producing), and edema (eg, cirrhosis, heart failure, nephrotic syndrome).

(c) ketabton.com: The Digital Library

Neuroendocrine tumors

Heterogeneous group of neoplasms originating from neuroendocrine cells (which have traits similar to nerve cells and hormone-producing cells).

Most neoplasms occur in the GI system (eg, carcinoid, gastrinoma), pancreas (eg, insulinoma, glucagonoma), and lungs (eg, small cell carcinoma). Also in thyroid (eg, medullary carcinoma) and adrenals (eg, pheochromocytoma).

Neuroendocrine cells (eg, pancreatic β cells, enterochromaffin cells) share a common biologic function through amine precursor uptake decarboxylase (APUD) despite differences in embryologic origin, anatomic site, and secretory products (eg, chromogranin A, neuron-specific enolase [NSE], synaptophysin, serotonin, histamine, calcitonin). Treatment: surgical resection, somatostatin analogs.

Neuroblastoma



Most common tumor of the adrenal medulla in **children,** usually < 4 years old. Originates from **N**eural crest cells. Occurs anywhere along the sympathetic chain.

Most common presentation is abdominal distension and a firm, irregular mass that can cross the midline (vs Wilms tumor, which is smooth and unilateral). Less likely to develop hypertension than with pheochromocytoma (**N**euroblastoma is **N**ormotensive). Can also present with opsoclonus-myoclonus syndrome ("dancing eyes-dancing feet").

↑ HVA and VMA (catecholamine metabolites) in urine. Homer-Wright rosettes (neuroblasts surrounding a central lumen A) characteristic of neuroblastoma and medulloblastoma. Bombesin and **N**SE ⊕. Associated with amplification of **N**-*myc* oncogene.

Pheochromocytoma

ETIOLOGY



Most common tumor of the adrenal medulla in **adults** A. Derived from chromaffin cells (arise from neural crest).

May be associated with germline mutations (eg, *NF-1*, *VHL*, *RET* [MEN 2A, 2B]).

Rule of 10's:
- 10% malignant
- 10% bilateral
- 10% extra-adrenal (eg, bladder wall, organ of Zuckerkandl)
- 10% calcify
- 10% kids

SYMPTOMS

Most tumors secrete epinephrine, norepinephrine, and dopamine, which can cause episodic hypertension. May also secrete EPO → polycythemia.

Symptoms occur in "spells"—relapse and remit.

Episodic hyperadrenergic symptoms (**5 P**'s):
- **P**ressure (↑ BP)
- **P**ain (headache)
- **P**erspiration
- **P**alpitations (tachycardia)
- **P**allor

FINDINGS

↑ catecholamines and metanephrines (eg, homovanillic acid, vanillylmandelic acid) in urine and plasma.

Chromogranin, synaptophysin and NSE ⊕.

TREATMENT

Irreversible α-antagonists (eg, phenoxybenzamine) followed by β-blockers prior to tumor resection. α-blockade must be achieved before giving β-blockers to avoid a hypertensive crisis. **A** before **B**.

Phenoxybenzamine for **phe**ochromocytoma.

(c) ketabton.com: The Digital Library

Multiple endocrine neoplasias

All **MEN** syndromes have autosomal **dominant** inheritance.
"All **MEN** are **dominant**" (or so they think).

SUBTYPE	CHARACTERISTICS
MEN 1	**P**ituitary tumors (prolactin or GH) **P**ancreatic endocrine tumors—Zollinger-Ellison syndrome, insulinomas, VIPomas, glucagonomas (rare) **P**arathyroid adenomas Associated with mutation of *MEN1* (menin, a tumor suppressor, chromosome 11), angiofibromas, collagenomas, meningiomas
MEN 2A	**P**arathyroid hyperplasia Medullary thyroid carcinoma—neoplasm of parafollicular C cells; secretes calcitonin; prophylactic thyroidectomy required **P**heochromocytoma (secretes catecholamines) Associated with mutation in *RET* (codes for receptor tyrosine kinase)
MEN 2B  	Medullary thyroid carcinoma **P**heochromocytoma Mucosal neuromas **A** (oral/intestinal ganglioneuromatosis) Associated with marfanoid habitus; mutation in *RET* gene



MEN 1 = **3 P**'s: **P**ituitary, **P**arathyroid, and **P**ancreas
MEN 2A = **2 P**'s: **P**arathyroid and **P**heochromocytoma
MEN 2B = **1 P**: **P**heochromocytoma

Pancreatic islet cell tumors

Insulinoma	Tumor of pancreatic β cells → overproduction of insulin → hypoglycemia. May see Whipple triad: low blood glucose, symptoms of hypoglycemia (eg, lethargy, syncope, diplopia), and resolution of symptoms after normalization of plasma glucose levels. Symptomatic patients have ↓ blood glucose and ↑ C-peptide levels (vs exogenous insulin use). ~ 10% of cases associated with MEN 1 syndrome. Treatment: surgical resection.
Glucagonoma	Tumor of pancreatic α cells → overproduction of glucagon. Presents with **6 D**'s: **D**ermatitis (necrolytic migratory erythema), **D**iabetes (hyperglycemia), **D**VT, **D**eclining weight, **D**epression, **D**iarrhea. Treatment: octreotide, surgical resection.
Somatostatinoma	Tumor of pancreatic δ cells → overproduction of somatostatin → ↓ secretion of secretin, cholecystokinin, glucagon, insulin, gastrin, gastric inhibitory peptide (GIP). May present with diabetes/glucose intolerance, steatorrhea, gallstones, achlorhydria. Treatment: surgical resection; somatostatin analogs (eg, octreotide) for symptom control.

(c) ketabton.com: The Digital Library

Carcinoid syndrome



Carcinoid tumors arise from neuroendocrine cells most commonly in the intestine or lung. Rare and does not occur if tumor is limited to the GI tract.
Prominent rosettes (arrow in A), chromogranin A ⊕ and synaptophysin ⊕).
Neuroendocrine cells secrete 5-HT → recurrent diarrhea, wheezing, right-sided valvular heart disease (eg, tricuspid regurgitation, pulmonic stenosis), niacin deficiency (pellagra). 5-HT undergoes hepatic first-pass metabolism and enzymatic breakdown by MAO in the lung.
Treatment: surgical resection, somatostatin analog (eg, octreotide, telotristat) for symptom control.
Rule of thirds:
- 1/3 metastasize
- 1/3 present with 2nd malignancy
- 1/3 are multiple

Zollinger-Ellison syndrome

Gastrin-secreting tumor (gastrinoma) of pancreas or duodenum. Acid hypersecretion causes recurrent ulcers in duodenum and jejunum. Presents with abdominal pain (peptic ulcer disease, distal ulcers), diarrhea (malabsorption). Positive secretin stimulation test: gastrin levels remain elevated after administration of secretin, which normally inhibits gastrin release. May be associated with MEN 1.

▸ ENDOCRINE—PHARMACOLOGY

Diabetes mellitus therapy

All patients with diabetes mellitus should receive education on diet, exercise, blood glucose monitoring, and complication management. Treatment differs based on the type of diabetes and glycemic control:
- Type 1 DM—insulin replacement
- Type 2 DM—oral agents (metformin is first line), non-insulin injectables, insulin replacement; weight loss particularly helpful in lowering blood glucose
- Gestational DM—insulin replacement if nutrition therapy and exercise alone fail

Regular (short-acting) insulin is preferred for DKA (IV), hyperkalemia (+ glucose), stress hyperglycemia.
To **N**ormalize **P**ancreatic **F**unction (-gli**T**s, -gli**N**s, -gli**P**s, -gli**F**s).

DRUG CLASS	MECHANISM	ADVERSE EFFECTS
Insulin preparations		
Rapid acting (1-hr peak): **L**ispro, **A**spart, **G**lulisine (no **LAG**) Short acting (2–3 hr peak): regular Intermediate acting (4–10 hr peak): NPH Long acting (no real peak): detemir, glargine	Bind insulin receptor (tyrosine kinase activity) Liver: ↑ glucose storage as glycogen Muscle: ↑ glycogen, protein synthesis Fat: ↑ TG storage Cell membrane: ↑ K^+ uptake	Hypoglycemia, lipodystrophy, hypersensitivity reactions (rare), weight gain



(c) ketabton.com: The Digital Library

Diabetes mellitus therapy *(continued)*

DRUG CLASS	MECHANISM	ADVERSE EFFECTS
Increase insulin sensitivity		
Biguanides Metformin	Inhibit mGPD → inhibition of hepatic gluconeogenesis and the action of glucagon. ↑ glycolysis, peripheral glucose uptake (↑ insulin sensitivity).	GI upset, lactic acidosis (use with caution in renal insufficiency), vitamin B_{12} deficiency. Weight loss (often desired).
Glitazones/ thiazolidinediones "-gliTs" Pioglitazone, rosiglitazone	Activate PPAR-γ (a nuclear receptor) → ↑ insulin sensitivity and levels of adiponectin → regulation of glucose metabolism and fatty acid storage.	Weight gain, edema, HF, ↑ risk of fractures. Delayed onset of action (several weeks). Rosiglitazone: ↑ risk of MI, cardiovascular death.
Increase insulin secretion		
Sulfonylureas (1st gen) Chlorpropamide, tolbutamide	Close K^+ channels in pancreatic B cell membrane → cell depolarizes → insulin release via ↑ Ca^{2+} influx.	DisulFIRam-like reaction (FIRst-generation only). Rarely used.
Sulfonylureas (2nd gen) Glipizide, glyburide		Hypoglycemia (↑ risk in renal insufficiency), weight gain.
Meglitinides "-gliNs" Nateglinide, repaglinide		
Increase glucose-induced insulin secretion		
GLP-1 analogs Exenatide, liraglutide	↓ glucagon release, ↓ gastric emptying, ↑ glucose-dependent insulin release.	Nausea, vomiting, pancreatitis. Weight loss (often desired). ↑ satiety (often desired).
DPP-4 inhibitors "-gliPs" Linagliptin, saxagliptin, sitagliptin	Inhibit DPP-4 enzyme that deactivates GLP-1 → ↓ glucagon release, ↓ gastric emptying. ↑ glucose-dependent insulin release.	Respiratory and urinary infections, weight neutral. ↑ satiety (often desired).
Decrease glucose absorption		
Sodium-glucose co-transporter 2 (SGLT2) inhibitors "-gliFs" Canagliflozin, dapagliflozin, empagliflozin	Block reabsorption of glucose in proximal convoluted tubule.	Glucosuria (UTIs, vulvovaginal candidiasis), dehydration (orthostatic hypotension), hyperkalemia, weight loss. Use with caution in renal insufficiency (↓ efficacy with ↓ GFR).
α-glucosidase inhibitors Acarbose, miglitol	Inhibit intestinal brush-border α-glucosidases → delayed carbohydrate hydrolysis and glucose absorption → ↓ postprandial hyperglycemia.	GI upset, bloating. Not recommended in renal insufficiency.
Others		
Amylin analogs Pramlintide	↓ glucagon release, ↓ gastric emptying.	Hypoglycemia, nausea. ↑ satiety (often desired).

(c) ketabton.com: The Digital Library

Thionamides

Propylthiouracil, methimazole.

MECHANISM	Block thyroid peroxidase, inhibiting the oxidation of iodide as well as the organification and coupling of iodine → inhibition of thyroid hormone synthesis. **P**TU also blocks 5′-deiodinase → ↓ **P**eripheral conversion of T_4 to T_3.
CLINICAL USE	Hyperthyroidism. PTU used in first trimester of pregnancy (due to methimazole teratogenicity); methimazole used in second and third trimesters of pregnancy (due to risk of PTU-induced hepatotoxicity). Not used to treat Graves ophthalmopathy (treated with corticosteroids).
ADVERSE EFFECTS	Skin rash, agranulocytosis (rare), aplastic anemia, hepatotoxicity. Methimazole is a possible teratogen (can cause aplasia cutis).

Levothyroxine, liothyronine

MECHANISM	Hormone replacement for T_4 (levothyroxine) or T_3 (liothyronine).
CLINICAL USE	Hypothyroidism, myxedema. May be abused for weight loss. Distinguish exogenous hyperthyroidism from endogenous hyperthyroidism by using a combination of TSH receptor antibodies, radioactive iodine uptake, and/or measurement of thyroid blood flow on ultrasound.
ADVERSE EFFECTS	Tachycardia, heat intolerance, tremors, arrhythmias.

Hypothalamic/pituitary drugs

DRUG	CLINICAL USE
Conivaptan, tolvaptan	ADH antagonists SIADH (block action of ADH at V_2-receptor)
Demeclocycline	ADH antagonist, a tetracycline SIADH
Desmopressin	Central DI, von Willebrand disease, sleep enuresis, hemophilia A
GH	GH deficiency, Turner syndrome
Oxytocin	Induction of labor (stimulates uterine contractions), control uterine hemorrhage
Somatostatin (octreotide)	Acromegaly, carcinoid syndrome, gastrinoma, glucagonoma, esophageal varices

Fludrocortisone

MECHANISM	Synthetic analog of aldosterone with little glucocorticoid effects.
CLINICAL USE	Mineralocorticoid replacement in 1° adrenal insufficiency.
ADVERSE EFFECTS	Similar to glucocorticoids; also edema, exacerbation of heart failure, hyperpigmentation.

(c) ketabton.com: The Digital Library

Cinacalcet

MECHANISM	Sensitizes Ca^{2+}-sensing receptor (CaSR) in parathyroid gland to circulating Ca^{2+} → ↓ PTH.
CLINICAL USE	2° hyperparathyroidism in patients with CKD receiving hemodialysis, hypercalcemia in 1° hyperparathyroidism (if parathyroidectomy fails), or in parathyroid carcinoma.
ADVERSE EFFECTS	Hypocalcemia.

Sevelamer

MECHANISM	Nonabsorbable phosphate binder that prevents phosphate absorption from the GI tract.
CLINICAL USE	Hyperphosphatemia in CKD.
ADVERSE EFFECTS	Hypophosphatemia, GI upset.

(c) ketabton.com: The Digital Library

▸ NOTES

(c) ketabton.com: The Digital Library

Gastrointestinal

"A good set of bowels is worth more to a man than any quantity of brains."
—Josh Billings

"Man should strive to have his intestines relaxed all the days of his life."
—Moses Maimonides

"All right, let's not panic. I'll make the money by selling one of my livers. I can get by with one."
—Homer Simpson

When studying the gastrointestinal system, be sure to understand the normal embryology, anatomy, and physiology and how it is affected in the various pathologic diseases. Study not only what a disease entails, but also its specific findings, so that you can differentiate between two similar diseases. For example, what specifically makes ulcerative colitis different than Crohn disease? Also, it is important to understand bile metabolism and which lab values increase or decrease depending on the disease process. Be comfortable with basic interpretation of abdominal x-rays, CT scans, and endoscopic images.

- Embryology 358
- Anatomy 360
- Physiology 371
- Pathology 376
- Pharmacology 398

FAS1_2019_09-Gastrointestinal.indd 357 11/7/19 4:42 PM

(c) ketabton.com: The Digital Library

▸ GASTROINTESTINAL—EMBRYOLOGY

Normal gastrointestinal embryology

Foregut—esophagus to duodenum at level of pancreatic duct and common bile duct insertion (ampulla of Vater).
Midgut—lower duodenum to proximal 2/3 of transverse colon.
Hindgut—distal 1/3 of transverse colon to anal canal above pectinate line.
Midgut development:

- 6th week—physiologic herniation of midgut through umbilical ring
- 10th week—returns to abdominal cavity + rotates around superior mesenteric artery (SMA), total 270° counterclockwise

Ventral wall defects

Developmental defects due to failure of rostral fold closure (eg, sternal defects [ectopia cordis]), lateral fold closure (eg, omphalocele, gastroschisis), or caudal fold closure (eg, bladder exstrophy).

	Gastroschisis	Omphalocele
ETIOLOGY	Extrusion of abdominal contents through abdominal folds (typically right of umbilicus)	Failure of lateral walls to migrate at umbilical ring → persistent midline herniation of abdominal contents into umbilical cord
COVERAGE	Not covered by peritoneum or amnion A; "the guts come out of the gap (schism) in the letter G"	Surrounded by peritoneum B (light gray shiny sac); "abdominal contents are sealed in the letter O"
ASSOCIATIONS	Not associated with chromosome abnormalities; favorable prognosis	Associated with congenital anomalies (eg, trisomies 13 and 18, Beckwith-Wiedemann syndrome) and other structural abnormalities (eg, cardiac, GU, neural tube)

Congenital umbilical hernia

Failure of umbilical ring to close after physiologic herniation of the midgut. Small defects usually close spontaneously.

(c) ketabton.com: The Digital Library

Tracheoesophageal anomalies

Esophageal atresia (EA) with distal tracheoesophageal fistula (TEF) is the most common (85%) and often presents as polyhydramnios in utero (due to inability of fetus to swallow amniotic fluid). Neonates drool, choke, and vomit with first feeding. TEFs allow air to enter stomach (visible on CXR). Cyanosis is 2° to laryngospasm (to avoid reflux-related aspiration). Clinical test: failure to pass nasogastric tube into stomach.

In **H**-type, the fistula resembles the letter **H**. In pure EA, CXR shows gasless abdomen.



Intestinal atresia



Presents with bilious vomiting and abdominal distension within first 1–2 days of life.

Duodenal atresia—failure to recanalize. Abdominal x-ray **A** shows "double bubble" (dilated stomach, proximal duodenum). Associated with **D**own syndrome.

Jejunal and ileal atresia—disruption of mesenteric vessels (typically SMA) → ischemic necrosis of fetal intestine → segmental resorption: bowel becomes discontinuous. X-ray shows dilated loops of small bowel with air-fluid levels.

Hypertrophic pyloric stenosis



Most common cause of gastric outlet obstruction in infants (1:600). Palpable olive-shaped mass in epigastric region, visible peristaltic waves, and nonbilious projectile vomiting at ~ 2–6 weeks old. More common in firstborn males; associated with exposure to macrolides.

Results in hypokalemic hypochloremic metabolic alkalosis (2° to vomiting of gastric acid and subsequent volume contraction).

Ultrasound shows thickened and lengthened pylorus **A**.

Treatment: surgical incision of pyloric muscles (pyloromyotomy).

(c) ketabton.com: The Digital Library

Pancreas and spleen embryology



Pancreas—derived from foregut. Ventral pancreatic bud contributes to uncinate process and main pancreatic duct. The dorsal pancreatic bud alone becomes the body, tail, isthmus, and accessory pancreatic duct. Both the ventral and dorsal buds contribute to pancreatic head.

Annular pancreas—abnormal rotation of ventral pancreatic bud forms a ring of pancreatic tissue → encircles 2nd part of duodenum; may cause duodenal narrowing (arrows in A) and vomiting.

Pancreas divisum—ventral and dorsal parts fail to fuse at 8 weeks. Common anomaly; mostly asymptomatic, but may cause chronic abdominal pain and/or pancreatitis.

Spleen—arises in mesentery of stomach (hence is mesodermal) but has foregut supply (celiac trunk → splenic artery).



▸ GASTROINTESTINAL—ANATOMY

Retroperitoneal structures

Retroperitoneal structures A are posterior to (and outside of) the peritoneal cavity. Injuries to retroperitoneal structures can cause blood or gas accumulation in retroperitoneal space.

SAD PUCKER:
- **S**uprarenal (adrenal) glands [not shown]
- **A**orta and IVC
- **D**uodenum (2nd through 4th parts)
- **P**ancreas (except tail)
- **U**reters [not shown]
- **C**olon (descending and ascending)
- **K**idneys
- **E**sophagus (thoracic portion) [not shown]
- **R**ectum (partially) [not shown]





(c) ketabton.com: The Digital Library

Important gastrointestinal ligaments



LIGAMENT	CONNECTS	STRUCTURES CONTAINED	NOTES
Falciform ligament	Liver to anterior abdominal wall	Ligamentum teres hepatis (derivative of fetal umbilical vein), patent paraumbilical veins	Derivative of ventral mesentery
Hepatoduodenal ligament	Liver to duodenum	Portal triad: proper hepatic artery, portal vein, common bile duct	Derivative of ventral mesentery Pringle maneuver—ligament is compressed manually or with a vascular clamp in omental foramen to control bleeding from hepatic inflow source Borders the omental foramen, which connects the greater and lesser sacs Part of lesser omentum
Gastrohepatic ligament	Liver to lesser curvature of stomach	Gastric vessels	Derivative of ventral mesentery Separates greater and lesser sacs on the right May be cut during surgery to access lesser sac Part of lesser omentum
Gastrocolic ligament (not shown)	Greater curvature and transverse colon	Gastroepiploic arteries	Derivative of dorsal mesentery Part of greater omentum
Gastrosplenic ligament	Greater curvature and spleen	Short gastrics, left gastroepiploic vessels	Derivative of dorsal mesentery Separates greater and lesser sacs on the left Part of greater omentum
Splenorenal ligament	Spleen to left pararenal space	Splenic artery and vein, tail of pancreas	Derivative of dorsal mesentery

(c) ketabton.com: The Digital Library

Digestive tract anatomy

Layers of gut wall A (inside to outside—**MSMS**):

- **M**ucosa—epithelium, lamina propria, muscularis mucosa
- **S**ubmucosa—includes **S**ubmucosal nerve plexus (Mei**ss**ner), **S**ecretes fluid
- **M**uscularis externa—includes **M**yenteric nerve plexus (Auerbach), **M**otility
- **S**erosa (when intraperitoneal), adventitia (when retroperitoneal)

Ulcers can extend into submucosa, inner or outer muscular layer. Erosions are in mucosa only.

Frequency of basal electric rhythm (slow waves), which originate in the interstitial cells of Cajal:

- Stomach—3 waves/min
- Duodenum—12 waves/min
- Ileum—8–9 waves/min



Digestive tract histology

Esophagus	Nonkeratinized stratified squamous epithelium. Upper 1/3, striated muscle; middle and lower 2/3 smooth muscle, with some overlap at the transition.
Stomach	Gastric glands A.
Duodenum	Villi B and microvilli ↑ absorptive surface. Brunner glands (HCO_3^--secreting cells of submucosa) and crypts of Lieberkühn (contain stem cells that replace enterocytes/goblet cells and Paneth cells that secrete defensins, lysozyme, and TNF).
Jejunum	Villi, crypts of Lieberkühn, and plicae circulares (also present in distal duodenum) C.
Ileum	Peyer patches (arrow in D; lymphoid aggregates in lamina propria, submucosa), plicae circulares (proximal ileum), and crypts of Lieberkühn. Largest number of goblet cells in the small intestine.
Colon	Crypts of Lieberkühn with abundant goblet cells, but no villi E.



(c) ketabton.com: The Digital Library

Abdominal aorta and branches



Arteries supplying GI structures are single and branch anteriorly.

Arteries supplying non-GI structures are paired and branch laterally and posteriorly.

Two areas of the colon have dual blood supply from distal arterial branches ("watershed regions") → susceptible in colonic ischemia:

- Splenic flexure—SMA and IMA
- Rectosigmoid junction—the last sigmoid arterial branch from the IMA and superior rectal artery

Nutcracker syndrome—compression of left renal vein between superior mesenteric artery and aorta. Characterized by abdominal (flank) pain and gross hematuria (from rupture of thin-walled renal varicosities).

Superior mesenteric artery syndrome—characterized by intermittent intestinal obstruction symptoms (primarily postprandial pain) when SMA and aorta compress transverse (third) portion of duodenum. Typically occurs in conditions associated with diminished mesenteric fat (eg, low body weight/malnutrition).



(c) ketabton.com: The Digital Library

Gastrointestinal blood supply and innervation

EMBRYONIC GUT REGION	ARTERY	PARASYMPATHETIC INNERVATION	VERTEBRAL LEVEL	STRUCTURES SUPPLIED
Foregut	Celiac	Vagus	T12/L1	Pharynx (vagus nerve only) and lower esophagus (celiac artery only) to proximal duodenum; liver, gallbladder, pancreas, spleen (mesoderm)
Midgut	SMA	Vagus	L1	Distal duodenum to proximal 2/3 of transverse colon
Hindgut	IMA	Pelvic	L3	Distal 1/3 of transverse colon to upper portion of anal canal

Celiac trunk

Branches of celiac trunk: common hepatic, splenic, and left gastric. These constitute the main blood supply of the foregut.

Strong anastomoses exist between:

- Left and right gastroepiploics
- Left and right gastrics



(c) ketabton.com: The Digital Library

Portosystemic anastomoses



SITE OF ANASTOMOSIS	CLINICAL SIGN	PORTAL ↔ SYSTEMIC
1 Esopha**gus**	Esophageal varices	Left gastric ↔ esophageal (drains into azygos)
2 Umbilicus	**Caput** medusae	Paraumbilical ↔ small epigastric veins of the anterior abdominal wall.
3 **Rectum**	Anorectal varices	Superior rectal ↔ middle and inferior rectal

Varices of **gut**, **butt**, and **caput** (medusae) are commonly seen with portal hypertension.

4 Treatment with a **t**ransjugular **i**ntrahepatic **p**ortosystemic **s**hunt (**TIPS**) between the portal vein and hepatic vein relieves portal hypertension by shunting blood to the systemic circulation, bypassing the liver. TIPS can precipitate hepatic encephalopathy due to ↓ clearance of ammonia from shunting.

(c) ketabton.com: The Digital Library

Pectinate line

Also called dentate line. Formed where endoderm (hindgut) meets ectoderm.



Above pectinate line: internal hemorrhoids, adenocarcinoma.
Internal hemorrhoids receive visceral innervation and are therefore **not painful**.

Below pectinate line: external hemorrhoids, anal fissures, squamous cell carcinoma.
External hemorrhoids receive somatic innervation (inferior rectal branch of pudendal nerve) and are therefore **painful** if thrombosed.

Anal fissure—tear in anal mucosa below Pectinate line. Pain while Pooping; blood on toilet Paper. Located Posteriorly because this area is Poorly Perfused. Innervated by Pudendal nerve. Associated with low-fiber diets and constipation.

(c) ketabton.com: The Digital Library

Liver tissue architecture





The functional unit of the liver is made up of hexagonally arranged lobules surrounding the central vein with portal triads on the edges (consisting of a portal vein, hepatic artery, bile ducts, as well as lymphatics) A.

Apical surface of hepatocytes faces bile canaliculi. Basolateral surface faces sinusoids.

Kupffer cells (specialized macrophages) located in sinusoids (black arrows in B; yellow arrows show hepatic venule) clear bacteria and damaged or senescent RBCs.

Hepatic stellate (Ito) cells in space of Disse store vitamin A (when quiescent) and produce extracellular matrix (when activated). Responsible for hepatic fibrosis.

Zone I—periportal zone:
- Affected 1st by viral hepatitis
- Best oxygenated, most resistant to circulatory compromise
- Ingested toxins (eg, cocaine)

Zone II—intermediate zone:
- Yellow fever

Zone III—pericentral vein (centrilobular) zone:
- Affected 1st by ischemia (least oxygenated)
- High concentration of cytochrome P-450
- Most sensitive to metabolic toxins (eg, ethanol, CCl_4, halothane, rifampin, acetaminophen)
- Site of alcoholic hepatitis



(c) ketabton.com: The Digital Library

Biliary structures



Gallstones that reach the confluence of the common bile and pancreatic ducts at the ampulla of Vater can block both the common bile and pancreatic ducts (double duct sign), causing both cholangitis and pancreatitis, respectively.

Tumors that arise in head of pancreas (usually ductal adenocarcinoma) can cause obstruction of common bile duct → enlarged gallbladder with painless jaundice (Courvoisier sign).

Cholangiography shows filling defects in gallbladder (blue arrow) and cystic duct (red arrow) A.



Femoral region

ORGANIZATION	**Lateral to medial**: **N**erve-**A**rtery-**Ve**in-**L**ymphatics.	You go from **lateral to medial** to find your **NAVeL**.
Femoral triangle	Contains femoral nerve, artery, vein.	**Venous** near the **penis**.
Femoral sheath	Fascial tube 3–4 cm below inguinal ligament. Contains femoral vein, artery, and canal (deep inguinal lymph nodes) but not femoral nerve.	



(c) ketabton.com: The Digital Library

Inguinal canal



Abdominal wall



Posterior (internal) abdominal wall **Anterior (external) abdominal wall**

(c) ketabton.com: The Digital Library

Hernias

Protrusion of peritoneum through an opening, usually at a site of weakness. Contents may be at risk for incarceration (not reducible back into abdomen/pelvis) and strangulation (ischemia and necrosis). Complicated hernias can present with tenderness, erythema, fever.

Diaphragmatic hernia



Abdominal structures enter the thorax A; may occur due to congenital defect of pleuroperitoneal membrane or from trauma. Commonly occurs on left side due to relative protection of right hemidiaphragm by liver. Most commonly a hiatal hernia, in which stomach herniates upward through the esophageal hiatus of the diaphragm.

Sliding hiatal hernia—gastroesophageal junction is displaced upward as gastric cardia slides into hiatus; "hourglass stomach." Most common type. Associated with GERD.

Paraesophageal hiatal hernia—gastroesophageal junction is usually normal but gastric fundus protrudes into the thorax.



Indirect inguinal hernia

Goes through the internal (deep) inguinal ring, external (superficial) inguinal ring, and into the groin. Enters internal inguinal ring lateral to inferior epigastric vessels. Caused by failure of processus vaginalis to close (can form hydrocele). May be noticed in infants or discovered in adulthood. Much more common in males B.

Follows the pathway of testicular descent. Covered by all 3 layers of spermatic fascia.



Direct inguinal hernia

Protrudes through inguinal (Hesselbach) triangle. Bulges directly through parietal peritoneum medial to the inferior epigastric vessels but lateral to the rectus abdominis. Goes through external (superficial) inguinal ring only. Covered by external spermatic fascia. Usually occurs in older men due to acquired weakness of transversalis fascia.

MDs don't LIe:

Medial to inferior epigastric vessels = Direct hernia.

Lateral to inferior epigastric vessels = Indirect hernia.



Femoral hernia

Protrudes below inguinal ligament through femoral canal below and lateral to pubic tubercle. More common in females, but overall inguinal hernias are the most common.

More likely to present with incarceration or strangulation (vs inguinal hernia).



(c) ketabton.com: The Digital Library

▸ GASTROINTESTINAL—PHYSIOLOGY

Gastrointestinal regulatory substances

REGULATORY SUBSTANCE	SOURCE	ACTION	REGULATION	NOTES
Gastrin	G cells (antrum of stomach, duodenum)	↑ gastric H^+ secretion ↑ growth of gastric mucosa ↑ gastric motility	↑ by stomach distention/ alkalinization, amino acids, peptides, vagal stimulation via gastrin-releasing peptide (GRP) ↓ by pH < 1.5	↑ by chronic PPI use ↑ in chronic atrophic gastritis (eg, *H pylori*) ↑↑ in Zollinger-Ellison syndrome (gastrinoma)
Somatostatin	D cells (pancreatic islets, GI mucosa)	↓ gastric acid and pepsinogen secretion ↓ pancreatic and small intestine fluid secretion ↓ gallbladder contraction ↓ insulin and glucagon release	↑ by acid ↓ by vagal stimulation	Inhibits secretion of various hormones (encourages **somato-sta**sis) Octreotide is an analog used to treat acromegaly, carcinoid syndrome, and variceal bleeding
Cholecystokinin	I cells (duodenum, jejunum)	↑ pancreatic secretion ↑ gallbladder contraction ↓ gastric emptying ↑ sphincter of Oddi relaxation	↑ by fatty acids, amino acids	Acts on neural muscarinic pathways to cause pancreatic secretion
Secretin	S cells (duodenum)	↑ pancreatic HCO_3^- secretion ↓ gastric acid secretion ↑ bile secretion	↑ by acid, fatty acids in lumen of duodenum	↑ HCO_3^- neutralizes gastric acid in duodenum, allowing pancreatic enzymes to function
Glucose-dependent insulinotropic peptide	K cells (duodenum, jejunum)	Exocrine: ↓ gastric H^+ secretion Endocrine: ↑ insulin release	↑ by fatty acids, amino acids, oral glucose	Also called gastric inhibitory peptide (GIP) Oral glucose load ↑ insulin compared to IV equivalent due to GIP secretion
Motilin	Small intestine	Produces migrating motor complexes (MMCs)	↑ in fasting state	Motilin receptor agonists (eg, erythromycin) are used to stimulate intestinal peristalsis.
Vasoactive intestinal polypeptide	Parasympathetic ganglia in sphincters, gallbladder, small intestine	↑ intestinal water and electrolyte secretion ↑ relaxation of intestinal smooth muscle and sphincters	↑ by distention and vagal stimulation ↓ by adrenergic input	VIPoma—non-α, non-β islet cell pancreatic tumor that secretes VIP; associated with **W**atery **D**iarrhea, **H**ypokalemia, **A**chlorhydria (**WDHA** syndrome)
Nitric oxide		↑ smooth muscle relaxation, including lower esophageal sphincter (LES)		Loss of NO secretion is implicated in ↑ LES tone of achalasia
Ghrelin	Stomach	↑ appetite ("**ghr**owlin' stomach")	↑ in fasting state ↓ by food	↑ in Prader-Willi syndrome ↓ after gastric bypass surgery

(c) ketabton.com: The Digital Library

Gastrointestinal secretory products

PRODUCT	SOURCE	ACTION	REGULATION	NOTES
Intrinsic factor	Parietal cells (stomach A)	Vitamin B_{12}–binding protein (required for B_{12} uptake in terminal ileum)		Autoimmune destruction of parietal cells → chronic gastritis and pernicious anemia.
Gastric acid	Parietal cells (stomach)	↓ stomach pH	↑ by histamine, vagal stimulation (ACh), gastrin ↓ by somatostatin, GIP, prostaglandin, secretin	
Pepsin	Chief cells (stomach)	Protein digestion	↑ by vagal stimulation (ACh), local acid	Pepsinogen (inactive) is converted to pepsin (active) in the presence of H^+.
Bicarbonate	Mucosal cells (stomach, duodenum, salivary glands, pancreas) and Brunner glands (duodenum)	Neutralizes acid	↑ by pancreatic and biliary secretion with secretin	Trapped in mucus that covers the gastric epithelium.



(c) ketabton.com: The Digital Library

Locations of gastrointestinal secretory cells



Gastrin ↑ acid secretion primarily through its effects on enterochromaffin-like (ECL) cells (leading to histamine release) rather than through its direct effect on parietal cells.

Pancreatic secretions

Isotonic fluid; low flow → high Cl^-, high flow → high HCO_3^-.

ENZYME	ROLE	NOTES
α-amylase	Starch digestion	Secreted in active form
Lipases	Fat digestion	
Proteases	Protein digestion	Includes trypsin, chymotrypsin, elastase, carboxypeptidases Secreted as proenzymes also called zymogens
Trypsinogen	Converted to active enzyme trypsin → activation of other proenzymes and cleaving of additional trypsinogen molecules into active trypsin (positive feedback loop)	Converted to trypsin by enterokinase/enteropeptidase, a brush-border enzyme on duodenal and jejunal mucosa

Carbohydrate absorption



Only monosaccharides (glucose, galactose, fructose) are absorbed by enterocytes. Glucose and galactose are taken up by SGLT1 (Na^+ dependent). **F**ructose is taken up via **F**acilitated diffusion by GLUT**5**. All are transported to blood by GLUT2.

D-xylose absorption test: simple sugar that requires intact mucosa for absorption, but does not require digestive enzymes. Helps distinguish GI mucosal damage from other causes of malabsorption.

(c) ketabton.com: The Digital Library

Vitamin and mineral absorption

Iron	Absorbed as Fe^{2+} in duodenum	Iron Fist, Bro Clinically relevant in patients with small bowel disease or after resection (eg, vitamin B_{12} deficiency following terminal ileum resection)
Folate	Absorbed in small bowel	
Vitamin B_{12}	Absorbed in terminal ileum along with bile salts, requires intrinsic factor	



Peyer patches



Unencapsulated lymphoid tissue **A** found in lamina propria and submucosa of ileum. Contain specialized **M** cells that sample and present antigens to immune cells.

B cells stimulated in germinal centers of Peyer patches differentiate into IgA-secreting plasma cells, which ultimately reside in lamina propria. IgA receives protective secretory component and is then transported across the epithelium to the gut to deal with intraluminal antigen.

Think of **IgA**, the **I**ntra-**g**ut **A**ntibody

Bile

Composed of bile salts (bile acids conjugated to glycine or taurine, making them water soluble), phospholipids, cholesterol, bilirubin, water, and ions. Cholesterol 7α-hydroxylase catalyzes rate-limiting step of bile acid synthesis.

Functions:

- Digestion and absorption of lipids and fat-soluble vitamins
- Cholesterol excretion (body's 1° means of eliminating cholesterol)
- Antimicrobial activity (via membrane disruption)

↓ absorption of enteric bile salts at distal ileum (as in short bowel syndrome, Crohn disease) prevents normal fat absorption

Calcium, which normally binds oxalate, binds fat instead, so free oxalate is absorbed by gut → ↑ frequency of calcium oxalate kidney stones

(c) ketabton.com: The Digital Library

Bilirubin

Heme is metabolized by heme oxygenase to biliverdin, which is subsequently reduced to bilirubin. Unconjugated bilirubin is removed from blood by liver, conjugated with glucuronate, and excreted in bile.

Direct bilirubin: conjugated with glucuronic acid; water soluble (**di**ssolves in water).

Indirect bilirubin: unconjugated; water **in**soluble.



(c) ketabton.com: The Digital Library

▸ GASTROINTESTINAL—PATHOLOGY

Sialolithiasis



Stone(s) in salivary gland duct A. Can occur in 3 major salivary glands (parotid, submandibular, sublingual). Single stone more common in submandibular gland (Wharton duct).
Presents as recurrent pre-/periprandial pain and swelling in affected gland.
Caused by dehydration or trauma.
Treat conservatively with NSAIDs, gland massage, warm compresses, sour candies (to promote salivary flow).

Sialadenitis—inflammation of salivary gland due to obstruction, infection, or immune-mediated mechanisms.

Salivary gland tumors



Most are benign and commonly affect parotid gland (80-85%). Nearly half of all submandibular gland neoplasms and most sublingual and minor salivary gland tumors are malignant. Typically present as painless mass/swelling. Facial paralysis or pain suggests malignant involvement.

- Pleomorphic adenoma (benign mixed tumor)—most common salivary gland tumor A. Composed of chondromyxoid stroma and epithelium and recurs if incompletely excised or ruptured intraoperatively. May undergo malignant transformation.
- Mucoepidermoid carcinoma—most common malignant tumor, has mucinous and squamous components.
- **War**thin tumor (papillary cystadenoma lymphomatosum)—benign cystic tumor with **germ**inal centers. Typically found in **smokers**. Bilateral in 10%; multifocal in 10%. "**War**riors from **Germ**any love **smoking**."

Achalasia



Failure of LES to relax due to degeneration of inhibitory neurons (containing NO and VIP) in the myenteric (Auerbach) plexus of the esophageal wall.
Manometry findings include uncoordinated or absent peristalsis with high LES resting pressure → progressive dysphagia to solids and liquids (vs obstruction—solids only). Barium swallow shows dilated esophagus with an area of distal stenosis ("bird's beak" A).
Associated with ↑ risk of esophageal cancer.

A-*chalasia* = absence of relaxation.
2° achalasia (pseudoachalasia) may arise from Chagas disease (*T cruzi* infection) or extraesophageal malignancies (mass effect or paraneoplastic).
Chagas disease can cause **acha**lasia.

(c) ketabton.com: The Digital Library

Esophageal pathologies

Diffuse esophageal spasm	Spontaneous, nonperistaltic (uncoordinated) contractions of the esophagus with normal LES pressure. Presents with dysphagia and angina-like chest pain. Barium swallow reveals "corkscrew" esophagus. Manometry is diagnostic. Treatment includes nitrates and CCBs.
Eosinophilic esophagitis	Infiltration of eosinophils in the esophagus often in atopic patients. Food allergens → dysphagia, food impaction. Esophageal rings and linear furrows often seen on endoscopy. Typically unresponsive to GERD therapy.
Esophageal perforation	Most commonly iatrogenic following esophageal instrumentation. Noniatrogenic causes include spontaneous rupture, foreign body ingestion, trauma, malignancy. May present with pneumomediastinum (arrows in A). Subcutaneous emphysema may be due to dissecting air (signs include crepitus in the neck region or chest wall). Boerhaave syndrome—transmural, usually distal esophageal rupture due to violent retching.
Esophageal strictures	Associated with caustic ingestion, acid reflux, and esophagitis.
Esophageal varices	Dilated submucosal veins (red arrows in B C) in lower 1/3 of esophagus 2° to portal hypertension. Common in cirrhotics, may be source of life-threatening hematemesis.
Esophagitis	Associated with reflux, infection in immunocompromised (*Candida:* white pseudomembrane D; HSV-1: punched-out ulcers; CMV: linear ulcers), caustic ingestion, or pill-induced esophagitis (eg, bisphosphonates, tetracycline, NSAIDs, iron, and potassium chloride).
Gastroesophageal reflux disease	Commonly presents as heartburn, regurgitation, dysphagia. May also present as chronic cough, hoarseness (laryngopharyngeal reflux). Associated with asthma. Transient decreases in LES tone.
Mallory-Weiss syndrome	Partial thickness, longitudinal lacerations of gastroesophageal junction, confined to mucosa/submucosa, due to severe vomiting. Often presents with hematemesis. Usually found in alcoholics and bulimics.
Plummer-Vinson syndrome	Triad of Dysphagia, Iron deficiency anemia, Esophageal webs. ↑ risk of esophageal Squamous cell carcinoma ("Plumber DIES"). May be associated with glossitis.
Schatzki rings	Rings formed at gastroesophageal junction, typically due to chronic acid reflux. Can present with dysphagia.
Sclerodermal esophageal dysmotility	Esophageal smooth muscle atrophy → ↓ LES pressure and dysmotility → acid reflux and dysphagia → stricture, Barrett esophagus, and aspiration. Part of CREST syndrome.



(c) ketabton.com: The Digital Library

Barrett esophagus





Specialized intestinal metaplasia **A**—replacement of nonkeratinized stratified squamous epithelium with intestinal epithelium (nonciliated columnar with goblet cells [stained blue in **B**]) in distal esophagus. Due to chronic gastroesophageal reflux disease (GERD). Associated with ↑ risk of esophageal adenocarcinoma.



Esophageal cancer

Typically presents with progressive dysphagia (first solids, then liquids) and weight loss. Aggressive course due to lack of serosa in esophageal wall, allowing rapid extension. Poor prognosis due to advanced disease at presentation.

CANCER	PART OF ESOPHAGUS AFFECTED	RISK FACTORS	PREVALENCE
Squamous cell carcinoma	Upper 2/3	Alcohol, hot liquids, caustic strictures, smoking, achalasia	More common worldwide
Adenocarcinoma	Lower 1/3	Chronic GERD, Barrett esophagus, obesity, smoking, achalasia	More common in America

(c) ketabton.com: The Digital Library

Gastritis

Acute gastritis	Erosions can be caused by: ▪ NSAIDs—↓ PGE_2 → ↓ gastric mucosa protection ▪ Burns (Curling ulcer)—hypovolemia → mucosal ischemia ▪ Brain injury (Cushing ulcer)—↑ vagal stimulation → ↑ ACh → ↑ H^+ production	Especially common among alcoholics and patients taking daily NSAIDs (eg, patients with rheumatoid arthritis) **Burn**ed by the **Curling** iron Always **Cush**ion the **brain**
Chronic gastritis	Mucosal inflammation, often leading to atrophy (hypochlorhydria → hypergastrinemia) and intestinal metaplasia (↑ risk of gastric cancers)	
H pylori	Most common. ↑ risk of peptic ulcer disease, MALT lymphoma	Affects antrum first and spreads to body of stomach
Autoimmune	Autoantibodies to the H^+/K^+ ATPase on parietal cells and to intrinsic factor. ↑ risk of pernicious anemia	Affects body/fundus of stomach

Ménétrier disease



Hyperplasia of gastric mucosa → hypertrophied rugae (look like brain gyri A). Causes excess mucus production with resultant protein loss and parietal cell atrophy with ↓ acid production.

Precancerous.

Presents with **W**eight loss, **A**norexia, **V**omiting, **E**pigastric pain, **E**dema (due to protein loss) (**WAVEE**).

Gastric cancer



Most commonly gastric adenocarcinoma; lymphoma, GI stromal tumor, carcinoid (rare). Early aggressive local spread with node/liver metastases. Often presents late, with weight loss, abdominal pain, early satiety, and in some cases acanthosis nigricans or Leser-Trélat sign. Associated with blood type A.

- Intestinal—associated with *H pylori*, dietary nitrosamines (smoked foods), tobacco smoking, achlorhydria, chronic gastritis. Commonly on lesser curvature; looks like ulcer with raised margins.
- Diffuse—not associated with *H pylori*; most cases due to E-cadherin mutation; signet ring cells (mucin-filled cells with peripheral nuclei) A; stomach wall grossly thickened and leathery (linitis plastica).

Virchow node—involvement of left supraclavicular node by metastasis from stomach.

Krukenberg tumor—bilateral metastases to ovaries. Abundant mucin-secreting, signet ring cells.

Sister Mary Joseph nodule—subcutaneous periumbilical metastasis.

Blumer shelf—palpable mass on digital rectal exam suggesting metastasis to rectouterine pouch (pouch of Douglas).

(c) ketabton.com: The Digital Library

Peptic ulcer disease

	Gastric ulcer	Duodenal ulcer
PAIN	Can be Greater with meals—weight loss	Decreases with meals—weight gain
H PYLORI INFECTION	~ 70%	~ 90%
MECHANISM	↓ mucosal protection against gastric acid	↓ mucosal protection or ↑ gastric acid secretion
OTHER CAUSES	NSAIDs	Zollinger-Ellison syndrome
RISK OF CARCINOMA	↑	Generally benign
OTHER	Biopsy margins to rule out malignancy	

Ulcer complications

Hemorrhage

Gastric, duodenal (posterior > anterior). Most common complication.
Ruptured gastric ulcer on the lesser curvature of stomach → bleeding from left gastric artery.
An ulcer on the posterior wall of duodenum → bleeding from gastroduodenal artery.

Obstruction

Pyloric channel, duodenal.

Perforation



Duodenal (anterior > posterior).
Anterior duodenal ulcers can perforate into the anterior abdominal cavity, potentially leading to pneumoperitoneum.
May see free air under diaphragm (pneumoperitoneum) A with referred pain to the shoulder via irritation of phrenic nerve.

(c) ketabton.com: The Digital Library

Malabsorption syndromes	Can cause diarrhea, steatorrhea, weight loss, weakness, vitamin and mineral deficiencies. Screen for fecal fat (eg, Sudan stain).	
Celiac disease 	Gluten-sensitive enteropathy, celiac sprue. Autoimmune-mediated intolerance of gliadin (gluten protein found in wheat) → malabsorption and steatorrhea. Associated with HLA-DQ2, HLA-DQ8, northern European descent, dermatitis herpetiformis, ↓ bone density. Findings: IgA anti-tissue transglutaminase (IgA tTG), anti-endomysial, anti-deamidated gliadin peptide antibodies; villous atrophy, crypt hyperplasia A, and intraepithelial lymphocytosis. Moderately ↑ risk of malignancy (eg, T-cell lymphoma).	↓ mucosal absorption primarily affects distal duodenum and/or proximal jejunum. D-xylose test: passively absorbed in proximal small intestine; blood and urine levels ↓ with mucosa defects or bacterial overgrowth, normal in pancreatic insufficiency. Treatment: gluten-free diet.
Lactose intolerance	Lactase deficiency. Normal-appearing villi, except when 2° to injury at tips of villi (eg, viral enteritis). Osmotic diarrhea with ↓ stool pH (colonic bacteria ferment lactose).	Lactose hydrogen breath test: ⊕ for lactose malabsorption if post-lactose breath hydrogen value rises > 20 ppm compared with baseline.
Pancreatic insufficiency	Due to chronic pancreatitis, cystic fibrosis, obstructing cancer. Causes malabsorption of fat and fat-soluble vitamins (A, D, E, K) as well as vitamin B_{12}.	↓ duodenal bicarbonate (and pH) and fecal elastase.
Tropical sprue	Similar findings as celiac sprue (affects small bowel), but responds to antibiotics. Cause is unknown, but seen in residents of or recent visitors to tropics.	↓ mucosal absorption affecting duodenum and jejunum but can involve ileum with time. Associated with megaloblastic anemia due to folate deficiency and, later, B_{12} deficiency.
Whipple disease 	Infection with *Tropheryma whipplei* (intracellular gram ⊕); PAS ⊕ foamy macrophages in intestinal lamina propria B, mesenteric nodes. Cardiac symptoms, Arthralgias, and Neurologic symptoms are common. Diarrhea/steatorrhea occur later in disease course. Most common in older men.	PAS the foamy Whipped cream in a CAN.

(c) ketabton.com: The Digital Library

Inflammatory bowel diseases

	Crohn disease	Ulcerative colitis
LOCATION	Any portion of the GI tract, usually the terminal ileum and colon. Skip lesions, rectal sparing.	Colitis = colon inflammation. Continuous colonic lesions, always with rectal involvement.
GROSS MORPHOLOGY	Transmural inflammation → fistulas. Cobblestone mucosa, creeping fat, bowel wall thickening ("string sign" on barium swallow x-ray A), linear ulcers, fissures.	Mucosal and submucosal inflammation only. Friable mucosa with superficial and/or deep ulcerations (compare normal B with diseased C). Loss of haustra → "lead pipe" appearance on imaging.
MICROSCOPIC MORPHOLOGY	Noncaseating granulomas and lymphoid aggregates. Th1 mediated.	Crypt abscesses and ulcers, bleeding, no granulomas. Th2 mediated.
COMPLICATIONS	Malabsorption/malnutrition, colorectal cancer (↑ risk with pancolitis).	
	Fistulas (eg, enterovesical fistulae, which can cause recurrent UTI and pneumaturia), phlegmon/abscess, strictures (causing obstruction), perianal disease.	Fulminant colitis, toxic megacolon, perforation.
INTESTINAL MANIFESTATION	Diarrhea that may or may not be bloody.	Bloody diarrhea.
EXTRAINTESTINAL MANIFESTATIONS	Rash (pyoderma gangrenosum, erythema nodosum), eye inflammation (episcleritis, uveitis), oral ulcerations (aphthous stomatitis), arthritis (peripheral, spondylitis).	
	Kidney stones (usually calcium oxalate), gallstones. May be ⊕ for anti-*Saccharomyces cerevisiae* antibodies (ASCA).	1° sclerosing cholangitis. Associated with p-ANCA.
TREATMENT	Corticosteroids, azathioprine, antibiotics (eg, ciprofloxacin, metronidazole), biologics (eg, infliximab, adalimumab).	5-aminosalicylic preparations (eg, mesalamine), 6-mercaptopurine, infliximab, colectomy.
	For Crohn, think of a fat granny and an old crone skipping down a cobblestone road away from the wreck (rectal sparing). Stones are more common in Crohns.	Ulcerative colitis causes ULCCCERS: Ulcers Large intestine Continuous, Colorectal carcinoma, Crypt abscesses Extends proximally Red diarrhea Sclerosing cholangitis



(c) ketabton.com: The Digital Library

Irritable bowel syndrome

Recurrent abdominal pain associated with ≥ 2 of the following:
- Related to defecation
- Change in stool frequency
- Change in form (consistency) of stool

No structural abnormalities. Most common in middle-aged women. Chronic symptoms may be diarrhea-predominant, constipation-predominant, or mixed. Pathophysiology is multifaceted.

First-line treatment is lifestyle modification and dietary changes.

Appendicitis



Acute inflammation of the appendix (yellow arrows in A), can be due to obstruction by fecalith (red arrow in A) (in adults) or lymphoid hyperplasia (in children).

Proximal obstruction of appendiceal lumen produces closed-loop obstruction → ↑ intraluminal pressure → stimulation of visceral afferent nerve fibers at T8-T10 → initial diffuse periumbilical pain → inflammation extends to serosa and irritates parietal peritoneum. Pain localized to RLQ/McBurney point (1/3 the distance from right anterior superior iliac spine to umbilicus). Nausea, fever; may perforate → peritonitis; may elicit psoas, obturator, and Rovsing signs, guarding and rebound tenderness on exam.

Differential: diverticulitis (elderly), ectopic pregnancy (use hCG to rule out), pseudoappendicitis.

Treatment: appendectomy.

Diverticula of the GI tract

Diverticulum	Blind pouch A protruding from the alimentary tract that communicates with the lumen of the gut. Most diverticula (esophagus, stomach, duodenum, colon) are acquired and are termed "false diverticula."	"True" diverticulum—all gut wall layers outpouch (eg, Meckel). "False" diverticulum or pseudodiverticulum—only mucosa and submucosa outpouch. Occur especially where vasa recta perforate muscularis externa.
Diverticulosis	Many false diverticula of the colon B, commonly sigmoid. Common (in ~ 50% of people > 60 years). Caused by ↑ intraluminal pressure and focal weakness in colonic wall. Associated with obesity and diets low in fiber, high in total fat/red meat.	Often asymptomatic or associated with vague discomfort. Complications include diverticular bleeding (painless hematochezia), diverticulitis.
Diverticulitis	Inflammation of diverticula with wall thickening (red arrows in C) classically causing LLQ pain, fever, leukocytosis. Treat with antibiotics.	Complications: abscess, fistula (colovesical fistula → pneumaturia), obstruction (inflammatory stenosis), perforation (white arrows in C) (→ peritonitis).



(c) ketabton.com: The Digital Library

Zenker diverticulum



Pharyngoesophageal **false** diverticulum A.
Esophageal dysmotility causes herniation of mucosal tissue at Killian triangle between the thyropharyngeal and cricopharyngeal parts of the inferior pharyngeal constrictor. Presenting symptoms: dysphagia, obstruction, gurgling, aspiration, foul breath, neck mass. Most common in elderly males.

Elder MIKE has bad breath:
- Elderly
- Males
- Inferior pharyngeal constrictor
- Killian triangle
- Esophageal dysmotility
- Halitosis



Meckel diverticulum



True diverticulum. Persistence of the vitelline (omphalomesenteric) duct. May contain ectopic acid–secreting gastric mucosa and/or pancreatic tissue. Most common congenital anomaly of GI tract. Can cause hematochezia/melena (less common), RLQ pain, intussusception, volvulus, or obstruction near terminal ileum.

Contrast with omphalomesenteric cyst = cystic dilation of vitelline duct.

Diagnosis: ^{99m}Tc-pertechnetate scan (aka Meckel scan) for uptake by heterotopic gastric mucosa.

The rule of 2's:
- 2 times as likely in males.
- 2 inches long.
- 2 feet from the ileocecal valve.
- 2% of population.
- Commonly presents in first 2 years of life.
- May have 2 types of epithelia (gastric/pancreatic).

Hirschsprung disease



Congenital megacolon characterized by lack of ganglion cells/enteric nervous plexuses (Auerbach and Meissner plexuses) in distal segment of colon. Due to failure of neural crest cell migration. Associated with loss of function mutations in *RET*.

Presents with bilious emesis, abdominal distention, and failure to pass meconium within 48 hours → chronic constipation. Normal portion of the colon proximal to the aganglionic segment is dilated, resulting in a "transition zone."

Risk ↑ with Down syndrome.

Explosive expulsion of feces (squirt sign) → empty rectum on digital exam.

Diagnosed by absence of ganglionic cells on rectal suction biopsy.

Treatment: resection.

RET mutation in the REcTum.

(c) ketabton.com: The Digital Library

Malrotation



Anomaly of midgut rotation during fetal development → improper positioning of bowel (small bowel clumped on the right side) A, formation of fibrous bands (Ladd bands).
Can lead to volvulus, duodenal obstruction.



Intussusception





Telescoping A of proximal bowel segment into a distal segment, commonly at the ileocecal junction. Most commonly idiopathic, but may be due to lead point.
Compromised blood supply → intermittent, severe, abdominal pain often with "currant jelly" dark red stools.
Majority of cases in infants, unusual in adults.
Most common pathologic lead point:
- Children—Meckel diverticulum
- Adults—intraluminal mass/tumor

On physical exam, patient may draw their legs to chest to ease pain, sausage shaped mass on palpation.
Imaging—Ultrasound/CT may show "target sign." B

May be associated with IgA vasculitis (HSP), recent viral infection (eg, adenovirus; Peyer patch hypertrophy creates lead point).

(c) ketabton.com: The Digital Library

Volvulus



Twisting of portion of bowel around its mesentery; can lead to obstruction and infarction. Can occur throughout the GI tract.

- **M**idgut volvulus more common in infants and children (**m**inors)
- **S**igmoid volvulus (coffee bean sign on x-ray A) more common in **s**eniors (elderly)





Other intestinal disorders

Acute mesenteric ischemia	Critical blockage of intestinal blood flow (often embolic occlusion of SMA) → small bowel necrosis A → abdominal pain out of proportion to physical findings. May see red "currant jelly" stools.
Adhesion	Fibrous band of scar tissue; commonly forms after surgery. Most common cause of small bowel obstruction, demonstrated by multiple dilated small bowel loops on x-ray (arrows in B).
Angiodysplasia	Tortuous dilation of vessels C → hematochezia. Most often found in the right-sided colon. More common in older patients. Confirmed by angiography. Associated with end-stage renal disease, von Willebrand disease, aortic stenosis.
Chronic mesenteric ischemia	"Intestinal angina": atherosclerosis of celiac artery, SMA, or IMA → intestinal hypoperfusion → postprandial epigastric pain → food aversion and weight loss.
Colonic ischemia	Reduction in intestinal blood flow causes ischemia. Crampy abdominal pain followed by hematochezia. Commonly occurs at watershed areas (splenic flexure, rectosigmoid junction). Typically affects elderly. Thumbprint sign on imaging due to mucosal edema/hemorrhage.
Ileus	Intestinal hypomotility without obstruction → constipation and ↓ flatus; distended/tympanic abdomen with ↓ bowel sounds. Associated with abdominal surgeries, opiates, hypokalemia, sepsis. Treatment: bowel rest, electrolyte correction, cholinergic drugs (stimulate intestinal motility).
Meconium ileus	Meconium plug obstructs intestine, prevents stool passage at birth. Associated with cystic fibrosis.
Necrotizing enterocolitis	Seen in premature, formula-fed infants with immature immune system. Necrosis of intestinal mucosa (most commonly terminal ileum and proximal colon) with possible perforation, which can lead to pneumatosis intestinalis (arrows in D), pneumoperitoneum, portal venous gas.



(c) ketabton.com: The Digital Library

Colonic polyps — Growths of tissue within the colon A. Grossly characterized as flat, sessile, or pedunculated on the basis of protrusion into colonic lumen. Generally classified by histologic type.

HISTOLOGIC TYPE	CHARACTERISTICS
Generally non-neoplastic	
Hamartomatous polyps	Solitary lesions do not have significant risk of transformation. Growths of normal colonic tissue with distorted architecture. Associated with Peutz-Jeghers syndrome and juvenile polyposis.
Hyperplastic polyps	Most common; generally smaller and predominantly located in rectosigmoid region. Occasionally evolves into serrated polyps and more advanced lesions.
Inflammatory pseudopolyps	Due to mucosal erosion in inflammatory bowel disease.
Mucosal polyps	Small, usually < 5 mm. Look similar to normal mucosa. Clinically insignificant.
Submucosal polyps	May include lipomas, leiomyomas, fibromas, and other lesions.
Malignant potential	
Adenomatous polyps	Neoplastic, via chromosomal instability pathway with mutations in *APC* and *KRAS*. Tubular B histology has less malignant potential than villous C ("**villous** histology is **vill**ain**ous**"); tubulovillous has intermediate malignant potential. Usually asymptomatic; may present with occult bleeding.
Serrated polyps	Neoplastic. Characterized by CpG island methylator phenotype (CIMP; cytosine base followed by guanine, linked by a phosphodiester bond). Defect may silence *MMR* gene (DNA mismatch repair) expression. Mutations lead to microsatellite instability and mutations in *BRAF*. "Saw-tooth" pattern of crypts on biopsy. Up to 20% of cases of sporadic CRC.







Polyposis syndromes

Familial adenomatous polyposis	Autosomal dominant mutation of *APC* tumor suppressor gene on chromosome 5q22. 2-hit hypothesis. Thousands of polyps arise starting after puberty; pancolonic; always involves rectum. Prophylactic colectomy or else 100% progress to CRC.
Gardner syndrome	FAP + osseous and soft tissue tumors (eg, osteomas of skull or mandible), congenital hypertrophy of retinal pigment epithelium, impacted/supernumerary teeth.
Turcot syndrome	FAP or Lynch syndrome + malignant CNS tumor (eg, medulloblastoma, glioma). **Tur**cot = **Tur**ban.
Peutz-Jeghers syndrome	Autosomal dominant syndrome featuring numerous hamartomas throughout GI tract, along with hyperpigmented macules on mouth, lips, hands, genitalia. Associated with ↑ risk of breast and GI cancers (eg, colorectal, stomach, small bowel, pancreatic).
Juvenile polyposis syndrome	Autosomal dominant syndrome in children (typically < 5 years old) featuring numerous hamartomatous polyps in the colon, stomach, small bowel. Associated with ↑ risk of CRC.

(c) ketabton.com: The Digital Library

Lynch syndrome

Previously called hereditary nonpolyposis colorectal cancer (HNPCC). Autosomal dominant mutation of mismatch repair genes (eg, *MLH1*, *MSH2*) with subsequent microsatellite instability. ~ 80% progress to CRC. Proximal colon is always involved. Associated with endometrial, ovarian, and skin cancers.

Colorectal cancer

DIAGNOSIS





Iron deficiency anemia in males (especially > 50 years old) and postmenopausal females raises suspicion.

Screening:

- Low risk: screen at age 50 with colonoscopy (polyp seen in A); alternatives include flexible sigmoidoscopy, fecal occult blood testing (FOBT), fecal immunochemical testing (FIT), FIT-fecal DNA, CT colonography
- Patients with a first-degree relative who has colon cancer: screen at age 40 with colonoscopy, or 10 years prior to the relative's presentation
- Patients with IBD: distinct screening protocol

"Apple core" lesion seen on barium enema x-ray B.

CEA tumor marker: good for monitoring recurrence, should not be used for screening.

EPIDEMIOLOGY

Most patients are > 50 years old. ~ 25% have a family history.

PRESENTATION

Rectosigmoid > ascending > descending.

Right side (cecal, ascending) associated with occult bleeding; left side (rectosigmoid) associated with hematochezia and obstruction (narrower lumen).

Ascending—exophytic mass, iron deficiency anemia, weight loss.

Descending—infiltrating mass, partial obstruction, colicky pain, hematochezia.

Can present with *S bovis* (*gallolyticus*) bacteremia/endocarditis or as an episode of diverticulitis.

RISK FACTORS

Adenomatous and serrated polyps, familial cancer syndromes, IBD, tobacco use, diet of processed meat with low fiber.

(c) ketabton.com: The Digital Library

Molecular pathogenesis of colorectal cancer

Chromosomal instability pathway: mutations in *APC* cause FAP and most sporadic cases of CRC via adenoma-carcinoma sequence; (**firing** order of events is "**AK-53**").

Microsatellite instability pathway: mutations or methylation of mismatch repair genes (eg, *MLH1*) cause Lynch syndrome and some sporadic CRC (via serrated polyp pathway).

Overexpression of COX-2 has been linked to colorectal cancer, NSAIDs may be chemopreventive.



Cirrhosis and portal hypertension





Cirrhosis—diffuse bridging fibrosis (via stellate cells) and regenerative nodules (red arrows in A; white arrows show splenomegaly) disrupt normal architecture of liver; ↑ risk for hepatocellular carcinoma (white arrow in B). Etiologies include alcohol, nonalcoholic steatohepatitis, chronic viral hepatitis, autoimmune hepatitis, biliary disease, genetic/metabolic disorders.

Portal hypertension—↑ pressure in portal venous system. Etiologies include cirrhosis (most common cause in Western countries), vascular obstruction (eg, portal vein thrombosis, Budd-Chiari syndrome), schistosomiasis.



(c) ketabton.com: The Digital Library

Spontaneous bacterial peritonitis

Also called 1° bacterial peritonitis. Common and potentially fatal bacterial infection in patients with cirrhosis and ascites. Often asymptomatic, but can cause fevers, chills, abdominal pain, ileus, or worsening encephalopathy. Commonly caused by gram ⊖ organisms (eg, *E coli*, *Klebsiella*) or less commonly gram ⊕ *Streptococcus*.

Diagnosis: paracentesis with ascitic fluid absolute neutrophil count (ANC) > 250 cells/mm^3. Empiric first-line treatment is 3rd generation cephalosporin (eg, cefotaxime).

Serum markers of liver pathology

ENZYMES RELEASED IN LIVER DAMAGE	
Aspartate aminotransferase and alanine aminotransferase	↑ in most liver disease: ALT > AST ↑ in alcoholic liver disease: AST > ALT (AST usually will not exceed 500 U/L in alcoholic hepatitis) AST > ALT in nonalcoholic liver disease suggests progression to advanced fibrosis or cirrhosis ↑↑↑ aminotransferases (>1000 U/L): differential includes drug-induced liver injury (eg, acetaminophen toxicity), ischemic hepatitis, acute viral hepatitis, autoimmune hepatitis
Alkaline phosphatase	↑ in cholestasis (eg, biliary obstruction), infiltrative disorders, bone disease
γ-glutamyl transpeptidase	↑ in various liver and biliary diseases (just as ALP can), but not in bone disease; associated with alcohol use
FUNCTIONAL LIVER MARKERS	
Bilirubin	↑ in various liver diseases (eg, biliary obstruction, alcoholic or viral hepatitis, cirrhosis), hemolysis
Albumin	↓ in advanced liver disease (marker of liver's biosynthetic function)
Prothrombin time	↑ in advanced liver disease (↓ production of clotting factors, thereby measuring the liver's biosynthetic function)
Platelets	↓ in advanced liver disease (↓ thrombopoietin, liver sequestration) and portal hypertension (splenomegaly/splenic sequestration)

Reye syndrome

Rare, often fatal childhood hepatic encephalopathy.

Associated with viral infection (especially VZV and influenza) that has been treated with aspirin. Aspirin metabolites ↓ β-oxidation by reversible inhibition of mitochondrial enzymes.

Findings: mitochondrial abnormalities, fatty liver (microvesicular fatty changes), hypoglycemia, vomiting, hepatomegaly, coma.

Avoid aspirin in children, except in those with Kawasaki disease.

Salicylates aren't a ray (**Reye**) of sun**SHINE** for kids:

Steatosis of liver/hepatocytes
Hypoglycemia/**H**epatomegaly
Infection (VZV, influenza)
Not awake (coma)
Encephalopathy

(c) ketabton.com: The Digital Library

Alcoholic liver disease

Hepatic steatosis	Macrovesicular fatty change A that may be reversible with alcohol cessation.	
Alcoholic hepatitis	Requires sustained, long-term consumption. Swollen and necrotic hepatocytes with neutrophilic infiltration. Mallory bodies B (intracytoplasmic eosinophilic inclusions of damaged keratin filaments).	Make a toAST with alcohol: AST > ALT (ratio usually > 2:1).
Alcoholic cirrhosis	Final and usually irreversible form. Sclerosis around central vein (arrows in C) may be seen in early disease. Regenerative nodules surrounded by fibrous bands in response to chronic liver injury → portal hypertension and end-stage liver disease.	



Nonalcoholic fatty liver disease



Metabolic syndrome (insulin resistance); obesity → fatty infiltration of hepatocytes A → cellular "ballooning" and eventual necrosis. May cause cirrhosis and HCC. Independent of alcohol use.

ALT > AST (Lipids)

Hepatic encephalopathy

Cirrhosis → portosystemic shunts → ↓ NH_3 metabolism → neuropsychiatric dysfunction. Reversible neuropsychiatric dysfunction ranging from disorientation/asterixis (mild) to difficult arousal or coma (severe).

Triggers:

- ↑ NH_3 production and absorption (due to GI bleed, constipation, infection).
- ↓ NH_3 removal (due to renal failure, diuretics, bypassed hepatic blood flow post-TIPS).

Treatment: lactulose (↑ NH_4^+ generation) and rifaximin (↓ NH_3-producing gut bacteria).

FAS1_2019_09-Gastrointestinal.indd 391 11/7/19 4:42 PM

(c) ketabton.com: The Digital Library

Hepatocellular carcinoma/hepatoma

Most common 1° malignant tumor of liver in adults A. Associated with HBV (+/– cirrhosis) and all other causes of cirrhosis (including HCV, alcoholic and nonalcoholic fatty liver disease, autoimmune disease, hemochromatosis, Wilson disease, α_1-antitrypsin deficiency) and specific carcinogens (eg, aflatoxin from *Aspergillus*). May lead to Budd-Chiari syndrome.

Findings: jaundice, tender hepatomegaly, ascites, polycythemia, anorexia. Spreads hematogenously.

Diagnosis: ↑ α-fetoprotein; ultrasound or contrast CT/MRI B, biopsy.



Other liver tumors

Angiosarcoma

Malignant tumor of endothelial origin; associated with exposure to arsenic, vinyl chloride.

Cavernous hemangioma

Most common benign liver tumor (venous malformation) A; typically occurs at age 30–50 years. Biopsy contraindicated because of risk of hemorrhage.



Hepatic adenoma

Rare, benign liver tumor, often related to oral contraceptive or anabolic steroid use; may regress spontaneously or rupture (abdominal pain and shock).

Metastases

GI malignancies, breast and lung cancer. Most common overall; metastases are rarely solitary.

Budd-Chiari syndrome

Thrombosis or compression of hepatic veins with centrilobular congestion and necrosis → congestive liver disease (hepatomegaly, ascites, varices, abdominal pain, liver failure). Absence of JVD. Associated with hypercoagulable states, polycythemia vera, postpartum state, HCC. May cause nutmeg liver (mottled appearance).

α_1-antitrypsin deficiency



Misfolded gene product protein aggregates in hepatocellular ER → cirrhosis with PAS ⊕ globules A in liver. Codominant trait.

Often presents in young patients with liver damage and dyspnea without a history of smoking.

In lungs, ↓ α_1-antitrypsin → uninhibited elastase in alveoli → ↓ elastic tissue → panacinar emphysema.

(c) ketabton.com: The Digital Library

Jaundice

Abnormal yellowing of the skin and/or sclera A due to bilirubin deposition. Hyperbilirubinemia 2° to ↑ production or ↓ clearance (impaired hepatic uptake, conjugation, excretion).

HOT Liver—common causes of ↑ bilirubin level:
Hemolysis
Obstruction
Tumor
Liver disease

Conjugated (direct) hyperbilirubinemia	Biliary tract obstruction: gallstones, cholangiocarcinoma, pancreatic or liver cancer, liver fluke. Biliary tract disease: ▪ 1° sclerosing cholangitis ▪ 1° biliary cholangitis Excretion defect: Dubin-Johnson syndrome, Rotor syndrome.
Unconjugated (indirect) hyperbilirubinemia	Hemolytic, physiologic (newborns), Crigler-Najjar, Gilbert syndrome.
Mixed (direct and indirect) hyperbilirubinemia	Hepatitis, cirrhosis.

Physiologic neonatal jaundice

At birth, immature UDP-glucuronosyltransferase → unconjugated hyperbilirubinemia → jaundice/kernicterus (deposition of unconjugated, lipid-soluble bilirubin in the brain, particularly basal ganglia).
Occurs after first 24 hours of life and usually resolves without treatment in 1–2 weeks.
Treatment: phototherapy (non-UV) isomerizes unconjugated bilirubin to water-soluble form.

Biliary atresia

Most common reason for pediatric liver transplantation.
Fibro-obliterative destruction of extrahepatic bile ducts → cholestasis.
Often presents as a newborn with persistent jaundice after 2 weeks of life, darkening urine, acholic stools, hepatomegaly.
Labs: ↑ direct bilirubin and GGT.

(c) ketabton.com: The Digital Library

Hereditary hyperbilirubinemias	All autosomal recessive.
❶ **Gilbert syndrome**	Mildly ↓ UDP-glucuronosyltransferase conjugation and impaired bilirubin uptake. Asymptomatic or mild jaundice usually with stress, illness, or fasting. ↑ unconjugated bilirubin without overt hemolysis. Relatively common, benign condition.
❷ **Crigler-Najjar syndrome, type I**	Absent UDP-glucuronosyltransferase. Presents early in life, but some patients may not have neurologic signs until later in life. Findings: jaundice, kernicterus (bilirubin deposition in brain), ↑ unconjugated bilirubin. Treatment: plasmapheresis and phototherapy (does not conjugate UCB; but does ↑ polarity and ↑ water solubility to allow excretion). Liver transplant is curative. Type II is less severe and responds to phenobarbital, which ↑ liver enzyme synthesis.
❸ **Dubin-Johnson syndrome**	Conjugated hyperbilirubinemia due to defective liver excretion. Grossly black (Dark) liver due to impaired excretion of epinephrine metabolites. Benign.
❹ **Rotor syndrome**	Similar to Dubin-Johnson syndrome, but milder in presentation without black (Regular) liver. Due to impaired hepatic uptake and excretion.



(c) ketabton.com: The Digital Library

Wilson disease



Also called hepatolenticular degeneration. Autosomal recessive mutations in hepatocyte copper-transporting ATPase (*ATP7B* gene; chromosome 13) → ↓ copper incorporation into apoceruloplasmin and excretion into bile → ↓ serum ceruloplasmin. Copper accumulates, especially in liver, brain, cornea, kidneys; ↑ urine copper.

Presents before age 40 with liver disease (eg, hepatitis, acute liver failure, cirrhosis), neurologic disease (eg, dysarthria, dystonia, tremor, parkinsonism), psychiatric disease, Kayser-Fleischer rings (deposits in Descemet membrane of cornea) A, hemolytic anemia, renal disease (eg, Fanconi syndrome).

Treatment: chelation with penicillamine or trientine, oral zinc. Liver transplant in acute liver failure related to Wilson disease.

Hemochromatosis



Autosomal recessive. On *HFE* gene, located on chromosome 6; associated with HLA-A3. Leads to abnormal **iron** sensing and ↑ intestinal absorption (↑ ferritin, ↑ iron, ↓ TIBC → ↑ transferrin saturation). Iron overload can also be 2° to chronic transfusion therapy (eg, β-thalassemia major). Iron accumulates, especially in liver, pancreas, skin, heart, pituitary, joints. Hemosiderin (iron) can be identified on liver MRI or biopsy with Prussian blue stain A.

Presents after age 40 when total body iron > 20 g; iron loss through menstruation slows progression in women. Classic triad of cirrhosis, diabetes mellitus, skin pigmentation ("bronze diabetes"). Also causes restrictive cardiomyopathy (classic) or dilated cardiomyopathy (reversible), hypogonadism, arthropathy (calcium pyrophosphate deposition; especially metacarpophalangeal joints). HCC is common cause of death.

Treatment: repeated phlebotomy, iron (**Fe**) chelation with de**fe**rasirox, de**fe**roxamine, de**fe**riprone.

Biliary tract disease

May present with pruritus, jaundice, dark urine, light-colored stool, hepatosplenomegaly. Typically with cholestatic pattern of LFTs (↑ conjugated bilirubin, ↑ cholesterol, ↑ ALP, ↑ GGT).

	PATHOLOGY	EPIDEMIOLOGY	ADDITIONAL FEATURES
Primary sclerosing cholangitis	Unknown cause of concentric "onion skin" bile duct fibrosis → alternating strictures and dilation with "beading" of intra- and extrahepatic bile ducts on ERCP, magnetic resonance cholangiopancreatography (MRCP).	Classically in middle-aged men with ulcerative colitis.	Associated with ulcerative colitis. p-ANCA ⊕. ↑ IgM. Can lead to 2° biliary cholangitis. ↑ risk of cholangiocarcinoma and gallbladder cancer.
Primary biliary cholangitis	Autoimmune reaction → lymphocytic infiltrate +/– granulomas → destruction of lobular bile ducts.	Classically in middle-aged women.	Anti-mitochondrial antibody ⊕, ↑ IgM. Associated with other autoimmune conditions (eg, Hashimoto thyroiditis, rheumatoid arthritis, celiac disease). Treatment: ursodiol.
Secondary biliary cirrhosis	Extrahepatic biliary obstruction → ↑ pressure in intrahepatic ducts → injury/ fibrosis and bile stasis.	Patients with known obstructive lesions (gallstones, biliary strictures, pancreatic carcinoma).	May be complicated by ascending cholangitis.

(c) ketabton.com: The Digital Library

Cholelithiasis and related pathologies



↑ cholesterol and/or bilirubin, ↓ bile salts, and gallbladder stasis all cause stones.

2 types of stones:

- Cholesterol stones (radiolucent with 10–20% opaque due to calcifications)—80% of stones. Associated with obesity, Crohn disease, advanced age, estrogen therapy, multiparity, rapid weight loss, Native American origin.
- Pigment stones A (black = radiopaque, Ca^{2+} bilirubinate, hemolysis; brown = radiolucent, infection). Associated with Crohn disease, chronic hemolysis, alcoholic cirrhosis, advanced age, biliary infections, total parenteral nutrition (TPN).

Risk factors (4 F's):

1. Female
2. Fat
3. Fertile (multiparity)
4. Forty

Most common complication is cholecystitis; can also cause acute pancreatitis, ascending cholangitis.

Diagnose with ultrasound. Treat with elective cholecystectomy if symptomatic.



RELATED PATHOLOGIES	CHARACTERISTICS
Biliary colic	Associated with nausea/vomiting and dull RUQ pain. Neurohormonal activation (eg, by CCK after a fatty meal) triggers contraction of gallbladder, forcing stone into cystic duct. Labs are normal, ultrasound shows cholelithiasis.
Choledocholithiasis	Presence of gallstone(s) in common bile duct, often leading to elevated ALP, GGT, direct bilirubin, and/or AST/ALT.
Cholecystitis 	Acute or chronic inflammation of gallbladder. Calculous cholecystitis—most common type; due to gallstone impaction in the cystic duct resulting in inflammation and gallbladder wall thickening (arrows in B); can produce 2° infection. Acalculous cholecystitis—due to gallbladder stasis, hypoperfusion, or infection (CMV); seen in critically ill patients. Murphy sign: inspiratory arrest on RUQ palpation due to pain. Pain may radiate to right shoulder (due to irritation of phrenic nerve). ↑ ALP if bile duct becomes involved (eg, ascending cholangitis). Diagnose with ultrasound or cholescintigraphy (HIDA scan). Failure to visualize gallbladder on HIDA scan suggests obstruction. Gallstone ileus—fistula between gallbladder and GI tract → stone enters GI lumen → obstructs at ileocecal valve (narrowest point); can see air in biliary tree (pneumobilia). Rigler triad: radiographic findings of pneumobilia, small bowel obstruction, gallstone (usually in iliac fossa).

(c) ketabton.com: The Digital Library

Cholelithiasis and related pathologies *(continued)*

Porcelain gallbladder



Calcified gallbladder due to chronic cholecystitis; usually found incidentally on imaging C.
Treatment: prophylactic cholecystectomy generally recommended due to ↑ risk of gallbladder cancer (mostly adenocarcinoma).

Ascending cholangitis

Infection of biliary tree usually due to obstruction that leads to stasis/bacterial overgrowth.
Charcot triad of cholangitis includes jaundice, fever, RUQ pain.
Reynolds pentad is Charcot triad plus altered mental status and shock (hypotension).

Acute pancreatitis





Autodigestion of pancreas by pancreatic enzymes (A shows pancreas [yellow arrows] surrounded by edema [red arrows]).
Causes: **I**diopathic, **G**allstones, **E**thanol, **T**rauma, **S**teroids, **M**umps, **A**utoimmune disease, **S**corpion sting, **H**ypercalcemia/**H**ypertriglyceridemia (> 1000 mg/dL), **E**RCP, **D**rugs (eg, sulfa drugs, NRTIs, protease inhibitors). **I GET SMASHED**.
Diagnosis by 2 of 3 criteria: acute epigastric pain often radiating to the back, ↑ serum amylase or lipase (more specific) to 3× upper limit of normal, or characteristic imaging findings.
Complications: pseudocyst B (lined by granulation tissue, not epithelium), abscess, necrosis, hemorrhage, infection, organ failure (ALI/ARDS, shock, renal failure), hypocalcemia (precipitation of Ca^{2+} soaps).

Chronic pancreatitis



Chronic inflammation, atrophy, calcification of the pancreas A. Major causes include alcohol abuse and genetic predisposition (ie, cystic fibrosis); can be idiopathic. Complications include pancreatic insufficiency and pseudocysts.
Pancreatic insufficiency (typically when <10% pancreatic function) may manifest with steatorrhea, fat-soluble vitamin deficiency, diabetes mellitus.
Amylase and lipase may or may not be elevated (almost always elevated in acute pancreatitis).

(c) ketabton.com: The Digital Library

Pancreatic adenocarcinoma





Very aggressive tumor arising from pancreatic ducts (disorganized glandular structure with cellular infiltration A); often metastatic at presentation, with average survival ~ 1 year after diagnosis. Tumors more common in pancreatic head B (lead to obstructive jaundice). Associated with CA 19-9 tumor marker (also CEA, less specific).

Risk factors:

- Tobacco use
- Chronic pancreatitis (especially > 20 years)
- Diabetes
- Age > 50 years
- Jewish and African-American males

Often presents with:

- Abdominal pain radiating to back
- Weight loss (due to malabsorption and anorexia)
- Migratory thrombophlebitis—redness and tenderness on palpation of extremities (Trousseau syndrome)
- Obstructive jaundice with palpable, nontender gallbladder (Courvoisier sign)

Treatment: Whipple procedure (pancreaticoduodenectomy), chemotherapy, radiation therapy.

▸ GASTROINTESTINAL—PHARMACOLOGY

Acid suppression therapy



(c) ketabton.com: The Digital Library

Histamine-2 blockers	Cimetidine, ranitidine, famotidine, nizatidine.	Take H_2 blockers before you dine. Think "table for 2" to remember H_2.
MECHANISM	Reversible block of histamine H_2-receptors → ↓ H^+ secretion by parietal cells.	
CLINICAL USE	Peptic ulcer, gastritis, mild esophageal reflux.	
ADVERSE EFFECTS	Cimetidine is a potent inhibitor of cytochrome P-450 (multiple drug interactions); it also has antiandrogenic effects (prolactin release, gynecomastia, impotence, ↓ libido in males); can cross blood-brain barrier (confusion, dizziness, headaches) and placenta. Both cimetidine and ranitidine ↓ renal excretion of creatinine. Other H_2 blockers are relatively free of these effects.	

Proton pump inhibitors	Omeprazole, lansoprazole, esomeprazole, pantoprazole, dexlansoprazole.
MECHANISM	Irreversibly inhibit H^+/K^+ ATPase in stomach parietal cells.
CLINICAL USE	Peptic ulcer, gastritis, esophageal reflux, Zollinger-Ellison syndrome, component of therapy for *H pylori*, stress ulcer prophylaxis.
ADVERSE EFFECTS	↑ risk of *C difficile* infection, pneumonia, acute interstitial nephritis. Vitamin B_{12} malabsorption; ↓ serum Mg^{2+} and ↓ Ca^{2+} absorption (potentially leading to increased fracture risk in elderly).

Antacids	Can affect absorption, bioavailability, or urinary excretion of other drugs by altering gastric and urinary pH or by delaying gastric emptying. All can cause hypokalemia. Overuse can also cause the following problems:	
Aluminum hydroxide	Constipation, Hypophosphatemia, Osteodystrophy, Proximal muscle weakness, Seizures	Aluminimum amount of feces CHOPS
Calcium carbonate	Hypercalcemia (milk-alkali syndrome), rebound acid ↑	Can chelate and ↓ effectiveness of other drugs (eg, tetracycline)
Magnesium hydroxide	Diarrhea, hyporeflexia, hypotension, cardiac arrest	Mg^{2+} = Must go 2 the bathroom

Bismuth, sucralfate	
MECHANISM	Bind to ulcer base, providing physical protection and allowing HCO_3^- secretion to reestablish pH gradient in the mucous layer. Sucralfate requires acidic environment, not given with PPIs/H_2 blockers.
CLINICAL USE	↑ ulcer healing, travelers' diarrhea (bismuth). Bismuth also used in quadruple therapy for *H pylori* gastritis.

Misoprostol	
MECHANISM	PGE_1 analog. ↑ production and secretion of gastric mucous barrier, ↓ acid production.
CLINICAL USE	Prevention of NSAID-induced peptic ulcers (NSAIDs block PGE_1 production). Also used off-label for induction of labor (ripens cervix).
ADVERSE EFFECTS	Diarrhea. Contraindicated in women of childbearing potential (abortifacient).

(c) ketabton.com: The Digital Library

Octreotide

MECHANISM	Long-acting somatostatin analog; inhibits secretion of various splanchnic vasodilatory hormones.
CLINICAL USE	Acute variceal bleeds, acromegaly, VIPoma, carcinoid tumors.
ADVERSE EFFECTS	Nausea, cramps, steatorrhea. ↑ risk of cholelithiasis due to CCK inhibition.

Sulfasalazine

MECHANISM	A combination of sulfapyridine (antibacterial) and 5-aminosalicylic acid (anti-inflammatory). Activated by colonic bacteria.
CLINICAL USE	Ulcerative colitis, Crohn disease (colitis component).
ADVERSE EFFECTS	Malaise, nausea, sulfonamide toxicity, reversible oligospermia.

Loperamide

MECHANISM	Agonist at μ-opioid receptors; slows gut motility. Poor CNS penetration (low addictive potential).
CLINICAL USE	Diarrhea.
ADVERSE EFFECTS	Constipation, nausea.

Ondansetron

MECHANISM	5-HT_3 antagonist; ↓ vagal stimulation. Powerful central-acting antiemetic.
CLINICAL USE	Control vomiting postoperatively and in patients undergoing cancer chemotherapy.
ADVERSE EFFECTS	Headache, constipation, QT interval prolongation, serotonin syndrome.

Metoclopramide

MECHANISM	D_2 receptor antagonist. ↑ resting tone, contractility, LES tone, motility, promotes gastric emptying. Does not influence colon transport time.
CLINICAL USE	Diabetic and postoperative gastroparesis, antiemetic, persistent GERD.
ADVERSE EFFECTS	↑ parkinsonian effects, tardive dyskinesia. Restlessness, drowsiness, fatigue, depression, diarrhea. Drug interaction with digoxin and diabetic agents. Contraindicated in patients with small bowel obstruction, Parkinson disease (due to D_2-receptor blockade), ↓ seizure threshold.

Orlistat

MECHANISM	Inhibits gastric and pancreatic lipase → ↓ breakdown and absorption of dietary fats. Taken with fat-containing meals.
CLINICAL USE	Weight loss.
ADVERSE EFFECTS	Abdominal pain, flatulence, bowel urgency/frequent bowel movements, steatorrhea; ↓ absorption of fat-soluble vitamins.

(c) ketabton.com: The Digital Library

Laxatives

Indicated for constipation or patients on opiates requiring a bowel regimen.

	EXAMPLES	MECHANISM	ADVERSE EFFECTS
Bulk-forming laxatives	Psyllium, methylcellulose	Soluble fibers draw water into gut lumen, forming a viscous liquid that promotes peristalsis	Bloating
Osmotic laxatives	Magnesium hydroxide, magnesium citrate, polyethylene glycol, lactulose	Provides osmotic load to draw water into GI lumen Lactulose also treats hepatic encephalopathy: gut flora degrade lactulose into metabolites (lactic acid, acetic acid) that promote nitrogen excretion as NH_4^+	Diarrhea, dehydration; may be abused by bulimics
Stimulants	Senna	Enteric nerve stimulation → colonic contraction	Diarrhea, melanosis coli
Emollients	Docusate	Promotes incorporation of water and fat into stool	Diarrhea

Aprepitant

MECHANISM	Substance P antagonist. Blocks NK_1 (neurokinin-1) receptors in brain.
CLINICAL USE	Antiemetic for chemotherapy-induced nausea and vomiting.

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

Hematology and Oncology

"You're always somebody's type! (blood type, that is)"

—BloodLink

"All the soarings of my mind begin in my blood."

—Rainer Maria Rilke

"The best blood will at some time get into a fool or a mosquito."

—Austin O'Malley

When studying hematology, pay close attention to the many cross connections to immunology. Make sure you master the different types of anemias. Be comfortable interpreting blood smears. When reviewing oncologic drugs, focus on mechanisms and adverse effects rather than details of clinical uses, which may be lower yield.

Please note that solid tumors are covered in their respective organ system chapters.

- Embryology 404
- Anatomy 406
- Physiology 410
- Pathology 414
- Pharmacology 435

FAS1_2019_10-HemaOncol.indd 403 11/7/19 5:04 PM

(c) ketabton.com: The Digital Library

▸ HEMATOLOGY AND ONCOLOGY—EMBRYOLOGY

Fetal erythropoiesis

Fetal erythropoiesis occurs in:
- Yolk sac (3–8 weeks)
- Liver (6 weeks–birth)
- Spleen (10–28 weeks)
- Bone marrow (18 weeks to adult)

Young Liver Synthesizes Blood.

Hemoglobin development

Embryonic globins: ζ and ε.
Fetal hemoglobin (HbF) = $\alpha_2\gamma_2$.
Adult hemoglobin (HbA_1) = $\alpha_2\beta_2$.
HbF has higher affinity for O_2 due to less avid binding of 2,3-BPG, allowing HbF to extract O_2 from maternal hemoglobin (HbA_1 and HbA_2) across the placenta. HbA_2 ($\alpha_2\delta_2$) is a form of adult hemoglobin present in small amounts.

From fetal to adult hemoglobin:
Alpha Always; Gamma Goes, Becomes Beta.



(c) ketabton.com: The Digital Library

Blood groups

	ABO classification				Rh classification	
	A	B	AB	O	Rh⊕	Rh⊖
RBC type						
Group antigens on RBC surface	A	B	A & B	None	Rh (D)	None
Antibodies in plasma	Anti-B IgM	Anti-A IgM	None	Anti-A Anti-B IgM, IgG	None	Anti-D IgG
Clinical relevance	Receive B or AB → hemolytic reaction	Receive A or AB → hemolytic reaction	Universal recipient of RBCs; universal donor of plasma	Receive any non-O → hemolytic reaction Universal donor of RBCs; universal recipient of plasma	Can receive either Rh⊕ or Rh⊖ blood	Treat mother with anti-D IgG during and after each pregnancy to prevent anti-D IgG formation

Hemolytic disease of the newborn

Also known as erythroblastosis fetalis.

	Rh hemolytic disease of the newborn	ABO hemolytic disease of the newborn
INTERACTION	Rh ⊖ mother; Rh ⊕ fetus.	Type O mother; type A or B fetus.
MECHANISM	First pregnancy: mother exposed to fetal blood (often during delivery) → formation of maternal anti-D IgG. Subsequent pregnancies: anti-D IgG crosses the placenta → attacks fetal RBCs → hemolysis in the fetus.	Pre-existing maternal anti-A and/or anti-B IgG antibodies cross placenta → hemolysis in the fetus.
PRESENTATION	Hydrops fetalis, jaundice shortly after birth, kernicterus.	Mild jaundice in the neonate within 24 hours of birth. Unlike Rh HDN, can occur in firstborn babies and is usually less severe.
TREATMENT/PREVENTION	Prevent by administration of anti-D IgG to Rh ⊖ pregnant women during third trimester and early postpartum period (if fetus Rh ⊕). Prevents maternal anti-D IgG production.	Treatment: phototherapy or exchange transfusion.

(c) ketabton.com: The Digital Library

▸ HEMATOLOGY AND ONCOLOGY—ANATOMY

Hematopoiesis



Neutrophils



Acute inflammatory response cells. Numbers ↑ in bacterial infections. Phagocytic. Multilobed nucleus A. Specific granules contain leukocyte alkaline phosphatase (LAP), collagenase, lysozyme, and lactoferrin. Azurophilic granules (lysosomes) contain proteinases, acid phosphatase, myeloperoxidase, and β-glucuronidase.

Hypersegmented neutrophils (nucleus has 6+ lobes) are seen in vitamin B_{12}/ folate deficiency.

A left shift with ↑ band cells (immature neutrophils) reflects states of ↑ myeloid proliferation (eg, bacterial infections, CML).

Important neutrophil chemotactic agents: C5a, IL-8, LTB_4, kallikrein, platelet-activating factor.

(c) ketabton.com: The Digital Library

Erythrocytes



Carry O_2 to tissues and CO_2 to lungs. Anucleate and lack organelles; biconcave A, with large surface area-to-volume ratio for rapid gas exchange. Life span of 120 days. Source of energy is glucose (90% used in glycolysis, 10% used in HMP shunt). Membranes contain Cl^-/HCO_3^- antiporter, which allow RBCs to export HCO_3^- and transport CO_2 from the periphery to the lungs for elimination.

Eryth = red; *cyte* = cell.

Erythrocytosis = polycythemia = ↑ Hct.
Anisocytosis = varying sizes.
Poikilocytosis = varying shapes.

Reticulocyte = immature RBC; reflects erythroid proliferation.
Bluish color (polychromasia) on Wright-Giemsa stain of reticulocytes represents residual ribosomal RNA.

Thrombocytes (platelets)



Involved in 1° hemostasis. Small cytoplasmic fragments A derived from megakaryocytes. Life span of 8–10 days. When activated by endothelial injury, aggregate with other platelets and interact with fibrinogen to form platelet plug. Contain dense granules (**C**a^{2+}, **A**DP, **S**erotonin, **H**istamine; **CASH**) and α granules (vWF, fibrinogen, fibronectin, platelet factor 4). Approximately ⅓ of platelet pool is stored in the spleen.

Thrombocytopenia or ↓ platelet function results in petechiae.
vWF receptor: GpIb.
Fibrinogen receptor: GpIIb/IIIa.
Thrombopoietin stimulates megakaryocyte proliferation.
Al**f**a granules contain vW**F**, **f**ibrinogen, **f**ibronectin, platelet **f**actor **f**our.

Monocytes



Found in blood, differentiate into macrophages in tissues.
Large, kidney-shaped nucleus A. Extensive "frosted glass" cytoplasm.

Mono = one (nucleus); *cyte* = cell.

Macrophages



Phagocytose bacteria, cellular debris, and senescent RBCs. Long life in tissues. Differentiate from circulating blood monocytes A. Activated by γ-interferon. Can function as antigen-presenting cell via MHC II. Important cellular component of granulomas (eg, TB, sarcoidosis).

Macro = large; *phage* = eater.
Macrophage naming varies by specific tissue type (eg, Kupffer cells in liver, histiocytes in connective tissue, Langerhans cells in skin, osteoclasts in bone, microglial cells in brain).
Lipid A from bacterial LPS binds CD14 on macrophages to initiate septic shock.

(c) ketabton.com: The Digital Library

Eosinophils



Defend against helminthic infections (major basic protein). Bilobate nucleus. Packed with large eosinophilic granules of uniform size A. Highly phagocytic for antigen-antibody complexes.
Produce histaminase, major basic protein (MBP, a helminthotoxin), eosinophil peroxidase, eosinophil cationic protein, and eosinophil-derived neurotoxin.

Eosin = pink dye; *philic* = loving.
Causes of eosinophilia = **PACCMAN**:
Parasites
Asthma
Eosinophilic granulomatosis with polyangiitis (**C**hurg-Strauss syndrome)
Chronic adrenal insufficiency
Myeloproliferative disorders
Allergic processes
Neoplasia (eg, Hodgkin lymphoma)

Basophils



Mediate allergic reaction. Densely basophilic granules A contain heparin (anticoagulant) and histamine (vasodilator). Leukotrienes synthesized and released on demand.

Basophilic—stains readily with **basic** stains.
Basophilia is uncommon, but can be a sign of myeloproliferative disorders, particularly CML.

Mast cells



Mediate local tissue allergic reactions. Contain basophilic granules A. Originate from same precursor as basophils but are not the same cell type. Can bind the Fc portion of IgE to membrane. Activated by tissue trauma, C3a and C5a, surface IgE cross-linking by antigen (IgE receptor aggregation) → degranulation → release of histamine, heparin, tryptase, and eosinophil chemotactic factors.

Involved in type I hypersensitivity reactions.
Cromolyn sodium prevents mast cell degranulation (used for asthma prophylaxis).
Vancomycin, opioids, and radiocontrast dye can elicit IgE-independent mast cell degranulation.

Dendritic cells



Highly phagocytic antigen-presenting cells (APCs) A. Function as link between innate and adaptive immune systems. Express MHC class II and Fc receptors on surface.

(c) ketabton.com: The Digital Library

Lymphocytes



Refer to B cells, T cells, and NK cells. B cells and T cells mediate adaptive immunity. NK cells are part of the innate immune response. Round, densely staining nucleus with small amount of pale cytoplasm A.

Natural killer cells



Important in innate immunity, especially against intracellular pathogens. Larger than B and T cells, with distinctive cytoplasmic lytic granules (containing perforin and granzymes) that, when released, act on target cells to induce apoptosis. Distinguish between healthy and infected cells by identifying cell surface proteins (induced by stress, malignant transformation, or microbial infections).

B cells



Mediate humoral immune response. Originate from stem cells in bone marrow and matures in marrow. Migrate to peripheral lymphoid tissue (follicles of lymph nodes, white pulp of spleen, unencapsulated lymphoid tissue). When antigen is encountered, B cells differentiate into plasma cells (which produce antibodies) and memory cells. Can function as an APC.

B = Bone marrow.

T cells



Mediate cellular immune response. Originate from stem cells in the bone marrow, but mature in the thymus. Differentiate into cytotoxic T cells (express CD8, recognize MHC I), helper T cells (express CD4, recognize MHC II), and regulatory T cells. CD28 (costimulatory signal) necessary for T-cell activation. Most circulating lymphocytes are T cells (80%).

T = Thymus.

CD4+ helper T cells are the primary target of HIV.

Rule of 8: MHC II × CD4 = 8; MHC I × CD8 = 8.

Plasma cells



Produce large amounts of antibody specific to a particular antigen. "Clock-face" chromatin distribution and eccentric nucleus, abundant RER, and well-developed Golgi apparatus (arrows in A). Found in bone marrow and normally do not circulate in peripheral blood.

Multiple myeloma is a plasma cell dyscrasia.

(c) ketabton.com: The Digital Library

▸ HEMATOLOGY AND ONCOLOGY—PHYSIOLOGY

Hemoglobin electrophoresis



On a gel, hemoglobin migrates from the negatively charged cathode to the positively charged anode. HbA migrates the farthest, followed by HbF, HbS, and HbC. This is because the missense mutations in HbS and HbC replace glutamic acid ⊖ with valine (neutral) and lysine ⊕, respectively, making HbC and HbS more positively charged than HbA.

A Fat Santa Claus can't (cathode → anode) go far.

(c) ketabton.com: The Digital Library

Platelet plug formation (primary hemostasis)



Thrombogenesis



Formation of insoluble fibrin mesh.

Aspirin irreversibly inhibits cyclooxygenase, thereby inhibiting TXA_2 synthesis.

Clopidogrel, prasugrel, and ticlopidine inhibit ADP-induced expression of GpIIb/IIIa by irreversibly blocking $P2Y_{12}$ receptor.

Abciximab, eptifibatide, and tirofiban inhibit GpIIb/IIIa directly.

Ristocetin activates vWF to bind GpIb. Failure of aggregation with ristocetin assay occurs in von Willebrand disease and Bernard-Soulier syndrome.

vWF carries/protects factor VIII; volksWagen Factories make gr8 cars.

(c) ketabton.com: The Digital Library

Coagulation and kinin pathways



(c) ketabton.com: The Digital Library

Vitamin K–dependent coagulation

Procoagulation



Anticoagulation



Vitamin K deficiency: ↓ synthesis of factors II, VII, IX, X, protein C, protein S.

Warfarin inhibits vitamin K epoxide reductase. Vitamin K administration can potentially reverse inhibitory effect of warfarin on clotting factor synthesis (delayed). FFP or PCC administration reverses action of warfarin immediately and can be given with vitamin K in cases of severe bleeding.

Neonates lack enteric bacteria, which produce vitamin K. Early administration of vitamin K overcomes neonatal deficiency/coagulopathy.

Factor VII (seven)—shortest half-life.

Factor II (two)—longest (tallest) half-life.

Antithrombin inhibits thrombin (factor IIa) and factors VIIa, IXa, Xa, XIa, XIIa.

Heparin enhances the activity of antithrombin.

Principal targets of antithrombin: thrombin and factor Xa.

Factor V Leiden mutation produces a factor V resistant to inhibition by activated protein C.

tPA is used clinically as a thrombolytic.

(c) ketabton.com: The Digital Library

▸ HEMATOLOGY AND ONCOLOGY—PATHOLOGY

RBC morphology

TYPE	EXAMPLE	ASSOCIATED PATHOLOGY	NOTES
Acanthocytes ("spur cells")		Liver disease, abetalipoproteinemia	Projections of varying size at irregular intervals.
Echinocytes ("burr cells")		Liver disease, ESRD, pyruvate kinase deficiency	Smaller and more uniform projections than acanthocytes
Dacrocytes ("teardrop cells")		Bone marrow infiltration (eg, myelofibrosis)	RBC "sheds a tear" because it's mechanically squeezed out of its home in the bone marrow
Schistocytes (eg, "helmet" cells)		MAHAs (eg, DIC, TTP/HUS, HELLP syndrome), mechanical hemolysis (eg, heart valve prosthesis)	Fragmented RBCs
Degmacytes ("bite cells")		G6PD deficiency	Due to removal of Heinz bodies by splenic macrophages
Elliptocytes		Hereditary elliptocytosis	Caused by mutation in genes encoding RBC membrane proteins (eg, spectrin)

(c) ketabton.com: The Digital Library

RBC morphology *(continued)*

TYPE	EXAMPLE	ASSOCIATED PATHOLOGY	NOTES
Spherocytes		Hereditary spherocytosis, autoimmune hemolytic anemia	Small, spherical cells without central pallor
Macro-ovalocytes		Megaloblastic anemia (also hypersegmented PMNs)	
Target cells		**H**bC disease, **A**splenia, **L**iver disease, **T**halassemia	"**HALT,**" said the hunter to his **target**
Sickle cells		Sickle cell anemia	Sickling occurs with low O_2 conditions (eg, high altitude, acidosis)

(c) ketabton.com: The Digital Library

RBC inclusions

TYPE	EXAMPLE	ASSOCIATED PATHOLOGY	NOTES
Bone marrow			
Iron granules (eg, in ringed sideroblasts)		Sideroblastic anemias (eg, lead poisoning, myelodysplastic syndromes, alcoholism)	Perinuclear mitochondria with excess iron (forming ring in ringed sideroblasts) Require Prussian blue stain to be visualized
Peripheral smear			
Howell-Jolly bodies		Functional hyposplenia (eg, sickle cell disease), asplenia	Basophilic nuclear remnants (do not contain iron) Usually removed by splenic macrophages
Basophilic stippling		Sideroblastic anemias, thalassemias	Basophilic ribosomal precipitates (do not contain iron)
Pappenheimer bodies		Sideroblastic anemia	Basophilic granules (contain iron)
Heinz bodies		G6PD deficiency	Denatured and precipitated hemoglobin (contain iron) Phagocytic removal of Heinz bodies → bite cells Requires supravital stain (eg, crystal violet) to be visualized

(c) ketabton.com: The Digital Library

Anemias



Reticulocyte index Also called corrected reticulocyte count. Used to correct falsely elevated reticulocyte count in anemia. Measures appropriate bone marrow response to anemic conditions (effective erythropoiesis). High reticulocyte index (RI) indicates compensatory RBC production; low RI indicates inadequate response to correct anemia. Calculated as:

$$\text{RI} = \text{reticulocyte \%} \times \text{actual Hct/normal Hct}$$

[normal Hct ≈ 45%]

(c) ketabton.com: The Digital Library

Microcytic, hypochromic anemias

MCV < 80 fL.

Iron deficiency

↓ iron due to chronic bleeding (eg, GI loss, menorrhagia), malnutrition, absorption disorders, GI surgery (eg, gastrectomy), or ↑ demand (eg, pregnancy) → ↓ final step in heme synthesis.

Labs: ↓ iron, ↑ TIBC, ↓ ferritin, ↑ free erythrocyte protoporphyrin, ↑ RDW, ↓ RI. Microcytosis and hypochromasia (↑ central pallor) A.

Symptoms: fatigue, conjunctival pallor B, pica (persistent craving and compulsive eating of nonfood substances), spoon nails (koilonychia).

May manifest as glossitis, cheilosis, Plummer-Vinson syndrome (triad of iron deficiency anemia, esophageal webs, and dysphagia).

α-thalassemia

α-globin gene deletions on chromosome 16 → ↓ α-globin synthesis. *cis* deletion (deletions occur on same chromosome) prevalent in Asian populations; *trans* deletion (deletions occur on separate chromosomes) prevalent in African populations. Normal is αα/αα.

NUMBER OF α-GLOBIN GENES DELETED	DISEASE	CLINICAL OUTCOME
1 (α α/α –)	α-thalassemia minima	No anemia (silent carrier)
2 (α –/α –; *trans*) or (α α/– –; *cis*)	α-thalassemia minor	Mild microcytic, hypochromic anemia; *cis* deletion may worsen outcome for the carrier's offspring
3 (– –/– α)	Hemoglobin H disease (HbH); excess β-globin forms β_4	Moderate to severe microcytic hypochromic anemia
4 (– –/– –)	Hemoglobin Barts disease; no α-globin, excess γ-globin forms γ_4	Hydrops fetalis; incompatible with life

β-thalassemia

Point mutations in splice sites and promoter sequences on chromosome 11 → ↓ β-globin synthesis. Prevalent in Mediterranean populations.

β-thalassemia minor (heterozygote): β chain is underproduced. Usually asymptomatic. Diagnosis confirmed by ↑ HbA_2 (> 3.5%) on electrophoresis.

β-thalassemia major (homozygote): β chain is absent → severe microcytic, hypochromic anemia with target cells and increased anisopoikilocytosis C requiring blood transfusion (2° hemochromatosis). Marrow expansion ("crew cut" on skull x-ray) → skeletal deformities (eg, "chipmunk" facies). Extramedullary hematopoiesis → hepatosplenomegaly. ↑ risk of parvovirus B19–induced aplastic crisis. ↑ HbF ($\alpha_2\gamma_2$), HbA_2 ($\alpha_2\delta_2$). HbF is protective in the infant and disease becomes symptomatic only after 6 months, when fetal hemoglobin declines.

HbS/β-thalassemia heterozygote: mild to moderate sickle cell disease depending on amount of β-globin production.

(c) ketabton.com: The Digital Library

Microcytic, hypochromic anemias *(continued)*

Lead poisoning

Lead inhibits ferrochelatase and ALA dehydratase → ↓ heme synthesis and ↑ RBC protoporphyrin. Also inhibits rRNA degradation → RBCs retain aggregates of rRNA (basophilic stippling).

Symptoms of **LEAD** poisoning:

- **L**ead **L**ines on gingivae (Burton lines) and on metaphyses of long bones **D** on x-ray.
- **E**ncephalopathy and **E**rythrocyte basophilic stippling.
- **A**bdominal colic and sideroblastic **A**nemia.
- **D**rops—wrist and foot drop. **D**imercaprol and **E**D**T**A are 1st line of treatment.

Succimer used for chelation for kids (It "**suc**ks" to be a kid who eats lead).

Exposure risk ↑ in old houses with chipped paint.

Sideroblastic anemia

Causes: genetic (eg, X-linked defect in ALA synthase gene), acquired (myelodysplastic syndromes), and reversible (alcohol is most common; also lead poisoning, vitamin B_6 deficiency, copper deficiency, drugs [eg, isoniazid, linezolid]).

Lab findings: ↑ iron, normal/↓ TIBC, ↑ ferritin. Ringed sideroblasts (with iron-laden, Prussian blue–stained mitochondria) seen in bone marrow **E**. Peripheral blood smear: basophilic stippling of RBCs. Some acquired variants may be normocytic or macrocytic.

Treatment: pyridoxine (B_6, cofactor for ALA synthase).



Interpretation of iron studies

	Iron deficiency	Chronic disease	Hemochromatosis	Pregnancy/ OCP use
Serum iron	↓	↓	↑	—
Transferrin or TIBC	↑	↓[a]	↓	↑
Ferritin	↓	↑	↑	—
% transferrin saturation (serum iron/TIBC)	↓↓	—/↓	↑↑	↓

↑↓ = 1° disturbance.

Transferrin—**transport**s iron in blood.

TIBC—indirectly measures transferrin.

Ferritin—1° iron storage protein of body.

[a]Evolutionary reasoning—pathogens use circulating iron to thrive. The body has adapted a system in which iron is stored within the cells of the body and prevents pathogens from acquiring circulating iron.

(c) ketabton.com: The Digital Library

Macrocytic anemias

MCV > 100 fL.

	DESCRIPTION	FINDINGS
Megaloblastic anemia	Impaired DNA synthesis → maturation of nucleus of precursor cells in bone marrow delayed relative to maturation of cytoplasm. Causes: vitamin B_{12} deficiency, folate deficiency, medications (eg, hydroxyurea, phenytoin, methotrexate, sulfa drugs).	RBC macrocytosis, hypersegmented neutrophils (arrow in A), glossitis.
Folate deficiency	Causes: malnutrition (eg, alcoholics), malabsorption, drugs (eg, methotrexate, trimethoprim, phenytoin), ↑ requirement (eg, hemolytic anemia, pregnancy).	↑ homocysteine, normal methylmalonic acid. **No neurologic symptoms** (vs B_{12} deficiency).
Vitamin B_{12} (cobalamin) deficiency	Causes: pernicious anemia, malabsorption (eg, Crohn disease), pancreatic insufficiency, gastrectomy, insufficient intake (eg, veganism), *Diphyllobothrium latum* (fish tapeworm).	↑ homocysteine, ↑ methylmalonic acid. **Neurologic symptoms:** reversible dementia, subacute combined degeneration (due to involvement of B_{12} in fatty acid pathways and myelin synthesis): spinocerebellar tract, lateral corticospinal tract, dorsal column dysfunction. Folate supplementation in vitamin B_{12} deficiency can correct the anemia, but worsens neurologic symptoms. Historically diagnosed with the Schilling test, a test that determines if the cause is dietary insufficiency vs malabsorption. Anemia 2° to insufficient intake may take several years to develop due to liver's ability to store B_{12} (as opposed to folate deficiency).
Orotic aciduria	Inability to convert orotic acid to UMP (de novo pyrimidine synthesis pathway) because of defect in UMP synthase. Autosomal recessive. Presents in children as failure to thrive, developmental delay, and megaloblastic anemia refractory to folate and B_{12}. No hyperammonemia (vs ornithine transcarbamylase deficiency—↑ orotic acid with hyperammonemia).	Orotic acid in urine. Treatment: uridine monophosphate or uridine triacetate to bypass mutated enzyme.
Nonmegaloblastic anemia	Macrocytic anemia in which DNA synthesis is normal. Causes: alcoholism, liver disease.	RBC macrocytosis without hypersegmented neutrophils.
Diamond-Blackfan anemia	A congenital form of pure red cell aplasia. Rapid-onset anemia within 1st year of life due to intrinsic defect in erythroid progenitor cells.	↑ % HbF (but ↓ total Hb). Short stature, craniofacial abnormalities, and upper extremity malformations (triphalangeal thumbs) in up to 50% of cases.

(c) ketabton.com: The Digital Library

Normocytic, normochromic anemias	Normocytic, normochromic anemias are classified as nonhemolytic or hemolytic. The hemolytic anemias are further classified according to the cause of the hemolysis (intrinsic vs extrinsic to the RBC) and by the location of the hemolysis (intravascular vs extravascular). Hemolysis can lead to increases in LDH, reticulocytes, unconjugated bilirubin, pigmented gallstones, and urobilinogen in urine.
Intravascular hemolysis	Findings: ↓ haptoglobin, ↑ schistocytes on blood smear. Characteristic hemoglobinuria, hemosiderinuria, and urobilinogen in urine. Notable causes are mechanical hemolysis (eg, prosthetic valve), paroxysmal nocturnal hemoglobinuria, microangiopathic hemolytic anemias.
Extravascular hemolysis	Mechanism: macrophages in spleen clear RBCs. Findings: spherocytes in peripheral smear (most commonly due to hereditary spherocytosis and autoimmune hemolytic anemia), no hemoglobinuria/hemosiderinuria. Can present with urobilinogen in urine.

Nonhemolytic, normocytic anemias

	DESCRIPTION	FINDINGS
Anemia of chronic disease	Inflammation (eg, ↑ IL-6) → ↑ hepcidin (released by liver, binds ferroportin on intestinal mucosal cells and macrophages, thus inhibiting iron transport) → ↓ release of iron from macrophages and ↓ iron absorption from gut. Associated with conditions such as chronic infections, neoplastic disorders, chronic kidney disease, and autoimmune diseases (eg, SLE, rheumatoid arthritis).	↓ iron, ↓ TIBC, ↑ ferritin. Normocytic, but can become microcytic. Treatment: address underlying cause of inflammation, judicious use of blood transfusion, consider erythropoiesis-stimulating agents such as EPO (eg, in chronic kidney disease).
Aplastic anemia  	Caused by failure or destruction of hematopoietic stem cells due to: ▪ Radiation and drugs (eg, benzene, chloramphenicol, alkylating agents, antimetabolites) ▪ Viral agents (eg, EBV, HIV, hepatitis viruses) ▪ Fanconi anemia (autosomal recessive DNA repair defect → bone marrow failure); normocytosis or macrocytosis on CBC ▪ Idiopathic (immune mediated, 1° stem cell defect); may follow acute hepatitis	↓ reticulocyte count, ↑ EPO. Pancytopenia characterized by anemia, leukopenia, and thrombocytopenia (not to be confused with aplastic crisis, which causes anemia only). Normal cell morphology, but hypocellular bone marrow with fatty infiltration A (dry bone marrow tap). Symptoms: fatigue, malaise, pallor, purpura, mucosal bleeding, petechiae, infection. Treatment: withdrawal of offending agent, immunosuppressive regimens (eg, antithymocyte globulin, cyclosporine), bone marrow allograft, RBC/platelet transfusion, bone marrow stimulation (eg, GM-CSF).

(c) ketabton.com: The Digital Library

Intrinsic hemolytic anemias

	DESCRIPTION	FINDINGS
Hereditary spherocytosis	Primarily autosomal dominant. Due to defect in proteins interacting with RBC membrane skeleton and plasma membrane (eg, ankyrin, band 3, protein 4.2, spectrin). Small, round RBCs with less surface area and no central pallor (↑ MCHC) → premature removal by spleen (extravascular hemolysis).	Splenomegaly, aplastic crisis (parvovirus B19 infection). Labs: ↓ mean fluorescence of RBCs in eosin 5-maleimide (EMA) binding test, ↑ fragility in osmotic fragility test. Normal to ↓ MCV with abundance of RBCs. Treatment: splenectomy.
G6PD deficiency	X-linked recessive. G6PD defect → ↓ NADPH → ↓ reduced glutathione → ↑ RBC susceptibility to oxidative stress (eg, sulfa drugs, antimalarials, **fava beans**) → hemolysis. Causes extravascular and intravascular hemolysis.	Back pain, hemoglobinuria a few days after oxidant **stress**. Labs: blood smear shows RBCs with **Heinz** bodies and **bite** cells. "**Stress** makes me eat **bites** of **fava beans** with **Heinz** ketchup."
Pyruvate kinase deficiency	Autosomal recessive. Pyruvate kinase defect → ↓ ATP → rigid RBCs → extravascular hemolysis. Increases levels of 2,3-BPG → ↓ hemoglobin affinity for O_2.	Hemolytic anemia in a newborn.
Paroxysmal nocturnal hemoglobinuria	Hematopoietic stem cell mutation → ↑ complement-mediated intravascular hemolysis, especially at night. Acquired *PIGA* mutation → impaired GPI anchor synthesis for decay-accelerating factor (DAF/CD55) and membrane inhibitor of reactive lysis (MIRL/CD59), which protect RBC membrane from complement.	Triad: Coombs ⊖ hemolytic anemia, pancytopenia, venous thrombosis (eg, Budd-Chiari syndrome). Pink/red urine in morning. Associated with aplastic anemia, acute leukemias. Labs: CD55/59 ⊖ RBCs on flow cytometry. Treatment: eculizumab (targets terminal complement protein C5).
Sickle cell anemia  A	Point mutation in β-globin gene → single amino acid substitution (glutamic acid → valine). Mutant HbA is termed HbS. Causes extravascular and intravascular hemolysis. Pathogenesis: low O_2, high altitude, or acidosis precipitates sickling (deoxygenated HbS polymerizes) → anemia, vaso-occlusive disease. Newborns are initially asymptomatic because of ↑ HbF and ↓ HbS. Heterozygotes (sickle cell trait) have resistance to malaria. 8% of African Americans carry an HbS allele. Sickle cells are crescent-shaped RBCs **A**. "Crew cut" on skull x-ray due to marrow expansion from ↑ erythropoiesis (also seen in thalassemias).	Complications in sickle cell disease: ▪ Aplastic crisis (transient arrest of erythropoiesis due to parvovirus B19). ▪ Autosplenectomy (Howell-Jolly bodies) → ↑ risk of infection by encapsulated organisms (eg, *S pneumoniae*). ▪ Splenic infarct/sequestration crisis. ▪ *Salmonella* osteomyelitis. ▪ Painful vaso-occlusive crises: dactylitis (painful swelling of hands/feet), priapism, acute chest syndrome (respiratory distress, new pulmonary infiltrates on CXR, common cause of death), avascular necrosis, stroke. ▪ Sickling in renal medulla (↓ Po_2) → renal papillary necrosis → hematuria. Hb electrophoresis: ↓↓ HbA, ↑ HbF, ↑↑ HbS. Treatment: hydroxyurea (↑ HbF), hydration.
HbC disease	Glutamic acid–to-lyCine (lysine) mutation in β-globin. Causes extravascular hemolysis.	Patients with HbSC (1 of each mutant gene) have milder disease than HbSS patients. Blood smear in homozygotes: hemoglobin Crystals inside RBCs, target cells.

(c) ketabton.com: The Digital Library

Extrinsic hemolytic anemias

	DESCRIPTION	FINDINGS
Autoimmune hemolytic anemia 	A normocytic anemia that is usually idiopathic and Coombs ⊕. Two types: ▪ **Warm** AIHA–chronic anemia in which Ig**G** causes RBC agglutination. Seen in SLE and CLL and with certain drugs (eg, α-methyldopa). "**Warm** weather is **G**ood." ▪ Cold AIHA–acute anemia in which Ig**M** + complement causes RBC agglutination upon exposure to cold → painful, blue fingers and toes. Seen in CLL, ***M**ycoplasma pneumoniae* infections, infectious **M**ononucleosis.	Spherocytes and agglutinated RBCs **A** on peripheral blood smear. Warm AIHA treatment: steroids, rituximab, splenectomy (if refractory). Cold AIHA treatment: cold avoidance, rituximab. Direct Coombs test—anti-Ig antibody (Coombs reagent) added to patient's RBCs. RBCs agglutinate if RBCs are coated with Ig. For comparison, Indirect Coombs test—normal RBCs added to patient's serum. If serum has anti-RBC surface Ig, RBCs agglutinate when Coombs reagent added.
Microangiopathic hemolytic anemia	RBCs are damaged when passing through obstructed or narrowed vessels. Causes intravascular hemolysis. Seen in DIC, TTP/HUS, SLE, HELLP syndrome, hypertensive emergency.	**Schisto**cytes (eg, "helmet cells") are seen on peripheral blood smear due to mechanical destruction (***schisto*** = to split) of RBCs.
Macroangiopathic hemolytic anemia	Prosthetic heart valves and aortic stenosis may also cause hemolytic anemia 2° to mechanical destruction of RBCs.	Schistocytes on peripheral blood smear.
Hemolytic anemia due to infection	↑ destruction of RBCs (eg, malaria, *Babesia*).	



(c) ketabton.com: The Digital Library

Leukopenias

CELL TYPE	CELL COUNT	CAUSES
Neutropenia	Absolute neutrophil count < 1500 cells/mm^3 Severe infections typical when < 500 cells/mm^3	Sepsis/postinfection, drugs (including chemotherapy), aplastic anemia, SLE, radiation
Lymphopenia	Absolute lymphocyte count < 1500 cells/mm^3 (< 3000 cells/mm^3 in children)	HIV, DiGeorge syndrome, SCID, SLE, corticosteroids[a], radiation, sepsis, postoperative
Eosinopenia	Absolute eosinophil count < 30 cells/mm^3	Cushing syndrome, corticosteroids[a]

[a]Corticosteroids cause neutrophilia, despite causing eosinopenia and lymphopenia. Corticosteroids ↓ activation of neutrophil adhesion molecules, impairing migration out of the vasculature to sites of inflammation. In contrast, corticosteroids sequester eosinophils in lymph nodes and cause apoptosis of lymphocytes.

Neutrophil left shift	↑ neutrophil precursors, such as band cells and metamyelocytes, in peripheral blood. Usually seen with neutrophilia in the acute response to infection or inflammation. Called leukoerythroblastic reaction when left shift is seen with immature RBCs. Occurs with severe anemia (physiologic response) or marrow response (eg, fibrosis, tumor taking up space in marrow).	A left shift is a shift to a more immature cell in the maturation process.

(c) ketabton.com: The Digital Library

Heme synthesis, porphyrias, and lead poisoning

The porphyrias are hereditary or acquired conditions of defective heme synthesis that lead to the accumulation of heme precursors. Lead inhibits specific enzymes needed in heme synthesis, leading to a similar condition.

CONDITION	AFFECTED ENZYME	ACCUMULATED SUBSTRATE	PRESENTING SYMPTOMS
Lead poisoning 	Ferrochelatase and ALA dehydratase	Protoporphyrin, ALA (blood)	Microcytic anemia (basophilic stippling in peripheral smear A, ringed sideroblasts in bone marrow), GI and kidney disease. Children—exposure to lead paint → mental deterioration. Adults—environmental exposure (eg, batteries, ammunition) → headache, memory loss, demyelination (peripheral neuropathy).
Acute intermittent porphyria	Porphobilinogen deaminase, previously called uroporphyrinogen I synthase (autosomal dominant mutation)	Porphobilinogen, ALA	Symptoms (5 P's): ▪ Painful abdomen ▪ Port wine–colored Pee ▪ Polyneuropathy ▪ Psychological disturbances ▪ Precipitated by drugs (eg, cytochrome P-450 inducers), alcohol, starvation Treatment: hemin and glucose.
Porphyria cutanea tarda	Uroporphyrinogen decarboxylase	Uroporphyrin (tea-colored urine)	Blistering cutaneous photosensitivity and hyperpigmentation B. Most common porphyria. Exacerbated with alcohol consumption. Causes: familial, hepatitis C. Treatment: phlebotomy, sun avoidance, antimalarials (eg, hydroxychloroquine).



(c) ketabton.com: The Digital Library

Iron poisoning

	Acute	Chronic
FINDINGS	High mortality rate associated with accidental ingestion by children (adult iron tablets may look like candy).	Seen in patients with 1° (hereditary) or 2° (eg, chronic blood transfusions for thalassemia or sickle cell disease) hemochromatosis.
MECHANISM	Cell death due to formation of free radicals and peroxidation of membrane lipids.	
SYMPTOMS/SIGNS	Abdominal pain, vomiting, GI bleeding. Radiopaque pill seen on x-ray. May progress to anion gap metabolic acidosis and multiorgan failure. Leads to scarring with GI obstruction.	Arthropathy, cirrhosis, cardiomyopathy, diabetes mellitus and skin pigmentation ("bronze diabetes"), hypogonadism.
TREATMENT	Chelation (eg, deferoxamine, deferasirox), gastric lavage.	Phlebotomy (patients without anemia) or chelation.

Coagulation disorders

PT—tests function of common and extrinsic pathway (factors I, II, V, VII, and X). Defect → ↑ **PT** (**P**lay **T**ennis outside [extrinsic pathway]).

INR (international normalized ratio) = patient PT/control PT. 1 = normal, > 1 = prolonged. Most common test used to follow patients on warfarin, which prolongs INR.

PTT—tests function of common and **in**trinsic pathway (all factors except VII and XIII). Defect → ↑ **PTT** (**P**lay **T**able **T**ennis **in**side).

Coagulation disorders can be due to clotting factor deficiencies or acquired factor inhibitors. Diagnosed with a mixing study, in which normal plasma is added to patient's plasma. Clotting factor deficiencies should correct (the PT or PTT returns to within the appropriate normal range), whereas factor inhibitors will not correct.

DISORDER	PT	PTT	MECHANISM AND COMMENTS
Hemophilia A, B, or C	—	↑	Intrinsic pathway coagulation defect (↑ PTT). ▪ A: deficiency of factor VIII; X-linked recessive. ▪ B: deficiency of factor IX; X-linked recessive. ▪ C: deficiency of factor XI; autosomal recessive. Hemorrhage in hemophilia—hemarthroses (bleeding into joints, eg, knee A), easy bruising, bleeding after trauma or surgery (eg, dental procedures). Treatment: desmopressin + factor VIII concentrate (A); factor IX concentrate (B); factor XI concentrate (C).
Vitamin K deficiency	↑	↑	General coagulation defect. Bleeding time normal. ↓ activity of factors II, VII, IX, X, protein C, protein S.

(c) ketabton.com: The Digital Library

Platelet disorders

All platelet disorders have ↑ bleeding time (BT), mucous membrane bleeding, and microhemorrhages (eg, petechiae, epistaxis). Platelet count (PC) is usually low, but may be normal in qualitative disorders.

DISORDER	PC	BT	NOTES
Bernard-Soulier syndrome	–/↓	↑	Defect in adhesion. ↓ GpIb → ↓ platelet-to-vWF adhesion. Labs: abnormal ristocetin test, large platelets.
Glanzmann thrombasthenia	–	↑	Defect in aggregation. ↓ GpIIb/IIIa (↓ integrin $\alpha_{IIb}\beta_3$) → ↓ platelet-to-platelet aggregation and defective platelet plug formation. Labs: blood smear shows no platelet clumping.
Immune thrombocytopenia	↓	↑	Destruction of platelets in spleen. Anti-GpIIb/IIIa antibodies → splenic macrophages phagocytose platelets. May be idiopathic or 2° to autoimmune disorders (eg, SLE), viral illness (eg, HIV, HCV), malignancy (eg, CLL), or drug reactions. Labs: ↑ megakaryocytes on bone marrow biopsy, ↓ platelet count. Treatment: steroids, IVIG, rituximab, TPO receptor agonists (eg, eltrombopag, romiplostim), or splenectomy for refractory ITP.

Thrombotic microangiopathies

Disorders overlap significantly in symptomatology.

	Thrombotic thrombocytopenic purpura	Hemolytic-uremic syndrome
EPIDEMIOLOGY	Typically females	Typically children
PATHOPHYSIOLOGY	Inhibition or deficiency of ADAMTS13 (a vWF metalloprotease) → ↓ degradation of vWF multimers → ↑ large vWF multimers → ↑ platelet adhesion and aggregation (microthrombi formation)	Commonly caused by Shiga-like toxin from EHEC (serotype O157:H7) infection
PRESENTATION	Triad of thrombocytopenia (↓ platelets), microangiopathic hemolytic anemia (↓ Hb, schistocytes, ↑ LDH), acute kidney injury (↑ Cr)	
DIFFERENTIATING SYMPTOMS	Triad + fever + neurologic symptoms	Triad + bloody diarrhea
LABS	Normal PT and PTT helps distinguish TTP and HUS (coagulation pathway is not activated) from DIC (coagulation pathway is activated)	
TREATMENT	Plasmapheresis, steroids, rituximab	Supportive care

(c) ketabton.com: The Digital Library

Mixed platelet and coagulation disorders

DISORDER	PC	BT	PT	PTT	NOTES
von Willebrand disease	—	↑	—	—/↑	Intrinsic pathway coagulation defect: ↓ vWF → ↑ PTT (vWF carries/protects factor VIII). Defect in platelet plug formation: ↓ vWF → defect in platelet-to-vWF adhesion. Autosomal dominant. Mild but most common inherited bleeding disorder. No platelet aggregation with ristocetin cofactor assay. Treatment: desmopressin, which releases vWF stored in endothelium.
Disseminated intravascular coagulation	↓	↑	↑	↑	Widespread clotting factor activation → deficiency in clotting factors → bleeding state. Causes: Snake bites, Sepsis (gram ⊖), Trauma, Obstetric complications, acute Pancreatitis, Malignancy, Nephrotic syndrome, Transfusion (SSTOP Making New Thrombi). Labs: schistocytes, ↑ fibrin degradation products (D-dimers), ↓ fibrinogen, ↓ factors V and VIII.

Hereditary thrombosis syndromes leading to hypercoagulability

DISEASE	DESCRIPTION
Antithrombin deficiency	Autosomal dominant inherited deficiency of antithrombin: has no direct effect on the PT, PTT, or thrombin time but diminishes the increase in PTT following heparin administration. Can also be acquired: renal failure/nephrotic syndrome → antithrombin loss in urine → ↓ inhibition of factors IIa and Xa.
Factor V Leiden	Autosomal dominant, most common cause of inherited hypercoagulability in Caucasians. Production of mutant factor V (guanine → adenine DNA point mutation → Arg506Gln mutation near the cleavage site) that is resistant to degradation by activated protein C. Complications include DVT, cerebral vein thrombosis, recurrent pregnancy loss.
Protein C or S deficiency	↓ ability to inactivate factors Va and VIIIa. ↑ risk of thrombotic skin necrosis with hemorrhage after administration of warfarin. If this occurs, think protein C deficiency. Together, protein C Cancels, and protein S Stops, coagulation.
Prothrombin gene mutation	Mutation in 3′ untranslated region → ↑ production of prothrombin → ↑ plasma levels and venous clots.

(c) ketabton.com: The Digital Library

Blood transfusion therapy

COMPONENT	DOSAGE EFFECT	CLINICAL USE
Packed RBCs	↑ Hb and O_2 carrying capacity	Acute blood loss, severe anemia
Platelets	↑ platelet count (↑ ~ 5000/mm^3/unit)	Stop significant bleeding (thrombocytopenia, qualitative platelet defects)
Fresh frozen plasma/prothrombin complex concentrate	↑ coagulation factor levels; FFP contains all coagulation factors and plasma proteins; PCC generally contains factors II, VII, IX, and X, as well as protein C and S	Cirrhosis, immediate anticoagulation reversal
Cryoprecipitate	Contains fibrinogen, factor VIII, factor XIII, vWF, and fibronectin	Coagulation factor deficiencies involving fibrinogen and factor VIII

Blood transfusion risks include infection transmission (low), transfusion reactions, iron overload (may lead to 2° hemochromatosis), hypocalcemia (citrate is a Ca^{2+} chelator), and hyperkalemia (RBCs may lyse in old blood units).

Leukemia vs lymphoma

Leukemia	Lymphoid or myeloid neoplasm with widespread involvement of bone marrow. Tumor cells are usually found in peripheral blood.
Lymphoma	Discrete tumor mass arising from lymph nodes. Presentations often blur definitions.

Hodgkin vs non-Hodgkin lymphoma

Hodgkin	Non-Hodgkin
Both may present with constitutional ("B") signs/symptoms: low-grade fever, night sweats, weight loss.	
Localized, single group of nodes with contiguous spread (stage is strongest predictor of prognosis). Better prognosis.	Multiple lymph nodes involved; extranodal involvement common; noncontiguous spread. Worse prognosis.
Characterized by Reed-Sternberg cells.	Majority involve B cells; a few are of T-cell lineage.
Bimodal distribution: young adulthood and > 55 years; more common in men except for nodular sclerosing type.	Can occur in children and adults.
Associated with EBV.	May be associated with autoimmune diseases and viral infections (eg, HIV, EBV, HTLV).

Hodgkin lymphoma



Contains Reed-Sternberg cells: distinctive tumor giant cells; binucleate or bilobed with the 2 halves as mirror images ("owl eyes" A). RS cells are CD15+ and CD30+ B-cell origin. 2 owl eyes × 15 = 30.

SUBTYPE	NOTES
Nodular sclerosis	Most common
Lymphocyte rich	Best prognosis
Mixed cellularity	Eosinophilia, seen in immunocompromised patients
Lymphocyte depleted	Seen in immunocompromised patients

(c) ketabton.com: The Digital Library

Non-Hodgkin lymphoma

TYPE	OCCURS IN	GENETICS	COMMENTS
Neoplasms of mature B cells			
Burkitt lymphoma	Adolescents or young adults	t(8;14)—translocation of c-*myc* (8) and heavy-chain Ig (14)	"Starry sky" appearance, sheets of lymphocytes with interspersed "tingible body" macrophages (arrows in A). Associated with EBV. Jaw lesion B in endemic form in Africa; pelvis or abdomen in sporadic form.
Diffuse large B-cell lymphoma	Usually older adults, but 20% in children	Mutations in *BCL-2*, *BCL-6*	Most common type of non-Hodgkin lymphoma in adults.
Follicular lymphoma	Adults	t(14;18)—translocation of heavy-chain Ig (14) and *BCL-2* (18)	Indolent course with painless "waxing and waning" lymphadenopathy. Bcl-2 normally inhibits apoptosis.
Mantle cell lymphoma	Adult **males** >> adult females	t(11;14)—translocation of cyclin D1 (11) and heavy-chain Ig (14), CD5+	Very aggressive, patients typically present with late-stage disease.
Marginal zone lymphoma	Adults	t(11;18)	Associated with chronic inflammation (eg, Sjögren syndrome, chronic gastritis [MALT lymphoma]).
Primary central nervous system lymphoma	Adults	EBV related; associated with HIV/AIDS	Considered an AIDS-defining illness. Variable presentation: confusion, memory loss, seizures. CNS mass (often single, ring-enhancing lesion on MRI) in immunocompromised patients C, needs to be distinguished from toxoplasmosis via CSF analysis or other lab tests.
Neoplasms of mature T cells			
Adult T-cell lymphoma	Adults	Caused by HTLV (associated with IV drug abuse)	Adults present with cutaneous lesions; common in Japan (**T**-cell in **T**okyo), West Africa, and the Caribbean. Lytic bone lesions, hypercalcemia.
Mycosis fungoides/ Sézary syndrome	Adults		Mycosis fungoides: skin patches and plaques D (cutaneous T-cell lymphoma), characterized by atypical CD4+ cells with "cerebriform" nuclei and intraepidermal neoplastic cell aggregates (Pautrier microabscess). May progress to Sézary syndrome (T-cell leukemia).



(c) ketabton.com: The Digital Library

Plasma cell dyscrasias



Characterized by monoclonal immunoglobulin (Ig) overproduction due to plasma cell disorder.
Labs: serum protein electrophoresis (SPEP) or free light chain (FLC) assay for initial tests (M spike on SPEP represents overproduction of a monoclonal Ig fragment). For urinalysis, use 24-hr urine protein electrophoresis (UPEP) to detect light chain, as routine urine dipstick detects only albumin.
Confirm with bone marrow biopsy.

Multiple myeloma

Overproduction of IgG (55% of cases) > IgA.
Clinical features: **CRAB**
- Hyper**C**alcemia
- **R**enal involvement
- **A**nemia
- Bone lytic lesions ("punched out" on X-ray **A**) → **B**ack pain.

Peripheral blood smear shows Rouleaux formation **B** (RBCs stacked like poker chips).
Urinalysis shows Ig light chains (Bence Jones proteinuria) with ⊖ urine dipstick.
Bone marrow analysis shows > 10% monoclonal plasma cells with clock-face chromatin **C** and intracytoplasmic inclusions containing IgG.
Complications: ↑ infection risk, 1° amyloidosis (AL).

Waldenstrom macroglobulinemia

Overproduction of Ig**M** (macroglobulinemia because Ig**M** is the **largest** Ig).
Clinical features:
- Peripheral neuropathy
- No CRAB findings
- Hyperviscosity syndrome:
  - Headache
  - Blurry vision
  - Raynaud phenomenon
  - Retinal hemorrhages

Bone marrow analysis shows >10% small lymphocytes with IgM-containing vacuoles (lymphoplasmacytic lymphoma).
Complication: thrombosis.

Monoclonal gammopathy of undetermined significance

Overproduction of any Ig type.
Usually asymptomatic. No CRAB findings.
Bone marrow analysis shows < 10% monoclonal plasma cells.
Complication: 1-2% risk per year of transitioning to multiple myeloma.







(c) ketabton.com: The Digital Library

Myelodysplastic syndromes



Stem cell disorders involving ineffective hematopoiesis → defects in cell maturation of nonlymphoid lineages. Caused by de novo mutations or environmental exposure (eg, radiation, benzene, chemotherapy). Risk of transformation to AML.

Pseudo-Pelger-Huet anomaly—neutrophils with bilobed ("duet") nuclei A. Typically seen after chemotherapy.

Leukemias

Unregulated growth and differentiation of WBCs in bone marrow → marrow failure → anemia (↓ RBCs), infections (↓ mature WBCs), and hemorrhage (↓ platelets). Usually presents with ↑ circulating WBCs (malignant leukocytes in blood); rare cases present with normal/↓ WBCs. Leukemic cell infiltration of liver, spleen, lymph nodes, and skin (leukemia cutis) possible.

TYPE	NOTES
Lymphoid neoplasms	
Acute lymphoblastic leukemia/lymphoma	Most frequently occurs in children; less common in adults (worse prognosis). T-cell ALL can present as mediastinal mass (presenting as SVC-like syndrome). Associated with Down syndrome. Peripheral blood and bone marrow have ↑↑↑ lymphoblasts A. TdT+ (marker of pre-T and pre-B cells), CD10+ (marker of pre-B cells). Most responsive to therapy. May spread to CNS and testes. t(12;21) → better prognosis.
Chronic lymphocytic leukemia/small lymphocytic lymphoma	Age > 60 years. Most common adult leukemia. CD20+, CD23+, CD5+ B-cell neoplasm. Often asymptomatic, progresses slowly; smudge cells B in peripheral blood smear; autoimmune hemolytic anemia. CLL = Crushed Little Lymphocytes (smudge cells). Richter transformation—CLL/SLL transformation into an aggressive lymphoma, most commonly diffuse large B-cell lymphoma (DLBCL).
Hairy cell leukemia	Adult males. Mature B-cell tumor. Cells have filamentous, hair-like projections (fuzzy appearing on LM C). Peripheral lymphadenopathy is uncommon. Causes marrow fibrosis → dry tap on aspiration. Patients usually present with massive splenomegaly and pancytopenia. Stains TRAP (tartrate-resistant acid phosphatase) ⊕ (trapped in a hairy situation). TRAP stain largely replaced with flow cytometry. Associated with *BRAF* mutations. Treatment: cladribine, pentostatin.
Myeloid neoplasms	
Acute myelogenous leukemia	Median onset 65 years. Auer rods D; myeloperoxidase ⊕ cytoplasmic inclusions seen mostly in APL (formerly M3 AML); ↑↑↑ circulating myeloblasts on peripheral smear. Risk factors: prior exposure to alkylating chemotherapy, radiation, myeloproliferative disorders, Down syndrome. APL: t(15;17), responds to all-*trans* retinoic acid (vitamin A) and arsenic, which induce differentiation of promyelocytes; DIC is a common presentation.

(c) ketabton.com: The Digital Library

Leukemias *(continued)*

Chronic myelogenous leukemia

Peak incidence: 45—85 years; median age: 64 years. Defined by the Philadelphia chromosome (t[9;22], *BCR-ABL*) and myeloid stem cell proliferation. Presents with dysregulated production of mature and maturing granulocytes (eg, neutrophils, metamyelocytes, myelocytes, basophils E) and splenomegaly. May accelerate and transform to AML or ALL ("blast crisis").

Very low leukocyte alkaline phosphatase (LAP) as a result of low activity in malignant neutrophils, vs benign neutrophilia (leukemoid reaction) in which LAP is ↑ due to ↑ leukocyte count with neutrophilia in response to stressors (eg, infections, medications, severe hemorrhage).

Responds to bcr-abl tyrosine kinase inhibitors (eg, imatinib).



Chronic myeloproliferative disorders

Malignant hematopoietic neoplasms with varying impacts on WBCs and myeloid cell lines.

Polycythemia vera

Primary polycythemia. Disorder of ↑ RBCs, usually due to acquired *JAK2* mutation. May present as intense itching after shower (aquagenic pruritus). Rare but classic symptom is erythromelalgia (severe, burning pain and red-blue coloration) due to episodic blood clots in vessels of the extremities A.

↓ EPO (vs 2° polycythemia, which presents with endogenous or artificially ↑ EPO).

Treatment: phlebotomy, hydroxyurea, ruxolitinib (JAK1/2 inhibitor).

Essential thrombocythemia

Characterized by massive proliferation of megakaryocytes and platelets. Symptoms include bleeding and thrombosis. Blood smear shows markedly increased number of platelets, which may be large or otherwise abnormally formed B. Erythromelalgia may occur.

Myelofibrosis

Obliteration of bone marrow with fibrosis C due to ↑ fibroblast activity. Associated with massive splenomegaly and "teardrop" RBCs D. "Bone marrow cries because it's fibrosed and is a dry tap."

	RBCs	WBCs	PLATELETS	PHILADELPHIA CHROMOSOME	*JAK2* MUTATIONS
Polycythemia vera	↑	↑	↑	⊖	⊕
Essential thrombocythemia	–	–	↑	⊖	⊕ (30–50%)
Myelofibrosis	↓	Variable	Variable	⊖	⊕ (30–50%)
CML	↓	↑	↑	⊕	⊖



(c) ketabton.com: The Digital Library

Polycythemia

	PLASMA VOLUME	RBC MASS	O_2 SATURATION	EPO LEVELS	ASSOCIATIONS
Relative	↓	–	–	–	Dehydration, burns.
Appropriate absolute	–	↑	↓	↑	Lung disease, congenital heart disease, high altitude.
Inappropriate absolute	–	↑	–	↑	Exogenous EPO: athlete abuse ("blood doping"). Inappropriate EPO secretion: malignancy (eg, renal cell carcinoma, hepatocellular carcinoma).
Polycythemia vera	↑	↑↑	–	↓	EPO ↓ in PCV due to negative feedback suppressing renal EPO production.

↑↓ = 1° disturbance

Chromosomal translocations

TRANSLOCATION	ASSOCIATED DISORDER	NOTES
t(8;14)	Burkitt (Burk-8) lymphoma (c-*myc* activation)	The Ig heavy chain genes on chromosome 14 are constitutively expressed. When other genes (eg, *c-myc* and *BCL-2*) are translocated next to this heavy chain gene region, they are overexpressed.
t(11;14)	Mantle cell lymphoma (cyclin D1 activation)	
t(11;18)	Marginal zone lymphoma	
t(14;18)	Follicular lymphoma (*BCL-2* activation)	
t(15;17)	APL (M3 type of AML; responds to all-trans retinoic acid)	
t(9;22) (Philadelphia chromosome)	CML (*BCR-ABL* hybrid), ALL (less common, poor prognostic factor); Philadelphia CreaML cheese	

Langerhans cell histiocytosis

Collective group of proliferative disorders of Langerhans cells. Presents in a child as lytic bone lesions A and skin rash or as recurrent otitis media with a mass involving the mastoid bone. Cells are functionally immature and do not effectively stimulate primary T cells via antigen presentation. Cells express S-100 (mesodermal origin) and CD1a. Birbeck granules ("tennis rackets" or rod shaped on EM) are characteristic B.



(c) ketabton.com: The Digital Library

Tumor lysis syndrome



Oncologic emergency triggered by massive tumor cell lysis, most often in lymphomas/leukemias. Release of K^+ → hyperkalemia, release of PO_4^{3-} → hyperphosphatemia, hypocalcemia due to Ca^{2+} sequestration by PO_4^{3-}. ↑ nucleic acid breakdown → hyperuricemia → acute kidney injury. Prevention and treatment include aggressive hydration, allopurinol, rasburicase.

Hemophagocytic lymphohistiocytosis



Systemic overactivation of macrophages and cytotoxic T cells → fever, pancytopenia, hepatosplenomegaly, ↑↑↑ serum ferritin levels. Can be inherited or 2° to strong immunologic activation (eg, after EBV infection, malignancy). Bone marrow biopsy shows macrophages phagocytosing marrow elements A.

▸ HEMATOLOGY AND ONCOLOGY—PHARMACOLOGY

Direct thrombin inhibitors

Bivalirudin, Argatroban, Dabigatran (only oral agent in class).

MECHANISM	Directly inhibits activity of free and clot-associated thrombin.
CLINICAL USE	Venous thromboembolism, atrial fibrillation. Can be used in HIT, when heparin is BAD for the patient. Does not require lab monitoring.
ADVERSE EFFECTS	Bleeding; can reverse dabigatran with idarucizumab. Consider PCC and/or antifibrinolytics (eg, tranexamic acid) if no reversal agent available.

(c) ketabton.com: The Digital Library

Heparin

MECHANISM	Activates antithrombin, which ↓ action of IIa (thrombin) and factor Xa. Short half-life.
CLINICAL USE	Immediate anticoagulation for pulmonary embolism (PE), acute coronary syndrome, MI, deep venous thrombosis (DVT). Used during pregnancy (does not cross placenta). Follow PTT.
ADVERSE EFFECTS	Bleeding, thrombocytopenia (HIT), osteoporosis, drug-drug interactions. For rapid reversal (antidote), use protamine sulfate (positively charged molecule that binds negatively charged heparin).
NOTES	Low-molecular-weight heparins (eg, enoxaparin, dalteparin)—act predominantly on factor Xa. Fondaparinux acts only on factor Xa. Have better bioavailability and 2–4× longer half life than unfractionated heparin; can be administered subcutaneously and without laboratory monitoring. LMWHs undergo renal clearance (vs hepatic clearance of unfractionated heparin) and are contraindicated in renal insufficiency. Not easily reversible. **Heparin-induced thrombocytopenia (HIT) type 2**—development of IgG antibodies against heparin-bound platelet factor 4 (PF4). Antibody-heparin-PF4 complex activates platelets → thrombosis and thrombocytopenia. Highest risk with unfractionated heparin. **HIT type 1** characterized by nonimmunologic milder drop in platelet count, usually asymptomatic.

Warfarin

MECHANISM	Inhibits epoxide reductase, which interferes with γ-carboxylation of vitamin K–dependent clotting factors II, VII, IX, X, and proteins C, S. Metabolism affected by polymorphisms in the gene for vitamin K epoxide reductase complex (*VKORC1*). In laboratory assay, has effect on **EX**trinsic pathway and ↑ **PT**. Long half-life.	The **EX-P**residen**T** went to **war**(farin).
CLINICAL USE	Chronic anticoagulation (eg, venous thromboembolism prophylaxis, and prevention of stroke in atrial fibrillation). Not used in pregnant women (because warfarin, unlike heparin, crosses placenta). Follow PT/INR.	
ADVERSE EFFECTS A	Bleeding, teratogenic, skin/tissue necrosis A, drug-drug interactions. Initial risk of hypercoagulation: protein C has a shorter half-life than factors II and X. Existing protein C depletes before existing factors II and X deplete, and before warfarin can reduce factors II and X production → hypercoagulation. Skin/tissue necrosis within first few days of large doses believed to be due to small vessel microthrombosis.	For reversal of warfarin, give vitamin K. For rapid reversal, give fresh frozen plasma (FFP) or PCC. Heparin "bridging": heparin frequently used when starting warfarin. Heparin's activation of antithrombin enables anticoagulation during initial, transient hypercoagulable state caused by warfarin. Initial heparin therapy reduces risk of recurrent venous thromboembolism and skin/tissue necrosis. Metabolized by cytochrome P-450.

(c) ketabton.com: The Digital Library

Heparin vs warfarin

	Heparin	Warfarin
ROUTE OF ADMINISTRATION	Parenteral (IV, SC)	Oral
SITE OF ACTION	Blood	Liver
ONSET OF ACTION	Rapid (seconds)	Slow, limited by half-lives of normal clotting factors
MECHANISM OF ACTION	Activates antithrombin, which ↓ the action of IIa (thrombin) and factor Xa	Impairs synthesis of vitamin K–dependent clotting factors II, VII, IX, and X, and anti-clotting proteins C and S
DURATION OF ACTION	Hours	Days
AGENTS FOR REVERSAL	Protamine sulfate	Vitamin K, FFP, PCC
MONITORING	PTT (intrinsic pathway)	PT/INR (extrinsic pathway)
CROSSES PLACENTA	No	Yes (teratogenic)

Direct factor Xa inhibitors	ApiXaban, rivaroXaban.
MECHANISM	Bind to and directly inhibit factor Xa.
CLINICAL USE	Treatment and prophylaxis for DVT and PE; stroke prophylaxis in patients with atrial fibrillation. Oral agents do not usually require coagulation monitoring.
ADVERSE EFFECTS	Bleeding. Reverse with andeXanet alfa.

Thrombolytics	Alteplase (tPA), reteplase (rPA), streptokinase, tenecteplase (TNK-tPA).
MECHANISM	Directly or indirectly aid conversion of plasminogen to plasmin, which cleaves thrombin and fibrin clots. ↑ PT, ↑ PTT, no change in platelet count.
CLINICAL USE	Early MI, early ischemic stroke, direct thrombolysis of severe PE.
ADVERSE EFFECTS	Bleeding. Contraindicated in patients with active bleeding, history of intracranial bleeding, recent surgery, known bleeding diatheses, or severe hypertension. Nonspecific reversal with antifibrinolytics (eg, aminocaproic acid, tranexamic acid), platelet transfusions, and factor corrections (eg, cryoprecipitate, FFP, PCC).

ADP receptor inhibitors	Clopidogrel, prasugrel, ticagrelor (reversible), ticlopidine.
MECHANISM	Irreversibly block ADP ($P2Y_{12}$) receptor, which prevents subsequent platelet aggregation. Prevent expression of glycoproteins IIb/IIIa on platelet surface.
CLINICAL USE	Acute coronary syndrome; coronary stenting. ↓ incidence or recurrence of thrombotic stroke.
ADVERSE EFFECTS	Neutropenia (ticlopidine). TTP may be seen.

(c) ketabton.com: The Digital Library

Glycoprotein IIb/IIIa inhibitors	Abciximab, eptifibatide, tirofiban.
MECHANISM	Bind to the glycoprotein receptor IIb/IIIa (fibrinogen receptor) on activated platelets, preventing aggregation. Abciximab is made from monoclonal antibody Fab fragments.
CLINICAL USE	Unstable angina, percutaneous coronary intervention.
ADVERSE EFFECTS	Bleeding, thrombocytopenia.

Cancer therapy—cell cycle



Cancer therapy—targets



FAS1_2019_10-HemaOncol.indd 438 11/7/19 5:05 PM

(c) ketabton.com: The Digital Library

Antitumor antibiotics

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Bleomycin	Induces free radical formation → breaks in DNA strands.	Testicular cancer, Hodgkin lymphoma.	Pulmonary fibrosis, skin hyperpigmentation. Minimal myelosuppression.
Dactinomycin (actinomycin D)	Intercalates into DNA, preventing RNA synthesis.	Wilms tumor, Ewing sarcoma, rhabdomyosarcoma. Used for childhood tumors.	Myelosuppression.
Anthracyclines (eg, doxorubicin, daunorubicin)	Generate free radicals. Intercalate in DNA → breaks in DNA → ↓ replication. Interferes with topoisomerase II enzyme.	Solid tumors, leukemias, lymphomas.	Cardiotoxicity (dilated cardiomyopathy), myelosuppression, alopecia. Dexrazoxane (iron chelating agent) used to prevent cardiotoxicity.

(c) ketabton.com: The Digital Library

Antimetabolites

DRUG	MECHANISM[a]	CLINICAL USE	ADVERSE EFFECTS
Azathioprine, 6-mercaptopurine	Purine (thiol) analogs → ↓ de novo purine synthesis. Activated by HGPRT. Azathioprine is metabolized into 6-MP.	Preventing organ rejection, rheumatoid arthritis, IBD, SLE; used to wean patients off steroids in chronic disease and to treat steroid-refractory chronic disease.	Myelosuppression; GI, liver toxicity. Azathioprine and 6-MP are metabolized by xanthine oxidase; thus both have ↑ risk of toxicity with allopurinol or febuxostat.
Cladribine	Purine analog → multiple mechanisms (eg, inhibition of DNA polymerase, DNA strand breaks).	Hairy cell leukemia.	Myelosuppression, nephrotoxicity, and neurotoxicity.
Cytarabine (arabinofuranosyl cytidine)	Pyrimidine analog → DNA chain termination. At higher concentrations, inhibits DNA polymerase.	Leukemias (AML), lymphomas.	Myelosuppression with megaloblastic anemia. CYTarabine causes panCYTopenia.
5-fluorouracil	Pyrimidine analog bioactivated to 5-FdUMP, which covalently complexes with thymidylate synthase and folic acid. Capecitabine is a prodrug. This complex inhibits thymidylate synthase → ↓ dTMP → ↓ DNA synthesis.	Colon cancer, pancreatic cancer, actinic keratosis, basal cell carcinoma (topical). Effects enhanced with the addition of leucovorin.	Myelosuppression, palmar-plantar erythrodysesthesia (hand-foot syndrome).
Methotrexate	Folic acid analog that competitively inhibits dihydrofolate reductase → ↓ dTMP → ↓ DNA synthesis.	Cancers: leukemias (ALL), lymphomas, choriocarcinoma, sarcomas. Non-neoplastic: ectopic pregnancy, medical abortion (with misoprostol), rheumatoid arthritis, psoriasis, IBD, vasculitis.	Myelosuppression, which is reversible with leucovorin (folinic acid) "rescue." Hepatotoxicity. Mucositis (eg, mouth ulcers). Pulmonary fibrosis. Folate deficiency, which may be teratogenic (neural tube defects) without supplementation. Nephrotoxicity.

[a]All are S-phase specific except cladribine, which is cell cycle nonspecific.

(c) ketabton.com: The Digital Library

Alkylating agents

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Busulfan	Cross-links DNA.	Used to ablate patient's bone marrow before bone marrow transplantation.	Severe myelosuppression (in almost all cases), pulmonary fibrosis, hyperpigmentation.
Cyclophosphamide, ifosfamide	Cross-link DNA at guanine. Require bioactivation by liver. A nitrogen mustard.	Solid tumors, leukemia, lymphomas, rheumatic disease (eg, SLE, granulomatosis with polyangiitis).	Myelosuppression; SIADH; Fanconi syndrome (ifosfamide); hemorrhagic cystitis and bladder cancer, prevented with mesna (sulfhydryl group of mesna binds toxic metabolites) and adequate hydration.
Nitrosoureas (eg, carmustine, lomustine)	Require bioactivation. Cross blood-brain barrier → CNS. Cross-link DNA.	Brain tumors (including glioblastoma multiforme).	CNS toxicity (convulsions, dizziness, ataxia).
Procarbazine	Cell cycle phase–nonspecific alkylating agent, mechanism unknown. Also a weak MAO inhibitor.	Hodgkin lymphoma, brain tumors.	Bone marrow suppression, pulmonary toxicity, leukemia, disulfiram-like reaction, tyramine-induced hypertensive crisis with consumption of tyramine-rich foods (eg, aged cheese, wine, fava beans).

Microtubule inhibitors

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Paclitaxel, other taxanes	Hyperstabilize polymerized microtubules in M phase so that mitotic spindle cannot break down (anaphase cannot occur).	Ovarian and breast carcinomas.	Myelosuppression, neuropathy, hypersensitivity. Taxes stabilize society.
Vincristine, vinblastine	Vinca alkaloids that bind β-tubulin and inhibit its polymerization into microtubules → prevent mitotic spindle formation (M-phase arrest).	Solid tumors, leukemias, Hodgkin and non-Hodgkin lymphomas.	Vincristine: neurotoxicity (areflexia, peripheral neuritis), constipation (including paralytic ileus). Crisps the nerves. Vinblastine: bone marrow suppression. Blasts the bone marrow.

(c) ketabton.com: The Digital Library

Cisplatin, carboplatin, oxaliplatin

MECHANISM	Cross-link DNA.
CLINICAL USE	Testicular, bladder, ovary, GI, and lung carcinomas.
ADVERSE EFFECTS	Nephrotoxicity (including Fanconi syndrome), peripheral neuropathy, ototoxicity. Prevent nephrotoxicity with amifostine (free radical scavenger) and chloride (saline) diuresis.

Etoposide, teniposide

MECHANISM	Inhibit topoisomerase II → ↑ DNA degradation (cell cycle arrest in G_2 and S phases).
CLINICAL USE	Solid tumors (particularly testicular and small cell lung cancer), leukemias, lymphomas.
ADVERSE EFFECTS	Myelosuppression, alopecia.

Irinotecan, topotecan

MECHANISM	Inhibit topoisomerase I and prevent DNA unwinding and replication.
CLINICAL USE	Colon cancer (irinotecan); ovarian and small cell lung cancers (topotecan).
ADVERSE EFFECTS	Severe myelosuppression, diarrhea.

Hydroxyurea

MECHANISM	Inhibits ribonucleotide reductase → ↓ DNA **S**ynthesis (**S**-phase specific).
CLINICAL USE	Myeloproliferative disorders (eg, CML, polycythemia vera), sickle cell disease (↑ HbF).
ADVERSE EFFECTS	Severe myelosuppression, megaloblastic anemia.

Bevacizumab

MECHANISM	Monoclonal antibody against VEGF. Inhibits angiogenesis (**Be**-**V**acizumab inhibits **B**lood **V**essel formation).
CLINICAL USE	Solid tumors (eg, colorectal cancer, renal cell carcinoma), wet age-related macular degeneration.
ADVERSE EFFECTS	Hemorrhage, blood clots, and impaired wound healing.

Erlotinib

MECHANISM	EGFR tyrosine kinase inhibitor.
CLINICAL USE	Non-small cell lung cancer.
ADVERSE EFFECTS	Rash, diarrhea.

Cetuximab, panitumumab

MECHANISM	Monoclonal antibodies against EGFR.
CLINICAL USE	Stage IV colorectal cancer (wild-type *KRAS*), head and neck cancer.
ADVERSE EFFECTS	Rash, elevated LFTs, diarrhea.

(c) ketabton.com: The Digital Library

Imatinib, dasatinib, nilotinib

MECHANISM	Tyrosine kinase inhibitors of bcr-abl (encoded by Philadelphia chromosome fusion gene in CML) and c-*kit* (common in GI stromal tumors).
CLINICAL USE	CML, GI stromal tumors (GISTs).
ADVERSE EFFECTS	Fluid retention.

Rituximab

MECHANISM	Monoclonal antibody against CD20, which is found on most B-cell neoplasms.
CLINICAL USE	Non-Hodgkin lymphoma, CLL, ITP, rheumatoid arthritis, TTP, AIHA.
ADVERSE EFFECTS	↑ risk of progressive multifocal leukoencephalopathy.

Bortezomib, carfilzomib

MECHANISM	Proteasome inhibitors, induce arrest at G2-M phase and apoptosis.
CLINICAL USE	Multiple myeloma, mantle cell lymphoma.
ADVERSE EFFECTS	Peripheral neuropathy, herpes zoster reactivation.

Tamoxifen, raloxifene

MECHANISM	Selective estrogen receptor modulators (SERMs)—receptor antagonists in breast and agonists in bone. Block the binding of estrogen to ER ⊕ cells.
CLINICAL USE	Breast cancer treatment (tamoxifen only) and prevention. Raloxifene also useful to prevent osteoporosis.
ADVERSE EFFECTS	Tamoxifen—partial agonist in endometrium, which ↑ the risk of endometrial cancer. **Raloxifene**—no ↑ in endometrial carcinoma (so you can **relax**!), because it is an estrogen receptor antagonist in endometrial tissue. Both ↑ risk of thromboembolic events (eg, DVT, PE) and "hot flashes."

Trastuzumab

MECHANISM	Monoclonal antibody against HER-2 *(c-erbB2)*, a tyrosine kinase receptor. Helps kill cancer cells that overexpress HER-2 through inhibition of HER-2 initiated cellular signaling and antibody-dependent cytotoxicity.
CLINICAL USE	HER-**2** ⊕ breast cancer and gastric cancer (tras**2**zumab).
ADVERSE EFFECTS	Dilated cardiomyopathy. "**Heart**ceptin" damages the **heart**.

(c) ketabton.com: The Digital Library

Dabrafenib, vemurafenib

MECHANISM	Small molecule inhibitors of *BRAF* oncogene ⊕ melanoma. **VEmuRAF-enib** is for **V**600**E**-**mu**tated ***BRAF* inhib**ition. Often co-administered with MEK inhibitors (eg, trametinib).
CLINICAL USE	Metastatic melanoma.

Rasburicase

MECHANISM	Recombinant uricase that catalyzes metabolism of uric acid to allantoin.
CLINICAL USE	Prevention and treatment of tumor lysis syndrome.

Key chemotoxicities



Cisplatin/Carboplatin → ototoxicity

Vincristine → peripheral neuropathy
Bleomycin, Busulfan → pulmonary fibrosis
Doxorubicin → cardiotoxicity
Trastuzumab → cardiotoxicity
Cisplatin/Carboplatin → nephrotoxicity

CYclophosphamide → hemorrhagic cystitis

Nonspecific common toxicities of nearly all cytotoxic chemotherapies include myelosuppression (neutropenia, anemia, thrombocytopenia), GI toxicity (nausea, vomiting, mucositis), alopecia.

(c) ketabton.com: The Digital Library

Musculoskeletal, Skin, and Connective Tissue

"Rigid, the skeleton of habit alone upholds the human frame."
—Virginia Woolf

"Beauty may be skin deep, but ugly goes clear to the bone."
—Redd Foxx

"The function of muscle is to pull and not to push, except in the case of the genitals and the tongue."
—Leonardo da Vinci

"To thrive in life you need three bones. A wishbone. A backbone. And a funny bone."
—Reba McEntire

This chapter provides information you will need to understand certain anatomical dysfunctions, rheumatic diseases, and dermatologic conditions. Be able to interpret 3D anatomy in the context of radiologic imaging. For the rheumatic diseases, create instructional cases or personas that include the most likely presentation and symptoms: risk factors, gender, important markers (eg, autoantibodies), and other epidemiologic factors. Doing so will allow you to answer the higher order questions that are likely to be asked on the exam.

- Anatomy and Physiology 446
- Pathology 459
- Dermatology 473
- Pharmacology 485

FAS1_2019_11-Musculo.indd 445 11/7/19 5:23 PM

(c) ketabton.com: The Digital Library

Rotator cuff muscles



Shoulder muscles that form the rotator cuff:

- **S**upraspinatus (suprascapular nerve)—abducts arm initially (before the action of the deltoid); most common rotator cuff injury (trauma or degeneration and impingement → tendinopathy or tear [arrow in A]), assessed by "empty/full can" test
- **I**nfraspinatus (suprascapular nerve)—externally rotates arm; pitching injury
- **t**eres minor (axillary nerve)—adducts and externally rotates arm
- **S**ubscapularis (upper and lower subscapular nerves)—internally rotates and adducts arm

Innervated primarily by C5-C6.

SItS (**small** t is for teres **minor**).



Arm abduction

DEGREE	MUSCLE	NERVE
0°–15°	Supraspinatus	Suprascapular
15°–100°	Deltoid	Axillary
> 90°	Trapezius	Accessory
> 100°	**S**erratus **A**nterior	**L**ong **T**horacic (**SALT**)

(c) ketabton.com: The Digital Library

Upper extremity nerves

NERVE	CAUSES OF INJURY	PRESENTATION
Axillary (C5-C6)	Fractured surgical neck of humerus Anterior dislocation of humerus	Flattened deltoid Loss of arm abduction at shoulder (> 15°) Loss of sensation over deltoid and lateral arm
Musculocutaneous (C5-C7)	Upper trunk compression	↓ biceps (C5-6) reflex Weakness of forearm flexion and supination Loss of sensation over lateral forearm
Radial (C5-T1)	Compression of axilla, eg, due to crutches or sleeping with arm over chair ("Saturday night palsy") Midshaft fracture of humerus Repetitive pronation/supination of forearm, eg, due to screwdriver use ("finger drop")	Wrist drop: loss of elbow, wrist, and finger extension ↓ grip strength (wrist extension necessary for maximal action of flexors) Loss of sensation over posterior arm/forearm and dorsal hand
Median (C5-T1)	Supracondylar fracture of humerus → proximal lesion of the nerve Carpal tunnel syndrome and wrist laceration → distal lesion of the nerve	"Ape hand" and "Pope's blessing" Loss of wrist flexion, flexion of lateral fingers, thumb opposition, lumbricals of index and middle fingers Loss of sensation over thenar eminence and dorsal and palmar aspects of lateral 3½ fingers with proximal lesion
Ulnar (C8-T1)	Fracture of medial epicondyle of humerus "funny bone" (proximal lesion) Fractured hook of hamate (distal lesion) from fall on outstretched hand	"Ulnar claw" on digit extension Radial deviation of wrist upon flexion (proximal lesion) Loss of wrist flexion, flexion of medial fingers, abduction and adduction of fingers (interossei), actions of medial 2 lumbrical muscles Loss of sensation over medial 1½ fingers including hypothenar eminence
Recurrent branch of median nerve (C5-T1)	Superficial laceration of palm	"Ape hand" Loss of thenar muscle group: opposition, abduction, and flexion of thumb No loss of sensation

Humerus fractures, proximally to distally, follow the **ARM** (**A**xillary → **R**adial → **M**edian)



(c) ketabton.com: The Digital Library

Brachial plexus lesions

1. Erb palsy ("waiter's tip")
2. Klumpke palsy (claw hand)
3. Wrist drop
4. Winged scapula
5. Deltoid paralysis
6. "Saturday night palsy" (wrist drop)
7. Difficulty flexing elbow, variable sensory loss
8. Decreased thumb function, "Pope's blessing"
9. Intrinsic muscles of hand, claw hand



Randy Travis Drinks Cold Beer

CONDITION	INJURY	CAUSES	MUSCLE DEFICIT	FUNCTIONAL DEFICIT	PRESENTATION
Erb palsy ("waiter's tip")	Traction or tear of **upper** trunk: C5-C6 roots	Infants—lateral traction on neck during delivery Adults—trauma	Deltoid, supraspinatus	Abduction (arm hangs by side)	
			Infraspinatus	Lateral rotation (arm medially rotated)	
			Biceps brachii Herb gets DIBs on tips	Flexion, supination (arm extended and pronated)	
Klumpke palsy	Traction or tear of **lower** trunk: C8-T1 roots	Infants—upward force on arm during delivery Adults—trauma (eg, grabbing a tree branch to break a fall)	Intrinsic hand muscles: lumbricals, interossei, thenar, hypothenar	Total claw hand: lumbricals normally flex MCP joints and extend DIP and PIP joints	
Thoracic outlet syndrome	Compression of **lower** trunk and subclavian vessels, most commonly within the scalene triangle	Cervical rib (arrows in A), Pancoast tumor	Same as Klumpke palsy	Atrophy of intrinsic hand muscles; ischemia, pain, and edema due to vascular compression	
Winged scapula	Lesion of long thoracic nerve, roots C5-C7 ("wings of heaven")	Axillary node dissection after mastectomy, stab wounds	Serratus anterior	Inability to anchor scapula to thoracic cage → cannot abduct arm above horizontal position B	

(c) ketabton.com: The Digital Library

Wrist region

Scaphoid, Lunate, Triquetrum, Pisiform, Hamate, Capitate, Trapezoid, Trapezium A. (So Long To Pinky, Here Comes The Thumb)

Scaphoid (palpable in anatomic snuff box B) is the most commonly fractured carpal bone, typically due to a fall on an outstretched hand. Complications of proximal scaphoid fractures include avascular necrosis and nonunion due to retrograde blood supply from a branch of the radial artery. Fracture not always seen on initial x-ray.

Dislocation of lunate may cause acute carpal tunnel syndrome.





(c) ketabton.com: The Digital Library

Hand muscles



Thenar (median)—**O**pponens pollicis, **A**bductor pollicis brevis, **F**lexor pollicis brevis, superficial head (deep head by ulnar nerve).

Hypothenar (ulnar)—**O**pponens digiti minimi, **A**bductor digiti minimi, **F**lexor digiti minimi brevis.

Dorsal interossei (ulnar)—abduct the fingers.

Palmar interossei (ulnar)—adduct the fingers.

Lumbricals (1st/2nd, median; 3rd/4th, ulnar)—flex at the MCP joint, extend PIP and DIP joints.

Both groups perform the same functions: **O**ppose, **A**bduct, and **F**lex (**OAF**).

DAB = **D**orsals **AB**duct.
PAD = **P**almars **AD**duct.

(c) ketabton.com: The Digital Library

Distortions of the hand

At rest, a balance exists between the extrinsic flexors and extensors of the hand, as well as the intrinsic muscles of the hand—particularly the lumbrical muscles (flexion of MCP, extension of DIP and PIP joints).

"Clawing"—seen best with **distal** lesions of median or ulnar nerves. Remaining extrinsic flexors of the digits exaggerate the loss of the lumbricals → fingers extend at MCP, flex at DIP and PIP joints.

Deficits less pronounced in **proximal** lesions; deficits present during voluntary flexion of the digits.

SIGN	"Ulnar claw"	"Pope's blessing"	"Median claw"	"OK gesture"
PRESENTATION				
CONTEXT	Extending fingers/at rest	Making a fist	Extending fingers/at rest	Making a fist
LOCATION OF LESION	Distal ulnar nerve	Proximal median nerve	Distal median nerve	Proximal ulnar nerve

Note: Atrophy of the thenar eminence (unopposable thumb → "ape hand") can be seen in median nerve lesions, while atrophy of the hypothenar eminence can be seen in ulnar nerve lesions.

Actions of hip muscles

ACTION	MUSCLES
Abductors	Gluteus medius, gluteus minimus
Adductors	Adductor magnus, adductor longus, adductor brevis
Extensors	Gluteus maximus, semitendinosus, semimembranosus
Flexors	Iliopsoas, rectus femoris, tensor fascia lata, pectineus, sartorius
Internal rotation	Gluteus medius, gluteus minimus, tensor fascia latae
External rotation	Iliopsoas, gluteus maximus, piriformis, obturator

(c) ketabton.com: The Digital Library

Lower extremity nerves

NERVE	INNERVATION	CAUSE OF INJURY	PRESENTATION/COMMENTS
Iliohypogastric (T12-L1)	Sensory—suprapubic region Motor—transversus abdominis and internal oblique	Abdominal surgery	Burning or tingling pain in surgical incision site radiating to inguinal and suprapubic region
Genitofemoral nerve (L1-L2)	Sensory—scrotum/labia majora, medial thigh Motor—cremaster	Laparoscopic surgery	↓ upper medial thigh and anterior thigh sensation beneath the inguinal ligament (lateral part of the femoral triangle); absent cremasteric reflex
Lateral femoral cutaneous (L2-L3)	Sensory—anterior and lateral thigh	Tight clothing, obesity, pregnancy, pelvic procedures	↓ thigh sensation (anterior and lateral)
Obturator (L2-L4)	Sensory—medial thigh Motor—obturator externus, adductor longus, adductor brevis, gracilis, pectineus, adductor magnus	Pelvic surgery	↓ thigh sensation (medial) and adduction
Femoral (L2-L4) 	Sensory—anterior thigh, medial leg Motor—quadriceps, iliacus, pectineus, sartorius	Pelvic fracture	↓ leg extension (↓ patellar reflex)
Sciatic (L4-S3)	Motor—semitendinosus, semimembranosus, biceps femoris, adductor magnus	Herniated disc, posterior hip dislocation	Splits into common peroneal and tibial nerves

(c) ketabton.com: The Digital Library

Lower extremity nerves *(continued)*

NERVE	INNERVATION	CAUSE OF INJURY	PRESENTATION/COMMENTS
Common (fibular) peroneal (L4-S2) 	Superficial peroneal nerve: ▪ Sensory—dorsum of foot (except webspace between hallux and 2nd digit) ▪ Motor—peroneus longus and brevis Deep peroneal nerve: ▪ Sensory—webspace between hallux and 2nd digit ▪ Motor—tibialis anterior	Trauma or compression of lateral aspect of leg, fibular neck fracture	PED = Peroneal Everts and Dorsiflexes; if injured, foot dropPED Loss of sensation on dorsum of foot Foot drop—inverted and plantarflexed at rest, loss of eversion and dorsiflexion; "steppage gait"
Tibial (L4-S3) 	Sensory—sole of foot Motor—biceps femoris (long head), triceps surae, plantaris, popliteus, flexor muscles of foot	Knee trauma, Baker cyst (proximal lesion); tarsal tunnel syndrome (distal lesion)	TIP = Tibial Inverts and Plantarflexes; if injured, can't stand on TIPtoes Inability to curl toes and loss of sensation on sole; in proximal lesions, foot everted at rest with loss of inversion and plantar flexion
Superior gluteal (L4-S1) 	Motor—gluteus medius, gluteus minimus, tensor fascia latae	Iatrogenic injury during intramuscular injection to superomedial gluteal region (prevent by choosing superolateral quadrant, preferably anterolateral region)	Trendelenburg sign/gait—pelvis tilts because weight-bearing leg cannot maintain alignment of pelvis through hip abduction Lesion is contralateral to the side of the hip that drops, ipsilateral to extremity on which the patient stands
Inferior gluteal (L5-S2)	Motor—gluteus maximus	Posterior hip dislocation	Difficulty climbing stairs, rising from seated position; loss of hip extension
Pudendal (S2-S4)	Sensory—perineum Motor—external urethral and anal sphincters	Stretch injury during childbirth, prolonged cycling, horseback riding	↓ sensation in perineum and genital area; can cause fecal and/or urinary incontinence Can be blocked with local anesthetic during childbirth using ischial spine as a landmark for injection

(c) ketabton.com: The Digital Library

Knee exam

Lateral femoral condyle to anterior tibia: **A**CL.
Medial femoral condyle to posterior tibia: **P**CL.
LAMP.



TEST	PROCEDURE	
Anterior drawer sign	Bending knee at 90° angle, ↑ anterior gliding of tibia (relative to femur) due to ACL injury Lachman test also tests ACL, but is more sensitive (↑ anterior gliding of tibia [relative to femur] with knee bent at 30° angle)	
Posterior drawer sign	Bending knee at 90° angle, ↑ posterior gliding of tibia due to PCL injury	
Abnormal passive abduction	Knee either extended or at ~ 30° angle, lateral (valgus) force → medial space widening of tibia → MCL injury	
Abnormal passive adduction	Knee either extended or at ~ 30° angle, medial (varus) force → lateral space widening of tibia → LCL injury	
McMurray test	During flexion and extension of knee with rotation of tibia/foot (**LIME**): ▪ Pain, "popping" on internal rotation and varus force → **L**ateral meniscal tear (**I**nternal rotation stresses lateral meniscus) ▪ Pain, "popping" on external rotation and valgus force → **M**edial meniscal tear (**E**xternal rotation stresses medial meniscus)	

(c) ketabton.com: The Digital Library

Ankle sprains

Anterior **T**alo**F**ibular ligament—most common ankle sprain overall, classified as a low ankle sprain. Due to overinversion/supination of foot.
Anterior inferior tibiofibular ligament—most common high ankle sprain.
Always **T**ears **F**irst.



Signs of lumbosacral radiculopathy

Paresthesia and weakness related to specific lumbosacral spinal nerves. Intervertebral disc (nucleus pulposus) herniates posterolaterally through annulus fibrosus (outer ring) into central canal due to thin posterior longitudinal ligament and thicker anterior longitudinal ligament along midline of vertebral bodies. Nerve affected is usually below the level of herniation.

Disc level herniation	L3-L4	L4-L5	L5-S1
Nerve root affected	L4	L5	S1
Dermatome affected			
Clinical findings	Weakness of knee extension ↓ patellar reflex	Weakness of dorsiflexion Difficulty in heel walking	Weakness of plantar flexion Difficulty in toe walking ↓ Achilles reflex



Neurovascular pairing

Nerves and arteries are frequently named together by the bones/regions with which they are associated. The following are exceptions to this naming convention.

LOCATION	NERVE	ARTERY
Axilla/lateral thorax	Long thoracic	Lateral thoracic
Surgical neck of humerus	Axillary	Posterior circumflex
Midshaft of humerus	Radial	Deep brachial
Distal humerus/cubital fossa	Median	Brachial
Popliteal fossa	Tibial	Popliteal
Posterior to medial malleolus	Tibial	Posterior tibial

(c) ketabton.com: The Digital Library

Motoneuron action potential to muscle contraction

T-tubules are extensions of plasma membrane in contact with the sarcoplasmic reticulum, allowing for coordinated contraction of striated muscles.





1. Action potential opens presynaptic voltage-gated Ca^{2+} channels, inducing acetylcholine (ACh) release.
2. Postsynaptic ACh binding leads to muscle cell depolarization at the motor end plate.
3. Depolarization travels over the entire muscle cell and deep into the muscle via the T-tubules.
4. Membrane depolarization induces conformational changes in the voltage-sensitive dihydropyridine receptor (DHPR) and its mechanically coupled ryanodine receptor (RR) → Ca^{2+} release from the sarcoplasmic reticulum into the cytoplasm.
5. Tropomyosin is blocking myosin-binding sites on the actin filament. Released Ca^{2+} binds to troponin C (TnC), shifting tropomyosin to expose the myosin-binding sites.
6. The myosin head binds strongly to actin, forming a crossbridge. P_i is then released, initiating the power stroke.
7. During the power stroke, force is produced as myosin pulls on the thin filament **A**. Muscle shortening occurs, with shortening of **H** and **I** bands and between **Z** lines (**HIZ** shrinkage). The A band remains the same length (**A** band is **A**lways the same length). ADP is released at the end of the power stroke.
8. Binding of new ATP molecule causes detachment of myosin head from actin filament. Ca^{2+} is resequestered.
9. ATP hydrolysis into ADP and P_i results in myosin head returning to high-energy position (cocked). The myosin head can bind to a new site on actin to form a crossbridge if Ca^{2+} remains available.

FAS1_2019_11-Musculo.indd 456 11/7/19 5:23 PM

(c) ketabton.com: The Digital Library

Types of muscle fibers

	Type I	Type II
CONTRACTION VELOCITY	Slow	Fast
FIBER COLOR	Red	White
PREDOMINANT METABOLISM	Oxidative phosphorylation → sustained contraction	Anaerobic glycolysis
MITOCHONDRIA, MYOGLOBIN	↑	↓
TYPE OF TRAINING	Endurance training	Weight/resistance training, sprinting
NOTES	Think "1 slow red ox"	

Vascular smooth muscle contraction and relaxation



(c) ketabton.com: The Digital Library

Muscle proprioceptors

Specialized sensory receptors that relay information about muscle dynamics.

	Muscle spindle	Golgi tendon organ
PATHWAY	❶ ↑ length and speed of stretch → ❷ via dorsal root ganglion (DRG) → ❸ activation of inhibitory interneuron and α motor neuron → ❹ simultaneous inhibition of antagonist muscle (prevents overstretching) and activation of agonist muscle (contraction).	❶ ↑ tension → ❷ via DRG → ❸ activation of inhibitory interneuron → ❹ inhibition of agonist muscle (reduced tension within muscle and tendon)
LOCATION	Body of muscle/type Ia and II sensory axons	Tendons/type Ib sensory axons
ACTIVATION BY	↑ muscle stretch	↑ muscle force



Bone formation

Endochondral ossification	Bones of axial skeleton, appendicular skeleton, and base of skull. Cartilaginous model of bone is first made by chondrocytes. Osteoclasts and osteoblasts later replace with woven bone and then remodel to lamellar bone. In adults, woven bone occurs after fractures and in Paget disease. Defective in achondroplasia.
Membranous ossification	Bones of calvarium, facial bones, and clavicle. Woven bone formed directly without cartilage. Later remodeled to lamellar bone.

(c) ketabton.com: The Digital Library

Cell biology of bone

Osteoblast	Builds bone by secreting collagen and catalyzing mineralization in alkaline environment via ALP. Differentiates from mesenchymal stem cells in periosteum. Osteoblastic activity measured by bone ALP, osteocalcin, propeptides of type I procollagen.
Osteoclast	Dissolves ("crushes") bone by secreting H^+ and collagenases. Differentiates from a fusion of monocyte/macrophage lineage precursors. RANK receptors on osteoclasts are stimulated by RANKL (RANK ligand, expressed on osteoblasts). OPG (osteoprotegerin, a RANKL decoy receptor) binds RANKL to prevent RANK-RANKL interaction → ↓ osteoclast activity.
Parathyroid hormone	At low, intermittent levels, exerts anabolic effects (building bone) on osteoblasts and osteoclasts (indirect). Chronically ↑ PTH levels (1° hyperparathyroidism) cause catabolic effects (osteitis fibrosa cystica).
Estrogen	Inhibits apoptosis in bone-forming osteoblasts and induces apoptosis in bone-resorbing osteoclasts. Causes closure of epiphyseal plate during puberty. Estrogen deficiency (surgical or postmenopausal) → ↑ cycles of remodeling and bone resorption → ↑ risk of osteoporosis.

▸ MUSCULOSKELETAL, SKIN, AND CONNECTIVE TISSUE—PATHOLOGY

Overuse injuries of the elbow

Medial epicondylitis (golfer's elbow)	Repetitive flexion (forehand shots) or idiopathic → pain near medial epicondyle.
Lateral epicondylitis (tennis elbow)	Repetitive extension (backhand shots) or idiopathic → pain near lateral epicondyle.

Wrist and hand injuries

Metacarpal neck fracture	Also called boxer's fracture. Common fracture caused by direct blow with a closed fist (eg, from punching a wall). Most commonly seen in 4th and 5th metacarpals A.	
Carpal tunnel syndrome	Entrapment of median nerve in carpal tunnel (between transverse carpal ligament and carpal bones) → nerve compression → paresthesia, pain, and numbness in distribution of median nerve. Thenar eminence atrophies B but sensation spared, because palmar cutaneous branch enters hand external to carpal tunnel.	Suggested by ⊕ Tinel sign (percussion of wrist causes tingling) and Phalen maneuver (90° flexion of wrist causes tingling). Associated with pregnancy (due to edema), rheumatoid arthritis, hypothyroidism, diabetes, acromegaly, dialysis-related amyloidosis; may be associated with repetitive use.
Guyon canal syndrome	Compression of ulnar nerve at wrist. Classically seen in cyclists due to pressure from handlebars.	

(c) ketabton.com: The Digital Library.

Clavicle fractures

Common in children and as birth trauma. Usually caused by a fall on outstretched hand or by direct trauma to shoulder. Weakest point at the junction of middle and lateral thirds; fractures at the middle third segment are most common. Presents as shoulder drop, shortened clavicle (lateral fragment is depressed due to arm weight and medially rotated by arm adductors [eg, pectoralis major]).

Common hip and knee conditions

"Unhappy triad"

Common injury in contact sports due to lateral force applied to a planted foot. Consists of damage to the ACL A, MCL, and medial meniscus (attached to MCL). However, lateral meniscus involvement is more common than medial meniscus involvement in conjunction with ACL and MCL injury. Presents with acute pain and signs of joint instability.



Anterior view of left knee

Prepatellar bursitis

Inflammation of the prepatellar bursa in front of the kneecap (red arrow in B). Can be caused by repeated trauma or pressure from excessive kneeling (also called "housemaid's knee").

Baker cyst

Popliteal fluid collection (red arrow in C) in gastrocnemius-semimembranosus bursa commonly communicating with synovial space and related to chronic joint disease (eg, osteoarthritis, rheumatoid arthritis).



(c) ketabton.com: The Digital Library

Common musculoskeletal conditions

De Quervain tenosynovitis	Noninflammatory thickening of abductor pollicis longus and extensor pollicis brevis tendons → pain or tenderness at radial styloid. ⊕ Finkelstein test (pain at radial styloid with active or passive stretch of thumb tendons). ↑ risk in new mothers, golfers, racquet sport players, "thumb" texters.
Ganglion cyst	Fluid-filled swelling overlying joint or tendon sheath, most commonly at dorsal side of wrist. Arises from herniation of dense connective tissue.
Iliotibial band syndrome	Overuse injury of lateral knee that occurs primarily in runners. Pain develops 2° to friction of iliotibial band against lateral femoral epicondyle.
Limb compartment syndrome	↑ pressure within fascial compartment of a limb → venous outflow obstruction and arteriolar collapse → anoxia and necrosis. Causes include significant long bone fractures, reperfusion injury, animal venoms. Presents with severe pain and tense, swollen compartments with passive stretch of muscles in the affected compartment. Motor deficits are late sign of irreversible muscle and nerve damage.
Medial tibial stress syndrome	Also called shin splints. Common cause of shin pain and diffuse tenderness in runners and military recruits. Caused by bone resorption that outpaces bone formation in tibial cortex.
Plantar fasciitis	Inflammation of plantar aponeurosis characterized by heel pain (worse with first steps in the morning or after period of inactivity) and tenderness.

Childhood musculoskeletal conditions

Developmental dysplasia of the hip	Abnormal acetabulum development in newborns. Major risk factor includes breech presentation. Results in hip instability/dislocation. Commonly tested with Ortolani and Barlow maneuvers (manipulation of newborn hip reveals a "clunk"). Confirmed via ultrasound (x-ray not used until ~4–6 months because cartilage is not ossified).
Legg-Calvé-Perthes disease	Idiopathic avascular necrosis of femoral head. Commonly presents between 5–7 years with insidious onset of hip pain that may cause child to limp. More common in males (4:1 ratio). Initial x-ray often normal.
Osgood-Schlatter disease 	Also called traction apophysitis. Overuse injury caused by repetitive strain and chronic avulsion of the secondary ossification center of proximal tibial tubercle. Occurs in adolescents after growth spurt. Common in running and jumping athletes. Presents with progressive anterior knee pain.
Patellofemoral syndrome	Overuse injury that commonly presents in young, female athletes as anterior knee pain. Exacerbated by prolonged sitting or weight-bearing on a flexed knee. Treatment: NSAIDs, thigh muscle strengthening.
Radial head subluxation	Also called nursemaid's elbow. Common elbow injury in children < 5 years. Caused by a sudden pull on the arm → immature annular ligament slips over head of radius. Injured arm held in extended/slightly flexed and pronated position.
Slipped capital femoral epiphysis	Classically presents in an obese young adolescent with hip/knee pain and altered gait. Increased axial force on femoral head → epiphysis displaces relative to femoral neck (like a scoop of ice cream slipping off a cone). Diagnosed via x-ray.

(c) ketabton.com: The Digital Library

Common pediatric fractures

Greenstick fracture — Incomplete fracture extending partway through width of bone A following bending stress; bone fails on tension side; compression side intact (compare to torus fracture). Bone is bent like a green twig.

Torus (buckle) fracture — Axial force applied to immature bone → cortex buckles on compression (concave) side and fractures B. Tension (convex) side remains solid (intact).



Achondroplasia

Failure of longitudinal bone growth (endochondral ossification) → short limbs. Membranous ossification is not affected → large head relative to limbs. Constitutive activation of fibroblast growth factor receptor (FGFR3) actually inhibits chondrocyte proliferation. > 85% of mutations occur sporadically; autosomal dominant with full penetrance (homozygosity is lethal). Associated with ↑ paternal age. Most common cause of short-limbed dwarfism.

Osteoporosis



Trabecular (spongy) and cortical bone lose mass despite normal bone mineralization and lab values (serum Ca^{2+} and PO_4^{3-}).

Most commonly due to ↑ bone resorption related to ↓ estrogen levels and old age. Can be 2° to drugs (eg, steroids, alcohol, anticonvulsants, anticoagulants, thyroid replacement therapy) or other conditions (eg, hyperparathyroidism, hyperthyroidism, multiple myeloma, malabsorption syndromes, anorexia).

Diagnosed by bone mineral density measurement by DEXA (dual-energy X-ray absorptiometry) at the lumbar spine, total hip, and femoral neck, with a T-score of ≤ –2.5 or by a fragility fracture (eg, fall from standing height, minimal trauma) at hip or vertebra. One time screening recommended in women ≥ 65 years old.

Prophylaxis: regular weight-bearing exercise and adequate Ca^{2+} and vitamin D intake throughout adulthood.

Treatment: bisphosphonates, teriparatide, SERMs, rarely calcitonin; denosumab (monoclonal antibody against RANKL).

Can lead to vertebral compression fractures A—acute back pain, loss of height, kyphosis. Also can present with fractures of femoral neck, distal radius (Colles fracture).





(c) ketabton.com: The Digital Library

Osteopetrosis



Failure of normal bone resorption due to defective osteoclasts → thickened, dense bones that are prone to fracture. Mutations (eg, carbonic anhydrase II) impair ability of osteoclast to generate acidic environment necessary for bone resorption. Overgrowth of cortical bone fills marrow space → pancytopenia, extramedullary hematopoiesis. Can result in cranial nerve impingement and palsies due to narrowed foramina.

X-rays show diffuse symmetric sclerosis (bone-in-bone, "stone bone" A). Bone marrow transplant is potentially curative as osteoclasts are derived from monocytes.

Osteomalacia/rickets

Defective mineralization of osteoid (osteomalacia) or cartilaginous growth plates (rickets, only in children). Most commonly due to vitamin D deficiency.

X-rays show osteopenia and "Looser zones" (pseudofractures) in osteomalacia, epiphyseal widening and metaphyseal cupping/fraying in rickets. Children with rickets have pathologic bow legs (genu varum A), bead-like costochondral junctions (rachitic rosary B), craniotabes (soft skull).

↓ vitamin D → ↓ serum Ca^{2+} → ↑ PTH secretion → ↓ serum PO_4^{3-}.

Hyperactivity of osteoblasts → ↑ ALP.



Osteitis deformans



Also called Paget disease of bone. Common, localized disorder of bone remodeling caused by ↑ osteoclastic activity followed by ↑ osteoblastic activity that forms poor-quality bone. Serum Ca^{2+}, phosphorus, and PTH levels are normal. ↑ ALP. Mosaic pattern of woven and lamellar bone (osteocytes within lacunae in chaotic juxtapositions); long bone chalk-stick fractures. ↑ blood flow from ↑ arteriovenous shunts may cause high-output heart failure. ↑ risk of osteosarcoma.

Hat size can be increased due to skull thickening A; hearing loss is common due to auditory foramen narrowing.

Stages of Paget disease:

- Lytic—osteoclasts
- Mixed—osteoclasts + osteoblasts
- Sclerotic—osteoblasts
- Quiescent—minimal osteoclast/osteoblast activity

Treatment: bisphosphonates.

Avascular necrosis of bone



Infarction of bone and marrow, usually very painful. Most common site is femoral head (watershed zone) A (due to insufficiency of medial circumflex femoral artery). Causes include Corticosteroids, Alcoholism, Sickle cell disease, Trauma, SLE, "the Bends" (caisson/decompression disease), LEgg-Calvé-Perthes disease (idiopathic), Gaucher disease, Slipped capital femoral epiphysis—CASTS Bend LEGS.



(c) ketabton.com: The Digital Library

Lab values in bone disorders

DISORDER	SERUM Ca^{2+}	PO_4^{3-}	ALP	PTH	COMMENTS
Osteoporosis	—	—	—	—	↓ bone mass
Osteopetrosis	—/↓	—	—	—	Dense, brittle bones. Ca^{2+} ↓ in severe, malignant disease
Paget disease of bone	—	—	↑	—	Abnormal "mosaic" bone architecture
Osteitis fibrosa cystica Primary hyperparathyroidism	↑	↓	↑	↑	"Brown tumors" due to fibrous replacement of bone, subperiosteal thinning Idiopathic or parathyroid hyperplasia, adenoma, carcinoma
Secondary hyperparathyroidism	↓	↑	↑	↑	Often as compensation for CKD (↓ PO_4^{3-} excretion and production of activated vitamin D)
Osteomalacia/rickets	↓	↓	↑	↑	Soft bones; vitamin D deficiency also causes 2° hyperparathyroidism
Hypervitaminosis D	↑	↑	—	↓	Caused by oversupplementation or granulomatous disease (eg, sarcoidosis)

↑ ↓ = 1° change.

Primary bone tumors

Metastatic disease is more common than 1° bone tumors. Benign bone tumors that start with O are more common in boys.

TUMOR TYPE	EPIDEMIOLOGY	LOCATION	CHARACTERISTICS
Benign tumors			
Osteochondroma	Most common benign bone tumor Males < 25 years old	Metaphysis of long bones	Lateral bony projection of growth plate (continuous with marrow space) covered by cartilaginous cap A Rarely transforms to chondrosarcoma
Osteoma	Middle age	Surface of facial bones	Associated with Gardner syndrome
Osteoid osteoma	Adults < 25 years old Males > females	Cortex of long bones	Presents as bone pain (worse at night) that is relieved by NSAIDs Bony mass (< 2 cm) with radiolucent osteoid core B
Osteoblastoma	Males > females	Vertebrae	Similar histology to osteoid osteoma Larger size (> 2 cm), pain unresponsive to NSAIDs
Chondroma		Medulla of small bones of hand and feet	Benign tumor of cartilage
Giant cell tumor	20–40 years old	Epiphysis of long bones (often in knee region)	Locally aggressive benign tumor Neoplastic mononuclear cells that express RANKL and reactive multinucleated giant (osteoclast-like) cells. "Osteoclastoma" "Soap bubble" appearance on x-ray C

(c) ketabton.com: The Digital Library

Primary bone tumors *(continued)*

TUMOR TYPE	EPIDEMIOLOGY	LOCATION	CHARACTERISTICS
Malignant tumors			
Osteosarcoma (osteogenic sarcoma)	Accounts for 20% of 1° bone cancers. Peak incidence of 1° tumor in males < 20 years. Less common in elderly; usually 2° to predisposing factors, such as Paget disease of bone, bone infarcts, radiation, familial retinoblastoma, Li-Fraumeni syndrome.	Metaphysis of long bones (often in knee region).	Pleomorphic osteoid-producing cells (malignant osteoblasts). Presents as painful enlarging mass or pathologic fractures. Codman triangle D (from elevation of periosteum) or sunburst pattern on x-ray E (think of an osteocod (bone fish) swimming in the sun). Aggressive. 1° usually responsive to treatment (surgery, chemotherapy), poor prognosis for 2°.
Chondrosarcoma		Medulla of pelvis, proximal femur and humerus.	Tumor of malignant chondrocytes.
Ewing sarcoma	Most common in Caucasians. Generally boys < 15 years old.	Diaphysis of long bones (especially femur), pelvic flat bones.	Anaplastic small blue cells of neuroectodermal origin (resemble lymphocytes) F. Differentiate from conditions with similar morphology (eg, lymphoma, chronic osteomyelitis) by testing for t(11;22) (fusion protein EWS-FLI1). "Onion skin" periosteal reaction in bone. Aggressive with early metastases, but responsive to chemotherapy. 11 + 22 = 33 (Patrick Ewing's jersey number).





(c) ketabton.com: The Digital Library

Osteoarthritis vs rheumatoid arthritis

	Osteoarthritis	Rheumatoid arthritis
PATHOGENESIS	Mechanical—wear and tear destroys articular cartilage (degenerative joint disorder) → inflammation with inadequate repair. Chondrocytes mediate degradation and inadequate repair.	Autoimmune—inflammation A induces formation of pannus (proliferative granulation tissue), which erodes articular cartilage and bone.
PREDISPOSING FACTORS	Age, female, obesity, joint trauma.	Female, HLA-DR4 (4-walled "rheum"), smoking. ⊕ rheumatoid factor (IgM antibody that targets IgG Fc region; in 80%), anti-cyclic citrullinated peptide antibody (more specific).
PRESENTATION	Pain in weight-bearing joints after use (eg, at the end of the day), improving with rest. Asymmetric joint involvement. Knee cartilage loss begins medially ("bowlegged"). No systemic symptoms.	Pain, swelling, and morning stiffness lasting > 1 hour, improving with use. Symmetric joint involvement. Systemic symptoms (fever, fatigue, weight loss). Extraarticular manifestations common.*
JOINT FINDINGS	Osteophytes (bone spurs), joint space narrowing, subchondral sclerosis and cysts. Synovial fluid noninflammatory (WBC < 2000/mm^3). Development of Heberden nodes B (at DIP) and Bouchard nodes C (at PIP), and 1st CMC; not MCP.	Erosions, juxta-articular osteopenia, soft tissue swelling, subchondral cysts, joint space narrowing. Deformities: cervical subluxation, ulnar finger deviation, swan neck D, boutonniere E. Involves MCP, PIP, wrist; not DIP or 1st CMC.
TREATMENT	Activity modification, acetaminophen, NSAIDs, intra-articular glucocorticoids.	NSAIDs, glucocorticoids, disease-modifying agents (eg, methotrexate, sulfasalazine), biologic agents (eg, TNF-α inhibitors).

*Extraarticular manifestations include rheumatoid nodules (fibrinoid necrosis with palisading histiocytes) in subcutaneous tissue and lung (+ pneumoconiosis → Caplan syndrome), interstitial lung disease, pleuritis, pericarditis, anemia of chronic disease, neutropenia + splenomegaly (Felty syndrome), AA amyloidosis, Sjögren syndrome, scleritis, carpal tunnel syndrome.



FAS1_2019_11-Musculo.indd 466 11/7/19 5:23 PM

(c) ketabton.com: The Digital Library

Gout

FINDINGS

Acute inflammatory monoarthritis caused by precipitation of monosodium urate crystals in joints A. Risk factors: male sex, hypertension, obesity, diabetes, dyslipidemia, alcohol use. Strongest risk factor is hyperuricemia, which can be caused by:

- Underexcretion of uric acid (90% of patients)—largely idiopathic, potentiated by renal failure; can be exacerbated by certain medications (eg, thiazide diuretics).
- Overproduction of uric acid (10% of patients)—Lesch-Nyhan syndrome, PRPP excess, ↑ cell turnover (eg, tumor lysis syndrome), von Gierke disease.

Crystals are needle shaped and ⊖ birefringent under polarized light (yellow under parallel light, blue under perpendicular light B). Serum uric acid levels may be normal during an acute attack.

SYMPTOMS

Asymmetric joint distribution. Joint is swollen, red, and painful. Classic manifestation is painful MTP joint of big toe (podagra). Tophus formation C (often on external ear, olecranon bursa, or Achilles tendon). Acute attack tends to occur after a large meal with foods rich in purines (eg, red meat, seafood), trauma, surgery, dehydration, diuresis, or alcohol consumption (alcohol metabolites compete for same excretion sites in kidney as uric acid → ↓ uric acid secretion and subsequent buildup in blood).

TREATMENT

Acute: NSAIDs (eg, indomethacin), glucocorticoids, colchicine.
Chronic (preventive): xanthine oxidase inhibitors (eg, allopurinol, febuxostat).



Calcium pyrophosphate deposition disease



Previously called pseudogout. Deposition of calcium pyrophosphate crystals within the joint space. Occurs in patients > 50 years old; both sexes affected equally. Usually idiopathic, sometimes associated with hemochromatosis, hyperparathyroidism, joint trauma.

Pain and swelling with acute inflammation (pseudogout) and/or chronic degeneration (pseudo-osteoarthritis). Most commonly affected joint is the knee.

Chondrocalcinosis (cartilage calcification) on x-ray.

Crystals are rhomboid and weakly ⊕ birefringent under polarized light (blue when parallel to light) A.

Acute treatment: NSAIDs, colchicine, glucocorticoids.

Prophylaxis: colchicine.

The blue P's—blue (when Parallel), Positive birefringence, calcium Pyrophosphate, Pseudogout

(c) ketabton.com: The Digital Library

Systemic juvenile idiopathic arthritis

Systemic arthritis seen in < 16 year olds. Usually presents with daily spiking fevers, salmon-pink macular rash, arthritis (commonly 2+ joints). Associated with anterior uveitis. Frequently presents with leukocytosis, thrombocytosis, anemia, ↑ ESR, ↑ CRP. Treatment: NSAIDs, steroids, methotrexate, TNF inhibitors.

Sjögren syndrome



Autoimmune disorder characterized by destruction of exocrine glands (especially lacrimal and salivary) by lymphocytic infiltrates A. Predominantly affects women 40–60 years old.

Findings:

- Inflammatory joint pain
- Keratoconjunctivitis sicca (↓ tear production and subsequent corneal damage)
- Xerostomia (↓ saliva production) → mucosal atrophy, fissuring of the tongue B
- Presence of antinuclear antibodies, rheumatoid factor (can be positive in the absence of rheumatoid arthritis), antiribonucleoprotein antibodies: SS-A (anti-Ro) and/or SS-B (anti-La)
- Bilateral parotid enlargement

Anti-SSA and anti-SSB may also be seen in SLE.

A common 1° disorder or a 2° syndrome associated with other autoimmune disorders (eg, rheumatoid arthritis, SLE, systemic sclerosis).

Complications: dental caries; mucosa-associated lymphoid tissue (MALT) lymphoma (may present as parotid enlargement).

Focal lymphocytic sialadenitis on labial salivary gland biopsy can confirm diagnosis.

Septic arthritis

S aureus, *Streptococcus*, and *Neisseria gonorrhoeae* are common causes. Affected joint is swollen A, red, and painful. Synovial fluid purulent (WBC > 50,000/mm^3).

Gonococcal arthritis—STI that presents as either purulent arthritis (eg, knee) or triad of polyarthralgia, tenosynovitis (eg, hand), dermatitis (eg, pustules).

(c) ketabton.com: The Digital Library

Seronegative spondyloarthritis	Arthritis without rheumatoid factor (no anti-IgG antibody). Strong association with HLA-B27 (MHC class I serotype). Subtypes (**PAIR**) share variable occurrence of inflammatory back pain (associated with morning stiffness, improves with exercise), peripheral arthritis, enthesitis (inflamed insertion sites of tendons, eg, Achilles), dactylitis ("sausage fingers"), uveitis.	
Psoriatic arthritis	Associated with skin psoriasis and nail lesions. Asymmetric and patchy involvement **A**. Dactylitis and "pencil-in-cup" deformity of DIP on x-ray **B**.	Seen in fewer than ⅓ of patients with psoriasis.
Ankylosing spondylitis	Symmetric involvement of spine and sacroiliac joints → ankylosis (joint fusion), uveitis, aortic regurgitation.	Bamboo spine (vertebral fusion) **C**. Costovertebral and costosternal ankylosis may cause restrictive lung disease. Monitor degree of reduced chest wall expansion to assess disease severity. More common in males.
Inflammatory bowel disease	Crohn disease and ulcerative colitis are often associated with spondyloarthritis.	
Reactive arthritis	Formerly called Reiter syndrome. Classic triad: ▪ **Conjunctivitis** ▪ **Urethritis** ▪ **Arthritis**	**"Can't see, can't pee, can't bend my knee."** ***Sh*igella, *Y*ersinia, *Ch*lamydia, *C*ampylobacter, *S*almonella (ShY ChiCS).**



(c) ketabton.com: The Digital Library

Systemic lupus erythematosus

Systemic, remitting, and relapsing autoimmune disease. Organ damage primarily due to a type III hypersensitivity reaction and, to a lesser degree, a type II hypersensitivity reaction. Associated with deficiency of early complement proteins (eg, C1q, C4, C2) → ↓ clearance of immune complexes. Classic presentation: rash, joint pain, and fever in a female of reproductive age (especially of African-American or Hispanic descent).



Libman-**S**acks **E**ndocarditis—nonbacterial, verrucous thrombi usually on mitral or aortic valve and can be present on either surface of the valve (but usually on undersurface). **LSE** in **SLE**.

Lupus nephritis (glomerular deposition of DNA-anti-DNA immune complexes) can be nephritic or nephrotic (causing hematuria or proteinuria). Most common and severe type is diffuse proliferative.

Common causes of death in SLE: **R**enal disease (most common), **I**nfections, **C**ardiovascular disease (accelerated CAD).

In an anti-SSA ⊕ pregnant woman, ↑ risk of newborn developing neonatal lupus → congenital heart block, periorbital/diffuse rash, transaminitis, and cytopenias at birth.

RASH OR PAIN:
- **R**ash (malar A or discoid B)
- **A**rthritis (nonerosive)
- **S**erositis (eg, pleuritis, pericarditis)
- **H**ematologic disorders (eg, cytopenias)
- **O**ral/nasopharyngeal ulcers (usually painless)
- **R**enal disease
- **P**hotosensitivity
- **A**ntinuclear antibodies
- **I**mmunologic disorder (anti-dsDNA, anti-Sm, antiphospholipid)
- **N**eurologic disorders (eg, seizures, psychosis)

Lupus patients die with **R**edness **I**n their **C**heeks.

Mixed connective tissue disease

Features of SLE, systemic sclerosis, and/or polymyositis. Associated with anti-U1 RNP antibodies (speckled ANA).

Antiphospholipid syndrome

1° or 2° autoimmune disorder (most commonly in SLE).

Diagnosed based on clinical criteria including history of thrombosis (arterial or venous) or spontaneous abortion along with laboratory findings of lupus anticoagulant, anticardiolipin, anti-β_2 glycoprotein I antibodies.

Treatment: systemic anticoagulation.

Anticardiolipin antibodies can cause false-positive VDRL/RPR.

Lupus anticoagulant can cause prolonged PTT that is not corrected by the addition of normal platelet-free plasma.

Polymyalgia rheumatica

SYMPTOMS	Pain and stiffness in proximal muscles (eg, shoulders, hips), often with fever, malaise, weight loss. Does not cause muscular weakness. More common in women > 50 years old; associated with giant cell (temporal) arteritis.
FINDINGS	↑ ESR, ↑ CRP, normal CK.
TREATMENT	Rapid response to low-dose corticosteroids.

(c) ketabton.com: The Digital Library

Fibromyalgia

Most common in women 20–50 years old. Chronic, widespread musculoskeletal pain associated with "tender points," stiffness, paresthesias, poor sleep, fatigue, cognitive disturbance ("fibro fog"). Treatment: regular exercise, antidepressants (TCAs, SNRIs), neuropathic pain agents (eg, gabapentin).

Polymyositis/ dermatomyositis

Nonspecific: ⊕ ANA, ↑ CK. Specific: ⊕ anti-Jo-1 (histidyl-tRNA synthetase), ⊕ anti-SRP (signal recognition particle), ⊕ anti-Mi-2 (helicase).

Polymyositis

Progressive symmetric proximal muscle weakness, characterized by endomysial inflammation with CD8+ T cells. Most often involves shoulders.

Dermatomyositis

Clinically similar to polymyositis, but also involves Gottron papules A, photodistributed facial erythema (eg, heliotrope [violaceous] edema of the eyelids B), "shawl and face" rash C, darkening and thickening of fingertips and sides resulting in irregular, "dirty"-appearing marks. ↑ risk of occult malignancy. Perimysial inflammation and atrophy with CD4+ T cells.

(c) ketabton.com: The Digital Library

Neuromuscular junction diseases

	Myasthenia gravis	Lambert-Eaton myasthenic syndrome
FREQUENCY	Most common NMJ disorder	Uncommon
PATHOPHYSIOLOGY	Autoantibodies to postsynaptic ACh receptor	Autoantibodies to presynaptic Ca^{2+} channel → ↓ ACh release
CLINICAL	Fatigable muscle weakness—ptosis; diplopia; proximal weakness; respiratory muscle involvement → dyspnea; bulbar muscle involvement → dysphagia, difficulty chewing Spared reflexes Worsens with muscle use	Proximal muscle weakness, autonomic symptoms (dry mouth, constipation, impotence) Hyporeflexia Improves with muscle use
ASSOCIATED WITH	Thymoma, thymic hyperplasia	Small cell lung cancer
ACHE INHIBITOR ADMINISTRATION	Reverses symptoms (pyridostigmine for treatment)	Minimal effect



Raynaud phenomenon

↓ blood flow to skin due to arteriolar (small vessel) vasospasm in response to cold or stress: color change from white (ischemia) to blue (hypoxia) to red (reperfusion). Most often in the fingers A and toes. Called Raynaud disease when 1° (idiopathic), Raynaud syndrome when 2° to a disease process such as mixed connective tissue disease, SLE, or CREST syndrome (limited form of systemic sclerosis). Digital ulceration (critical ischemia) seen in 2° Raynaud syndrome. Treat with calcium^{2+} channel blockers.

FAS1_2019_11-Musculo.indd 472 11/7/19 5:23 PM

(c) ketabton.com: The Digital Library

Scleroderma

Systemic sclerosis. Triad of autoimmunity, noninflammatory vasculopathy, and collagen deposition with fibrosis. Commonly sclerosis of skin, manifesting as puffy, taut skin A without wrinkles, fingertip pitting B. Can involve other systems, eg, renal (scleroderma renal crisis; treat with ACE inhibitors), pulmonary (interstitial fibrosis, pulmonary HTN), GI (esophageal dysmotility and reflux), cardiovascular. 75% female. 2 major types:

- **Diffuse scleroderma**—widespread skin involvement, rapid progression, early visceral involvement. Associated with anti-Scl-70 antibody (anti-DNA topoisomerase-I antibody) and anti-RNA polymerase III.
- **Limited scleroderma**—limited skin involvement confined to fingers and face. Also with CREST syndrome: Calcinosis cutis C, anti-Centromere antibody, Raynaud phenomenon, Esophageal dysmotility, Sclerodactyly, and Telangiectasia. More benign clinical course.

▸ MUSCULOSKELETAL, SKIN, AND CONNECTIVE TISSUE—DERMATOLOGY

Skin layers

Skin has 3 layers: epidermis, dermis, subcutaneous fat (hypodermis, subcutis).
Epidermal layers: Come, Let's Get Sun Burned.



(c) ketabton.com: The Digital Library

Epithelial cell junctions



Tight junctions (zonula occludens) **A**–prevents paracellular movement of solutes; composed of claudins and occludins.

Adherens junction (belt desmosome, zonula adherens) **B**–forms "belt" connecting actin cytoskeletons of adjacent cells with **CAD**herins (**Ca**$^{2+}$-dependent **ad**hesion proteins). Loss of E-cadherin promotes metastasis.

Desmosome (spot desmosome, macula adherens) **C**–structural support via intermediate filament interactions. Autoantibodies to desmoglein 1 and/or 3 → pemphigus vulgaris.

Gap junction **D**–channel proteins called connexons permit electrical and chemical communication between cells.

Hemidesmosome **E**–connects keratin in basal cells to underlying basement membrane. Autoantibodies → **bullo**us pemphigoid. (Hemidesmosomes are down "**bullo**w.")

Integrins–membrane proteins that maintain **integri**ty of basolateral membrane by binding to collagen, laminin, and fibronectin in basement membrane.

FAS1_2019_11-Musculo.indd 474 11/7/19 5:24 PM

(c) ketabton.com: The Digital Library

Dermatologic macroscopic terms

LESION	CHARACTERISTICS	EXAMPLES
Macule	Flat lesion with well-circumscribed change in skin color < 1 cm	Freckle (ephelide), labial macule A
Patch	Macule > 1 cm	Large birthmark (congenital nevus) B
Papule	Elevated solid skin lesion < 1 cm	Mole (nevus) C, acne
Plaque	Papule > 1 cm	Psoriasis D
Vesicle	Small fluid-containing blister < 1 cm	Chickenpox (varicella), shingles (zoster) E
Bulla	Large fluid-containing blister > 1 cm	Bullous pemphigoid F
Pustule	Vesicle containing pus	Pustular psoriasis G
Wheal	Transient smooth papule or plaque	Hives (urticaria) H
Scale	Flaking off of stratum corneum	Eczema, psoriasis, SCC I
Crust	Dry exudate	Impetigo J



Dermatologic microscopic terms

LESION	CHARACTERISTICS	EXAMPLES
Hyperkeratosis	↑ thickness of stratum corneum	Psoriasis, calluses
Parakeratosis	Retention of nuclei in stratum corneum	Psoriasis, actinic keratosis
Hypergranulosis	↑ thickness of stratum granulosum	Lichen planus
Spongiosis	Epidermal accumulation of edematous fluid in intercellular spaces	Eczematous dermatitis
Acantholysis	Separation of epidermal cells	Pemphigus vulgaris
Acanthosis	Epidermal hyperplasia (↑ spinosum)	Acanthosis nigricans, psoriasis

(c) ketabton.com: The Digital Library

Pigmented skin disorders

Albinism	Normal melanocyte number with ↓ melanin production A due to ↓ tyrosinase activity or defective tyrosine transport. ↑ risk of skin cancer.
Melasma (chloasma)	Acquired hyperpigmentation associated with pregnancy ("mask of pregnancy" B) or OCP use. More common in women with darker complexions.
Vitiligo	Irregular patches of complete depigmentation C. Caused by destruction of melanocytes (believed to be autoimmune). Associated with other autoimmune disorders.

Seborrheic dermatitis

Erythematous, well-demarcated plaques A with greasy yellow scales in areas rich in sebaceous glands, such as scalp, face, and periocular region. Common in both infants (cradle cap) and adults, associated with Parkinson disease. Sebaceous glands are not inflamed, but play a role in disease development. Possibly associated with *Malassezia* spp. Treatment: topical antifungals and corticosteroids.

FAS1_2019_11-Musculo.indd 476 11/7/19 5:24 PM

(c) ketabton.com: The Digital Library

Common skin disorders

Acne	Multifactorial etiology—↑ sebum/androgen production, abnormal keratinocyte desquamation, *Cutibacterium acnes* colonization of the pilosebaceous unit (comedones), and inflammation (papules/pustules A, nodules, cysts). Treatment: retinoids, benzoyl peroxide, and antibiotics.
Atopic dermatitis (eczema)	Type I hypersensitivity reaction. Pruritic eruption, commonly on skin flexures. Associated with other atopic diseases (asthma, allergic rhinitis, food allergies); ↑ serum IgE. Mutations in filaggrin gene predispose (via skin barrier dysfunction). Often appears on face in infancy B and then in antecubital fossa C in children and adults.
Allergic contact dermatitis	Type IV hypersensitivity reaction secondary to contact allergen (eg, nickel D, poison ivy, neomycin E).
Melanocytic nevus	Common mole. Benign, but melanoma can arise in congenital or atypical moles. Intradermal nevi are papular F. Junctional nevi are flat macules G.
Pseudofolliculitis barbae	Foreign body inflammatory facial skin disorder characterized by firm, hyperpigmented papules and pustules that are painful and pruritic. Located on cheeks, jawline, and neck. Commonly occurs as a result of shaving ("razor bumps"), primarily affects African-American males.
Psoriasis	Papules and plaques with silvery scaling H, especially on knees and elbows. Acanthosis with parakeratotic scaling (nuclei still in stratum corneum), Munro microabscesses. ↑ stratum spinosum, ↓ stratum granulosum. Auspitz sign (I)—pinpoint bleeding spots from exposure of dermal papillae when scales are scraped off. Associated with nail pitting and psoriatic arthritis.
Rosacea	Inflammatory facial skin disorder characterized by erythematous papules and pustules J, but no comedones. May be associated with facial flushing in response to external stimuli (eg, alcohol, heat). Phymatous rosacea can cause rhinophyma (bulbous deformation of nose).
Seborrheic keratosis	Flat, greasy, pigmented squamous epithelial proliferation of immature keratinocytes with keratin-filled cysts (horn cysts) K. Looks "stuck on." Lesions occur on head, trunk, and extremities. Common benign neoplasm of older persons. Leser-Trélat sign L—rapid onset of multiple seborrheic keratoses, indicates possible malignancy (eg, GI adenocarcinoma).
Verrucae	Warts; caused by low-risk HPV strains. Soft, tan-colored, cauliflower-like papules M. Epidermal hyperplasia, hyperkeratosis, koilocytosis. Condyloma acuminatum on anus or genitals N.
Urticaria	Hives. Pruritic wheals that form after mast cell degranulation O. Characterized by superficial dermal edema and lymphatic channel dilation.

(c) ketabton.com: The Digital Library

Vascular tumors of skin

Angiosarcoma	Rare blood vessel malignancy typically occurring in the head, neck, and breast areas. Usually in elderly, on sun-exposed areas. Associated with radiation therapy and chronic postmastectomy lymphedema. Hepatic angiosarcoma associated with vinyl chloride and arsenic exposures. Very aggressive and difficult to resect due to delay in diagnosis.
Bacillary angiomatosis	Benign capillary skin papules A found in AIDS patients. Caused by *Bartonella* infections. Frequently mistaken for Kaposi sarcoma, but has neutrophilic infiltrate.
Cherry hemangioma	Benign capillary hemangioma B commonly appearing in middle-aged adults. Does not regress. Frequency ↑ with age.
Glomus tumor	Benign, painful, red-blue tumor, commonly under fingernails C. Arises from modified smooth muscle cells of the thermoregulatory glomus body.
Kaposi sarcoma	Endothelial malignancy most commonly affecting the skin, mouth, GI tract, respiratory tract. Classically seen in older Eastern European males, patients with AIDS, and organ transplant patients. Associated with HHV-8 and HIV. Rarely mistaken for bacillary angiomatosis, but has lymphocytic infiltrate.
Pyogenic granuloma	Polypoid lobulated capillary hemangioma D that can ulcerate and bleed. Associated with trauma and pregnancy.
Strawberry hemangioma	Benign capillary hemangioma of infancy E. Appears in first few weeks of life (1/200 births); grows rapidly and regresses spontaneously by 5–8 years old.



(c) ketabton.com: The Digital Library

Skin infections

Bacterial infections	
Impetigo	Very superficial skin infection. Usually from *S aureus* or *S pyogenes*. Highly contagious. Honey-colored crusting A. Bullous impetigo B has bullae and is usually caused by *S aureus*.
Erysipelas	Infection involving upper dermis and superficial lymphatics, usually from *S pyogenes*. Presents with well-defined, raised demarcation between infected and normal skin C.
Cellulitis	Acute, painful, spreading infection of deeper dermis and subcutaneous tissues. Usually from *S pyogenes* or *S aureus*. Often starts with a break in skin from trauma or another infection D.
Abscess	Collection of pus from a walled-off infection within deeper layers of skin E. Offending organism is almost always *S aureus*.
Necrotizing fasciitis	Deeper tissue injury, usually from anaerobic bacteria or *S pyogenes*. Pain may be out of proportion to exam findings. Results in crepitus from methane and CO_2 production. "Flesh-eating bacteria." Causes bullae and skin necrosis → violaceous color of bullae, surrounding skin F. Surgical emergency.
Staphylococcal scalded skin syndrome	Exotoxin destroys keratinocyte attachments in stratum granulosum only (vs toxic epidermal necrolysis, which destroys epidermal-dermal junction). Characterized by fever and generalized erythematous rash with sloughing of the upper layers of the epidermis G that heals completely. ⊕ Nikolsky sign (separation of epidermis upon manual stroking of skin). Commonly seen in newborns and children/adults with renal insufficiency.
Viral infections	
Herpes	Herpes virus infections (HSV1 and HSV2) of skin can occur anywhere from mucosal surfaces to normal skin. These include herpes labialis, herpes genitalis, herpetic whitlow H (finger).
Molluscum contagiosum	Umbilicated papules I caused by a poxvirus. While frequently seen in children, it may be sexually transmitted in adults.
Varicella zoster virus	Causes varicella (chickenpox) and zoster (shingles). Varicella presents with multiple crops of lesions in various stages from vesicles to crusts. Zoster is a reactivation of the virus in dermatomal distribution (unless it is disseminated).
Hairy leukoplakia	Irregular, white, painless plaques on lateral tongue that cannot be scraped off J. EBV mediated. Occurs in HIV-positive patients, organ transplant recipients. Contrast with thrush (scrapable) and leukoplakia (precancerous).

(c) ketabton.com: The Digital Library

Autoimmune blistering skin disorders

	Pemphigus vulgaris	Bullous pemphigoid
PATHOPHYSIOLOGY	Potentially fatal. Most commonly seen in older adults. Type II hypersensitivity reaction. IgG antibodies against desmoglein-1 and/or desmoglein-3 (component of desmosomes, which connect keratinocytes in the stratum spinosum).	Less severe than pemphigus vulgaris. Most commonly seen in older adults. Type II hypersensitivity reaction. IgG antibodies against hemidesmosomes (epidermal basement membrane; antibodies are "bullow" the epidermis).
GROSS MORPHOLOGY	Flaccid intraepidermal bullae A caused by acantholysis (separation of keratinocytes, "row of tombstones" on H&E stain); oral mucosa is involved. Nikolsky sign ⊕.	Tense blisters C containing eosinophils; oral mucosa spared. Nikolsky sign ⊖.
IMMUNOFLUORESCENCE	Reticular pattern around epidermal cells B.	Linear pattern at epidermal-dermal junction D.



(c) ketabton.com: The Digital Library

Other blistering skin disorders

Dermatitis herpetiformis	Pruritic papules, vesicles, and bullae (often found on elbows, knees, buttocks) A. Deposits of IgA at tips of dermal papillae. Associated with celiac disease. Treatment: dapsone, gluten-free diet.
Erythema multiforme	Associated with infections (eg, *Mycoplasma pneumoniae*, HSV), drugs (eg, sulfa drugs, β-lactams, phenytoin). Presents with multiple types of lesions—macules, papules, vesicles, target lesions (look like targets with multiple rings and dusky center showing epithelial disruption) B.
Stevens-Johnson syndrome	Characterized by fever, bullae formation and necrosis, sloughing of skin at dermal-epidermal junction (⊕ Nikolsky), high mortality rate. Typically mucous membranes are involved C D. Targetoid skin lesions may appear, as seen in erythema multiforme. Usually associated with adverse drug reaction. Toxic epidermal necrolysis (TEN) E F is more severe form of SJS involving > 30% body surface area. 10–30% involvement denotes SJS-TEN.

(c) ketabton.com: The Digital Library

Miscellaneous skin disorders

Acanthosis nigricans	Epidermal hyperplasia causing symmetric, hyperpigmented thickening of skin, especially in axilla or on neck A B. Associated with insulin resistance (eg, diabetes, obesity, Cushing syndrome, PCOS), visceral malignancy (eg, gastric adenocarcinoma).
Actinic keratosis	Premalignant lesions caused by sun exposure. Small, rough, erythematous or brownish papules or plaques C D. Risk of squamous cell carcinoma is proportional to degree of epithelial dysplasia.
Erythema nodosum	Painful, raised inflammatory lesions of subcutaneous fat (panniculitis), usually on anterior shins. Often idiopathic, but can be associated with sarcoidosis, coccidioidomycosis, histoplasmosis, TB, streptococcal infections E, leprosy F, inflammatory bowel disease.
Lichen Planus	Pruritic, Purple, Polygonal Planar Papules and Plaques are the 6 P's of lichen Planus G H. Mucosal involvement manifests as Wickham striae (reticular white lines) and hypergranulosis. Sawtooth infiltrate of lymphocytes at dermal-epidermal junction. Associated with hepatitis C.
Pityriasis rosea	"Herald patch" I followed days later by other scaly erythematous plaques, often in a "Christmas tree" distribution on trunk J. Multiple pink plaques with collarette scale. Self-resolving in 6–8 weeks.
Sunburn	Acute cutaneous inflammatory reaction due to excessive UV irradiation. Causes DNA mutations, inducing apoptosis of keratinocytes. UVB is dominant in sunBurn, UVA in tAnning and photoAging. Exposure to UVA and UVB ↑ risk of skin cancer.



(c) ketabton.com: The Digital Library

Rule of 9's

The extent of a burn injury can be estimated as a percentage of the body surface area.



Burn classification

DEPTH	INVOLVEMENT	APPEARANCE	SENSATION
Superficial burn	Epidermis only	Similar to sunburn; localized, painful, dry, blanching redness with no blisters	Painful
Superficial partial-thickness	All of epidermis and some dermis	Blisters, blanches with pressure, swollen, warm	Painful to temperature and air
Deep partial-thickness burn	All of epidermis and some dermis	Blisters (easily unroofed), does not blanch with pressure	Painless; perception of pressure only
Full-thickness burn	All of skin (epidermis and dermis)	White, waxy, dry, inelastic, leathery, does not blanch with pressure	Painless; perception of deep pressure only
Deeper injury burn	All of skin and at least partial involvement of muscle and/or fascia	White, dry, inelastic, does not blanch with pressure	Painless; some perception of deep pressure

(c) ketabton.com: The Digital Library

Skin cancer

Basal cell carcinoma more common above upper lip

Squamous cell carcinoma more common below lower lip

Sun exposure strongly predisposes to skin cancer.

Basal cell carcinoma

Most common skin cancer. Found in sun-exposed areas of body (eg, face). Locally invasive, but rarely metastasizes. Waxy, pink, pearly nodules, commonly with telangiectasias, rolled borders A, central crusting or ulceration. BCCs also appear as nonhealing ulcers with infiltrating growth B or as a scaling plaque (superficial BCC) C. Basal cell tumors have "palisading" (aligned) nuclei D.

Keratoacanthoma

Seen in middle-aged and elderly individuals. Rapidly growing, resembles squamous cell carcinoma. Presents as dome-shaped nodule with keratin-filled center. Grows rapidly (4-6 weeks) and may spontaneously regress E.

Melanoma

Common tumor with significant risk of metastasis. S-100 tumor marker. Associated with dysplastic nevi; fair-skinned persons are at ↑ risk. Depth of tumor (Breslow thickness) correlates with risk of metastasis. Look for the ABCDEs: Asymmetry, Border irregularity, Color variation, Diameter > 6 mm, and Evolution over time. At least 4 different types of melanoma, including superficial spreading F, nodular G, lentigo maligna H, and acral lentiginous (highest prevalence in African-Americans and Asians) I. Often driven by activating mutation in BRAF kinase. Primary treatment is excision with appropriately wide margins. Metastatic or unresectable melanoma in patients with *BRAF* V600E mutation may benefit from vemurafenib, a BRAF kinase inhibitor.

Squamous cell carcinoma

Second most common skin cancer. Associated with immunosuppression, chronic nonhealing wounds, and occasionally arsenic exposure. Commonly appears on face J, lower lip K, ears, hands. Locally invasive, may spread to lymph nodes, and will rarely metastasize. Ulcerative red lesions. Histopathology: keratin "pearls" L.

Actinic keratosis, a scaly plaque, is a precursor to squamous cell carcinoma.

(c) ketabton.com: The Digital Library

Arachidonic acid pathways



LTB_4 is a **neutrophil** chemotactic agent.
Neutrophils arrive "**B4**" others.

PGI_2 inhibits platelet aggregation and promotes vasodilation.
Platelet-**G**athering **I**nhibitor.

Acetaminophen

MECHANISM	Reversibly inhibits cyclooxygenase, mostly in CNS. Inactivated peripherally.
CLINICAL USE	Antipyretic, analgesic, but not anti-inflammatory. Used instead of aspirin to avoid Reye syndrome in children with viral infection.
ADVERSE EFFECTS	Overdose produces hepatic necrosis; acetaminophen metabolite (NAPQI) depletes glutathione and forms toxic tissue byproducts in liver. *N*-acetylcysteine is antidote—regenerates glutathione.

(c) ketabton.com: The Digital Library

Aspirin

MECHANISM	NSAID that irreversibly inhibits cyclooxygenase (both COX-1 and COX-2) by covalent acetylation → ↓ synthesis of TXA_2 and prostaglandins. ↑ bleeding time. No effect on PT, PTT. Effect lasts until new platelets are produced.
CLINICAL USE	Low dose (< 300 mg/day): ↓ platelet aggregation. Intermediate dose (300–2400 mg/day): antipyretic and analgesic. High dose (2400–4000 mg/day): anti-inflammatory.
ADVERSE EFFECTS	Gastric ulceration, tinnitus (CN VIII), allergic reactions (especially in patients with asthma or nasal polyps). Chronic use can lead to acute kidney injury, interstitial nephritis, GI bleeding. Risk of Reye syndrome in children treated with aspirin for viral infection. Toxic doses cause respiratory alkalosis early, but transitions to mixed metabolic acidosis-respiratory alkalosis. Treatment of overdose: $NaHCO_3$.

Celecoxib

MECHANISM	Reversibly and **sele**ctively inh**ib**its the cyclooxygenase (**COX**) isoform 2 ("**Selecoxib**"), which is found in inflammatory cells and vascular endothelium and mediates inflammation and pain; spares COX-1, which helps maintain gastric mucosa. Thus, does not have the corrosive effects of other NSAIDs on the GI lining. Spares platelet function as TXA_2 production is dependent on COX-1.
CLINICAL USE	Rheumatoid arthritis, osteoarthritis.
ADVERSE EFFECTS	↑ risk of thrombosis, sulfa allergy.

Nonsteroidal anti-inflammatory drugs

Ibuprofen, naproxen, indomethacin, ketorolac, diclofenac, meloxicam, piroxicam.

MECHANISM	Reversibly inhibit cyclooxygenase (both COX-1 and COX-2). Block prostaglandin synthesis.
CLINICAL USE	Antipyretic, analgesic, anti-inflammatory. Indomethacin is used to close a PDA.
ADVERSE EFFECTS	Interstitial nephritis, gastric ulcer (prostaglandins protect gastric mucosa), renal ischemia (prostaglandins vasodilate afferent arteriole), aplastic anemia.

Leflunomide

MECHANISM	Reversibly inhibits dihydroorotate dehydrogenase, preventing pyrimidine synthesis. Suppresses T-cell proliferation.
CLINICAL USE	Rheumatoid arthritis, psoriatic arthritis.
ADVERSE EFFECTS	Diarrhea, hypertension, hepatotoxicity, teratogenicity.

Bisphosphonates

Alendronate, ibandronate, risedronate, zoledronate.

MECHANISM	Pyrophosphate analogs; bind hydroxyapatite in bone, inhibiting osteoclast activity.
CLINICAL USE	Osteoporosis, hypercalcemia, Paget disease of bone, metastatic bone disease, osteogenesis imperfecta.
ADVERSE EFFECTS	Esophagitis (if taken orally, patients are advised to take with water and remain upright for 30 minutes), osteonecrosis of jaw, atypical femoral stress fractures.

(c) ketabton.com: The Digital Library

Teriparatide

MECHANISM	Recombinant PTH analog. ↑ osteoblastic activity when administered in pulsatile fashion.
CLINICAL USE	Osteoporosis. Causes ↑ bone growth compared to antiresorptive therapies (eg, bisphosphonates).
ADVERSE EFFECTS	↑ risk of osteosarcoma (avoid use in patients with Paget disease of the bone or unexplained elevation of alkaline phosphatase). Avoid in patients who have had prior cancers or radiation therapy. Transient hypercalcemia.

Gout drugs

Chronic gout drugs (preventive)	
Probenecid	Inhibits reabsorption of uric acid in proximal convoluted tubule (also inhibits secretion of penicillin). Can precipitate uric acid calculi.
Allopurinol	Competitive inhibitor of xanthine oxidase → ↓ conversion of hypoxanthine and xanthine to urate. Also used in lymphoma and leukemia to prevent tumor lysis–associated urate nephropathy. ↑ concentrations of xanthine oxidase active metabolites, azathioprine, and 6-MP.
Pegloticase	Recombinant uricase catalyzing uric acid to allantoin (a more water-soluble product).
Febuxostat	Inhibits xanthine oxidase.
Acute gout drugs	
NSAIDs	Any NSAID. Use salicylates with caution (may decrease uric acid excretion, particularly at low doses).
Glucocorticoids	Oral, intra-articular, or parenteral.
Colchicine	Binds and stabilizes tubulin to inhibit microtubule polymerization, impairing neutrophil chemotaxis and degranulation. Acute and prophylactic value. GI, neuromyopathic side effects.

Prevent A Painful Flare.



TNF-α inhibitors

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Etanercept	Fusion protein (decoy receptor for TNF-α + IgG_1 Fc), produced by recombinant DNA. Etanercept intercepts TNF.	Rheumatoid arthritis, psoriasis, ankylosing spondylitis	Predisposition to infection, including reactivation of latent TB, since TNF is important in granuloma formation and stabilization. Can also lead to drug-induced lupus.
Infliximab, adalimumab, certolizumab, golimumab	Anti-TNF-α monoclonal antibody.	Inflammatory bowel disease, rheumatoid arthritis, ankylosing spondylitis, psoriasis	

FAS1_2019_11-Musculo.indd 487 11/7/19 5:24 PM

(c) ketabton.com: The Digital Library

▸ NOTES

(c) ketabton.com: The Digital Library

Neurology and Special Senses

"We are all now connected by the Internet, like neurons in a giant brain."
—Stephen Hawking

"Anything's possible if you've got enough nerve."
—J.K. Rowling, *Harry Potter and the Order of the Phoenix*

"I like nonsense; it wakes up the brain cells."
—Dr. Seuss

"I believe in an open mind, but not so open that your brains fall out."
—Arthur Hays Sulzberger

"The chief function of the body is to carry the brain around."
—Thomas Edison

"Exactly how [the brain] operates remains one of the biggest unsolved mysteries, and it seems the more we probe its secrets, the more surprises we find."
—Neil deGrasse Tyson

Understand the difference between upper motor neuron (UMN) and lower motor neuron (LMN) findings and the underlying anatomy. Know the major motor, sensory, cerebellar and visual pathways and their respective locations in the CNS. Connect key neurological associations with certain pathologies (eg, cerebellar lesions, stroke manifestations, Brown-Séquard syndrome). Recognize common findings on MRI/CT (eg, ischemic and hemorrhagic stroke) and on neuropathology (eg, neurofibrillary tangles and Lewy bodies). High-yield medications include those used to treat epilepsy, Parkinson disease, migraine, and pain (eg, opioids).

- Embryology 490
- Anatomy and Physiology 493
- Pathology 511
- Otology 533
- Ophthalmology 534
- Pharmacology 544

FAS1_2019_12-Neurol.indd 489 11/8/19 7:39 AM

(c) ketabton.com: The Digital Library

▸ NEUROLOGY—EMBRYOLOGY

Neural development



Notochord induces overlying ectoderm to differentiate into neuroectoderm and form neural plate.
Neural plate gives rise to neural tube and neural crest cells.
Notochord becomes nucleus pulposus of intervertebral disc in adults.

Alar plate (dorsal): sensory; regulated by TGF-β (including bone morphogenetic protein [*BMP*])
Basal plate (ventral): motor; regulated by sonic hedgehog gene (*SHH*)

] Same orientation as spinal cord

Regional specification of developing brain

Telencephalon is the 1st part. **Di**encephalon is the **2nd** part. The rest are arranged alphabetically: **mes**encephalon, **met**encephalon, **my**elencephalon.



Central and peripheral nervous systems origins

Neuroepithelia in neural tube—CNS neurons, ependymal cells (inner lining of ventricles, make CSF), oligodendrocytes, astrocytes.
Neural crest—PNS neurons, Schwann cells, glia, melanocytes, adrenal medulla.
Mesoderm—**M**icroglia (like **M**acrophages).

(c) ketabton.com: The Digital Library

Neural tube defects — Neuropores fail to fuse (4th week) → persistent connection between amniotic cavity and spinal canal. Associated with maternal diabetes and folate deficiency. ↑ α-fetoprotein (AFP) in amniotic fluid and maternal serum (except spina bifida occulta = normal AFP). ↑ acetylcholinesterase (AChE) in amniotic fluid is a helpful confirmatory test.

Spina bifida occulta — Failure of caudal neuropore to close, but no herniation. Usually seen at lower vertebral levels. Dura is intact. Associated with tuft of hair or skin dimple at level of bony defect.

Meningocele — Meninges (but no neural tissue) herniate through bony defect.

Myelomeningocele — Meninges and neural tissue (eg, cauda equina) herniate through bony defect.

Myeloschisis — Also called rachischisis. Exposed, unfused neural tissue without skin/meningeal covering.

Anencephaly — Failure of rostral neuropore to close → no forebrain, open calvarium. Clinical findings: polyhydramnios (no swallowing center in brain).



Holoprosencephaly — Failure of the embryonic forebrain (prosencephalon) to separate into 2 cerebral hemispheres; usually occurs during weeks 5–6. May be related to mutations in sonic hedgehog signaling pathway. Associated with other midline defects including cleft lip/palate (moderate form) and cyclopia (severe form). ↑ risk for pituitary dysfunction (eg, diabetes insipidus). Can be seen with Patau syndrome (trisomy 13).



MRI reveals monoventricle **A** and fusion of basal ganglia (star in **A**).

Lissencephaly — Failure of neuronal migration resulting in a "smooth brain" that lacks sulci and gyri. May be associated with microcephaly, ventriculomegaly.

(c) ketabton.com: The Digital Library

Posterior fossa malformations

Chiari I malformation
Ectopia of cerebellar **tonsils** inferior to foramen magnum (1 structure) A. Congenital, usually asymptomatic in childhood, manifests in adulthood with headaches and cerebellar symptoms. Associated with spinal cavitations (eg, syringomyelia).

Chiari II malformation
Herniation of cerebellar **vermis** and **tonsils** (2 structures) through foramen magnum with aqueductal stenosis → noncommunicating hydrocephalus. Usually associated with lumbosacral myelomeningocele (may present as paralysis/sensory loss at and below the level of the lesion). More severe than Chiari I, usually presents early in life.

Dandy-Walker malformation
Agenesis of cerebellar vermis → cystic enlargement of 4th ventricle (arrow in B) that fills the enlarged posterior fossa. Associated with noncommunicating hydrocephalus, spina bifida.

Syringomyelia



Cystic cavity (syrinx) within central canal of spinal cord (yellow arrows in A). Fibers crossing in anterior white commissure (spinothalamic tract) are typically damaged first. Results in a "cape-like," bilateral, symmetrical loss of pain and temperature sensation in upper extremities (fine touch sensation is preserved).

Associated with Chiari I malformation (red arrow in A shows low-lying cerebellar tonsils), scoliosis and other congenital malformations; acquired causes include trauma and tumors. Most common location cervical > thoracic >> lumbar. Syrinx = tube, as in "syringe."



(c) ketabton.com: The Digital Library

Tongue development



1st and 2nd pharyngeal arches form anterior 2/3 (thus sensation via CN V_3, taste via CN VII).
3rd and 4th pharyngeal arches form posterior 1/3(thus sensation and taste mainly via CN IX, extreme posterior via CN X).
Motor innervation is via CN XII to hyoglossus (retracts and depresses tongue), **genio**glossus (**protrudes** tongue), and **styl**oglossus (draws sides of tongue upward to create a trough for swallowing).
Motor innervation is via CN X to palatoglossus (elevates posterior tongue during swallowing).

Taste—CN VII, IX, X (solitary nucleus).
Pain—CN V_3, IX, X.
Motor—CN X, XII.

The **Geni**e comes **out** of the lamp in **styl**e.

▸ NEUROLOGY—ANATOMY AND PHYSIOLOGY

Neurons

Signal-transmitting cells of the nervous system. Permanent cells—do not divide in adulthood.
Signal-relaying cells with dendrites (receive input), cell bodies, and axons (send output). Cell bodies and dendrites can be seen on Nissl staining (stains RER). RER is not present in the axon. Neuron markers: neurofilament protein, synaptophysin.

Astrocytes



Most common glial cell type in CNS. Physical support, repair, extracellular K^+ buffer, removal of excess neurotransmitter, component of blood-brain barrier, glycogen fuel reserve buffer. Reactive gliosis in response to neural injury.

Derived from neuroectoderm.
Astrocyte marker: GFAP.

Microglia



Phagocytic scavenger cells of CNS (mesodermal, mononuclear origin). Activation in response to tissue damage → release of inflammatory mediators (eg, nitric oxide, glutamate). Not readily discernible by Nissl stain.

HIV-infected microglia fuse to form multinucleated giant cells in CNS seen in HIV-associated dementia.

Ependymal cells

Ciliated simple columnar glial cells line the ventricles and central canal of spinal cord. Apical surfaces are covered in cilia (which circulate CSF) and microvilli (which help with CSF absorption). Specialized ependymal cells (choroid plexus) produce CSF.

(c) ketabton.com: The Digital Library

Myelin



↑ conduction velocity of signals transmitted down axons → saltatory conduction of action potential at the nodes of Ranvier, where there are high concentrations of Na^+ channels.
In CNS (including CN II), myelin is synthesized by oligodendrocytes; in PNS (including CN III-XII), myelin is synthesized by Schwann cells.

Wraps and insulates axons (arrow in A): ↑ space constant and ↑ conduction velocity.

COPS: CNS = Oligodendrocytes, PNS = Schwann cells.

Schwann cells



Promote axonal regeneration. Derived from neural crest.

Each "Schwone" cell myelinates only 1 PNS axon.
Injured in Guillain-Barré syndrome.

Oligodendrocytes



Myelinate axons of neurons in CNS. Each oligodendrocyte can myelinate many axons (~ 30). Predominant type of glial cell in white matter.

Derived from neuroectoderm.
"Fried egg" appearance histologically.
Injured in multiple sclerosis, progressive multifocal leukoencephalopathy (PML), leukodystrophies.

Sensory receptors

RECEPTOR TYPE	SENSORY NEURON FIBER TYPE	LOCATION	SENSES
Free nerve endings	Aδ—fast, myelinated fibers C—slow, unmyelinated A Delta plane is fast, but a taxC is slow	All skin, epidermis, some viscera	Pain, temperature
Meissner corpuscles	Large, myelinated fibers; adapt quickly	Glabrous (hairless) skin	Dynamic, fine/light touch, position sense, low-frequency vibration
Pacinian corpuscles	Large, myelinated fibers; adapt quickly	Deep skin layers, ligaments, joints	High-frequency vibration, pressure
Merkel discs	Large, myelinated fibers; adapt slowly	Finger tips, superficial skin	Pressure, deep static touch (eg, shapes, edges), position sense
Ruffini corpuscles	Dendritic endings with capsule; adapt slowly	Finger tips, joints	Pressure, slippage of objects along surface of skin, joint angle change

(c) ketabton.com: The Digital Library

Peripheral nerve



Endoneurium—thin, supportive connective tissue that ensheathes and supports individual myelinated nerve fibers.

Perineurium (blood-nerve Permeability barrier)—surrounds a fascicle of nerve fibers.

Epineurium—dense connective tissue that surrounds entire nerve (fascicles and blood vessels).

Endo = inner
Peri = around
Epi = outer

Chromatolysis



Reaction of neuronal cell body to axonal injury. Changes reflect ↑ protein synthesis in effort to repair the damaged axon. Characterized by:

- Round cellular swelling A
- Displacement of the nucleus to the periphery
- Dispersion of Nissl substance throughout cytoplasm

Wallerian degeneration—disintegration of the axon and myelin sheath distal to site of axonal injury with macrophages removing debris.

Proximal to the injury, the axon retracts, and the cell body sprouts new protrusions that grow toward other neurons for potential reinnervation. Serves as a preparation for axonal regeneration and functional recovery.



Neurotransmitter changes with disease

	LOCATION OF SYNTHESIS	ANXIETY	DEPRESSION	SCHIZOPHRENIA	ALZHEIMER DISEASE	HUNTINGTON DISEASE	PARKINSON DISEASE
Acetylcholine	Basal nucleus of Meynert				↓	↓	↑
Dopamine	Ventral tegmentum, SNc		↓	↑		↑	↓
GABA	Nucleus accumbens	↓				↓	
Norepinephrine	Locus ceruleus (pons)	↑	↓				
Serotonin	Raphe nuclei (medulla, pons)	↓	↓				↓

(c) ketabton.com: The Digital Library

Meninges



Three membranes that surround and protect the brain and spinal cord:

- Dura mater—thick outer layer closest to skull. Derived from mesoderm.
- Arachnoid mater—middle layer, contains web-like connections. Derived from neural crest.
- Pia mater—thin, fibrous inner layer that firmly adheres to brain and spinal cord. Derived from neural crest.

CSF flows in the subarachnoid space, located between arachnoid and pia mater.

Epidural space—potential space between the dura mater and skull/vertebral column containing fat and blood vessels. Site of blood collection associated with middle meningeal artery injury.

Blood-brain barrier



Prevents circulating blood substances (eg, bacteria, drugs) from reaching the CSF/CNS. Formed by 3 structures:

- Tight junctions between nonfenestrated capillary endothelial cells
- Basement membrane
- Astrocyte foot processes

Glucose and amino acids cross slowly by carrier-mediated transport mechanisms.

Nonpolar/lipid-soluble substances cross rapidly via diffusion.

Circumventricular organs with fenestrated capillaries and no blood-brain barrier allow molecules in blood to affect brain function (eg, area postrema—vomiting after chemotherapy; OVLT [organum vasculosum lamina terminalis]—osmoreceptors) or neurosecretory products to enter circulation (eg, neurohypophysis—ADH release).

Infarction and/or neoplasm destroys endothelial cell tight junctions → vasogenic edema.

Hyperosmolar agents (eg, mannitol) can disrupt the BBB → ↑ permeability of medications.

Vomiting center

Coordinated by nucleus tractus solitarius (NTS) in the medulla, which receives information from the chemoreceptor trigger zone (CTZ, located within area postrema in 4th ventricle), GI tract (via vagus nerve), vestibular system, and CNS.

CTZ and adjacent vomiting center nuclei receive input from 5 major receptors: muscarinic (M_1), dopamine (D_2), histamine (H_1), serotonin (5-HT_3), and neurokinin (NK-1) receptors.

- 5-HT_3, D_2, and NK-1 antagonists used to treat chemotherapy-induced vomiting.
- H_1 and M_1 antagonists treat motion sickness; H_1 antagonists treat hyperemesis gravidarum.

(c) ketabton.com: The Digital Library

Sleep physiology

Sleep cycle is regulated by the circadian rhythm, which is driven by suprachiasmatic nucleus (SCN) of the hypothalamus. Circadian rhythm controls nocturnal release of ACTH, prolactin, melatonin, norepinephrine: SCN → norepinephrine release → pineal gland → ↑ melatonin. SCN is regulated by environment (eg, light).

Two stages: rapid-eye movement (REM) and non-REM.

Alcohol, benzodiazepines, and barbiturates are associated with ↓ REM sleep and N3 sleep; norepinephrine also ↓ REM sleep.

Benzodiazepines are useful for night terrors and sleepwalking by ↓ N3 and REM sleep.

SLEEP STAGE (% OF TOTAL SLEEP TIME IN YOUNG ADULTS)	DESCRIPTION	EEG WAVEFORM AND NOTES
Awake (eyes open)	Alert, active mental concentration.	**B**eta (highest frequency, lowest amplitude)
Awake (eyes closed)		**A**lpha
Non-REM sleep		
Stage N1 (5%)	Light sleep.	**T**heta
Stage N2 (45%)	Deeper sleep; when bruxism ("**two**th" [tooth] grinding) occurs.	**S**leep spindles and K complexes
Stage N3 (25%)	Deepest non-REM sleep (slow-wave sleep); **sleepwalking**, night terrors, and **bedwetting** occur (**wee** and **flee** in N3).	**D**elta (lowest frequency, highest amplitude)
REM sleep (25%)	Loss of motor tone, ↑ brain O_2 use, variable pulse/BP, ↑ ACh. REM is when dreaming, nightmares, and penile/clitoral tumescence occur; may serve memory processing function. Extraocular movements due to activity of PPRF (paramedian pontine reticular formation/conjugate gaze center). Occurs every 90 minutes, and duration ↑ through the night.	**B**eta Changes in elderly: ↓ REM sleep time, ↓ N3. Changes in depression: ↑ REM sleep time, ↓ REM latency, ↓ N3, repeated nighttime awakenings, early morning awakening (terminal insomnia). Changes in narcolepsy: ↓ REM latency. At night, **BATS** **D**rink **B**lood

(c) ketabton.com: The Digital Library

Hypothalamus	Maintains homeostasis by regulating **T**hirst and water balance, controlling **A**denohypophysis (anterior pituitary) and **N**eurohypophysis (posterior pituitary) release of hormones produced in the hypothalamus, and regulating **H**unger, **A**utonomic nervous system, **T**emperature, and **S**exual urges (**TAN HATS**). Inputs (areas not protected by blood-brain barrier): OVLT (senses change in osmolarity), area postrema (found in dorsal medulla, responds to emetics).	
Lateral nucleus	Hunger. Destruction → anorexia, failure to thrive (infants). Stimulated by ghrelin, inhibited by leptin.	**L**ateral injury makes you **L**ean.
Ventromedial nucleus	Satiety. Destruction (eg, craniopharyngioma) → hyperphagia. Stimulated by leptin.	**V**entro**M**edial injury makes you **V**ery **M**assive.
Anterior nucleus	Cooling, parasympathetic.	**A**/**C** = **A**nterior **C**ooling.
Posterior nucleus	Heating, sympathetic.	**Heat**ing controlled by **Post**erior nucleus ("**Hot Pot**").
Suprachiasmatic nucleus	Circadian rhythm.	**SCN** is a **S**un-**C**ensing **N**ucleus.
Supraoptic and paraventricular nuclei	Synthesize ADH and oxytocin.	**SAD POX**: **S**upraoptic = **AD**H, **P**araventricular = **OX**ytocin ADH and oxytocin are carried by neurophysins down axons to posterior pituitary, where these hormones are stored and released.
Preoptic nucleus	Thermoregulation, sexual behavior. Releases GnRH.	Failure of GnRH-producing neurons to migrate from olfactory pit → Kallmann syndrome.

Thalamus Major relay for all ascending sensory information except olfaction.

NUCLEI	INPUT	SENSES	DESTINATION	MNEMONIC
Ventral **P**ostero-**L**ateral nucleus	Spinothalamic and dorsal columns/medial lemniscus	**V**ibration, **P**ain, **P**ressure, **P**roprioception, **L**ight touch, temperature	1° somatosensory cortex	
Ventral **P**ostero-**M**edial nucleus	Trigeminal and gustatory pathway	**Face** sensation, taste	1° somatosensory cortex	**M**akeup goes on the **face**
Lateral geniculate nucleus	CN II, optic chiasm, optic tract	Vision	1° visual cortex (calcarine sulcus)	**L**ateral = **L**ight
Medial geniculate nucleus	Superior olive and inferior colliculus of tectum	Hearing	Auditory cortex of temporal lobe	**M**edial = **M**usic
Ventral **l**ateral nucleus	Cerebellum, basal ganglia	Motor	Motor cortex	

(c) ketabton.com: The Digital Library

Limbic system



Collection of neural structures involved in emotion, long-term memory, olfaction, behavior modulation, ANS function.

Consists of hippocampus (red arrows in A), amygdalae, mammillary bodies, anterior thalamic nuclei, cingulate gyrus (yellow arrows in A), entorhinal cortex. Responsible for Feeding, Fleeing, Fighting, Feeling, and Sex.

The famous 5 F's.

Dopaminergic pathways

Commonly altered by drugs (eg, antipsychotics) and movement disorders (eg, Parkinson disease).

PATHWAY	SYMPTOMS OF ALTERED ACTIVITY	NOTES
Mesocortical	↓ activity → "negative" symptoms (eg, anergia, apathy, lack of spontaneity)	Antipsychotic drugs have limited effect
Mesolimbic	↑ activity → "positive" symptoms (eg, delusions, hallucinations)	1° therapeutic target of antipsychotic drugs → ↓ positive symptoms (eg, in schizophrenia)
Nigrostriatal	↓ activity → extrapyramidal symptoms (eg, dystonia, akathisia, parkinsonism, tardive dyskinesia)	Major dopaminergic pathway in brain Significantly affected by movement disorders and antipsychotic drugs
Tuberoinfundibular	↓ activity → ↑ prolactin → ↓ libido, sexual dysfunction, galactorrhea, gynecomastia (in men)	

Cerebellum



Modulates movement; aids in coordination and balance A.

Input:
- Contralateral cortex via middle cerebellar peduncle
- Ipsilateral proprioceptive information via inferior cerebellar peduncle from spinal cord

Output:
- The only output of cerebellar cortex = Purkinje cells (always inhibitory) → deep nuclei of cerebellum → contralateral cortex via superior cerebellar peduncle
- Deep nuclei (lateral → medial)—Dentate, Emboliform, Globose, Fastigial

Lateral lesions—affect voluntary movement of extremities (lateral structures); when injured, propensity to fall toward injured (ipsilateral) side.

Medial lesions (eg, vermis, fastigial nuclei, flocculonodular lobe)—truncal ataxia (wide-based cerebellar gait), nystagmus, head tilting. Generally result in bilateral motor deficits affecting axial and proximal limb musculature (medial structures).

Don't Eat Greasy Foods

(c) ketabton.com: The Digital Library

Basal ganglia



Important in voluntary movements and adjusting posture A.
Receives cortical input, provides negative feedback to cortex to modulate movement.
Striatum = putamen (motor) + Caudate (cognitive).
Lentiform = putamen + globus pallidus.

D_1 Receptor = D1Rect pathway.
Indirect (D_2) = Inhibitory.

Direct (excitatory) pathway—SNc input to the striatum via the nigrostriatal dopaminergic pathway releases GABA, which inhibits GABA release from the GPi, disinhibiting the Thalamus via the GPi (↑ motion).
Indirect (inhibitory) pathway—SNc input to the striatum via the nigrostriatal dopaminergic pathway releases GABA that disinhibits STN via GPe inhibition, and STN stimulates GPi to inhibit the thalamus (↓ motion).

Dopamine binds to D_1, stimulating the excitatory pathway, and to D_2, inhibiting the inhibitory pathway → ↑ motion.





(c) ketabton.com: The Digital Library

Cerebral cortex regions



Cerebral perfusion

Relies on tight autoregulation. Primarily driven by P_{CO_2} (P_{O_2} also modulates perfusion in severe hypoxia).

Also relies on a pressure gradient between mean arterial pressure (MAP) and intracranial pressure (ICP). ↓ blood pressure or ↑ ICP → ↓ cerebral perfusion pressure (CPP).

Therapeutic hyperventilation → ↓ P_{CO_2} → vasoconstriction → ↓ cerebral blood flow → ↓ ICP. May be used to treat acute cerebral edema (eg, 2° to stroke) unresponsive to other interventions.

CPP = MAP – ICP. If CPP = 0, there is no cerebral perfusion → brain death.

Hypoxemia increases CPP only if P_{O_2} < 50 mm Hg.

CPP is directly proportional to P_{CO_2} until P_{CO_2} > 90 mm Hg.



(c) ketabton.com: The Digital Library

Homunculus

Topographic representation of motor and sensory areas in the cerebral cortex. Distorted appearance is due to certain body regions being more richly innervated and thus having ↑ cortical representation.



Cerebral arteries—cortical distribution



Watershed zones

Cortical border zones occur between anterior and middle cerebral arteries and posterior and middle cerebral arteries (blue areas in A).

Internal border zones occur between the superficial and deep vascular territories of the middle cerebral artery (red areas in A).

Infarct due to severe hypoperfusion → proximal upper and lower extremity weakness ("man-in-the-barrel syndrome"), higher order visual dysfunction (if posterior cerebral/middle cerebral cortical border zone stroke).

(c) ketabton.com: The Digital Library

Circle of Willis

System of anastomoses between anterior and posterior blood supplies to brain.



Dural venous sinuses



Large venous channels A that run through the periosteal and meningeal layers of the dura mater. Drain blood from cerebral veins (arrow) and receive CSF from arachnoid granulations. Empty into internal jugular vein.

Venous sinus thrombosis—presents with signs/symptoms of ↑ ICP (eg, headache, seizures, papilledema, focal neurologic deficits). May lead to venous hemorrhage. Associated with hypercoagulable states (eg, pregnancy, OCP use, factor V Leiden).



(c) ketabton.com: The Digital Library

Ventricular system



Lateral ventricles → 3rd ventricle via right and left interventricular foramina of Monro.
3rd ventricle → 4th ventricle via cerebral aqueduct of Sylvius.
4th ventricle → subarachnoid space via:
- Foramina of **L**uschka = **L**ateral.
- Foramen of **M**agendie = **M**edial.

CSF made by choroid plexuses located in the lateral and fourth ventricles. Travels to subarachnoid space via foramina of Luschka and Magendie, is reabsorbed by arachnoid granulations, and then drains into dural venous sinuses.

Brain stem—ventral view



4 CN are above pons (I, II, III, IV).
4 CN exit the pons (V, VI, VII, VIII).
4 CN are in medulla (IX, X, XI, XII).
4 CN nuclei are medial (III, IV, VI, XII).
"Factors of 12, except 1 and 2."

Brain stem—dorsal view (cerebellum removed)



Pineal gland—melatonin secretion, circadian rhythms.
Superior colliculi—direct eye movements to stimuli (noise/movements) or objects of interest.
Inferior colliculi—auditory.
Your eyes are **above** your ears, and the superior colliculus (visual) is **above** the inferior colliculus (auditory).

(c) ketabton.com: The Digital Library

Cranial nerve nuclei

Located in tegmentum portion of brain stem (between dorsal and ventral portions):

- Midbrain—nuclei of CN III, IV
- Pons—nuclei of CN V, VI, VII, VIII
- Medulla—nuclei of CN IX, X, XII
- Spinal cord—nucleus of CN XI

Lateral nuclei = sensory (a**L**ar plate).
—Sulcus limitans—
Medial nuclei = **M**otor (basal plate).

Cranial nerve and vessel pathways



Divisions of CN V exit owing to **S**tanding **R**oom **O**nly

FAS1_2019_12-Neurol.indd 505 11/8/19 7:39 AM

(c) ketabton.com: The Digital Library

Cranial nerves

NERVE	CN	FUNCTION	TYPE	MNEMONIC
Olfactory	I	Smell (only CN without thalamic relay to cortex)	Sensory	Some
Optic	II	Sight	Sensory	Say
Oculomotor	III	Eye movement (SR, IR, MR, IO), pupillary constriction (sphincter pupillae: Edinger-Westphal nucleus, muscarinic receptors), accommodation, eyelid opening (levator palpebrae)	Motor	Marry
Trochlear	IV	Eye movement (SO)	Motor	Money
Trigeminal	V	Mastication, facial sensation (ophthalmic, maxillary, mandibular divisions), somatosensation from anterior 2/3 of tongue, dampening of loud noises (tensor tympani)	Both	But
Abducens	VI	Eye movement (LR)	Motor	My
Facial	VII	Facial movement, taste from anterior 2/3 of tongue (chorda tympani), lacrimation, salivation (submandibular and sublingual glands are innervated by CN seven), eye closing (orbicularis oculi), auditory volume modulation (stapedius)	Both	Brother
Vestibulocochlear	VIII	Hearing, balance	Sensory	Says
Glossopharyngeal	IX	Taste and sensation from posterior 1/3 of tongue, swallowing, salivation (parotid gland), monitoring carotid body and sinus chemo- and baroreceptors, and elevation of pharynx/larynx (stylopharyngeus)	Both	Big
Vagus	X	Taste from supraglottic region, swallowing, soft palate elevation, midline uvula, talking, cough reflex, parasympathetics to thoracoabdominal viscera, monitoring aortic arch chemo- and baroreceptors	Both	Brains
Accessory	XI	Head turning, shoulder shrugging (SCM, trapezius)	Motor	Matter
Hypoglossal	XII	Tongue movement	Motor	Most

Vagal nuclei

NUCLEUS	FUNCTION	CRANIAL NERVES
Nucleus tractus Solitarius	Visceral Sensory information (eg, taste, baroreceptors, gut distention)	VII, IX, X
Nucleus aMbiguus	Motor innervation of pharynx, larynx, upper esophagus (eg, swallowing, palate elevation)	IX, X, XI (cranial portion)
Dorsal motor nucleus	Sends autonomic (parasympathetic) fibers to heart, lungs, upper GI	X

(c) ketabton.com: The Digital Library

Cranial nerve reflexes

REFLEX	AFFERENT	EFFERENT
Corneal	V_1 ophthalmic (nasociliary branch)	Bilateral VII (temporal branch—orbicularis oculi)
Lacrimation	V_1 (loss of reflex does not preclude emotional tears)	VII
Jaw jerk	V_3 (sensory—muscle spindle from masseter)	V_3 (motor—masseter)
Pupillary	II	III
Gag	IX	X
Cough	X	X

Mastication muscles

3 muscles close jaw: **M**asseter, te**M**poralis, **M**edial pterygoid. 1 opens: **L**ateral pterygoid. All are innervated by trigeminal nerve (V_3).

M's **M**unch.
Lateral **L**owers (when speaking of pterygoids with respect to jaw motion).
"It takes more muscle to keep your mouth shut."

Spinal nerves

There are 31 pairs of spinal nerves: 8 cervical, 12 thoracic, 5 lumbar, 5 sacral, 1 coccygeal.
Nerves C1–C7 exit above the corresponding vertebrae (eg, C3 exits above the 3rd cervical vertebra). C8 spinal nerve exits below C7 and above T1. All other nerves exit below (eg, L2 exits below the 2nd lumbar vertebra).

Spinal cord—lower extent

In adults, spinal cord ends at lower border of L1–L2 vertebrae. **S**ubarachnoid **S**pace (which contains the CSF) extends to lower border of **S2** vertebra. Lumbar puncture is usually performed between L3–L4 or L4–L5 (level of cauda equina).
Goal of lumbar puncture is to obtain sample of CSF without damaging spinal cord. To **keep** the cord **alive**, keep the spinal needle between **L3** and **L5**.
Needle passes through:

1. skin
2. fascia and fat
3. supraspinous ligament
4. interspinous ligament
5. ligamentum flavum
6. epidural space
   (epidural anesthesia needle stops here)
7. dura mater
8. arachnoid mater
9. subarachnoid space
   (CSF collection occurs here)



(c) ketabton.com: The Digital Library

Spinal cord and associated tracts

Legs (**L**umbosacral) are **L**ateral in **L**ateral corticospinal, spinothalamic tracts. Thoracic spinal cord section in **A**.

Dorsal columns are organized as you are, with hands at sides. "Arms outside, legs inside."





(c) ketabton.com: The Digital Library

Spinal tract anatomy and functions Ascending tracts synapse and then cross.

TRACT	FUNCTION	1ST-ORDER NEURON	SYNAPSE 1	2ND-ORDER NEURON	SYNAPSE 2 + PROJECTIONS
Ascending tracts					
Dorsal column	Pressure, vibration, fine touch, proprioception	Sensory nerve ending → bypasses pseudounipolar cell body in dorsal root ganglion → enters spinal cord → ascends ipsilaterally in dorsal columns	Nucleus gracilis, nucleus cuneatus (ipsilateral medulla)	Decussates in medulla → ascends contralaterally as the medial lemniscus	VPL (thalamus) → sensory cortex
Spinothalamic tract	Lateral: pain, temperature Anterior: crude touch, pressure	Sensory nerve ending (Aδ and C fibers) → bypasses pseudounipolar cell body in dorsal root ganglion → enters spinal cord	Ipsilateral gray matter (spinal cord)	Decussates in spinal cord as the anterior white commissure → ascends contralaterally	
Descending tract					
Lateral corticospinal tract	Voluntary movement of contralateral limbs	UMN: cell body in 1° motor cortex → descends ipsilaterally (through posterior limb of internal capsule and cerebral peduncle), most fibers decussate at caudal medulla (pyramidal decussation) → descends contralaterally	Cell body of anterior horn (spinal cord)	LMN: leaves spinal cord	NMJ → muscle fibers

(c) ketabton.com: The Digital Library

Clinical reflexes



Reflexes count up in order (main nerve root in **bold**):

- **Achilles reflex** = **S1**, S2 ("buckle my shoe")
- **Patellar reflex** = L2-**L4** ("kick the door")
- **Biceps and brachioradialis reflexes** = **C5**, C6 ("pick up sticks")
- **Triceps reflex** = C6, **C7**, C8 ("lay them straight")

Additional reflexes:

- **Cremasteric reflex** = L1, L2 ("testicles move")
- **Anal wink reflex** = S3, S4 ("winks galore")

Primitive reflexes

CNS reflexes that are present in a healthy infant, but are absent in a neurologically intact adult. Normally disappear within 1st year of life. These primitive reflexes are inhibited by a mature/developing frontal lobe. They may reemerge in adults following frontal lobe lesions → loss of inhibition of these reflexes.

Moro reflex	"Hang on for life" reflex—abduct/extend arms when startled, and then draw together
Rooting reflex	Movement of head toward one side if cheek or mouth is stroked (nipple seeking)
Sucking reflex	Sucking response when roof of mouth is touched
Palmar reflex	Curling of fingers if palm is stroked
Plantar reflex	Dorsiflexion of large toe and fanning of other toes with plantar stimulation Babinski sign—presence of this reflex in an adult, which may signify a UMN lesion
Galant reflex	Stroking along one side of the spine while newborn is in ventral suspension (face down) causes lateral flexion of lower body toward stimulated side

Landmark dermatomes

DERMATOME	CHARACTERISTICS
C2	Posterior half of skull
C3	High turtleneck shirt Diaphragm and gallbladder pain referred to the right shoulder via phrenic nerve C3, 4, 5 keeps the diaphragm alive
C4	Low-collar shirt
C6	Includes thumbs Thumbs up sign on left hand looks like a 6
T4	At the nipple T4 at the teat pore
T7	At the xiphoid process 7 letters in xiphoid
T10	At the umbilicus (belly button) Point of referred pain in early appendicitis
L1	At the Inguinal Ligament
L4	Includes the kneecaps Down on ALL 4's
S2, S3, S4	Sensation of penile and anal zones S2, 3, 4 keep the penis off the floor



(c) ketabton.com: The Digital Library

▸ NEUROLOGY—PATHOLOGY

Common brain lesions

AREA OF LESION	CONSEQUENCE	EXAMPLES/COMMENTS
Frontal lobe	Disinhibition and deficits in concentration, orientation, judgment; may have reemergence of primitive reflexes	
Frontal eye fields	Destructive lesions (eg, MCA stroke): eyes look toward brain lesion (ie, away from side of hemiplegia)	
Paramedian pontine reticular formation	Eyes look away from brain lesion (ie, toward side of hemiplegia)	
Medial longitudinal fasciculus	Internuclear ophthalmoplegia (impaired adduction of ipsilateral eye; nystagmus of contralateral eye with abduction)	Multiple sclerosis
Dominant parietal cortex	Agraphia, acalculia, finger agnosia, left-right disorientation	Gerstmann syndrome
Nondominant parietal cortex	Agnosia of the contralateral side of the world	Hemispatial neglect syndrome
Hippocampus (bilateral)	Anterograde amnesia—inability to make new memories	
Basal ganglia	May result in tremor at rest, chorea, athetosis	Parkinson disease, Huntington disease, Wilson disease
Subthalamic nucleus	Contralateral hemiballismus	
Mammillary bodies (bilateral)	Wernicke-Korsakoff syndrome—**C**onfusion, **A**taxia, **N**ystagmus, **O**phthalmoplegia, memory loss (anterograde and retrograde amnesia), confabulation, personality changes	Wernicke problems come in a **CAN O**' beer and other conditions associated with thiamine deficiency
Amygdala (bilateral)	Klüver-Bucy syndrome—disinhibited behavior (eg, hyperphagia, hypersexuality, hyperorality)	HSV-1 encephalitis
Dorsal midbrain	Parinaud syndrome—vertical gaze palsy, pupillary light-near dissociation, lid retraction, convergence-retraction nystagmus	Stroke, hydrocephalus, pinealoma
Reticular activating system (midbrain)	Reduced levels of arousal and wakefulness	Coma
Cerebellar hemisphere	Intention tremor, limb ataxia, loss of balance; damage to cerebellum → ipsilateral deficits; fall toward side of lesion	Cerebellar hemispheres are **laterall**y located—affect **lateral** limbs
Cerebellar vermis	Truncal ataxia (wide-based, "drunken sailor" gait), nystagmus	Vermis is **centrall**y located—affects **central** body Degeneration associated with chronic alcohol use
Red nucleus (midbrain)	Decorticate (flexor) posturing—lesion above red nucleus, presents with flexion of upper extremities and extension of lower extremities Decerebrate (extensor) posturing—lesion at or below red nucleus, presents with extension of upper and lower extremities	Worse prognosis with decerebrate posturing In de**cor**ticate posturing, your hands are near the **cor** (heart)

(c) ketabton.com: The Digital Library

Ischemic brain disease/stroke

Irreversible neuronal injury begins after 5 minutes of hypoxia. Most **vulnerable**: **hippo**campus, **n**eocortex, cerebellum (**Pur**kinje cells), **water**shed areas ("**vulnerable hippos n**eed **pur**e **water**").

Stroke imaging: noncontrast CT to exclude hemorrhage (before tPA can be given). CT detects ischemic changes in 6–24 hr. Diffusion-weighted MRI can detect ischemia within 3–30 min.

TIME SINCE ISCHEMIC EVENT	12–24 HOURS	24–72 HOURS	3–5 DAYS	1–2 WEEKS	> 2 WEEKS
Histologic features	Eosinophilic cytoplasm + pyknotic nuclei (red neurons)	Necrosis + neutrophils	Macrophages (microglia)	Reactive gliosis (astrocytes) + vascular proliferation	Glial scar

Ischemic stroke



Acute blockage of vessels → disruption of blood flow and subsequent ischemia → infarction → liquefactive necrosis.

3 types:

- Thrombotic—due to a clot forming directly at site of infarction (commonly the MCA A), usually over a ruptured atherosclerotic plaque.
- Embolic—embolus from another part of the body obstructs vessel. Can affect multiple vascular territories. Examples: atrial fibrillation, carotid artery stenosis, DVT with patent foramen ovale, infective endocarditis.
- Hypoxic—due to hypoperfusion or hypoxemia. Common during cardiovascular surgeries, tends to affect watershed areas.

Treatment: tPA (if within 3–4.5 hr of onset and no hemorrhage/risk of hemorrhage) and/or thrombectomy (if large artery occlusion). Reduce risk with medical therapy (eg, aspirin, clopidogrel); optimum control of blood pressure, blood sugars, lipids; smoking cessation; and treat conditions that ↑ risk (eg, atrial fibrillation, carotid artery stenosis).

Transient ischemic attack

Brief, reversible episode of focal neurologic dysfunction without acute infarction (⊖ MRI), with the majority resolving in < 15 minutes; ischemia (eg, embolus, small vessel stenosis).

Neonatal intraventricular hemorrhage



Bleeding into ventricles (arrow in coronal transcranial ultrasound A shows blood in right intraventricular space, extending into periventricular white matter). Increased risk in premature and low-birth-weight infants. Originates in germinal matrix, a highly vascularized layer within the subventricular zone. Due to reduced glial fiber support and impaired autoregulation of BP in premature infants. Can present with altered level of consciousness, bulging fontanelle, hypotension, seizures, coma.

(c) ketabton.com: The Digital Library

Intracranial hemorrhage

Epidural hematoma

Rupture of middle meningeal artery (branch of maxillary artery), often 2° to skull fracture (circle in A) involving the pterion (thinnest area of the lateral skull). Might present with transient loss of consciousness → recovery ("lucid interval") → rapid deterioration due to hematoma expansion.

Scalp hematoma (arrows in A) and rapid intracranial expansion (arrows in B) under systemic arterial pressure → transtentorial herniation, CN III palsy.

CT shows biconvex (lentiform), hyperdense blood collection B **not crossing suture lines**.



Subdural hematoma

Rupture of bridging veins. Can be acute (traumatic, high-energy impact → hyperdense on CT) or chronic (associated with mild trauma, cerebral atrophy, elderly, alcoholism → hypodense on CT). Also seen in shaken babies. Predisposing factors: brain atrophy, trauma.

Crescent-shaped hemorrhage (red arrows in C and D) that **crosses suture lines**. Can cause midline shift (yellow arrow in C), findings of "acute on chronic" hemorrhage (blue arrows in D).



Subarachnoid hemorrhage

Bleeding E F due to trauma, or rupture of an aneurysm (such as a saccular aneurysm E) or arteriovenous malformation. Rapid time course. Patients complain of "worst headache of my life." Bloody or yellow (xanthochromic) lumbar puncture.

Vasospasm can occur due to blood breakdown or rebleed 3–10 days after hemorrhage → ischemic infarct; nimodipine used to prevent/reduce vasospasm. ↑ risk of developing communicating and/or obstructive hydrocephalus.



Intraparenchymal hemorrhage

Most commonly caused by systemic hypertension. Also seen with amyloid angiopathy (recurrent lobar hemorrhagic stroke in elderly), vasculitis, neoplasm. May be 2° to reperfusion injury in ischemic stroke.

Hypertensive hemorrhages (Charcot-Bouchard microaneurysm) most often occur in putamen of basal ganglia (lenticulostriate vessels G), followed by thalamus, pons, and cerebellum H.



(c) ketabton.com: The Digital Library

Effects of strokes

ARTERY	AREA OF LESION	SYMPTOMS	NOTES
Anterior circulation			
Middle cerebral artery	Motor and sensory cortices A—upper limb and face. Temporal lobe (Wernicke area); frontal lobe (Broca area).	Contralateral paralysis and sensory loss—face and upper limb. Aphasia if in dominant (usually left) hemisphere. Hemineglect if lesion affects nondominant (usually right) hemisphere.	Wernicke aphasia is associated with right superior quadrant visual field defect due to temporal lobe involvement.
Anterior cerebral artery	Motor and sensory cortices—lower limb.	Contralateral paralysis and sensory loss—lower limb, urinary incontinence.	
Lenticulostriate artery	Striatum, internal capsule.	Contralateral paralysis. Absence of cortical signs (eg, neglect, aphasia, visual field loss).	Pure motor stroke. Common location of lacunar infarcts B, due to hyaline arteriosclerosis (lipohyalinosis) 2° to unmanaged hypertension.
Posterior circulation			
Anterior spinal artery	Corticospinal tract. Medial lemniscus. Caudal medulla—hypoglossal nerve.	Contralateral paralysis—upper and lower limbs. ↓ contralateral proprioception. Ipsilateral hypoglossal dysfunction (tongue deviates ipsilaterally).	Medial medullary syndrome—caused by infarct of paramedian branches of ASA and/or vertebral arteries.
Posterior inferior cerebellar artery	Lateral medulla: Nucleus ambiguus (CN IX, X, XI) Vestibular nuclei Lateral spinothalamic tract, spinal trigeminal nucleus Sympathetic fibers Inferior cerebellar peduncle	 **Dysphagia**, **hoarseness**, ↓ **gag reflex**, hiccups. Vomiting, vertigo, nystagmus ↓ pain and temperature sensation from contralateral body, ipsilateral face. Ipsilateral Horner syndrome. Ipsilateral ataxia, dysmetria.	Lateral medullary (Wallenberg) syndrome. Nucleus ambiguus effects are specific to PICA lesions C. "Don't **pick a** (**PICA**) **horse** (hoarseness) that **can't eat** (dysphagia)." Also supplies inferior cerebellar peduncle (part of cerebellum).
Anterior inferior cerebellar artery	Lateral pons: Facial nucleus Vestibular nuclei Spinothalamic tract, spinal trigeminal nucleus Sympathetic fibers Middle and inferior cerebellar peduncles Labyrinthine artery	 **Paralysis of face** (LMN lesion vs UMN lesion in cortical stroke), ↓ lacrimation, ↓ salivation, ↓ taste from anterior 2/3 of tongue. Vomiting, vertigo, nystagmus ↓ pain and temperature sensation from contralateral body, ipsilateral face. Ipsilateral Horner syndrome. Ipsilateral ataxia, dysmetria. Ipsilateral sensorineural deafness, vertigo.	Lateral pontine syndrome. Facial nucleus effects are specific to AICA lesions. "**Facial droop** means AICA's **pooped**." Also supplies middle and inferior cerebellar peduncles (part of cerebellum).

(c) ketabton.com: The Digital Library

Effects of strokes *(continued)*

ARTERY	AREA OF LESION	SYMPTOMS	NOTES
Basilar artery	Pons, medulla, lower midbrain.	If RAS spared, consciousness is preserved.	Locked-in syndrome (locked in the basement).
	Corticospinal and corticobulbar tracts.	Quadriplegia; loss of voluntary facial, mouth, and tongue movements.	
	Ocular cranial nerve nuclei, paramedian pontine reticular formation.	Loss of horizontal, but not vertical, eye movements.	
Posterior cerebral artery	Occipital lobe D.	Contralateral hemianopia with macular sparing; alexia without agraphia (dominant hemisphere).	



Central poststroke pain syndrome

Neuropathic pain due to thalamic lesions. Initial paresthesias followed in weeks to months by allodynia (ordinarily painless stimuli cause pain) and dysesthesia (altered sensation) on the contralateral side. Occurs in 10% of stroke patients.

Diffuse axonal injury



Caused by traumatic shearing forces during rapid acceleration and/or deceleration of the brain (eg, motor vehicle accident). Usually results in devastating neurologic injury, often causing coma or persistent vegetative state. MRI A shows multiple lesions (punctate hemorrhages) involving the white matter tracts.

(c) ketabton.com: The Digital Library

Aphasia

Aphasia—higher-order language deficit (inability to understand/produce/use language appropriately); caused by pathology in dominant cerebral hemisphere (usually left).

Dysarthria—motor inability to produce speech (movement deficit).



TYPE	COMMENTS
Broca (expressive)	Broca area in inferior frontal gyrus of frontal lobe. Patient appears frustrated, insight intact. Broca = Broken Boca (*boca* = mouth in Spanish).
Wernicke (receptive)	Wernicke area in superior temporal gyrus of temporal lobe. Patients do not have insight. Wernicke is a Word salad and makes no sense.
Conduction	Can be caused by damage to arCuate fasciculus.
Global	Broca and Wernicke areas affected.
Transcortical motor	Affects frontal lobe around Broca area, but Broca area is spared.
Transcortical sensory	Affects temporal lobe around Wernicke area, but Wernicke area is spared.
Transcortical mixed	Broca and Wernicke areas and arcuate fasciculus remain intact; surrounding watershed areas affected.

Aneurysms

Abnormal dilation of an artery due to weakening of vessel wall.

Saccular aneurysm



Also called berry aneurysm A. Occurs at bifurcations in the circle of Willis. Most common site is junction of ACom and ACA. Associated with ADPKD, Ehlers-Danlos syndrome. Other risk factors: advanced age, hypertension, smoking, race (↑ risk in African-Americans).

Usually clinically silent until rupture (most common complication) → subarachnoid hemorrhage ("worst headache of my life" or "thunderclap headache") → focal neurologic deficits. Can also cause symptoms via direct compression of surrounding structures by growing aneurysm.

- ACom—compression → bitemporal hemianopia (compression of optic chiasm); visual acuity deficits; rupture → ischemia in ACA distribution → contralateral lower extremity hemiparesis, sensory deficits.
- MCA—rupture → ischemia in MCA distribution → contralateral upper extremity and lower facial hemiparesis, sensory deficits.
- PCom—compression → ipsilateral CN III palsy → mydriasis ("blown pupil"); may also see ptosis, "down and out" eye.

Charcot-Bouchard microaneurysm

Common, associated with chronic hypertension; affects small vessels (eg, lenticulostriate arteries in basal ganglia, thalamus) and can cause hemorrhagic intraparenchymal strokes. Not visible on angiography.

(c) ketabton.com: The Digital Library

Seizures

Characterized by synchronized, high-frequency neuronal firing. Variety of forms.

Partial (focal) seizures

Affect single area of the brain. Most commonly originate in medial temporal lobe. Types:

- Simple partial (consciousness intact)—motor, sensory, autonomic, psychic
- Complex partial (impaired consciousness, automatisms)

Generalized seizures

Diffuse. Types:

- Absence (petit mal)—3 Hz spike-and-wave discharges, no postictal confusion, blank stare
- Myoclonic—quick, repetitive jerks
- Tonic-clonic (grand mal)—alternating stiffening and movement, postictal confusion, urinary incontinence, tongue biting
- Tonic—stiffening
- Atonic—"drop" seizures (falls to floor); commonly mistaken for fainting

Epilepsy—disorder of recurrent, unprovoked seizures (febrile seizures are not epilepsy).

Status epilepticus—continuous (≥ 5 min) or recurring seizures that may result in brain injury.

Causes of seizures by age:

- Children—genetic, infection (febrile), trauma, congenital, metabolic
- Adults—tumor, trauma, stroke, infection
- Elderly—stroke, tumor, trauma, metabolic, infection



Fever vs heat stroke

	Fever	Heat stroke
PATHOPHYSIOLOGY	Cytokine activation during inflammation (eg, infection)	Inability of body to dissipate heat (eg, exertion)
TEMPERATURE	Usually < 40 °C	Usually > 40 °C
COMPLICATIONS	Febrile seizure (benign, usually self-limiting)	CNS dysfunction (eg, confusion), end-organ damage, acute respiratory distress syndrome, rhabdomyolysis
MANAGEMENT	Acetaminophen or ibuprofen for comfort (does not prevent future febrile seizures), antibiotic therapy if indicated	Rapid external cooling, rehydration and electrolyte correction

(c) ketabton.com: The Digital Library

Headaches

Pain due to irritation of structures such as the dura, cranial nerves, or extracranial structures. More common in females, except cluster headaches.

CLASSIFICATION	LOCALIZATION	DURATION	DESCRIPTION	TREATMENT
Cluster[a]	Unilateral	15 min–3 hr; repetitive	Excruciating periorbital pain ("suicide headache") with lacrimation and rhinorrhea. May present with Horner syndrome. More common in males.	Acute: sumatriptan, 100% O_2. Prophylaxis: verapamil.
Migraine 	Unilateral	4–72 hr	Pulsating pain with nausea, photophobia, or phonophobia. May have "aura." Due to irritation of CN V, meninges, or blood vessels (release of vasoactive neuropeptides [eg, substance P, calcitonin gene-related peptide]).	Acute: NSAIDs, triptans, dihydroergotamine. Prophylaxis: lifestyle changes (eg, sleep, exercise, diet), β-blockers, amitriptyline, topiramate, valproate, botulinum toxin, anti-CGRP monoclonal antibodies. **POUND**–**P**ulsatile, **O**ne-day duration, **U**nilateral, **N**ausea, **D**isabling.
Tension	Bilateral	> 30 min (typically 4–6 hr); constant	Steady, "band-like" pain. No photophobia or phonophobia. No aura.	Acute: analgesics, NSAIDs, acetaminophen. Prophylaxis: TCAs (eg, amitriptyline), behavioral therapy.

Other causes of headache include subarachnoid hemorrhage ("worst headache of my life"), meningitis, hydrocephalus, neoplasia, giant cell (temporal) arteritis.

[a] Compare with **trigeminal neuralgia**, which produces repetitive, unilateral, shooting/shock-like pain in the distribution of CN V. Triggered by chewing, talking, touching certain parts of the face. Lasts (typically) for seconds to minutes, but episodes often increase in intensity and frequency over time. First-line therapy: carbamazepine.

(c) ketabton.com: The Digital Library

Movement disorders

DISORDER	PRESENTATION	CHARACTERISTIC LESION	NOTES
Akathisia	Restlessness and intense urge to move.		Can be seen with neuroleptic use or as a side effect of Parkinson treatment.
Asterixis	Extension of wrists causes "flapping" motion.		Associated with hepatic encephalopathy, Wilson disease, and other metabolic derangements.
Athetosis	Slow, snake-like, writhing movements; especially seen in the fingers.	Basal ganglia.	Seen in Huntington disease.
Chorea	Sudden, jerky, purposeless movements.	Basal ganglia.	*Chorea* = dancing. Seen in Huntington disease and in acute rheumatic fever (Sydenham chorea).
Dystonia	Sustained, involuntary muscle contractions.		Writer's cramp, blepharospasm, torticollis. Treatment: botulinum toxin injection.
Essential tremor	High-frequency tremor with sustained posture (eg, outstretched arms), worsened with movement or when anxious.		Often familial. Patients often self-medicate with alcohol, which ↓ tremor amplitude. Treatment: nonselective β-blockers (eg, propranolol), primidone.
Intention tremor	Slow, zigzag motion when pointing/extending toward a target.	Cerebellar dysfunction.	
Resting tremor	Uncontrolled movement of distal appendages (most noticeable in hands); tremor alleviated by intentional movement.	Substantia nigra (Parkinson disease).	Occurs at rest; "pill-rolling tremor" of Parkinson disease. When you park your car, it is at rest.
Hemiballismus	Sudden, wild flailing of one side of the body.	Contralateral subthalamic nucleus (eg, lacunar stroke).	Pronounce "Half-of-body ballistic."
Myoclonus	Sudden, brief, uncontrolled muscle contraction.		Jerks; hiccups; common in metabolic abnormalities such as renal and liver failure.
Restless legs syndrome	Worse at rest/nighttime. Relieved by movement.		Associated with iron deficiency, CKD. Treatment: dopamine agonists (pramipexole, ropinirole).

(c) ketabton.com: The Digital Library

Neurodegenerative disorders

↓ in cognitive ability, memory, or function with intact consciousness.
Must rule out depression as cause of dementia (called pseudodementia). Other reversible causes of dementia: hypothyroidism, vitamin B_{12} deficiency, neurosyphilis, normal pressure hydrocephalus.

DISEASE	DESCRIPTION	HISTOLOGIC/GROSS FINDINGS
Parkinson disease	Parkinson **TRAPSS** your body: **T**remor (pill-rolling tremor at rest) **R**igidity (cogwheel) **A**kinesia (or bradykinesia) **P**ostural instability **S**huffling gait **S**mall handwriting (micrographia) MPTP, a contaminant in illegal drugs, is metabolized to MPP+, which is toxic to substantia nigra.	Loss of dopaminergic neurons (ie, depigmentation) of substantia nigra pars compacta. Lewy bodies: composed of α-synuclein (intracellular eosinophilic inclusions **A**).
Huntington disease	Autosomal dominant trinucleotide $(CAG)_n$ repeat expansion in the **hunt**ingtin *(HTT)* gene on chromosome **4** **(4 letters).** Symptoms manifest between ages 20 and 50: chorea, athetosis, aggression, depression, dementia (sometimes initially mistaken for substance abuse). Anticipation results from expansion of **CAG** repeats. **C**audate loses **A**Ch and **G**ABA.	Atrophy of caudate and putamen with ex vacuo ventriculomegaly. ↑ dopamine, ↓ GABA, ↓ ACh in brain. Neuronal death via NMDA-R binding and glutamate excitotoxicity.
Alzheimer disease	Most common cause of dementia in elderly. Down syndrome patients have ↑ risk of developing Alzheimer disease, as APP is located on chromosome 21. ↓ ACh. Associated with the following altered proteins: ▪ ApoE-2: ↓ risk of sporadic form ▪ ApoE-4: ↑ risk of sporadic form ▪ APP, presenilin-1, presenilin-2: familial forms (10%) with earlier onset	Widespread cortical atrophy (normal cortex **B**; cortex in Alzheimer disease **C**), especially hippocampus (arrows in **B** and **C**). Narrowing of gyri and widening of sulci. Senile plaques **D** in gray matter: extracellular β-amyloid core; may cause amyloid angiopathy → intracranial hemorrhage; Aβ (amyloid-β) synthesized by cleaving amyloid precursor protein (APP). Neurofibrillary tangles **E**: intracellular, hyperphosphorylated tau protein = insoluble cytoskeletal elements; number of tangles correlates with degree of dementia. Hirano bodies—intracellular eosinophilic proteinaceous rods in hippocampus.
Frontotemporal dementia	Formerly called Pick disease. Early changes in personality and behavior (behavioral variant), or aphasia (primary progressive aphasia). May have associated movement disorders.	Frontotemporal lobe degeneration **F**. Inclusions of hyperphosphorylated tau (round Pick bodies **G**) or ubiquitinated TDP-43.

(c) ketabton.com: The Digital Library

Neurodegenerative disorders *(continued)*

DISEASE	DESCRIPTION	HISTOLOGIC/GROSS FINDINGS
Lewy body dementia	Visual hallucinations ("haLewycinations"), dementia with fluctuating cognition/alertness, REM sleep behavior disorder, and parkinsonism. Called Lewy body dementia if cognitive and motor symptom onset < 1 year apart, otherwise considered dementia 2° to Parkinson disease.	Intracellular Lewy bodies A primarily in cortex.
Vascular dementia	Result of multiple arterial infarcts and/or chronic ischemia. Step-wise decline in cognitive ability with late-onset memory impairment. 2nd most common cause of dementia in elderly.	MRI or CT shows multiple cortical and/or subcortical infarcts.
Creutzfeldt-Jakob disease	Rapidly progressive (weeks to months) dementia with myoclonus ("startle myoclonus") and ataxia. Commonly see periodic sharp waves on EEG and ↑ 14-3-3 protein in CSF.	Spongiform cortex (vacuolization without inflammation). Prions (PrP^{c} → PrP^{sc} sheet [β-pleated sheet resistant to proteases]) H.



Idiopathic intracranial hypertension

Also called pseudotumor cerebri. ↑ ICP with no obvious findings on imaging. Risk factors include female sex, Tetracyclines, Obesity, vitamin A excess, Danazol (female TOAD). Associated with cerebral venous sinus stenosis. Findings: headache, tinnitus, diplopia (usually from CN VI palsy), no change in mental status. Impaired optic nerve axoplasmic flow → papilledema. Visual field testing shows enlarged blind spot and peripheral constriction. Lumbar puncture reveals ↑ opening pressure and provides temporary headache relief.

Treatment: weight loss, acetazolamide, invasive procedures for refractory cases (eg, CSF shunt placement, optic nerve sheath fenestration surgery for visual loss).

(c) ketabton.com: The Digital Library

Hydrocephalus	↑ CSF volume → ventricular dilation +/– ↑ ICP.
Communicating	
Communicating hydrocephalus	↓ CSF absorption by arachnoid granulations (eg, arachnoid scarring post-meningitis) → ↑ ICP, papilledema, herniation.
Normal pressure hydrocephalus	Affects the elderly; idiopathic; CSF pressure elevated only episodically; does not result in increased subarachnoid space volume. Expansion of ventricles A distorts the fibers of the corona radiata → triad of **urinary incontinence**, **gait apraxia** (magnetic gait), and **cognitive dysfunction**. "**Wet**, **wobbly**, and **wacky**." Symptoms potentially reversible with CSF drainage via lumbar puncture or shunt placement.
Noncommunicating (obstructive)	
Noncommunicating hydrocephalus	Caused by structural blockage of CSF circulation within ventricular system (eg, stenosis of aqueduct of Sylvius, colloid cyst blocking foramen of Monro, tumor B).
Hydrocephalus mimics	
Ex vacuo ventriculomegaly	Appearance of ↑ CSF on imaging C, but is actually due to ↓ brain tissue and neuronal atrophy (eg, Alzheimer disease, advanced HIV, Pick disease, Huntington disease). ICP is normal; NPH triad is not seen.

(c) ketabton.com: The Digital Library

Multiple sclerosis

Autoimmune inflammation and demyelination of CNS (brain and spinal cord) with subsequent axonal damage. Can present with:

- Acute optic neuritis (painful unilateral visual loss associated with Marcus Gunn pupil)
- Brain stem/cerebellar syndromes (eg, diplopia, ataxia, scanning speech, intention tremor, nystagmus/INO [bilateral > unilateral])
- Pyramidal tract demyelination (eg, weakness, spasticity)
- Spinal cord syndromes (eg, electric shock-like sensation along cervical spine on neck flexion, neurogenic bladder, paraparesis, sensory manifestations affecting the trunk or one or more extremity)

Symptoms may exacerbate with increased body temperature (eg, hot bath, exercise). Relapsing and remitting is most common clinical course. Most often affects women in their 20s and 30s; more common in individuals living farther from equator and with low serum vitamin D levels.

FINDINGS



↑ IgG level and myelin basic protein in CSF. Oligoclonal bands are diagnostic. MRI is gold standard. Periventricular plaques A (areas of oligodendrocyte loss and reactive gliosis). Multiple white matter lesions disseminated in space and time.

TREATMENT

Stop relapses and halt/slow progression with disease-modifying therapies (eg, β-interferon, glatiramer, natalizumab). Treat acute flares with IV steroids. Symptomatic treatment for neurogenic bladder (catheterization, muscarinic antagonists), spasticity (baclofen, $GABA_B$ receptor agonists), pain (TCAs, anticonvulsants).

(c) ketabton.com: The Digital Library

Other demyelinating and dysmyelinating disorders

Osmotic demyelination syndrome



Also called central pontine myelinolysis. Massive axonal demyelination in pontine white matter A 2° to rapid osmotic changes, most commonly iatrogenic correction of hyponatremia but also rapid shifts of other osmolytes (eg, glucose). Acute paralysis, dysarthria, dysphagia, diplopia, loss of consciousness. Can cause "locked-in syndrome."

Correcting serum Na^+ too fast:

- "From low to high, your pons will die" (osmotic demyelination syndrome)
- "From high to low, your brains will blow" (cerebral edema/herniation)

Acute inflammatory demyelinating polyradiculopathy

Most common subtype of **Guillain-Barré syndrome**.

Autoimmune condition that destroys Schwann cells via inflammation and demyelination of motor fibers, sensory fibers, peripheral nerves (including CN III-XII). Likely facilitated by molecular mimicry and triggered by inoculations or stress. Despite association with infections (eg, *Campylobacter jejuni*, viruses [eg, Zika]), no definitive causal link to any pathogen.

Results in symmetric ascending muscle weakness/paralysis and depressed/absent DTRs beginning in lower extremities. Facial paralysis (usually bilateral) and respiratory failure are common. May see autonomic dysregulation (eg, cardiac irregularities, hypertension, hypotension) or sensory abnormalities. Almost all patients survive; majority recover completely after weeks to months.

↑ CSF protein with normal cell count (albuminocytologic dissociation).

Respiratory support is critical until recovery. Disease-modifying treatment: plasmapheresis or IV immunoglobulins. No role for steroids.

Acute disseminated (postinfectious) encephalomyelitis

Multifocal inflammation and demyelination after infection or vaccination. Presents with rapidly progressive multifocal neurologic symptoms, altered mental status.

Charcot-Marie-Tooth disease

Also called hereditary motor and sensory neuropathy. Group of progressive hereditary nerve disorders related to the defective production of proteins involved in the structure and function of peripheral nerves or the myelin sheath. Typically autosomal dominant and associated with foot deformities (eg, pes cavus, hammer toe), lower extremity weakness (eg, foot drop), and sensory deficits. Most common type, CMT1A, is caused by *PMP22* gene duplication.

Progressive multifocal leukoencephalopathy

Demyelination of CNS B due to destruction of oligodendrocytes (2° to reactivation of latent JC virus infection). Seen in 2–4% of patients with AIDS. Rapidly progressive, usually fatal. Predominantly involves parietal and occipital areas; visual symptoms are common. ↑ risk associated with natalizumab.

Other disorders

Krabbe disease, metachromatic leukodystrophy, adrenoleukodystrophy.

(c) ketabton.com: The Digital Library

Neurocutaneous disorders

DISORDER	GENETICS	PRESENTATION	NOTES
Sturge-Weber syndrome	Congenital nonhereditary anomaly of neural crest derivatives. Somatic mosaicism of an activating mutation in one copy of the *GNAQ* gene.	Capillary vascular malformation → port-wine stain A (nevus flammeus or non-neoplastic birthmark) in CN V_1/V_2 distribution; ipsilateral leptomeningeal angioma B → seizures/epilepsy; intellectual disability; episcleral hemangioma → ↑ IOP → early-onset glaucoma.	Also called encephalotrigeminal angiomatosis. SSTURGGE-Weber: Sporadic, port-wine Stain, Tram track calcifications (opposing gyri), Unilateral, intellectual disability (Retardation), Glaucoma, *GNAQ* gene, Epilepsy.
Tuberous sclerosis	AD, variable expression. Mutation in tumor suppressor genes *TSC1* on chromosome 9 (hamartin), *TSC2* on chromosome 16 (tuberin).	Hamartomas in CNS and skin, Angiofibromas C, Mitral regurgitation, Ash-leaf spots D, cardiac Rhabdomyoma, (Tuberous sclerosis), autosomal dOminant; Mental retardation (intellectual disability), renal Angiomyolipoma E, Seizures, Shagreen patches.	HAMARTOMASS. ↑ incidence of Subependymal giant cell astrocytomas and ungual fibromas.
Neurofibromatosis type I	AD, 100% penetrance. Mutation in *NF1* tumor suppressor gene on chromosome 17 (encodes neurofibromin, a negative RAS regulator).	Café-au-lait spots F, Intellectual disability, Cutaneous neurofibromas G, Lisch nodules (pigmented iris hamartomas H), Optic gliomas, Pheochromocytomas, Seizures/focal neurologic Signs (often from meningioma), bone lesions (eg, sphenoid dysplasia).	Also called von Recklinghausen disease. 17 letters in "von Recklinghausen." CICLOPSS.
Neurofibromatosis type II	AD. Mutation in *NF2* tumor suppressor gene (merlin) on chromosome 22.	Bilateral vestibular schwannomas, juvenile cataracts, meningiomas, ependymomas.	NF2 affects 2 ears, 2 eyes.
von Hippel-Lindau disease	AD. Deletion of *VHL* gene on chromosome 3p. pVHL ubiquitinates hypoxia-inducible factor 1a.	Hemangioblastomas (high vascularity with hyperchromatic nuclei I) in retina, brain stem, cerebellum, spine J; Angiomatosis; bilateral Renal cell carcinomas; Pheochromocytomas.	Numerous tumors, benign and malignant. VHL = 3 letters. HARP.



(c) ketabton.com: The Digital Library

Adult primary brain tumors

TUMOR	DESCRIPTION	HISTOLOGY
Glioblastoma multiforme	Grade IV astrocytoma. Common, highly malignant 1° brain tumor with ~ 1-year median survival. Found in cerebral hemispheres. Can cross corpus callosum ("butterfly glioma" A).	Astrocyte origin, GFAP ⊕. "Pseudopalisading" pleomorphic tumor cells B border central areas of necrosis, hemorrhage, and/or microvascular proliferation.
Oligodendroglioma	Relatively rare, slow growing. Most often in frontal lobes C. Often calcified.	Oligodendrocyte origin. "Fried egg" cells—round nuclei with clear cytoplasm D. "Chicken-wire" capillary pattern.
Meningioma	Common, typically benign. Females > males. Most often occurs near surfaces of brain and in parasagittal region. Extra-axial (external to brain parenchyma) and may have a dural attachment ("tail" E). Often asymptomatic; may present with seizures or focal neurologic signs. Resection and/or radiosurgery.	Arachnoid cell origin. Spindle cells concentrically arranged in a whorled pattern F; psammoma bodies (laminated calcifications).
Hemangioblastoma	Most often cerebellar G. Associated with von Hippel-Lindau syndrome when found with retinal angiomas. Can produce erythropoietin → 2° polycythemia.	Blood vessel origin. Closely arranged, thin-walled capillaries with minimal intervening parenchyma H.



(c) ketabton.com: The Digital Library

Adult primary brain tumors *(continued)*

TUMOR	DESCRIPTION	HISTOLOGY
Pituitary adenoma	May be nonfunctioning (silent) or hyperfunctioning (hormone-producing). Nonfunctional tumors present with mass effect (eg, bitemporal hemianopia [due to pressure on optic chiasm I]). Pituitary apoplexy → hyper- or hypopituitarism. Prolactinoma classically presents as galactorrhea, amenorrhea, ↓ bone density due to suppression of estrogen in women and as ↓ libido, infertility in men. Treatment: dopamine agonists (eg, bromocriptine, cabergoline), transsphenoidal resection.	Hyperplasia of only one type of endocrine cells found in pituitary. Most commonly from lactotrophs (prolactin) J → hyperprolactinemia. Less commonly, from somatotrophs (GH) → acromegaly, gigantism; corticotrophs (ACTH) → Cushing disease. Rarely, from thyrotrophs (TSH), gonadotrophs (FSH, LH).
Schwannoma	Classically at the cerebellopontine angle K, benign, involving CNs V, VII, and VIII, but can be along any peripheral nerve. Often localized to CN VIII in internal acoustic meatus → vestibular schwannoma (can present as hearing loss and tinnitus). Bilateral vestibular schwannomas found in NF-2. Resection or stereotactic radiosurgery.	Schwann cell origin, S-100 ⊕. Biphasic, dense, hypercellular areas containing spindle cells alternating with hypocellular, myxoid areas L.





(c) ketabton.com: The Digital Library

Childhood primary brain tumors

TUMOR	DESCRIPTION	HISTOLOGY
Pilocytic astrocytoma	Low-grade astrocytoma. Most common 1° brain tumor in childhood. Usually well circumscribed. In children, most often found in posterior fossa A (eg, cerebellum). May be supratentorial. Benign; good prognosis.	Astrocyte origin, GFAP ⊕. Rosenthal fibers—eosinophilic, corkscrew fibers B. Cystic + solid (gross).
Medulloblastoma	Most common malignant brain tumor in childhood. Commonly involves cerebellum C. Can compress 4th ventricle, causing noncommunicating hydrocephalus → headaches, papilledema. Can involve the cerebellar vermis → truncal ataxia. Can send "drop metastases" to spinal cord.	Form of primitive neuroectodermal tumor (PNET). Homer-Wright rosettes, small blue cells D. Synaptophysin ⊕.
Ependymoma	Most commonly found in 4th ventricle E. Can cause hydrocephalus. Poor prognosis.	Ependymal cell origin. Characteristic perivascular pseudorosettes F. Rod-shaped blepharoplasts (basal ciliary bodies) found near the nucleus.
Craniopharyngioma	Most common childhood supratentorial tumor. May be confused with pituitary adenoma (both cause bitemporal hemianopia).	Derived from remnants of Rathke pouch (ectoderm). Calcification is common G H. Cholesterol crystals found in "motor oil"-like fluid within tumor.
Pinealoma	Tumor of pineal gland. Can cause Parinaud syndrome (compression of tectum → vertical gaze palsy); obstructive hydrocephalus (compression of cerebral aqueduct); precocious puberty in males (hCG production).	Similar to germ cell tumors (eg, testicular seminoma).



(c) ketabton.com: The Digital Library

Herniation syndromes



❶ Cingulate (subfalcine) herniation under falx cerebri	Can compress anterior cerebral artery.
❷ Central/downward transtentorial herniation	Caudal displacement of brain stem → rupture of paramedian basilar artery branches → Duret hemorrhages. Usually fatal.
❸ Uncal transtentorial herniation	Uncus = medial temporal lobe. Early herniation → ipsilateral blown pupil (unilateral CN III compression), contralateral hemiparesis. Late herniation → coma, Kernohan phenomenon (misleading contralateral blown pupil and ipsilateral hemiparesis due to contralateral compression against Kernohan notch).
❹ Cerebellar tonsillar herniation into the foramen magnum	Coma and death result when these herniations compress the brain stem.

Motor neuron signs

SIGN	UMN LESION	LMN LESION	COMMENTS
Weakness	+	+	Lower motor neuron = everything lowered (less muscle mass, ↓ muscle tone, ↓ reflexes, downgoing toes) Upper motor neuron = everything up (tone, DTRs, toes)
Atrophy	–	+	
Fasciculations	–	+	Fasciculations = muscle twitching
Reflexes	↑	↓	
Tone	↑	↓	
Babinski	+	–	Positive Babinski is normal in infants
Spastic paresis	+	–	
Flaccid paralysis	–	+	
Clasp knife spasticity	+	–	

(c) ketabton.com: The Digital Library

Spinal lesions

AREA AFFECTED	DISEASE	CHARACTERISTICS
	Spinal muscular atrophy	Congenital degeneration of anterior horns of spinal cord. LMN symptoms only, symmetric weakness. "**F**loppy baby" with marked hypotonia (**F**laccid paralysis) and tongue **F**asciculations. Autosomal recessive mutation in *SMN1* → defective snRNP assembly. SMA type 1 is called Werdnig-Hoffmann disease.
	**Amyotrophic lateral sclerosis**	Also called **Lou** Gehrig disease. Combined UMN (corticobulbar/corticospinal) and LMN (medullary and spinal cord) degeneration. No sensory or bowel/bladder deficits. Can be caused by defect in superoxide dismutase 1. LMN deficits: flaccid limb weakness, fasciculations, atrophy, bulbar palsy (dysarthria, dysphagia, tongue atrophy). UMN deficits: spastic limb weakness, hyperreflexia, clonus, pseudobulbar palsy (dysarthria, dysphagia, emotional lability). Fatal. Treatment: "ri**Lou**zole".
	**Complete occlusion of anterior spinal artery**	Spares dorsal columns and Lissauer tract; mid-thoracic ASA territory is watershed area, as artery of Adamkiewicz supplies ASA below T8. Can be caused by aortic aneurysm repair. Presents with UMN deficit below the lesion (corticospinal tract), LMN deficit at the level of the lesion (anterior horn), and loss of pain and temperature sensation below the lesion (spinothalamic tract).
	**Tabes dorsalis**	Caused by 3° syphilis. Results from degeneration/demyelination of dorsal columns and roots → progressive sensory ataxia (impaired proprioception → poor coordination). ⊕ Romberg sign and absent DTRs. Associated with Charcot joints, shooting pain, Argyll Robertson pupils.
	Syringomyelia	Syrinx expands and damages anterior white commissure of spinothalamic tract (2nd-order neurons) → bilateral symmetric loss of pain and temperature sensation in cape-like distribution. Seen with Chiari I malformation. Can affect other tracts.
	**Vitamin B_{12} deficiency**	Subacute combined degeneration (**SCD**)—demyelination of **S**pinocerebellar tracts, lateral **C**orticospinal tracts, and **D**orsal columns. Ataxic gait, paresthesia, impaired position/vibration sense, UMN symptoms.
	**Cauda equina syndrome**	Compression of spinal roots L2 and below, often due to intervertebral disc herniation or tumor. Radicular pain, absent knee and ankle reflexes, loss of bladder and anal sphincter control, saddle anesthesia.

(c) ketabton.com: The Digital Library

Poliomyelitis

Caused by poliovirus (fecal-oral transmission). Replicates in oropharynx and small intestine before spreading via bloodstream to CNS. Infection causes destruction of cells in anterior horn of spinal cord (LMN death).

Signs of LMN lesion: asymmetric weakness (vs symmetric weakness in spinal muscular atrophy), hypotonia, flaccid paralysis, fasciculations, hyporeflexia, muscle atrophy. Respiratory muscle involvement leads to respiratory failure. Signs of infection: malaise, headache, fever, nausea, etc.

CSF shows ↑ WBCs (lymphocytic pleocytosis) and slight ↑ of protein (with no change in CSF glucose). Virus recovered from stool or throat.

Brown-Séquard syndrome



Hemisection of spinal cord. Findings:

1. Ipsilateral loss of all sensation **at** level of lesion
2. Ipsilateral LMN signs (eg, flaccid paralysis) **at** level of lesion
3. Ipsilateral UMN signs **below** level of lesion (due to corticospinal tract damage)
4. Ipsilateral loss of proprioception, vibration, light (2-point discrimination) touch, and tactile sense **below** level of lesion (due to dorsal column damage)
5. Contralateral loss of pain, temperature, and crude (non-discriminative) touch **below** level of lesion (due to spinothalamic tract damage)

If lesion occurs above T1, patient may present with ipsilateral Horner syndrome due to damage of oculosympathetic pathway.



Friedreich ataxia

Autosomal recessive trinucleotide repeat disorder $(GAA)_n$ on chromosome 9 in gene that encodes frataxin (iron-binding protein). Leads to impairment in mitochondrial functioning. Degeneration of lateral corticospinal tract (spastic paralysis), spinocerebellar tract (ataxia), dorsal columns (↓ vibratory sense, proprioception), and dorsal root ganglia (loss of DTRs). **Staggering** gait, frequent **falling**, nystagmus, dysarthria, pes cavus, hammer toes, **diabetes** mellitus, **hypertrophic cardiomyopathy** (cause of death). Presents in childhood with kyphoscoliosis A B.

Friedreich is **F**ratastic (**frataxin**): he's your favorite **frat** brother, always **staggering** and **falling** but has a **sweet**, **big heart**.

Ataxic **GAA**it.

(c) ketabton.com: The Digital Library

Common cranial nerve lesions

CN V motor lesion	Jaw deviates **toward** side of lesion due to unopposed force from the opposite pterygoid muscle.
CN X lesion	Uvula deviates **away** from side of lesion. Weak side collapses and uvula points away.
CN XI lesion	Weakness turning head to contralateral side of lesion (SCM). Shoulder droop on side of lesion (trapezius). The left SCM contracts to help turn the head to the right.
CN XII lesion	LMN lesion. Tongue deviates **toward** side of lesion ("lick your wounds") due to weakened tongue muscles on affected side.

Facial nerve lesions

Bell palsy is the most common cause of peripheral facial palsy A. Usually develops after HSV reactivation. Treatment: corticosteroids +/– acyclovir. Most patients gradually recover function, but aberrant regeneration can occur. Other causes of peripheral facial palsy include Lyme disease, herpes zoster (Ramsay Hunt syndrome), sarcoidosis, tumors (eg, parotid gland), diabetes mellitus.

	Upper motor neuron lesion	Lower motor neuron lesion
LESION LOCATION	Motor cortex, connection from motor cortex to facial nucleus in pons	Facial nucleus, anywhere along CN VII
AFFECTED SIDE	Contralateral	Ipsilateral
MUSCLES INVOLVED	Lower muscles of facial expression	Upper and lower muscles of facial expression
FOREHEAD INVOLVED?	Spared, due to bilateral UMN innervation	Affected
OTHER SYMPTOMS	None	Incomplete eye closure (dry eyes, corneal ulceration), hyperacusis, loss of taste sensation to anterior tongue



FAS1_2019_12-Neurol.indd 532 11/8/19 7:39 AM

(c) ketabton.com: The Digital Library

▸ NEUROLOGY—OTOLOGY

Auditory physiology

Outer ear	Visible portion of ear (pinna), includes auditory canal and tympanic membrane. Transfers sound waves via vibration of tympanic membrane.
Middle ear	Air-filled space with three bones called the ossicles (malleus, incus, stapes). Ossicles conduct and amplify sound from tympanic membrane to inner ear.
Inner ear	Snail-shaped, fluid-filled cochlea. Contains basilar membrane that vibrates 2° to sound waves. Vibration transduced via specialized hair cells → auditory nerve signaling → brain stem. Each frequency leads to vibration at specific location on basilar membrane (tonotopy): ▪ Low frequency heard at apex near helicotrema (wide and flexible). ▪ High frequency heard best at base of cochlea (thin and rigid).

Diagnosing hearing loss



Types of hearing loss

Noise-induced hearing loss	Damage to stereociliated cells in organ of Corti. Loss of high-frequency hearing first. Sudden extremely loud noises can produce hearing loss due to tympanic membrane rupture.
Presbycusis	Aging-related progressive bilateral/symmetric sensorineural hearing loss (often of higher frequencies) due to destruction of hair cells at the cochlear base (preserved low-frequency hearing at apex).

Cholesteatoma

Overgrowth of desquamated keratin debris within the middle ear space (A, arrows); may erode ossicles, mastoid air cells → conductive hearing loss. Often presents with painless otorrhea.



(c) ketabton.com: The Digital Library

Vertigo	Sensation of spinning while actually stationary. Subtype of "dizziness," but distinct from "lightheadedness."
Peripheral vertigo	More common. Inner ear etiology (eg, semicircular canal debris, vestibular nerve infection, Ménière disease [triad: sensorineural hearing loss, vertigo, tinnitus; endolymphatic hydrops → ↑ endolymph within the inner ear], benign paroxysmal positional vertigo [BPPV]). Treatment: antihistamines, anticholinergics, antiemetics (symptomatic relief); low-salt diet +/– diuretics (Ménière disease); Epley maneuver (BPPV).
Central vertigo	Brain stem or cerebellar lesion (eg, stroke affecting vestibular nuclei, demyelinating disease, or posterior fossa tumor). Findings: directional or purely vertical nystagmus, skew deviation (vertical misalignment of the eyes), diplopia, dysmetria. Focal neurologic findings.

▸ NEUROLOGY—OPHTHALMOLOGY

Normal eye anatomy





Conjunctivitis



Inflammation of the conjunctiva → red eye A.
Allergic—itchy eyes, bilateral.
Bacterial—pus; treat with antibiotics.
Viral—most common, often adenovirus; sparse mucous discharge, swollen preauricular node, ↑ lacrimation; self-resolving.

(c) ketabton.com: The Digital Library

Refractive errors	Common cause of impaired vision, correctable with glasses.
Hyperopia	Also called "farsightedness." Eye too short for refractive power of cornea and lens → light focused behind retina. Correct with convex (converging) lenses.
Myopia	Also called "nearsightedness." Eye too long for refractive power of cornea and lens → light focused in front of retina. Correct with concave (diverging) lens.
Astigmatism	Abnormal curvature of cornea → different refractive power at different axes. Correct with cylindrical lens.

Presbyopia

Aging-related impaired accommodation (focusing on near objects), primarily due to ↓ lens elasticity, changes in lens curvature, ↓ strength of the ciliary muscle. Patients often need "reading glasses" (magnifiers).

Cataract



Painless, often bilateral, opacification of lens A, often resulting in glare and ↓ vision, especially at night. Acquired risk factors: ↑ age, smoking, excessive alcohol use, excessive sunlight, prolonged corticosteroid use, diabetes mellitus, trauma, infection. Congenital risk factors: classic galactosemia, galactokinase deficiency, trisomies (13, 18, 21), TORCH infections (eg, rubella), Marfan syndrome, Alport syndrome, myotonic dystrophy, neurofibromatosis 2.

Aqueous humor pathway



(c) ketabton.com: The Digital Library

Glaucoma

Optic disc atrophy with characteristic cupping (normal A versus thinning of outer rim of optic nerve head B), usually with elevated intraocular pressure (IOP) and progressive peripheral visual field loss if untreated. Treatment is through pharmacologic or surgical lowering of IOP.

Open-angle glaucoma

Associated with ↑ age, African-American race, family history. Painless, more common in US.
Primary—cause unclear.
Secondary—blocked trabecular meshwork from WBCs (eg, uveitis), RBCs (eg, vitreous hemorrhage), retinal elements (eg, retinal detachment).

Closed- or narrow-angle glaucoma

Primary—enlargement or anterior movement of lens against central iris (pupil margin) → obstruction of normal aqueous flow through pupil → fluid builds up behind iris, pushing peripheral iris against cornea C and impeding flow through trabecular meshwork.
Secondary—hypoxia from retinal disease (eg, diabetes mellitus, vein occlusion) induces vasoproliferation in iris that contracts angle.
Chronic closure—often asymptomatic with damage to optic nerve and peripheral vision.
Acute closure—true ophthalmic emergency. ↑ IOP pushes iris forward → angle closes abruptly. Very painful, red eye D, sudden vision loss, halos around lights, frontal headache, fixed and mid-dilated pupil, nausea and vomiting. Mydriatic agents contraindicated.



Uveitis



Inflammation of uvea; specific name based on location within affected eye. Anterior uveitis: iritis; posterior uveitis: choroiditis and/or retinitis. May have hypopyon (accumulation of pus in anterior chamber A) or conjunctival redness. Associated with systemic inflammatory disorders (eg, sarcoidosis, rheumatoid arthritis, juvenile idiopathic arthritis, HLA-B27–associated conditions).

Age-related macular degeneration



Degeneration of macula (central area of retina). Causes distortion (metamorphopsia) and eventual loss of central vision (scotomas).

- Dry (nonexudative, > 80%)—Deposition of yellowish extracellular material ("Drusen") in between Bruch membrane and retinal pigment epithelium A with gradual ↓ in vision. Prevent progression with multivitamin and antioxidant supplements.
- Wet (exudative, 10–15%)—rapid loss of vision due to bleeding 2° to choroidal neovascularization. Treat with anti-VEGF (vascular endothelial growth factor) injections (eg, bevacizumab, ranibizumab).

(c) ketabton.com: The Digital Library

Diabetic retinopathy



Retinal damage due to chronic hyperglycemia. Two types:

- Nonproliferative—damaged capillaries leak blood → lipids and fluid seep into retina → hemorrhages (arrows in A) and macular edema. Treatment: blood sugar control.
- Proliferative—chronic hypoxia results in new blood vessel formation with resultant traction on retina → retinal detachment. Treatment: anti-VEGF injections, peripheral retinal photocoagulation, surgery.

Hypertensive retinopathy

Retinal damage due to chronic uncontrolled HTN.

Flame-shaped retinal hemorrhages, arteriovenous nicking, microaneurysms, macular star (exudate, red arrow in A), cotton-wool spots (blue arrow in A). Presence of papilledema requires immediate lowering of BP.

Associated with ↑ risk of stroke, CAD, kidney disease.

Retinal vein occlusion



Blockage of central or branch retinal vein due to compression from nearby arterial atherosclerosis. Retinal hemorrhage and venous engorgement ("blood and thunder appearance"; arrows in A), edema in affected area.

Retinal detachment



Separation of neurosensory layer of retina (photoreceptor layer with rods and cones) from outermost pigmented epithelium (normally shields excess light, supports retina) → degeneration of photoreceptors → vision loss. May be 2° to retinal breaks, diabetic traction, inflammatory effusions. Visualized on fundoscopy as crinkling of retinal tissue A and changes in vessel direction.

Breaks more common in patients with high myopia and/or history of head trauma. Often preceded by posterior vitreous detachment ("flashes" and "floaters") and eventual monocular loss of vision like a "curtain drawn down." Surgical emergency.

(c) ketabton.com: The Digital Library

Central retinal artery occlusion



Acute, painless monocular vision loss. Retina cloudy with attenuated vessels and "cherry-red" spot at fovea (center of macula) A. Evaluate for embolic source (eg, carotid artery atherosclerosis, cardiac vegetations, patent foramen ovale).

Retinitis pigmentosa



Inherited progressive retinal degeneration. Nyctalopia (night blindness) → peripheral vision loss. Bone spicule-shaped deposits A.

Papilledema



Optic disc swelling (usually bilateral) due to ↑ ICP (eg, 2° to mass effect). Enlarged blind spot and elevated optic disc with blurred margins A.

Leukocoria

Loss (whitening) of the red reflex. Important causes in children include retinoblastoma A, congenital cataract, toxocariasis.

(c) ketabton.com: The Digital Library

Pupillary control

Miosis

Constriction, parasympathetic:

- 1st neuron: Edinger-Westphal nucleus to ciliary ganglion via CN III
- 2nd neuron: short ciliary nerves to sphincter pupillae muscles

Short ciliary nerves shorten the pupil diameter.

Pupillary light reflex

Light in either retina sends a signal via CN II to pretectal nuclei (dashed lines in image) in midbrain that activates bilateral Edinger-Westphal nuclei; pupils constrict bilaterally (direct and consensual reflex).

Result: illumination of 1 eye results in bilateral pupillary constriction.



Mydriasis

Dilation, sympathetic:

- 1st neuron: hypothalamus to ciliospinal center of Budge (C8–T2)
- 2nd neuron: exit at T1 to superior cervical ganglion (travels along cervical sympathetic chain near lung apex, subclavian vessels)
- 3rd neuron: plexus along internal carotid, through cavernous sinus; enters orbit as long ciliary nerve to pupillary dilator muscles. Sympathetic fibers also innervate smooth muscle of eyelids (minor retractors) and sweat glands of forehead and face.

Long ciliary nerves make the pupil diameter longer.

Marcus Gunn pupil

Also called relative afferent pupillary defect (RAPD). When the light shines into a normal eye, constriction of the ipsilateral (direct reflex) and contralateral eye (consensual reflex) is observed. When the light is then swung to the affected eye, both pupils dilate instead of constrict due to impaired conduction of light signal along the injured optic nerve. Associated with optic neuritis, early multiple sclerosis.

FAS1_2019_12-Neurol.indd 539 11/8/19 7:39 AM

(c) ketabton.com: The Digital Library

Horner syndrome

Sympathetic denervation of face →:

- **P**tosis (slight drooping of eyelid: superior tarsal muscle)
- **A**nhidrosis (absence of sweating) and flushing of affected side of face
- **M**iosis (pupil constriction)

Associated with lesions along the sympathetic chain:

- 1st neuron: pontine hemorrhage, lateral medullary syndrome, spinal cord lesion above T1 (eg, Brown-Séquard syndrome, late-stage syringomyelia)
- 2nd neuron: stellate ganglion compression by Pancoast tumor
- 3rd neuron: carotid dissection (painful)

PAM is **horn**y (**Horn**er).



Ocular motility



CN **VI** innervates the **L**ateral **R**ectus.
CN **IV** innervates the **S**uperior **O**blique.
CN **III** innervates the **R**est.
The "chemical formula" $LR_6SO_4R_3$.



Obliques go **O**pposite (left SO and IO tested with patient looking right).
IOU: **IO** tested looking **U**p.

(c) ketabton.com: The Digital Library

CN III, IV, VI palsies

CN III damage

CN III has both motor (central) and parasympathetic (peripheral) components. Common causes include:

- Ischemia → pupil sparing (motor fibers affected more than parasympathetic fibers)
- Uncal herniation → coma
- PCom aneurysm → sudden-onset headache
- Cavernous sinus thrombosis → proptosis, involvement of CNs IV, V_1/V_2, VI
- Midbrain stroke → contralateral hemiplegia



Motor output to extraocular muscles—affected primarily by vascular disease (eg, diabetes mellitus: glucose → sorbitol) due to ↓ diffusion of oxygen and nutrients to the interior fibers from compromised vasculature that resides on outside of nerve. Signs: ptosis, "down-and-out" gaze.

Parasympathetic output—fibers on the periphery are first affected by compression (eg, PCom aneurysm, uncal herniation). Signs: diminished or absent pupillary light reflex, "blown pupil" often with "down-and-out" gaze A.

Motor = Middle (central)
Parasympathetic = Peripheral

CN IV damage

Pupil is higher in the affected eye B. Characteristic head tilt to contralateral/unaffected side to compensate for lack of intorsion in affected eye.

Can't see the floor with CN IV damage (eg, difficulty going down stairs, reading).



CN VI damage

Affected eye unable to abduct and is displaced medially in primary position of gaze C.

(c) ketabton.com: The Digital Library

Visual field defects

1. Right anopia (monocular vision loss)
2. Bitemporal hemianopia (pituitary lesion, chiasm)
3. Left homonymous hemianopia
4. Left upper quadrantanopia (right temporal lesion, MCA)
5. Left lower quadrantanopia (right parietal lesion, MCA)
6. Left hemianopia with macular sparing (right occipital lesion, PCA)
7. Central scotoma (eg, macular degeneration)

Meyer **L**oop—**L**ower retina; **L**oops around inferior horn of **L**ateral ventricle.
Dorsal optic radiation—superior retina; takes shortest path via internal capsule.



Note: When an image hits 1° visual cortex, it is upside down and left-right reversed.

Cavernous sinus

Collection of venous sinuses on either side of pituitary. Blood from eye and superficial cortex → cavernous sinus → internal jugular vein.
CNs III, IV, V_1, V_2, and VI plus postganglionic sympathetic pupillary fibers en route to orbit all pass through cavernous sinus. Cavernous portion of internal carotid artery is also here.
Cavernous sinus syndrome—presents with variable ophthalmoplegia, ↓ corneal sensation, Horner syndrome and occasional decreased maxillary sensation. 2° to pituitary tumor mass effect, carotid-cavernous fistula, or cavernous sinus thrombosis related to infection.



FAS1_2019_12-Neurol.indd 542 11/8/19 7:39 AM

(c) ketabton.com: The Digital Library

Internuclear ophthalmoplegia

Medial longitudinal fasciculus (MLF): pair of tracts that allows for crosstalk between CN VI and CN III nuclei. Coordinates both eyes to move in same horizontal direction. Highly myelinated (must communicate quickly so eyes move at same time). Lesions may be unilateral or bilateral (latter classically seen in multiple sclerosis, stroke).

Lesion in MLF = internuclear ophthalmoplegia (INO), a conjugate horizontal gaze palsy. Lack of communication such that when CN VI nucleus activates ipsilateral lateral rectus, contralateral CN III nucleus does not stimulate medial rectus to contract. Abducting eye displays nystagmus (CN VI overfires to stimulate CN III). Convergence normal.



MLF in MS.

When looking left, the left nucleus of CN VI fires, which contracts the left lateral rectus and stimulates the contralateral (right) nucleus of CN III via the right MLF to contract the right medial rectus.

Directional term (eg, right INO, left INO) refers to the eye that is unable to adduct.

INO = Ipsilateral adduction failure, Nystagmus Opposite.



(c) ketabton.com: The Digital Library

▸ NEUROLOGY—PHARMACOLOGY

Epilepsy therapy

	GENERALIZED						
	PARTIAL (FOCAL)	TONIC-CLONIC	ABSENCE	STATUS EPILEPTICUS	MECHANISM	SIDE EFFECTS	NOTES
Benzodiazepines				** ✓	↑ $GABA_A$ action	Sedation, tolerance, dependence, respiratory depression	Also for eclampsia seizures (1st line is $MgSO_4$)
Carbamazepine	* ✓	✓			Blocks Na^+ channels	Diplopia, ataxia, blood dyscrasias (agranulocytosis, aplastic anemia), liver toxicity, teratogenesis (cleft lip/palate, spina bifida), induction of cytochrome P-450, SIADH, SJS	1st line for trigeminal neuralgia
Ethosuximide			* ✓		Blocks thalamic T-type Ca^{2+} channels	**EFGHIJ**—**E**thosuximide causes **F**atigue, **G**I distress, **H**eadache, **I**tching (and urticaria), S**J**S	**S**ucks to have **S**ilent (absence) **S**eizures
Gabapentin	✓				Primarily inhibits high-voltage-activated Ca^{2+} channels; designed as GABA analog	Sedation, ataxia	Also used for peripheral neuropathy, postherpetic neuralgia
Lamotrigine	✓	✓	✓		Blocks voltage-gated Na^+ channels, inhibits the release of glutamate	SJS (must be titrated slowly), hemophagocytic lymphohistiocytosis (black box warning)	
Levetiracetam	✓	✓			SV2A receptor blocker; may modulate GABA and glutamate release, inhibit voltage-gated Ca^{2+} channels	Neuropsychiatric symptoms (eg, personality change), fatigue, drowsiness, headache	
Phenobarbital	✓	✓		✓	↑ $GABA_A$ action	Sedation, tolerance, dependence, induction of cytochrome P-450, cardiorespiratory depression	1st line in **neonates** ("phenobabytal")
Phenytoin, fosphenytoin	✓	* ✓		*** ✓	Blocks Na^+ channels; zero-order kinetics	**PHENYTOIN**: cytochrome **P**-450 induction, **H**irsutism, **E**nlarged gums, **N**ystagmus, **Y**ellow-brown skin, **T**eratogenicity (fetal hydantoin syndrome), **O**steopenia, **I**nhibited folate absorption, **N**europathy. Rare: SJS, DRESS syndrome, SLE-like syndrome. Toxicity leads to diplopia, ataxia, sedation.	
Topiramate	✓	✓			Blocks Na^+ channels, ↑ GABA action	**S**edation, **s**low cognition, kidney **s**tones, **s**kinny (weight loss), **s**ight threatened (glaucoma), **s**peech (word-finding) difficulties	Also used for migraine prophylaxis
Valproic acid	✓	* ✓	✓		↑ Na^+ channel inactivation, ↑ GABA concentration by inhibiting GABA transaminase	GI distress, rare but fatal hepatotoxicity (measure LFTs), pancreatitis, neural tube defects, tremor, weight gain, contraindicated in pregnancy	Also used for myoclonic seizures, bipolar disorder, migraine prophylaxis
Vigabatrin	✓				↑ GABA. Irreversible GABA transaminase inhibitor	Permanent visual loss (black box warning)	**V**ision **g**one **a**ll **b**ad with **Vigab**atrin

* = Common use, ** = 1st line for acute, *** = 1st line for recurrent seizure prophylaxis.

(c) ketabton.com: The Digital Library

Epilepsy therapy *(continued)*



FAS1_2019_12-Neurol.indd 545 11/8/19 7:39 AM

(c) ketabton.com: The Digital Library

Barbiturates	Phenobarbital, pentobarbital, thiopental, secobarbital.
MECHANISM	Facilitate $GABA_A$ action by ↑ **duration** of Cl^- channel opening, thus ↓ neuron firing (barbi**durat**es ↑ **durat**ion).
CLINICAL USE	Sedative for anxiety, seizures, insomnia, induction of anesthesia (thiopental).
ADVERSE EFFECTS	Respiratory and cardiovascular depression (can be fatal); CNS depression (can be exacerbated by alcohol use); dependence; drug interactions (induces cytochrome P-450). Overdose treatment is supportive (assist respiration and maintain BP). Contraindicated in porphyria.
Benzodiazepines	Diazepam, lorazepam, triazolam, temazepam, oxazepam, midazolam, chlordiazepoxide, alprazolam.
MECHANISM	Facilitate $GABA_A$ action by ↑ **frequency** of Cl^- channel opening ("**fre**nzodiazepines" ↑ **fre**quency). ↓ REM sleep. Most have long half-lives and active metabolites (exceptions [**ATOM**]: **A**lprazolam, **T**riazolam, **O**xazepam, and **M**idazolam are short acting → higher addictive potential).
CLINICAL USE	Anxiety, panic disorder, spasticity, status epilepticus (lorazepam, diazepam, midazolam), eclampsia, detoxification (especially alcohol withdrawal– DTs), night terrors, sleepwalking, general anesthetic (amnesia, muscle relaxation), hypnotic (insomnia). **L**orazepam, **O**xazepam, and **T**emazepam can be used for those with liver disease who drink a **LOT** due to minimal first-pass metabolism.
ADVERSE EFFECTS	Dependence, additive CNS depression effects with alcohol and barbiturates (all bind the $GABA_A$ receptor). Less risk of respiratory depression and coma than with barbiturates. Treat overdose with flumazenil (competitive antagonist at GABA benzodiazepine receptor). Can precipitate seizures by causing acute benzodiazepine withdrawal.
Nonbenzodiazepine hypnotics	**Z**olpidem, **Z**aleplon, es**Z**opiclone. "These **ZZZ**s put you to sleep."
MECHANISM	Act via the BZ_1 subtype of the GABA receptor. Effects reversed by flumazenil. Sleep cycle less affected as compared with benzodiazepine hypnotics.
CLINICAL USE	Insomnia.
ADVERSE EFFECTS	Ataxia, headaches, confusion. Short duration because of rapid metabolism by liver enzymes. Unlike older sedative-hypnotics, cause only modest day-after psychomotor depression and few amnestic effects. ↓ dependence risk than benzodiazepines.

(c) ketabton.com: The Digital Library

Suvorexant

MECHANISM	Orexin (hypocretin) receptor antagonist.	Suvorexant is an orexin antagonist.
CLINICAL USE	Insomnia.	
ADVERSE EFFECTS	CNS depression (somnolence), headache, abnormal sleep-related activities. Contraindications: narcolepsy, combination with strong CYP3A4 inhibitors. Not recommended in patients with liver disease. Limited physical dependence or abuse potential.	

Ramelteon

MECHANISM	Melatonin receptor agonist; binds MT1 and MT2 in suprachiasmatic nucleus.	Ramelteon is a melatonin receptor agonist.
CLINICAL USE	Insomnia.	
ADVERSE EFFECTS	Dizziness, nausea, fatigue, headache. No dependence (not a controlled substance).	

Triptans — Sumatriptan

MECHANISM	5-$HT_{1B/1D}$ agonists. Inhibit trigeminal nerve activation, prevent vasoactive peptide release, induce vasoconstriction.	A sumo wrestler trips and falls on his head.
CLINICAL USE	Acute migraine, cluster headache attacks.	
ADVERSE EFFECTS	Coronary vasospasm (contraindicated in patients with CAD or vasospastic angina), mild paresthesia, serotonin syndrome (in combination with other 5-HT agonists).	

(c) ketabton.com: The Digital Library

Parkinson disease therapy

Parkinsonism is due to loss of dopaminergic neurons and excess cholinergic activity.
Bromocriptine, Amantadine, Levodopa (with carbidopa), Selegiline (and COMT inhibitors), Antimuscarinics (BALSA).

STRATEGY	AGENTS
Dopamine agonists	Ergot—Bromocriptine. Non-ergot (preferred)—pramipexole, ropinirole; toxicity includes nausea, impulse control disorder (eg, gambling), postural hypotension, hallucinations, confusion.
↑ dopamine availability	Amantadine (↑ dopamine release and ↓ dopamine reuptake); toxicity = peripheral edema, livedo reticularis, ataxia.
↑ L-DOPA availability	Agents prevent peripheral (pre-BBB) L-DOPA degradation → ↑ L-DOPA entering CNS → ↑ central L-DOPA available for conversion to dopamine. ▪ Levodopa (L-DOPA)/carbidopa—carbidopa blocks peripheral conversion of L-DOPA to dopamine by inhibiting DOPA decarboxylase. Also reduces side effects of peripheral L-DOPA conversion into dopamine (eg, nausea, vomiting). ▪ Entacapone and tolcapone prevent peripheral L-DOPA degradation to 3-O-methyldopa (3-OMD) by inhibiting COMT. Used in conjunction with levodopa.
Prevent dopamine breakdown	Agents act centrally (post-BBB) to inhibit breakdown of dopamine. ▪ Selegiline, rasagiline—block conversion of dopamine into DOPAC by selectively inhibiting MAO-B. ▪ Tolcapone—crosses BBB and blocks conversion of dopamine to 3-methoxytyramine (3-MT) in the brain by inhibiting central COMT.
Curb excess cholinergic activity	Benztropine, trihexyphenidyl (Antimuscarinic; improves tremor and rigidity but has little effect on bradykinesia in Parkinson disease). Park your Mercedes-Benz.



FAS1_2019_12-Neurol.indd 548 11/8/19 7:40 AM

(c) ketabton.com: The Digital Library

Carbidopa/levodopa

MECHANISM	↑ dopamine in brain. Unlike dopamine, L-DOPA can cross blood-brain barrier and is converted by dopa decarboxylase in the CNS to dopamine. Carbidopa, a peripheral DOPA decarboxylase inhibitor, is given with L-DOPA to ↑ bioavailability of L-DOPA in the brain and to limit peripheral side effects.
CLINICAL USE	Parkinson disease.
ADVERSE EFFECTS	Nausea, hallucinations, postural hypotension. With progressive disease, L-DOPA can lead to "on-off" phenomenon with improved mobility during "on" periods, then impaired motor function during "off" periods when patient responds poorly to L-DOPA or medication wears off.

Selegiline, rasagiline

MECHANISM	Selectively inhibit MAO-B (metabolize dopamine) → ↑ dopamine availability. **Sele**giline **sele**ctively inhibits MAO-**B** and is more commonly found in the **B**rain than in the periphery.
CLINICAL USE	Adjunctive agent to L-DOPA in treatment of Parkinson disease.
ADVERSE EFFECTS	May enhance adverse effects of L-DOPA.

Neurodegenerative disease therapy

DISEASE	AGENT	MECHANISM	NOTES
Alzheimer disease	**Don**epezil, **riva**stigmine, **gala**ntamine	AChE inhibitor	1st-line treatment Adverse effects: nausea, dizziness, insomnia **Don**a **Riva** dances at the **gala**
	Memantine	NMDA receptor antagonist; helps prevent excitotoxicity (mediated by Ca^{2+})	Used for moderate to advanced dementia Adverse effects: dizziness, confusion, hallucinations
Amyotrophic lateral sclerosis	Riluzole	↓ neuron glutamate excitotoxicity	↑ survival Treat **Lou** Gehrig disease with ril**ou**zole
Huntington disease	Tetrabenazine	Inhibit vesicular monoamine transporter (VMAT) → ↓ dopamine vesicle packaging and release	May be used for Huntington chorea and tardive dyskinesia

Anesthetics—general principles

CNS drugs must be lipid soluble (cross the blood-brain barrier) or be actively transported.
Drugs with ↓ solubility in blood = rapid induction and recovery times.

Drugs with ↑ solubility in lipids = ↑ potency = $\frac{1}{\text{MAC}}$

MAC = **M**inimum **A**lveolar **C**oncentration (of inhaled anesthetic) required to prevent 50% of subjects from moving in response to noxious stimulus (eg, skin incision).
Examples: nitrous oxide (N_2O) has ↓ blood and lipid solubility, and thus fast induction and low potency. Halothane has ↑ lipid and blood solubility, and thus high potency and slow induction.

(c) ketabton.com: The Digital Library

Inhaled anesthetics

Desflur**ane**, haloth**ane**, enflur**ane**, isoflur**ane**, sevoflur**ane**, methoxyflur**ane**, N_2O.

MECHANISM

Mechanism unknown.

EFFECTS

Myocardial depression, respiratory depression, postoperative nausea/vomiting, ↑ cerebral blood flow, ↓ cerebral metabolic demand.

ADVERSE EFFECTS

Hepatotoxicity (**h**alothane), **neph**rotoxicity (**meth**oxyflurane), proconvulsant (**e**nflurane, **e**pileptogenic), expansio**n** of trapped gas in a body cavity (N_2O).

Malignant hyperthermia—rare, life-threatening condition in which inhaled anesthetics or succinylcholine induce severe muscle contractions and hyperthermia. Susceptibility is often inherited as autosomal dominant with variable penetrance. Mutations in voltage-sensitive ryanodine receptor (*RYR1* gene) cause ↑ Ca^{2+} release from sarcoplasmic reticulum.

Treatment: dantrolene (a ryanodine receptor antagonist).

Intravenous anesthetics

AGENT	MECHANISM	ANESTHESIA USE	NOTES
Thiopental	Facilitates $GABA_A$ (barbiturate)	Anesthesia induction, short surgical procedures	↓ cerebral blood flow. High lipid solubility Effect terminated by rapid redistribution into tissue, fat
Midazolam	Facilitates $GABA_A$ (benzodiazepine)	Procedural sedation (eg, endoscopy), anesthesia induction	May cause severe postoperative respiratory depression, ↓ BP, anterograde amnesia
Propofol	Potentiates $GABA_A$	Rapid anesthesia induction, short procedures, ICU sedation	May cause respiratory depression, hypotension
Ketamine	NMDA receptor antagonist	Dissociative anesthesia Sympathomimetic	↑ cerebral blood flow Emergence reaction possible with disorientation, hallucination, vivid dreams

Local anesthetics

Esters—procaine, tetracaine, benzocaine, chloroprocaine.

Amides—l**I**doca**I**ne, mep**I**vaca**I**ne, bup**I**vaca**I**ne, rop**I**vaca**I**ne (am**I**des have **2 I**'s in name).

MECHANISM

Block Na^+ channels by binding to specific receptors on inner portion of channel. Most effective in rapidly firing neurons. 3° amine local anesthetics penetrate membrane in uncharged form, then bind to ion channels as charged form.

Can be given with vasoconstrictors (usually epinephrine) to enhance local action—↓ bleeding, ↑ anesthesia by ↓ systemic concentration.

In infected (acidic) tissue, alkaline anesthetics are charged and cannot penetrate membrane effectively → need more anesthetic.

Order of nerve blockade: small-diameter fibers > large diameter. Myelinated fibers > unmyelinated fibers. Overall, size factor predominates over myelination such that small myelinated fibers > small unmyelinated fibers > large myelinated fibers > large unmyelinated fibers.

Order of loss: (1) pain, (2) temperature, (3) touch, (4) pressure.

CLINICAL USE

Minor surgical procedures, spinal anesthesia. If allergic to esters, give amides.

ADVERSE EFFECTS

CNS excitation, severe cardiovascular toxicity (bupivacaine), hypertension, hypotension, arrhythmias (cocaine), methemoglobinemia (benzocaine).

(c) ketabton.com: The Digital Library

Neuromuscular blocking drugs	Muscle paralysis in surgery or mechanical ventilation. Selective for Nm nicotinic receptors at neuromuscular junction but not autonomic Nn receptors.
Depolarizing neuromuscular blocking drugs	Succinylcholine—strong ACh receptor agonist; produces sustained depolarization and prevents muscle contraction. Reversal of blockade: ▪ Phase I (prolonged depolarization)—no antidote. Block potentiated by cholinesterase inhibitors. ▪ Phase II (repolarized but blocked; ACh receptors are available, but desensitized)—may be reversed with cholinesterase inhibitors. Complications include hypercalcemia, hyperkalemia, malignant hyperthermia.
Nondepolarizing neuromuscular blocking drugs	Atracurium, cisatracurium, pancuronium, rocuronium, tubocurarine, vecuronium—competitive ACh antagonist. Reversal of blockade—cholinesterase inhibitors (eg, neostigmine, edrophonium) are given with anticholinergics (eg, atrophine, glycopyrrolate) to prevent muscarinic effects, such as bradycardia.

Spasmolytics, antispasmodics

DRUG	MECHANISM	CLINICAL USE	NOTES
Baclofen	$GABA_B$ receptor agonist in spinal cord.	Muscle spasticity, dystonia, multiple sclerosis.	Acts on the back (spinal cord).
Cyclobenzaprine	Acts within CNS, mainly at the brain stem.	Muscle spasticity.	Centrally acting. Structurally related to TCAs. May cause anticholinergic side effects, sedation.
Dantrolene	Prevents release of Ca^{2+} from sarcoplasmic reticulum of skeletal muscle by inhibiting the ryanodine receptor.	Malignant hyperthermia (toxicity of inhaled anesthetics and succinylcholine) and neuroleptic malignant syndrome (toxicity of antipsychotic drugs).	Acts Directly on muscle.
Tizanidine	α_2 agonist, acts centrally.	Muscle spasticity, multiple sclerosis, ALS, cerebral palsy.	

Opioid analgesics

MECHANISM	Act as agonists at opioid receptors (μ = β-endorphin, δ = enkephalin, κ = dynorphin) to modulate synaptic transmission—close presynaptic Ca^{2+} channels, open postsynaptic K^+ channels → ↓ synaptic transmission. Inhibit release of ACh, norepinephrine, 5-HT, glutamate, substance P.
EFFICACY	Full agonist: morphine, heroin, meperidine, methadone, codeine, fentanyl. Partial agonist: buprenorphine. Mixed agonist/antagonist: nalbuphine, pentazocine, butorphanol. Antagonist: naloxone, naltrexone, methylnaltrexone.
CLINICAL USE	Moderate to severe or refractory pain, diarrhea (loperamide, diphenoxylate), acute pulmonary edema, maintenance programs for heroin addicts (methadone, buprenorphine + naloxone).
ADVERSE EFFECTS	Nausea, vomiting, pruritus, addiction, respiratory depression, constipation, sphincter of Oddi spasm, miosis (except meperidine → mydriasis), additive CNS depression with other drugs. Tolerance does not develop to miosis and constipation. Treat toxicity with naloxone (competitive opioid receptor antagonist) and prevent relapse with naltrexone once detoxified.

(c) ketabton.com: The Digital Library

Mixed agonist and antagonist opioid analgesics

DRUG	MECHANISM	CLINICAL USE	NOTES
Pentazocine	κ-opioid receptor agonist and μ-opioid receptor weak antagonist or partial agonist.	Analgesia for moderate to severe pain.	Can cause opioid withdrawal symptoms if patient is also taking full opioid agonist (due to competition for opioid receptors).
Butorphanol	κ-opioid receptor agonist and μ-opioid receptor partial agonist.	Severe pain (eg, migraine, labor).	Causes less respiratory depression than full opioid agonists. Use with full opioid agonist can precipitate withdrawal. Not easily reversed with naloxone.

Tramadol

MECHANISM	Very weak opioid agonist; also inhibits the reuptake of norepinephrine and serotonin.
CLINICAL USE	Chronic pain.
ADVERSE EFFECTS	Similar to opioids; decreases seizure threshold; serotonin syndrome.

Tramadol is a **S**light opioid agonist, and a **S**erotonin and norepinephrine reuptake inhibitor. It is used for **S**tubborn pain, but can lower **S**eizure threshold, and may cause **S**erotonin **S**yndrome.

Glaucoma therapy

↓ IOP via ↓ amount of aqueous humor (inhibit synthesis/secretion or ↑ drainage).
BAD humor may not be **P**olitically **C**orrect.

DRUG CLASS	EXAMPLES	MECHANISM	ADVERSE EFFECTS
β-blockers	Timolol, betaxolol, carteolol	↓ aqueous humor synthesis	No pupillary or vision changes
α-agonists	Epinephrine (α_1), apraclonidine, brimonidine (α_2)	↓ aqueous humor synthesis via vasoconstriction (epinephrine) ↓ aqueous humor synthesis (apraclonidine, brimonidine)	Mydriasis (α_1); do not use in closed-angle glaucoma Blurry vision, ocular hyperemia, foreign body sensation, ocular allergic reactions, ocular pruritus
Diuretics	Acetazolamide	↓ aqueous humor synthesis via inhibition of carbonic anhydrase	No pupillary or vision changes
Prostaglandins	Bimatoprost, latanoprost ($PGF_{2\alpha}$)	↑ outflow of aqueous humor via ↓ resistance of flow through uveoscleral pathway	Darkens color of iris (browning), eyelash growth
Cholinomimetics (M_3)	Direct: pilocarpine, carbachol Indirect: physostigmine, echothiophate	↑ outflow of aqueous humor via contraction of ciliary muscle and opening of trabecular meshwork Use pilocarpine in acute angle closure glaucoma—very effective at opening meshwork into canal of Schlemm	Miosis (contraction of pupillary sphincter muscles) and cyclospasm (contraction of ciliary muscle)

(c) ketabton.com: The Digital Library

HIGH-YIELD PRINCIPLES IN

Psychiatry

"Words of comfort, skillfully administered, are the oldest therapy known to man."

—Louis Nizer

"All men should strive to learn before they die what they are running from, and to, and why."

—James Thurber

"The sorrow which has no vent in tears may make other organs weep."

—Henry Maudsley

"It's no use going back to yesterday, because I was a different person then."

—Lewis Carroll, Alice in Wonderland

This chapter encompasses overlapping areas in psychiatry, psychology, sociology, and psychopharmacology. High-yield topics include schizophrenia, mood disorders, eating disorders, personality disorders, somatic symptom disorders, substance abuse, and antipsychotic agents. Know the DSM-5 criteria for diagnosing common psychiatric disorders.

Psychology 554

Pathology 556

Pharmacology 572

FAS1_2019_13-Psych.indd 553 11/7/19 5:28 PM

(c) ketabton.com: The Digital Library

▸ PSYCHIATRY—PSYCHOLOGY

Classical conditioning	Learning in which a natural response (salivation) is elicited by a conditioned, or learned, stimulus (bell) that previously was presented in conjunction with an unconditioned stimulus (food).	Usually elicits **involuntary** responses. Pavlov's classical experiments with dogs—ringing the bell provoked salivation.

Operant conditioning	Learning in which a particular action is elicited because it produces a punishment or reward. Usually elicits **voluntary** responses.
Reinforcement	Target behavior (response) is followed by desired reward (positive reinforcement) or removal of aversive stimulus (negative reinforcement).
Punishment	Repeated application of aversive stimulus (positive punishment) or removal of desired reward (negative punishment) to extinguish unwanted behavior.
Extinction	Discontinuation of reinforcement (positive or negative) eventually eliminates behavior. Can occur in operant or classical conditioning.

Skinner operant conditioning quadrants:

	Increase behavior	Decrease behavior
Add a stimulus	Positive reinforcement	Positive punishment
Remove a stimulus	Negative reinforcement	Negative punishment

Transference and countertransference	
Transference	Patient projects feelings about formative or other important persons onto physician (eg, psychiatrist is seen as parent).
Countertransference	Doctor projects feelings about formative or other important persons onto patient (eg, patient reminds physician of younger sibling).

Ego defenses — Thoughts and behaviors (voluntary or involuntary) used to resolve conflict and prevent undesirable feelings (eg, anxiety, depression).

IMMATURE DEFENSES	DESCRIPTION	EXAMPLE
Acting out	Subconsciously coping with stressors or emotional conflict using actions rather than reflections or feelings.	A patient skips therapy appointments after deep discomfort from dealing with his past.
Denial	Avoiding the awareness of some painful reality.	A patient with cancer plans a full-time work schedule despite being warned of significant fatigue during chemotherapy.
Displacement	Redirection of emotions or impulses to a neutral person or object (vs projection).	After being reprimanded by her principal, a frustrated teacher returns home and criticizes her husband's cooking instead of confronting the principal directly.
Dissociation	Temporary, drastic change in personality, memory, consciousness, or motor behavior to avoid emotional stress. Patient has incomplete or no memory of traumatic event.	A victim of sexual abuse suddenly appears numb and detached when she is exposed to her abuser.

(c) ketabton.com: The Digital Library

Ego defenses *(continued)*

IMMATURE DEFENSES	DESCRIPTION	EXAMPLE
Fixation	Partially remaining at a more childish level of development (vs regression).	A surgeon throws a tantrum in the operating room because the last case ran very late.
Idealization	Expressing extremely positive thoughts of self and others while ignoring negative thoughts.	A patient boasts about his physician and his accomplishments while ignoring any flaws.
Identification	Largely unconscious assumption of the characteristics, qualities, or traits of another person or group.	A resident starts putting his stethoscope in his pocket like his favorite attending, instead of wearing it around his neck like before.
Intellectualization	Using facts and logic to emotionally distance oneself from a stressful situation.	A patient diagnosed with cancer discusses the pathophysiology of the disease.
Isolation (of affect)	Separating feelings from ideas and events.	Describing murder in graphic detail with no emotional response.
Passive aggression	Demonstrating hostile feelings in a nonconfrontational manner; showing indirect opposition.	A disgruntled employee is repeatedly late to work, but won't admit it is a way to get back at the manager.
Projection	Attributing an unacceptable internal impulse to an external source (vs displacement).	A man who wants to cheat on his wife accuses his wife of being unfaithful.
Rationalization	Asserting plausible explanations for events that actually occurred for other reasons, usually to avoid self-blame.	A man who was recently fired claims that the job was not important anyway.
Reaction formation	Replacing a warded-off idea or feeling with an emphasis on its opposite (vs sublimation).	A stepmother treats a child she resents with excessive nurturing and overprotection.
Regression	Involuntarily turning back the maturational clock to behaviors previously demonstrated under stress (vs fixation).	A previously toilet-trained child begins bedwetting again following the birth of a sibling.
Repression	Involuntarily withholding an idea or feeling from conscious awareness (vs suppression).	A 20-year-old does not remember going to counseling during his parents' divorce 10 years earlier.
Splitting	Believing that people are either all good or all bad at different times due to intolerance of ambiguity. Common in borderline personality disorder.	A patient says that all the nurses are cold and insensitive, but the doctors are warm and friendly.
MATURE DEFENSES		
Sublimation	Replacing an unacceptable wish with a course of action that is similar to the wish but socially acceptable (vs reaction formation).	A teenager's aggression toward his parents because of their high expectations is channeled into excelling in sports.
Altruism	Alleviating negative feelings via unsolicited generosity, which provides gratification (vs reaction formation).	A mafia boss makes a large donation to charity.
Suppression	Intentionally withholding an idea or feeling from conscious awareness (vs repression); temporary.	An athlete focuses on other tasks to prevent worrying about an important upcoming match.
Humor	Lightheartedly expressing uncomfortable feelings to shift the internal focus away from the distress.	A nervous medical student jokes about the boards.
	Mature adults wear a **SASH**.	

(c) ketabton.com: The Digital Library

▸ PSYCHIATRY—PATHOLOGY

Infant deprivation effects

Long-term deprivation of affection results in:
- Failure to thrive
- Poor language/socialization skills
- Lack of basic trust
- Reactive attachment disorder (infant withdrawn/unresponsive to comfort)
- Disinhibited social engagement (child indiscriminately attaches to strangers)

Deprivation for > 6 months can lead to irreversible changes.
Severe deprivation can result in infant death.

Child abuse

	Physical abuse	Sexual abuse	Emotional abuse
SIGNS	Fractures, bruises, or burns. Injuries often in different stages of healing or in patterns resembling possible implements of injury. Includes abusive head trauma (shaken baby syndrome), characterized by subdural hematomas or retinal hemorrhages. Caregivers may delay seeking medical attention for the child or provide explanations inconsistent with the child's developmental stage or pattern of injury.	STIs, UTIs, and genital, anal, or oral trauma. Most often, there are no physical signs; sexual abuse should not be excluded from a differential diagnosis in the absence of physical trauma. Children often exhibit sexual knowledge or behavior incongruent with their age.	Babies or young children may lack a bond with the caregiver but are overly affectionate with less familiar adults. They may be aggressive toward children and animals or unusually anxious. Older children are often emotionally labile and prone to angry outbursts. They may distance themselves from caregivers and other children. They can experience vague somatic symptoms for which a medical cause cannot be found.
EPIDEMIOLOGY	40% of deaths related to child abuse or neglect occur in children < 1 year old.	Peak incidence 9–12 years old.	~80% of young adult victims of child emotional abuse meet the criteria for ≥ 1 psychiatric illness by age 21.

Child neglect

Failure to provide a child with adequate food, shelter, supervision, education, and/or affection. Most common form of child maltreatment. Signs: poor hygiene, malnutrition, withdrawal, impaired social/emotional development, failure to thrive.
As with child abuse, suspected child neglect must be reported to local child protective services.

Vulnerable child syndrome

Parents perceive the child as especially susceptible to illness or injury (vs factitious disorder imposed on another). Usually follows a serious illness or life-threatening event. Can result in missed school or overuse of medical services.

(c) ketabton.com: The Digital Library

Childhood and early-onset disorders

Attention-deficit hyperactivity disorder	Onset before age 12. ≥ 6 months of limited attention span and/or poor impulse control. Characterized by hyperactivity, impulsivity, and/or inattention in ≥ 2 settings (eg, school, home, places of worship). Normal intelligence, but commonly coexists with difficulties in school. Often persists into adulthood. Commonly coexists with oppositional defiant disorder. Treatment: stimulants (eg, methylphenidate) +/– behavioral therapy; alternatives include atomoxetine, guanfacine, clonidine.
Autism spectrum disorder	Onset in early childhood. Social and communication deficits, repetitive/ritualized behaviors, restricted interests. May be accompanied by intellectual disability and/or above average abilities in specific skills (eg, music). More common in boys. Associated with ↑ head and/or brain size.
Conduct disorder	Repetitive, pervasive behavior violating societal norms or the basic rights of others (eg, aggression toward people and animals, destruction of property, theft). After age 18, often reclassified as antisocial personality disorder. Treatment: psychotherapy (eg, cognitive behavioral therapy [CBT]).
Disruptive mood dysregulation disorder	Onset before age 10. Severe, recurrent temper outbursts out of proportion to situation. Child is constantly angry and irritable between outbursts. Treatment: CBT, stimulants, antipsychotics.
Intellectual disability	Global cognitive deficits (vs specific learning disorder) that affect reasoning, memory, abstract thinking, judgment, language, learning. Adaptive functioning is impaired, leading to major difficulties with education, employment, communication, socialization, independence. Treatment: psychotherapy, occupational therapy, special education.
Oppositional defiant disorder	Enduring pattern of anger and irritability with argumentative, vindictive, and defiant behavior toward authority figures. Treatment: psychotherapy (eg, CBT).
Selective mutism	Onset before age 5. Anxiety disorder lasting ≥ 1 month involving refraining from speech in certain situations despite speaking in other, usually more comfortable situations. Development (eg, speech and language) not typically impaired. Interferes with social, academic, and occupational tasks. Commonly coexists with social anxiety disorder. Treatment: behavioral, family, and play therapy; SSRIs.
Separation anxiety disorder	Overwhelming fear of separation from home or attachment figure lasting ≥ 4 weeks. Can be normal behavior up to age 3–4. May lead to factitious physical complaints to avoid school. Treatment: CBT, play therapy, family therapy.
Specific learning disorder	Onset during school-age years. Inability to acquire or use information from a specific subject (eg, math, reading, writing) near age-expected proficiency for ≥ 6 months despite focused intervention. General functioning and intelligence are normal (vs intellectual disability). Treatment: academic support, counseling, extracurricular activities.
Tourette syndrome	Onset before age 18. Sudden, Sudden, recurrent, nonrhythmic, stereotyped motor and vocal tics that persist for > 1 year. Coprolalia (involuntary obscene speech) found in some patients. Associated with OCD and ADHD. Treatment: psychoeducation, behavioral therapy. For intractable and distressing tics, high-potency antipsychotics (eg, haloperidol, fluphenazine), tetrabenazine, α_2-agonists (eg, guanfacine, clonidine), or atypical antipsychotics.

Orientation	Patients' ability to know the date and time, where they are, and who they are (order of loss: time → place → person). Common causes of loss of orientation: alcohol, drugs, fluid/electrolyte imbalance, head trauma, hypoglycemia, infection, nutritional deficiencies, hypoxia.

(c) ketabton.com: The Digital Library

Amnesias

Retrograde amnesia	Inability to remember things that occurred before a CNS insult.
Anterograde amnesia	Inability to remember things that occurred after a CNS insult (↓ acquisition of new memory).
Korsakoff syndrome	Amnesia (anterograde > retrograde) and disorientation caused by vitamin B_1 deficiency. Associated with disruption and destruction of the limbic system, especially mammillary bodies and anterior thalamus. Seen in alcoholics as a late neuropsychiatric manifestation of Wernicke encephalopathy. Confabulations are characteristic.

Dissociative disorders

Depersonalization/ derealization disorder	Persistent feelings of detachment or estrangement from one's own body, thoughts, perceptions, and actions (depersonalization) or one's environment (derealization). Intact reality testing (vs psychosis).
Dissociative amnesia	Inability to recall important personal information, usually following severe trauma or stress. May be accompanied by dissociative fugue (abrupt, unexpected travelling away from home).
Dissociative identity disorder	Formerly called multiple personality disorder. Presence of ≥ 2 distinct identities or personality states. More common in women. Associated with history of sexual abuse, PTSD, depression, substance abuse, borderline personality, somatic symptom disorders.

Delirium

"Waxing and waning" level of consciousness with acute onset, ↓ attention span, ↓ level of arousal. Characterized by disorganized thinking, hallucinations (often visual), misperceptions (eg, illusions), disturbance in sleep-wake cycle, cognitive dysfunction, agitation. Reversible.

Usually 2° to other identifiable illness (eg, CNS disease, infection, trauma, substance abuse/ withdrawal, metabolic/electrolyte disturbances, hemorrhage, urinary/fecal retention), or medications (eg, anticholinergics), especially in the elderly.

Most common presentation of altered mental status in inpatient setting, especially in the ICU or during prolonged hospital stays. EEG may show diffuse background rhythm slowing.

Delirium = changes in sensorium.

Treatment: identification and management of underlying condition. Orientation protocols (eg, keeping a clock or calendar nearby), ↓ sleep disturbances, and ↑ cognitive stimulation to manage symptoms. Antipsychotics as needed. Avoid unnecessary restraints and drugs that may worsen delirium (eg, anticholinergics, benzodiazepines, opioids).

(c) ketabton.com: The Digital Library

Psychosis	Distorted perception of reality characterized by delusions, hallucinations, and/or disorganized thought/speech. Can occur in patients with medical illness, psychiatric illness, or both.
Delusions	False, fixed, idiosyncratic beliefs that persist despite evidence to the contrary and are not typical of a patient's culture or religion (eg, a patient who believes that others are reading his thoughts). Types include erotomanic, grandiose, jealous, persecutory, somatic, mixed, and unspecified.
Disorganized thought	Speech may be incoherent ("word salad"), tangential, or derailed ("loose associations").
Hallucinations	Perceptions in the absence of external stimuli (eg, seeing a light that is not actually present). Contrast with misperceptions (eg, illusions) of real external stimuli. Types include: ▪ Auditory—more commonly due to psychiatric illness (eg, schizophrenia) than medical illness. ▪ Visual—more commonly due to medical illness (eg, drug intoxication, delirium) than psychiatric illness. ▪ Tactile—common in alcohol withdrawal and stimulant use (eg, "cocaine crawlies," a type of delusional parasitosis). ▪ Olfactory—often occur as an aura of temporal lobe epilepsy (eg, burning rubber) and in brain tumors. ▪ Gustatory—rare, but seen in epilepsy. ▪ Hypnagogic—occurs while going to sleep. Sometimes seen in narcolepsy. ▪ Hypnopompic—occurs while waking from sleep ("get pomped up in the morning"). Sometimes seen in narcolepsy.

(c) ketabton.com: The Digital Library

Schizophrenia spectrum disorders

Schizophrenia

Chronic illness causing profound functional impairment. Symptom categories include:
- Positive—hallucinations, delusions, unusual thought processes, disorganized speech, bizarre behavior
- Negative—flat or blunted affect, apathy, anhedonia, alogia, social withdrawal
- Cognitive—reduced ability to understand or make plans, diminished working memory, inattention

Diagnosis requires ≥ 2 of the following active symptoms, including ≥ 1 from symptoms #1–3:
1. Delusions
2. Hallucinations, often auditory
3. Disorganized speech
4. Disorganized or catatonic behavior
5. Negative symptoms

Requires ≥ 1 month of active symptoms over the past 6 months; onset ≥ 6 months prior to diagnosis.

Associated with altered dopaminergic activity, ↑ serotonergic activity, and ↓ dendritic branching. Ventriculomegaly on brain imaging. Lifetime prevalence—1.5% (males > females). Presents earlier in men (late teens to early 20s) than in women (late 20s to early 30s). ↑ suicide risk.

Heavy cannabis use in adolescence is associated with ↑ incidence and worsened course of psychotic, mood, and anxiety disorders.

Treatment: atypical antipsychotics (eg, risperidone) are first line.

Negative symptoms often persist after treatment, despite resolution of positive symptoms.

Brief psychotic disorder—≥ 1 positive symptom(s) lasting < 1 month, usually stress-related.
Schizophreniform disorder—≥ 2 symptoms lasting 1–6 months.

Schizoaffective disorder

Shares symptoms with both schizophrenia and mood disorders (major depressive or bipolar disorder). To differentiate from a mood disorder with psychotic features, patient must have > 2 weeks of psychotic symptoms without a manic or depressive episode.

Delusional disorder

≥ 1 delusion(s) lasting > 1 month, but without a mood disorder or other psychotic symptoms. Daily functioning, including socialization, may be impacted by the pathological, fixed belief but is otherwise unaffected. Can be shared by individuals in close relationships (folie à deux).

Schizotypal personality disorder

Cluster A personality disorder that also falls on the schizophrenia spectrum. May include brief psychotic episodes (eg, delusions) that are less frequent and severe than in schizophrenia.

Mood disorder

Characterized by an abnormal range of moods or internal emotional states and loss of control over them. Severity of moods causes distress and impairment in social and occupational functioning. Includes major depressive, bipolar, dysthymic, and cyclothymic disorders. Episodic superimposed psychotic features (delusions, hallucinations, disorganized speech/behavior) may be present.

Manic episode

Distinct period of abnormally and persistently elevated, expansive, or irritable mood and ↑ activity or energy lasting ≥ 1 week. Diagnosis requires hospitalization or marked functional impairment with ≥ 3 of the following (manics **DIG FAST**):
- **D**istractibility
- **I**mpulsivity/**I**ndiscretion—seeks pleasure without regard to consequences (hedonistic)
- **G**randiosity—inflated self-esteem
- **F**light of ideas—racing thoughts
- ↑ goal-directed **A**ctivity/psychomotor **A**gitation
- ↓ need for **S**leep
- **T**alkativeness or pressured speech

(c) ketabton.com: The Digital Library

Hypomanic episode

Similar to a manic episode except mood disturbance is not severe enough to cause marked impairment in social and/or occupational functioning or to necessitate hospitalization. Abnormally ↑ activity or energy usually present. No psychotic features. Lasts ≥ 4 consecutive days.

Bipolar disorder

Bipolar I—≥ 1 manic episode +/– a hypomanic or depressive episode (may be separated by any length of time).

Bipolar II—a hypomanic and a depressive episode (no history of manic episodes).

Patient's mood and functioning usually normalize between episodes. Use of antidepressants can destabilize mood. High suicide risk. Treatment: mood stabilizers (eg, lithium, valproic acid, carbamazepine, lamotrigine), atypical antipsychotics.

Cyclothymic disorder—milder form of bipolar disorder fluctuating between mild depressive and hypomanic symptoms. Must last ≥ 2 years with symptoms present at least half of the time, with any remission lasting ≤ 2 months.

Major depressive disorder

Recurrent episodes lasting ≥ 2 weeks characterized by ≥ 5 of 9 diagnostic symptoms (must include depressed mood or anhedonia) (DIGS SPACE):

- **D**epressed mood (or irritability in children)
- ↓ **I**nterest (anhedonia)
- **G**uilt or feelings of worthlessness
- **S**leep disturbances
- **S**uicidal ideation
- **P**sychomotor retardation or agitation
- **A**ppetite/weight changes
- ↓ **C**oncentration
- ↓ **E**nergy

Screen for previous manic or hypomanic episodes to rule out bipolar disorder.

Treatment: CBT and SSRIs are first line. Also SNRIs, mirtazapine, bupropion, electroconvulsive therapy (ECT).

MDD with psychotic features

MDD + hallucinations or delusions. Psychotic features are typically mood congruent (eg, depressive themes of inadequacy, guilt, punishment, nihilism, disease, or death) and occur only in the context of major depressive episode (vs schizoaffective disorder). Treatment: antidepressant with atypical antipsychotic, ECT.

Persistent depressive disorder (dysthymia)

Often milder than MDD; ≥ 2 depressive symptoms lasting ≥ 2 years (≥ 1 year in children), with any remission lasting ≤ 2 months.

MDD with seasonal pattern

Formerly called seasonal affective disorder. Major depressive episodes occurring only during a particular season (usually winter) in ≥ 2 consecutive years and in most years across a lifetime. Atypical symptoms common.

Depression with atypical features

Characterized by mood reactivity (transient improvement in response to a positive event), hypersomnia, hyperphagia, leaden paralysis (heavy feeling in arms and legs), long-standing interpersonal rejection sensitivity. Most common subtype of depression. Treatment: CBT and SSRIs are first line. MAO inhibitors (MAOIs) are effective but not first line because of their risk profile.

(c) ketabton.com: The Digital Library

Peripartum mood disturbances

Onset during or shortly after pregnancy or within 4 weeks of delivery. ↑ risk with history of mood disorders.

Maternal (postpartum) blues	50–85% incidence rate. Characterized by depressed affect, tearfulness, and fatigue starting 2–3 days after delivery. Usually resolves within 2 weeks. Treatment: supportive. Follow up to assess for possible MDD with peripartum onset.
MDD with peripartum onset	10–15% incidence rate. Formerly called postpartum depression. Meets MDD criteria with onset no later than 1 year after delivery. Treatment: CBT and SSRIs are first line.
Postpartum psychosis	0.1–0.2% incidence rate. Characterized by mood-congruent delusions, hallucinations, and thoughts of harming the baby or self. Risk factors include first pregnancy, family history, bipolar disorder, psychotic disorder, recent medication change. Treatment: hospitalization and initiation of atypical antipsychotic; if insufficient, ECT may be used.

Grief

The five stages of grief per the Kübler-Ross model are denial, anger, bargaining, depression, and acceptance (may occur in any order). Other normal grief symptoms include shock, guilt, sadness, anxiety, yearning, and somatic symptoms that usually occur in waves. Simple hallucinations of the deceased person are common (eg, hearing the deceased speaking). Any thoughts of dying are limited to joining the deceased (vs complicated grief). Duration varies widely; usually resolves within 6–12 months.

Persistent complex bereavement disorder involves obsessive preoccupation with the deceased and causes functional impairment, lasting at least 12 months (6 months in children). Can also meet criteria for major depressive episode.

Electroconvulsive therapy

Rapid-acting method to treat refractory depression, depression with psychotic symptoms, catatonia, and acute suicidality. Induces tonic-clonic seizure under anesthesia and neuromuscular blockade. Adverse effects include disorientation, headache, partial anterograde/retrograde amnesia usually resolving in 6 months. No absolute contraindications. Safe in pregnant and elderly individuals.

Risk factors for suicide completion

Sex (male)
Age (young adult or elderly)
Depression
Previous attempt (highest risk factor)
Ethanol or drug use
Rational thinking loss (psychosis)
Sickness (medical illness)
Organized plan
No spouse or other social support
Stated future intent

SAD PERSONS are more likely to complete suicide.

Most common method in US is firearms; access to guns ↑ risk of suicide completion.

Women try more often; men complete more often.

Other risk factors include recent psychiatric hospitalization and family history of completed suicide.

Anxiety disorders

Inappropriate experiences of fear/worry and their physical manifestations incongruent with the magnitude of the stressors. Symptoms are not attributable to another psychiatric disorder, medical condition (eg, hyperthyroidism), or substance abuse. Includes panic disorder, phobias, generalized anxiety disorder, and selective mutism.

(c) ketabton.com: The Digital Library

Panic disorder

Recurrent panic attacks involving intense fear and discomfort +/– a known trigger. Attacks typically peak in 10 minutes with ≥ 4 of the following: palpitations, paresthesias, depersonalization or derealization, abdominal pain, nausea, intense fear of dying, intense fear of losing control, lightheadedness, chest pain, chills, choking, sweating, shaking, shortness of breath. Strong genetic component. ↑ risk of suicide.

Diagnosis requires attack followed by ≥ 1 month of ≥ 1 of the following:

- Persistent concern of additional attacks
- Worrying about consequences of attack
- Behavioral change related to attacks

Symptoms are systemic manifestations of fear.

Treatment: CBT, SSRIs, and venlafaxine are first line. Benzodiazepines occasionally used in acute setting.

Phobias

Severe, persistent (≥ 6 months) fear or anxiety due to presence or anticipation of a specific object or situation. Person often recognizes fear is excessive. Treatment: CBT with exposure therapy.

Social anxiety disorder—exaggerated fear of embarrassment in social situations (eg, public speaking, using public restrooms). Treatment: CBT, SSRIs, venlafaxine. For performance type (eg, anxiety restricted to public speaking), use β-blockers or benzodiazepines as needed.

Agoraphobia—irrational fear/anxiety while facing or anticipating ≥ 2 specific situations (eg, open/closed spaces, lines, crowds, public transport). If severe, patients may refuse to leave their homes. Associated with panic disorder. Treatment: CBT, SSRIs.

Generalized anxiety disorder

Excessive anxiety and worry about different aspects of daily life (eg, work, school, children) for most days of ≥ 6 months. Associated with ≥ 3 of the following for adults (≥ 1 for kids): restlessness, irritability, sleep disturbance, fatigue, muscle tension, difficulty concentrating. Treatment: CBT, SSRIs, SNRIs are first line. Buspirone, TCAs, benzodiazepines are second line.

Obsessive-compulsive disorders

Obsessions (recurring intrusive thoughts, feelings, or sensations) that cause severe distress, relieved in part by compulsions (performance of repetitive, often time-consuming actions). Ego-dystonic: behavior inconsistent with one's beliefs and attitudes (vs obsessive-compulsive personality disorder, ego-syntonic). Associated with Tourette syndrome. Treatment: CBT and SSRIs; clomipramine and venlafaxine are second line.

Body dysmorphic disorder—preoccupation with minor or imagined defects in appearance. Causes significant emotional distress and repetitive appearance-related behaviors (eg, mirror checking, excessive grooming). Common in eating disorders. Treatment: CBT.

Trichotillomania

Compulsively pulling out one's hair. Causes significant distress and persists despite attempts to stop. Presents with areas of thinning hair or baldness on any area of the body, most commonly the scalp A. Incidence highest in childhood but spans all ages. Treatment: psychotherapy.

(c) ketabton.com: The Digital Library

Trauma and stress-related disorders

Adjustment disorder	Emotional or behavioral symptoms (eg, anxiety, outbursts) that occur within 3 months of an identifiable psychosocial stressor (eg, divorce, illness) lasting < 6 months once the stressor has ended. If symptoms persist > 6 months after stressor ends, it is GAD. Symptoms do not meet criteria for MDD. Treatment: CBT is first line; antidepressants and anxiolytics may be considered.
Post-traumatic stress disorder	Experiencing, or discovering that a loved one has experienced, a life-threatening situation (eg, serious injury, rape, witnessing death) → persistent **H**yperarousal, **A**voidance of associated stimuli, intrusive **R**e-experiencing of the event (eg, nightmares, flashbacks), changes in cognition or mood (eg, fear, horror, **D**istress) (having PTSD is **HARD**). Disturbance lasts > 1 month with significant distress or impaired functioning. Treatment: CBT, SSRIs, and venlafaxine are first line. Prazosin can reduce nightmares. Acute stress disorder—lasts between 3 days and 1 month. Treatment: CBT; pharmacotherapy is usually not indicated.

Diagnostic criteria by symptom duration



(c) ketabton.com: The Digital Library

Personality

Personality trait	An enduring, repetitive pattern of perceiving, relating to, and thinking about the environment and oneself.
Personality disorder	Inflexible, maladaptive, and rigidly pervasive pattern of behavior causing subjective distress and/or impaired functioning; person is usually not aware of problem (ego-syntonic). Usually presents by early adulthood. Three clusters: **A**, **B**, **C**; remember as **Weird**, **Wild**, and **Worried**, respectively, based on symptoms.

Cluster A personality disorders	Odd or eccentric; inability to develop meaningful social relationships. No psychosis; genetic association with schizophrenia.	Cluster **A**: **A**ccusatory, **A**loof, **A**wkward. "**Weird**."
Paranoid	Pervasive distrust (**A**ccusatory), suspiciousness, hypervigilance, and a profoundly cynical view of the world.	
Schizoid	Voluntary social withdrawal (**A**loof), limited emotional expression, content with social isolation (vs avoidant).	
Schizotypal	Eccentric appearance, odd beliefs or magical thinking, interpersonal **A**wkwardness.	Included on the schizophrenia spectrum. Pronounce schiz**o**-**type**-al: **o**dd-**type** thoughts.

Cluster B personality disorders	Dramatic, emotional, or erratic; genetic association with mood disorders and substance abuse.	Cluster **B**: **B**ad, **B**orderline, flam**B**oyant, must be the **B**est. "**Wild**."
Antisocial	Disregard for the rights of others with lack of remorse. Involves criminality, impulsivity, hostility, and manipulation. Males > females. Must be ≥ 18 years old with evidence of conduct disorder onset before age 15. Diagnosis is conduct disorder if < 18 years old.	Anti**soc**ial = **soc**iopath. **B**ad.
Borderline	Unstable mood and interpersonal relationships, fear of abandonment, impulsivity, self-mutilation, suicidality, sense of emotional emptiness. Females > males. Splitting is a major defense mechanism.	Treatment: dialectical behavior therapy. **B**orderline.
Histrionic	Attention-seeking, dramatic speech and emotional expression, shallow and labile emotions, sexually provocative. May use physical appearance to draw attention.	Flam**B**oyant.
Narcissistic	Grandiosity, sense of entitlement; lacks empathy and requires excessive admiration; often demands the "best" and reacts to criticism with rage and/or defensiveness. Fragile self-esteem. Often envious of others.	Must be the **B**est.

(c) ketabton.com: The Digital Library

Cluster C personality disorders	Anxious or fearful; genetic association with anxiety disorders.	Cluster C: Cowardly, obsessive-Compulsive, Clingy. "Worried."
Avoidant	Hypersensitive to rejection and criticism, socially inhibited, timid, feelings of inadequacy, desires relationships with others (vs schizoid).	Cowardly.
Obsessive-Compulsive	Preoccupation with order, perfectionism, and control; ego-syntonic: behavior consistent with one's own beliefs and attitudes (vs OCD).	
Dependent	Excessive need for support, low self-confidence. Patients often get stuck in abusive relationships.	Submissive and Clingy.
Malingering	Symptoms are **intentional**, motivation is **intentional**. Patient **consciously** fakes, profoundly exaggerates, or claims to have a disorder in order to attain a specific **2° (external) gain** (eg, avoiding work, obtaining compensation). Poor compliance with treatment or follow-up of diagnostic tests. Complaints cease after gain (vs factitious disorder).	
Factitious disorders	Symptoms are **intentional**, motivation is **unconscious**. Patient **consciously** creates physical and/or psychological symptoms in order to assume "sick role" and to get medical attention and sympathy (**1° [internal] gain**).	
Factitious disorder imposed on self	Formerly called Munchausen syndrome. Chronic factitious disorder with predominantly physical signs and symptoms. Characterized by a history of multiple hospital admissions and willingness to undergo invasive procedures. More common in women and healthcare workers.	
Factitious disorder imposed on another	Formerly called Munchausen syndrome by proxy. Illness in a child or elderly patient is caused or fabricated by the caregiver. Motivation is to assume a sick role by proxy. Form of child/elder abuse.	
Somatic symptom and related disorders	Symptoms are **unconscious**, motivation is **unconscious**. Category of disorders characterized by physical symptoms causing significant distress and impairment. Symptoms not intentionally produced or feigned.	
Somatic symptom disorder	≥ 1 bodily complaints (eg, abdominal pain, fatigue) lasting months to years. Associated with excessive, persistent thoughts and anxiety about symptoms. May co-occur with medical illness. Treatment: regular office visits with the same physician in combination with psychotherapy.	
Conversion disorder	Also called functional neurologic symptom disorder. Loss of sensory or motor function (eg, paralysis, blindness, mutism), often following an acute stressor; patient may be aware of but indifferent toward symptoms (*la belle indifférence*); more common in females, adolescents, and young adults.	
Illness anxiety disorder	Preoccupation with acquiring or having a serious illness, often despite medical evaluation and reassurance; minimal to no somatic symptoms.	

(c) ketabton.com: The Digital Library

Eating disorders	Most common in young women.
Anorexia nervosa	Intense fear of weight gain, overvaluation of thinness, and body image distortion leading to calorie restriction and severe weight loss resulting in inappropriately low body weight. Binge-eating/purging type—recurring purging behaviors (eg, laxative or diuretic abuse, self-induced vomiting) or binge eating over the last 3 months. Restricting type—primary disordered behaviors include dieting, fasting, and/or over-exercising. No recurring purging behaviors or binge eating over the last 3 months. Refeeding syndrome—often occurs in significantly malnourished patients with sudden ↑ calorie intake → ↑ insulin → ↓ PO_4^{3-}, ↓ K^+, ↓ Mg^{2+} → cardiac complications, rhabdomyolysis, seizures. Treatment: psychotherapy, nutritional rehabilitation, antidepressants (eg, SSRIs).
Bulimia nervosa	Recurring episodes of binge eating with compensatory purging behaviors at least weekly over the last 3 months. BMI often normal or slightly overweight (vs anorexia). Associated with parotid gland hypertrophy (may see ↑ serum amylase), enamel erosion, Mallory-Weiss syndrome, electrolyte disturbances (eg, ↓ K^+, ↓ Cl^-), metabolic alkalosis, dorsal hand calluses from induced vomiting (Russell sign). Treatment: psychotherapy, nutritional rehabilitation, antidepressants (eg, SSRIs). Bupropion is contraindicated due to seizure risk.
Binge-eating disorder	Recurring episodes of binge eating without purging behaviors at least weekly over the last 3 months. ↑ diabetes risk. Most common eating disorder in adults. Treatment: psychotherapy (first line); SSRIs; lisdexamfetamine.
Pica	Recurring episodes of eating non-food substances (eg, dirt, hair, paint chips) over ≥ 1 month that are not culturally or developmentally recognized as normal. May provide temporary emotional relief. Common in children and during pregnancy. Associated with malnutrition, iron deficiency anemia, developmental disabilities, emotional trauma. Treatment: psychotherapy and nutritional rehabilitation (first line); SSRIs (second line).

Gender dysphoria	Significant incongruence between one's experienced gender and the gender assigned at birth, lasting > 6 months and leading to persistent distress. Individuals may self-identify as another gender, pursue surgery or hormone treatment to rid self of primary/secondary sex characteristics, and/or live as another gender. Gender nonconformity itself is not a mental disorder. **Transgender**—desiring and often making lifestyle changes to live as a different gender. Medical interventions (eg, hormone therapy, sex reassignment surgery) may be utilized during the transition to enable the individual's appearance to match their gender identity. **Transvestism**—deriving pleasure from wearing clothes (eg, a vest) of the opposite sex (cross-dressing). **Transvestic disorder**—transvestism that causes significant distress/functional impairment. It is a paraphilia (psychosexual disorder), not part of gender dysphoria.

Sexual dysfunction	Includes sexual desire disorders (hypoactive sexual desire or sexual aversion), sexual arousal disorders (erectile dysfunction), orgasmic disorders (anorgasmia, premature ejaculation), sexual pain disorders (dyspareunia, vaginismus). Differential diagnosis includes (PENIS): ▪ Psychological (if nighttime erections still occur) ▪ Endocrine (eg, diabetes, low testosterone) ▪ Neurogenic (eg, postoperative, spinal cord injury) ▪ Insufficient blood flow (eg, atherosclerosis) ▪ Substances (eg, antihypertensives, antidepressants, ethanol)

(c) ketabton.com: The Digital Library

Sleep terror disorder

Periods of inconsolable terror with screaming in the middle of the night. Most common in children. Occurs during slow-wave/deep (stage N3) non-REM sleep with no memory of the arousal episode, as opposed to nightmares that occur during **REM** sleep (**rem**embering a scary dream). Triggers include emotional stress, fever, and lack of sleep. Usually self limited.

Enuresis

Nighttime urinary incontinence ≥ 2 times/week for ≥ 3 months in person > 5 years old. First-line treatment: behavioral modification (eg, scheduled voids, nighttime fluid restriction) and positive reinforcement. For refractory cases: bedwetting alarm, oral desmopressin (ADH analog; preferred over imipramine due to fewer side effects).

Narcolepsy

Excessive daytime sleepiness (despite awakening well-rested) with recurrent episodes of rapid-onset, overwhelming sleepiness ≥ 3 times/week for the last 3 months. Due to ↓ orexin (hypocretin) production in lateral hypothalamus and dysregulated sleep-wake cycles. Associated with:

- Hypna**go**gic (just before **go**ing to sleep) or hypno**pomp**ic (just before awakening; get **pomp**ed up in the morning) hallucinations.
- Nocturnal and narcoleptic sleep episodes that start with REM sleep (sleep paralysis).
- Cataplexy (loss of all muscle tone following strong emotional stimulus, such as laughter).

Treatment: good sleep hygiene (scheduled naps, regular sleep schedule), daytime stimulants (eg, amphetamines, modafinil) and/or nighttime sodium oxybate (GHB).

Substance use disorder

Maladaptive pattern of substance use involving ≥ 2 of the following in the past year:

- Tolerance
- Withdrawal
- Intense, distracting cravings
- Using more, or longer, than intended
- Persistent desire but inability to cut down
- Time-consuming substance acquisition, use, or recovery
- Impaired functioning at work, school, or home
- Social or interpersonal conflicts
- Reduced recreational activities
- > 1 episode of use involving danger (eg, unsafe sex, driving while impaired)
- Continued use despite awareness of harm

Stages of change in overcoming addiction

1. **Precontemplation**—denying problem
2. **Contemplation**—acknowledging problem, but unwilling to change
3. **Preparation/determination**—preparing for behavioral changes
4. **Action/willpower**—changing behaviors
5. **Maintenance**—maintaining changes
6. **Relapse**—(if applicable) returning to old behaviors and abandoning changes



(c) ketabton.com: The Digital Library

Psychiatric emergencies

	CAUSE	MANIFESTATION	TREATMENT
Serotonin syndrome	Any drug that ↑ 5-HT. Psychiatric drugs: MAOIs, SSRIs, SNRIs, TCAs, vilazodone, vortioxetine, buspirone Nonpsychiatric drugs: tramadol, ondansetron, triptans, linezolid, MDMA, dextromethorphan, meperidine, St. John's wort	3 A's: ↑ Activity (neuromuscular; eg, clonus, hyperreflexia, hypertonia, tremor, seizure), Autonomic instability (eg, hyperthermia, diaphoresis, diarrhea), Altered mental status	Cyproheptadine (5-HT_2 receptor antagonist)
Hypertensive crisis	Eating tyramine-rich foods (eg, aged cheeses, cured meats, wine, chocolate) while taking MAOIs	Hypertensive crisis (tyramine displaces other neurotransmitters [eg, NE] in the synaptic cleft → ↑ sympathetic stimulation)	Phentolamine
Neuroleptic malignant syndrome	Antipsychotics (typical > atypical) + genetic predisposition	Malignant FEVER: Myoglobinuria, Fever, Encephalopathy, Vitals unstable, ↑ Enzymes (eg, CK), muscle Rigidity ("lead pipe")	Dantrolene, dopamine agonist (eg, bromocriptine), discontinue causative agent
Delirium tremens	Alcohol withdrawal; occurs 2–4 days after last drink Classically seen in hospital setting when inpatient cannot drink	Altered mental status, hallucinations, autonomic hyperactivity, anxiety, seizures, tremors, psychomotor agitation, insomnia, nausea	Benzodiazepines (eg, chlordiazepoxide, lorazepam, diazepam)
Acute dystonia	Typical antipsychotics, anticonvulsants (eg, carbamazepine), metoclopramide	Sudden onset of muscle spasms, stiffness, and/or oculogyric crisis occurring hours to days after medication use; can lead to laryngospasm requiring intubation	Benztropine or diphenhydramine
Lithium toxicity	↑ lithium dosage, ↓ renal elimination (eg, acute kidney injury), medications affecting clearance (eg, ACE inhibitors, thiazide diuretics, NSAIDs). Narrow therapeutic window.	Nausea, vomiting, slurred speech, hyperreflexia, seizures, ataxia, nephrogenic diabetes insipidus	Discontinue lithium, hydrate aggressively with isotonic sodium chloride, consider hemodialysis
Tricyclic antidepressant toxicity	TCA overdose	Respiratory depression, hyperpyrexia, prolonged QT Tri-CyCliC's: Convulsions, Coma, Cardiotoxicity (arrhythmia due to Na^+ channel inhibition)	Supportive treatment, monitor ECG, $NaHCO_3$ (prevents arrhythmia), activated charcoal

(c) ketabton.com: The Digital Library

Psychoactive drug intoxication and withdrawal

DRUG	INTOXICATION	WITHDRAWAL
Depressants		
	Nonspecific: mood elevation, ↓ anxiety, sedation, behavioral disinhibition, respiratory depression.	Nonspecific: anxiety, tremor, seizures, insomnia.
Alcohol	Emotional lability, slurred speech, ataxia, coma, blackouts. Serum γ-glutamyltransferase (GGT)—sensitive indicator of alcohol use. **AST** value is 2× **AL**T value ("To**AST 2 AL**cohol"). Treatment: benzodiazepines.	
Barbiturates	Low safety margin, marked respiratory depression. Treatment: symptom management (eg, assist respiration, ↑ BP).	Delirium, life-threatening cardiovascular collapse.
Benzodiazepines	Greater safety margin. Ataxia, minor respiratory depression. Treatment: flumazenil (benzodiazepine receptor antagonist, but rarely used as it can precipitate seizures).	Sleep disturbance, depression.
Opioids	Euphoria, respiratory and CNS depression, ↓ gag reflex, pupillary constriction (pinpoint pupils), seizures. Most common cause of drug overdose death. Treatment: naloxone.	Sweating, dilated pupils, piloerection ("cold turkey"), rhinorrhea, lacrimation, yawning, nausea, stomach cramps, diarrhea ("flu-like" symptoms). Treatment: symptom management, methadone, buprenorphine.
Inhalants	Disinhibition, euphoria, slurred speech, disturbed gait, disorientation, drowsiness.	Irritability, dysphoria, sleep disturbance, headache.
Stimulants		
	Nonspecific: mood elevation, ↓ appetite, psychomotor agitation, insomnia, cardiac arrhythmias, tachycardia, anxiety.	Nonspecific: post-use "crash," including depression, lethargy, ↑ appetite, sleep disturbance, vivid nightmares.
Amphetamines	Euphoria, grandiosity, pupillary dilation, prolonged wakefulness, hyperalertness, hypertension, paranoia, fever, fractured teeth. Skin excoriations with methamphetamine use. Severe: cardiac arrest, seizures. Treatment: benzodiazepines for agitation and seizures.	
Caffeine	Palpitation, agitation, tremor, insomnia.	Headache, difficulty concentrating, flu-like symptoms.

(c) ketabton.com: The Digital Library

Psychoactive drug intoxication and withdrawal *(continued)*

DRUG	INTOXICATION	WITHDRAWAL
Cocaine	Impaired judgment, pupillary dilation, hallucinations (including tactile), paranoia, angina, sudden cardiac death. Chronic use may lead to perforated nasal septum due to vasoconstriction and resulting ischemic necrosis. Treatment: benzodiazepines; consider mixed α-/β-blocker (eg, labetalol) for hypertension and tachycardia. Pure β-blocker usage is controversial as a first-line therapy.	
Nicotine	Restlessness.	Irritability, anxiety, restlessness, ↓ concentration, ↑ appetite/weight. Treatment: nicotine patch, gum, or lozenges; bupropion/varenicline.
Hallucinogens		
Lysergic acid diethylamide	Perceptual distortion (visual, auditory), depersonalization, anxiety, paranoia, psychosis, flashbacks (usually nondisturbing).	
Marijuana (cannabinoid)	Euphoria, anxiety, paranoid delusions, perception of slowed time, impaired judgment, social withdrawal, ↑ appetite, dry mouth, conjunctival injection, hallucinations. Pharmaceutical form is dronabinol: used as antiemetic (chemotherapy) and appetite stimulant (in AIDS).	Irritability, anxiety, depression, insomnia, restlessness, ↓ appetite.
MDMA (ecstasy)	Hallucinogenic stimulant: euphoria, hallucinations, disinhibition, hyperactivity, ↑ thirst, bruxism, distorted sensory and time perception. Life-threatening effects include hypertension, tachycardia, hyperthermia, hyponatremia, serotonin syndrome.	Depression, fatigue, change in appetite, difficulty concentrating, anxiety.
Phencyclidine	Violence, impulsivity, psychomotor agitation, nystagmus, tachycardia, hypertension, analgesia, psychosis, delirium, seizures. Trauma is most common complication.	

Alcohol use disorder

Physiologic tolerance and dependence on alcohol with symptoms of withdrawal when intake is interrupted.

Complications: vitamin B_1 (thiamine) deficiency, alcoholic cirrhosis, hepatitis, pancreatitis, peripheral neuropathy, testicular atrophy.

Treatment: naltrexone (reduces cravings), acamprosate, disulfiram (to condition the patient to abstain from alcohol use). Support groups such as Alcoholics Anonymous are helpful in sustaining abstinence and supporting patient and family.

Wernicke-Korsakoff syndrome

Results from vitamin B_1 deficiency. Symptoms can be precipitated by administering dextrose before vitamin B_1. Triad of confusion, ophthalmoplegia, ataxia (Wernicke encephalopathy). May progress to irreversible memory loss, confabulation, personality change (Korsakoff syndrome). Treatment: IV vitamin B_1 (before dextrose).

(c) ketabton.com: The Digital Library

▸ PSYCHIATRY—PHARMACOLOGY

Psychotherapy

Behavioral therapy	Teaches patients how to identify and change maladaptive behaviors or reactions to stimuli. Examples include systematic desensitization for treatment of phobia.
Cognitive behavioral therapy	Teaches patients to recognize distortions in their thought processes, develop constructive coping skills, and ↓ maladaptive coping behaviors → greater emotional control and tolerance of distress. Examples include recognizing triggers for alcohol consumption.
Dialectical behavioral therapy	Designed for use in borderline personality disorder, but can be used in other psychiatric conditions as well (eg, depression).
Interpersonal therapy	Focused on improving interpersonal relationships and communication skills.
Supportive therapy	Utilizes empathy to help individuals during a time of hardship to maintain optimism or hope.

Preferred medications for selected psychiatric conditions

PSYCHIATRIC CONDITION	PREFERRED DRUGS
ADHD	Stimulants (methylphenidate, amphetamines)
Alcohol withdrawal	Benzodiazepines (eg, chlordiazepoxide, lorazepam, diazepam)
Bipolar disorder	Lithium, valproic acid, carbamazepine, lamotrigine, atypical antipsychotics
Bulimia nervosa	SSRIs
Depression	SSRIs
Generalized anxiety disorder	SSRIs, SNRIs
Obsessive-compulsive disorder	SSRIs, venlafaxine, clomipramine
Panic disorder	SSRIs, venlafaxine, benzodiazepines
PTSD	SSRIs, venlafaxine
Schizophrenia	Atypical antipsychotics
Social anxiety disorder	SSRIs, venlafaxine Performance only: β-blockers, benzodiazepines
Tourette syndrome	Antipsychotics (eg, fluphenazine, risperidone), tetrabenazine

Central nervous system stimulants

Methylphenidate, dextroamphetamine, methamphetamine, lisdexamfetamine.

MECHANISM	↑ catecholamines in the synaptic cleft, especially norepinephrine and dopamine.
CLINICAL USE	ADHD, narcolepsy, binge-eating disorder.
ADVERSE EFFECTS	Nervousness, agitation, anxiety, insomnia, anorexia, tachycardia, hypertension, weight loss, tics, bruxism.

(c) ketabton.com: The Digital Library

Typical antipsychotics	Haloperidol, pimozide, trifluoperazine, fluphenazine, thioridazine, chlorpromazine.
MECHANISM	Block dopamine D_2 receptor (↑ cAMP).
CLINICAL USE	Schizophrenia (1° positive symptoms), psychosis, bipolar disorder, delirium, Tourette syndrome, Huntington disease, OCD. Use with caution in dementia.
POTENCY	**High** potency: **Hal**operidol, **Tri**fluoperazine, **Fl**uphenazine (**Hal Tri**es to **Fl**y **High**)—more neurologic side effects (eg, extrapyramidal symptoms [EPS]). **Low** potency: **Ch**lorpromazine, **Thi**oridazine (**Ch**eating **Thi**eves are **low**)—more anticholinergic, antihistamine, α_1-blockade effects.
ADVERSE EFFECTS	Lipid soluble → stored in body fat → slow to be removed from body. Endocrine: dopamine receptor antagonism → hyperprolactinemia → galactorrhea, oligomenorrhea, gynecomastia. Metabolic: dyslipidemia, weight gain, hyperglycemia. Antimuscarinic: dry mouth, constipation. Antihistamine: sedation. α_1-blockade: orthostatic hypotension. Cardiac: QT prolongation. Ophthalmologic: **C**hlorpromazine—**C**orneal deposits; **T**hioridazine—re**T**inal deposits. Neuroleptic malignant syndrome. Extrapyramidal symptoms— **ADAPT**: ▪ Hours to days: **A**cute **D**ystonia (muscle spasm, stiffness, oculogyric crisis). Treatment: benztropine, diphenhydramine. ▪ Days to months: ▪▪ **A**kathisia (restlessness). Treatment: β-blockers, benztropine, benzodiazepines. ▪▪ **P**arkinsonism (bradykinesia). Treatment: benztropine, amantadine. ▪ Months to years: **T**ardive dyskinesia (chorea, especially orofacial). Treatment: atypical antipsychotics (eg, clozapine), valbenazine, deutetrabenazine.

Atypical antipsychotics	Aripiprazole, asen**apine**, cloz**apine**, olanz**apine**, quetiapine, ilo**peridone**, pali**peridone**, ris**peridone**, lurasidone, ziprasidone.	
MECHANISM	Not completely understood. Most are 5-HT_2 and D_2 antagonists; aripiprazole is a D_2 partial agonist. Varied effects on α and H_1 receptors.	
CLINICAL USE	Schizophrenia—both positive and negative symptoms. Also used for bipolar disorder, OCD, anxiety disorders, depression, mania, Tourette syndrome.	Use clozapine for treatment-resistant schizophrenia or schizoaffective disorder and for suicidality in schizophrenia.
ADVERSE EFFECTS	All—prolonged QT, fewer EPS and anticholinergic side effects than typical antipsychotics.	
	"-apines"—metabolic syndrome (weight gain, diabetes, dyslipidemia).	**O**lanzapine, cl**O**zapine → **O**besity
	Clozapine—agranulocytosis (monitor WBCs frequently) and seizures (dose related).	Must watch bone marrow **cloz**ely with **cloz**apine.
	Risperidone—hyperprolactinemia (amenorrhea, galactorrhea, gynecomastia).	

(c) ketabton.com: The Digital Library

Lithium

MECHANISM	Not established; possibly related to inhibition of phosphoinositol cascade.
CLINICAL USE	Mood stabilizer for bipolar disorder; treats acute manic episodes and prevents relapse.
ADVERSE EFFECTS	Tremor, thyroid abnormalities (eg, hypothyroidism), polyuria (causes nephrogenic diabetes insipidus), teratogenesis. Causes Ebstein anomaly in newborn if taken by pregnant mother. Narrow therapeutic window requires close monitoring of serum levels. Almost exclusively excreted by kidneys; most is reabsorbed at PCT via Na^+ channels. Thiazides, NSAIDs, and other drugs affecting clearance are implicated in lithium toxicity.

LiTHIUM:
Low **T**hyroid (hypothyroidism)
Heart (Ebstein anomaly)
Insipidus (nephrogenic diabetes insipidus)
Unwanted **M**ovements (tremor)

Buspirone

MECHANISM	Stimulates 5-HT_{1A} receptors.
CLINICAL USE	Generalized anxiety disorder. Does not cause sedation, addiction, or tolerance. Begins to take effect after 1–2 weeks. Does not interact with alcohol (vs barbiturates, benzodiazepines).

I get **anxious** if the **bus** doesn't arrive at **one**, so I take **bus**pir**one**.

Antidepressants



(c) ketabton.com: The Digital Library

Selective serotonin reuptake inhibitors

Fluoxetine, fluvoxamine, paroxetine, sertraline, escitalopram, citalopram.

MECHANISM	Inhibit 5-HT reuptake.
CLINICAL USE	Depression, generalized anxiety disorder, panic disorder, OCD, bulimia, binge-eating disorder, social anxiety disorder, PTSD, premature ejaculation, premenstrual dysphoric disorder.
ADVERSE EFFECTS	Fewer than TCAs. Serotonin syndrome, GI distress, SIADH, sexual dysfunction (anorgasmia, ↓ libido).

It normally takes 4–8 weeks for antidepressants to show appreciable effect.

Serotonin-norepinephrine reuptake inhibitors

Venlafaxine, desvenlafaxine, duloxetine, levomilnacipran, milnacipran.

MECHANISM	Inhibit 5-HT and NE reuptake.
CLINICAL USE	Depression, generalized anxiety disorder, diabetic neuropathy. Venlafaxine is also indicated for social anxiety disorder, panic disorder, PTSD, OCD. Duloxetine and milnacipran are also indicated for fibromyalgia.
ADVERSE EFFECTS	↑ BP, stimulant effects, sedation, nausea.

Tricyclic antidepressants

Amitriptyline, nortriptyline, imipramine, desipramine, clomipramine, doxepin, amoxapine.

MECHANISM	TCAs inhibit 5-HT and NE reuptake.
CLINICAL USE	MDD, peripheral neuropathy, chronic neuropathic pain, migraine prophylaxis, OCD (clomipramine), nocturnal enuresis (imipramine, although adverse effects may limit use).
ADVERSE EFFECTS	Sedation, α_1-blocking effects including postural hypotension, and atropine-like (anticholinergic) side effects (tachycardia, urinary retention, dry mouth). 3° TCAs (**a**mitriptyline) have more **a**nticholinergic effects than 2° TCAs (nortriptyline). Can prolong QT interval. **Tri**-**Cy**C**li**C's: **C**onvulsions, **C**oma, **C**ardiotoxicity (arrhythmia due to Na^+ channel inhibition); also respiratory depression, hyperpyrexia. Confusion and hallucinations are more common in the elderly due to anticholinergic side effects (2° amines [eg, nortriptyline] better tolerated). Treatment: $NaHCO_3$ to prevent arrhythmia.

Monoamine oxidase inhibitors

Tranylcypromine, **P**henelzine, **I**socarboxazid, **S**elegiline (selective **MAO**-B inhibitor).
(**MAO** **T**akes **P**ride **I**n **S**hanghai).

MECHANISM	Nonselective MAO inhibition → ↑ levels of amine neurotransmitters (norepinephrine, 5-HT, dopamine).
CLINICAL USE	Atypical depression, anxiety. Parkinson disease (selegiline).
ADVERSE EFFECTS	CNS stimulation; hypertensive crisis, most notably with ingestion of tyramine. Contraindicated with SSRIs, TCAs, St. John's wort, meperidine, dextromethorphan, linezolid (to avoid precipitating serotonin syndrome). Wait 2 weeks after stopping MAOIs before starting serotonergic drugs or stopping dietary restrictions.

(c) ketabton.com: The Digital Library

Atypical antidepressants	
Bupropion	Inhibits NE and DA reuptake. Also used for smoking cessation. Toxicity: stimulant effects (tachycardia, insomnia), headache, seizures in patients with bulimia and anorexia nervosa. Favorable sexual side effect profile.
Mirtazapine	α_2-antagonist (↑ release of NE and 5-HT), potent 5-HT_2 and 5-HT_3 receptor antagonist, and H_1 antagonist. Toxicity: sedation (which may be desirable in depressed patients with insomnia), ↑ appetite, weight gain (which may be desirable in underweight patients), dry mouth.
Trazodone	Primarily blocks 5-HT_2, α_1-adrenergic, and H_1 receptors; also weakly inhibits 5-HT reuptake. Used primarily for insomnia, as high doses are needed for antidepressant effects. Toxicity: sedation, nausea, priapism, postural hypotension. Think tra**ZZZ**o**bone** due to sedative and male-specific side effects.
Varenicline	Nicotinic ACh receptor partial agonist. Used for smoking cessation. Toxicity: sleep disturbance, depressed mood, suicidal ideation. Vare**nicline** helps **ni**cotine cravings de**cline**.
Vilazodone	Inhibits 5-HT reuptake; 5-HT_{1A} receptor partial agonist. Used for MDD. Toxicity: headache, diarrhea, nausea, anticholinergic effects. May cause serotonin syndrome if taken with other serotonergic agents.
Vortioxetine	Inhibits 5-HT reuptake; 5-HT_{1A} receptor agonist and 5-HT_3 receptor antagonist. Used for MDD. Toxicity: nausea, sexual dysfunction, sleep disturbances, anticholinergic effects. May cause serotonin syndrome if taken with other serotonergic agents.

Opioid detoxification and relapse prevention	Intravenous drug users at ↑ risk for hepatitis, HIV, abscesses, bacteremia, right-heart endocarditis.
Methadone	Long-acting oral opiate used for heroin detoxification or long-term maintenance therapy.
Buprenorphine	Sublingual form (partial agonist) used to prevent relapse.
Naloxone	Short-acting opioid antagonist given IM, IV, or as a nasal spray to treat acute opioid overdose, particularly to reverse respiratory and CNS depression.
Naltrexone	Long-acting oral opioid antagonist used after detoxification to prevent relapse. Use nal**trex**one for the long **trex** back to sobriety.

(c) ketabton.com: The Digital Library

Renal

"But I know all about love already. I know precious little still about kidneys."

—Aldous Huxley, *Antic Hay*

"This too shall pass. Just like a kidney stone."

—Hunter Madsen

"I drink too much. The last time I gave a urine sample it had an olive in it."

—Rodney Dangerfield

Being able to understand and apply renal physiology will be critical for the exam. Important topics include electrolyte disorders, acid-base derangements, glomerular disorders (including histopathology), acute and chronic kidney disease, urine casts, diuretics, ACE inhibitors, and AT-II receptor blockers. Renal anomalies associated with various congenital defects are also high-yield associations to think about when evaluating pediatric vignettes.

- Embryology 578
- Anatomy 580
- Physiology 581
- Pathology 594
- Pharmacology 607

FAS1_2019_14-Renal.indd 577 11/7/19 5:42 PM

(c) ketabton.com: The Digital Library

▸ RENAL—EMBRYOLOGY

Kidney embryology

Pronephros—week 4; then degenerates.

Mesonephros—functions as interim kidney for 1st trimester; later contributes to male genital system.

Metanephros—permanent; first appears in 5th week of gestation; nephrogenesis continues through weeks 32–36 of gestation.

- Ureteric bud (metanephric diverticulum)—derived from caudal end of mesonephric duct; gives rise to ureter, pelvises, calyces, collecting ducts; fully canalized by 10th week
- Metanephric mesenchyme (ie, metanephric blastema)—ureteric bud interacts with this tissue; interaction induces differentiation and formation of glomerulus through to distal convoluted tubule (DCT)
- Aberrant interaction between these 2 tissues may result in several congenital malformations of the kidney (eg, renal agenesis, multicystic dysplastic kidney)

Ureteropelvic junction—last to canalize → congenital obstruction. Most common cause of prenatal hydronephrosis. Detected by prenatal ultrasound.



Potter sequence (syndrome)



Oligohydramnios → compression of developing fetus → limb deformities, facial anomalies (eg, low-set ears and retrognathia A, flattened nose), compression of chest and lack of amniotic fluid aspiration into fetal lungs → pulmonary hypoplasia (cause of death).

Causes include ARPKD, obstructive uropathy (eg, posterior urethral valves), bilateral renal agenesis, chronic placental insufficiency.

Babies who can't "Pee" in utero develop Potter sequence.

POTTER sequence associated with:

- Pulmonary hypoplasia
- Oligohydramnios (trigger)
- Twisted face
- Twisted skin
- Extremity defects
- Renal failure (in utero)

(c) ketabton.com: The Digital Library

Horseshoe kidney



Inferior poles of both kidneys fuse abnormally A. As they ascend from pelvis during fetal development, horseshoe kidneys get trapped under inferior mesenteric artery and remain low in the abdomen. Kidneys function normally. Associated with hydronephrosis (eg, ureteropelvic junction obstruction), renal stones, infection, ↑ risk of renal cancer.

Higher incidence in chromosomal aneuploidy (eg, Turner syndrome, trisomies 13, 18, 21).



Congenital solitary functioning kidney

Condition of being born with only one functioning kidney. Majority asymptomatic with compensatory hypertrophy of contralateral kidney, but anomalies in contralateral kidney are common. Often diagnosed prenatally via ultrasound.

Unilateral renal agenesis

Ureteric bud fails to develop and induce differentiation of metanephric mesenchyme → complete absence of kidney and ureter.

Multicystic dysplastic kidney

Ureteric bud fails to induce differentiation of metanephric mesenchyme → nonfunctional kidney consisting of cysts and connective tissue. Predominantly nonhereditary and usually unilateral; bilateral leads to Potter sequence.

Duplex collecting system

Bifurcation of ureteric bud before it enters the metanephric blastema creates a Y-shaped bifid ureter. Duplex collecting system can alternatively occur through two ureteric buds reaching and interacting with metanephric blastema. Strongly associated with vesicoureteral reflux and/or ureteral obstruction, ↑ risk for UTIs.

Posterior urethral valves

Membrane remnant in the posterior urethra in males; its persistence can lead to urethral obstruction. Can be diagnosed prenatally by bilateral hydronephrosis and dilated or thick-walled bladder on ultrasound. Most common cause of bladder outlet obstruction in male infants. Associated with oligohydramnios in cases of severe obstruction.



(c) ketabton.com: The Digital Library

▸ RENAL—ANATOMY

Kidney anatomy and glomerular structure



Cross-section of kidney



*Components of glomerular filtration barrier.

Cross-section of glomerulus

Left kidney is taken during living donor transplantation because it has a longer renal vein.

Afferent = Arriving.

Efferent = Exiting.

Renal blood flow: renal artery → segmental artery → interlobar artery → arcuate artery → interlobular artery → afferent arteriole → glomerulus A → efferent arteriole → vasa recta/peritubular capillaries → venous outflow.

Left renal vein receives two additional veins: left suprarenal and left gonadal veins.

Despite high overall renal blood flow, renal medulla receives significantly less blood flow than renal cortex → very sensitive to hypoxia → vulnerable to ischemic damage.



FAS1_2019_14-Renal.indd 580 11/7/19 5:42 PM

(c) ketabton.com: The Digital Library

Course of ureters



Course of ureter A: arises from renal pelvis, travels under gonadal arteries → **over** common iliac artery → **under** uterine artery/vas deferens (retroperitoneal).

Gynecologic procedures (eg, ligation of uterine or ovarian vessels) may damage ureter → ureteral obstruction or leak.

Bladder contraction compresses the intravesical ureter, preventing urine reflux.

Blood supply to ureter:

- Proximal—renal arteries
- Middle—gonadal artery, aorta, common and internal iliac arteries
- Distal—internal iliac and superior vesical arteries

3 common points of ureteral obstruction: ureteropelvic junction, pelvic inlet, ureterovesical junction.

Water (ureters) flows **over** the iliacs and **under** the bridge (uterine artery or vas deferens).



▸ RENAL—PHYSIOLOGY

Fluid compartments



HIKIN': **HI**gh **K**$^+$ **IN**tracellularly.

60–40–20 rule (% of body weight for average person):

- 60% total body water
- 40% ICF, mainly composed of K^+, Mg^{2+}, organic phosphates (eg, ATP)
- 20% ECF, mainly composed of Na^+, Cl^-, HCO_3^-, albumin

Plasma volume can be measured by radiolabeling albumin.

Extracellular volume can be measured by inulin or mannitol.

Serum osmolality = 285–295 mOsm/kg H_2O.

Plasma volume = TBV × (1 – Hct).

Glomerular filtration barrier



Responsible for filtration of plasma according to size and charge selectivity.

Composed of:

- Fenestrated capillary endothelium
- Basement membrane with type IV collagen chains and heparan sulfate
- Visceral epithelial layer consisting of podocyte foot processes (FPs) A

Charge barrier—all 3 layers contain ⊖ charged glycoproteins that prevent entry of ⊖ charged molecules (eg, albumin).

Size barrier—fenestrated capillary endothelium (prevents entry of > 100 nm molecules/blood cells); podocyte foot processes interpose with glomerular basement membrane (GBM); slit diaphragm (prevents entry of molecules > 50–60 nm).

(c) ketabton.com: The Digital Library

Renal clearance

$C_x = (U_x V)/P_x$ = volume of plasma from which the substance is completely cleared in the urine per unit time.

If $C_x < GFR$: net tubular reabsorption and/or not freely filtered.

If $C_x > GFR$: net tubular secretion of X.

If $C_x = GFR$: no net secretion or reabsorption.

C_x = clearance of X (mL/min).
U_x = urine concentration of X (eg, mg/mL).
P_x = plasma concentration of X (eg, mg/mL).
V = urine flow rate (mL/min).

Glomerular filtration rate

Inulin clearance can be used to calculate GFR because it is freely filtered and is neither reabsorbed nor secreted.

$$C_{inulin} = GFR = U_{inulin} \times V/P_{inulin}$$
$$= K_f [(P_{GC} - P_{BS}) - (\pi_{GC} - \pi_{BS})]$$

(GC = glomerular capillary; BS = Bowman space; π_{BS} normally equals zero; K_f = filtration coefficient).

Normal GFR ≈ 100 mL/min.

Creatinine clearance is an approximate measure of GFR. Slightly overestimates GFR because creatinine is moderately secreted by renal tubules.



Effective renal plasma flow

Effective renal plasma flow (eRPF) can be estimated using *para*-aminohippuric acid (PAH) clearance. Between filtration and secretion, there is nearly 100% excretion of all PAH that enters the kidney.

$eRPF = U_{PAH} \times V/P_{PAH} = C_{PAH}$.

Renal blood flow (RBF) = RPF/(1 − Hct). Usually 20–25% of cardiac output.

eRPF underestimates true renal plasma flow (RPF) slightly.

(c) ketabton.com: The Digital Library

Filtration

Filtration fraction (FF) = GFR/RPF.
Normal FF = 20%.
Filtered load (mg/min) = GFR (mL/min) × plasma concentration (mg/mL).

GFR can be estimated with creatinine clearance.
RPF is best estimated with PAH clearance.
Prostaglandins **D**ilate **A**fferent arteriole (**PDA**).
Angiotensin II **C**onstricts **E**fferent arteriole (**ACE**).



Changes in glomerular dynamics

	GFR	RPF	FF (GFR/RPF)
Afferent arteriole constriction	↓	↓	—
Efferent arteriole constriction	↑	↓	↑
↑ plasma protein concentration	↓	—	↓
↓ plasma protein concentration	↑	—	↑
Constriction of ureter	↓	—	↓
Dehydration	↓	↓↓	↑

(c) ketabton.com: The Digital Library

Calculation of reabsorption and secretion rate

Filtered load = GFR × P_x.
Excretion rate = V × U_x.
Reabsorption rate = filtered – excreted.
Secretion rate = excreted – filtered.
Fe_{Na} = fractional excretion of sodium.

$$Fe_{Na} = \frac{Na^+ \text{ excreted}}{Na^+ \text{ filtered}} = \frac{V \times U_{Na}}{GFR \times P_{Na}} = \frac{P_{Cr} \times U_{Na}}{U_{Cr} \times P_{Na}} \text{ where GFR} = \frac{U_{Cr} \times V}{P_{Cr}}$$

Glucose clearance

Glucose at a normal plasma level (range 60–120 mg/dL) is completely reabsorbed in proximal convoluted tubule (PCT) by Na^+/glucose cotransport.

In adults, at plasma glucose of ~ 200 mg/dL, glucosuria begins (threshold). At rate of ~ 375 mg/min, all transporters are fully saturated (T_m).

Normal pregnancy is associated with ↑ GFR. With ↑ filtration of all substances, including glucose, the glucose threshold occurs at lower plasma glucose concentrations → glucosuria at normal plasma glucose levels.

Sodium-glucose cotransporter 2 (SGLT2) inhibitors (eg, -flozin drugs) result in glucosuria at plasma concentrations < 200 mg/dL.

Glucosuria is an important clinical clue to diabetes mellitus.

Splay phenomenon—T_m for glucose is reached gradually rather than sharply due to the heterogeneity of nephrons (ie, different T_m points); represented by the portion of the titration curve between threshold and T_m.



(c) ketabton.com: The Digital Library

Nephron transport physiology



Early PCT—contains brush border. Reabsorbs all glucose and amino acids and most HCO_3^-, Na^+, Cl^-, PO_4^{3-}, K^+, H_2O, and uric acid. Isotonic absorption. Generates and secretes NH_3, which enables the kidney to secrete more H^+.
PTH—inhibits Na^+/PO_4^{3-} cotransport → ↑ PO_4^{3-} excretion.
AT II—stimulates Na^+/H^+ exchange → ↑ Na^+, H_2O, and HCO_3^- reabsorption (permitting contraction alkalosis).
65–80% Na^+ and H_2O reabsorbed.

Thin descending loop of Henle—passively reabsorbs H_2O via medullary hypertonicity (impermeable to Na^+). Concentrating segment. Makes urine hypertonic.



Thick ascending loop of Henle—reabsorbs Na^+, K^+, and Cl^-. Indirectly induces paracellular reabsorption of Mg^{2+} and Ca^{2+} through ⊕ lumen potential generated by K^+ backleak. Impermeable to H_2O. Makes urine less concentrated as it ascends.
10–20% Na^+ reabsorbed.



Early DCT—reabsorbs Na^+, Cl^-. Impermeable to H_2O. Makes urine fully dilute (hypotonic).
PTH—↑ Ca^{2+}/Na^+ exchange → ↑ Ca^{2+} reabsorption.
5–10% Na^+ reabsorbed.



Collecting tubule—reabsorbs Na^+ in exchange for secreting K^+ and H^+ (regulated by aldosterone).
Aldosterone—acts on mineralocorticoid receptor → mRNA → protein synthesis. In principal cells: ↑ apical K^+ conductance, ↑ Na^+/K^+ pump, ↑ epithelial Na^+ channel (ENaC) activity → lumen negativity → K^+ secretion. In α-intercalated cells: lumen negativity → ↑ H^+ ATPase activity → ↑ H^+ secretion → ↑ HCO_3^-/Cl^- exchanger activity.
ADH—acts at V_2 receptor → insertion of aquaporin H_2O channels on apical side.
3–5% Na^+ reabsorbed.

(c) ketabton.com: The Digital Library

Renal tubular defects

Order: Fanconi's BaGeLS



	DEFECTS	EFFECTS	CAUSES	NOTES
Fanconi syndrome	Generalized reabsorption defect in PCT → ↑ excretion of amino acids, glucose, HCO_3^-, and PO_4^{3-}, and all substances reabsorbed by the PCT	May lead to metabolic acidosis (proximal RTA), hypophosphatemia, osteopenia	Hereditary defects (eg, Wilson disease, tyrosinemia, glycogen storage disease), ischemia, multiple myeloma, nephrotoxins/drugs (eg, ifosfamide, cisplatin), lead poisoning	
Bartter syndrome	Reabsorption defect in thick ascending loop of Henle (affects $Na^+/K^+/2Cl^-$ cotransporter)	Metabolic alkalosis, hypokalemia, hypercalciuria	Autosomal recessive	Presents similarly to chronic loop diuretic use
Gitelman syndrome	Reabsorption defect of NaCl in DCT	Metabolic alkalosis, hypomagnesemia, hypokalemia, hypocalciuria	Autosomal recessive	Presents similarly to lifelong thiazide diuretic use Less severe than Bartter syndrome
Liddle syndrome	Gain of function mutation → ↓ Na^+ channel degradation → ↑ Na^+ reabsorption in collecting tubules	Metabolic alkalosis, hypokalemia, hypertension, ↓ aldosterone	Autosomal dominant	Presents similarly to hyperaldosteronism, but aldosterone is nearly undetectable Treatment: amiloride
Syndrome of Apparent Mineralocorticoid Excess	Cortisol activates mineralocorticoid receptors. 11β-HSD converts cortisol to cortisone (inactive on these receptors) Hereditary 11β-HSD deficiency → ↑ cortisol → ↑ mineralocorticoid receptor activity	Metabolic alkalosis, hypokalemia, hypertension ↓ serum aldosterone level; cortisol tries to be the SAME as aldosterone	Autosomal recessive Can acquire disorder from glycyrrhetinic acid (present in licorice), which blocks activity of 11β-hydroxysteroid dehydrogenase	Treatment: K^+-sparing diuretics (↓ mineralocorticoid effects) or corticosteroids (exogenous corticosteroid ↓ endogenous cortisol production → ↓ mineralocorticoid receptor activation)

(c) ketabton.com: The Digital Library

Relative concentrations along proximal convoluted tubules

[TF/P] > 1
when solute is reabsorbed less quickly than water or when solute is secreted

[TF/P] = 1
when solute and water are reabsorbed at the same rate

[TF/P] < 1
when solute is reabsorbed more quickly than water



Tubular inulin ↑ in concentration (but not amount) along the PCT as a result of water reabsorption. Cl^- reabsorption occurs at a slower rate than Na^+ in early PCT and then matches the rate of Na^+ reabsorption more distally. Thus, its relative concentration ↑ before it plateaus.

(c) ketabton.com: The Digital Library

Renin-angiotensin-aldosterone system



Renin	Secreted by JG cells in response to ↓ renal perfusion pressure (detected by renal baroreceptors in afferent arteriole), ↑ renal sympathetic discharge (β_1 effect), and ↓ NaCl delivery to macula densa cells.
AT II	Helps maintain blood volume and blood pressure. Affects baroreceptor function; limits reflex bradycardia, which would normally accompany its pressor effects.
ANP, BNP	Released from atria (ANP) and ventricles (BNP) in response to ↑ volume; inhibits renin-angiotensin-aldosterone system; relaxes vascular smooth muscle via cGMP → ↑ GFR, ↓ renin. Dilates afferent arteriole, promotes natriuresis.
ADH	Primarily regulates serum osmolality; also responds to low blood volume states. Stimulates reabsorption of water in collecting ducts. Also stimulates reabsorption of urea in collecting ducts to maximizes corticopapillary osmotic gradient.
Aldosterone	Primarily regulates ECF volume and Na^+ content; ↑ release in ↓ blood volume states. Responds to hyperkalemia by ↑ K^+ excretion.

(c) ketabton.com: The Digital Library

Juxtaglomerular apparatus	Consists of mesangial cells, JG cells (modified smooth muscle of afferent arteriole), and the macula densa (NaCl sensor, located at distal end of loop of Henle). JG cells secrete renin in response to ↓ renal blood pressure and ↑ sympathetic tone (β_1). Macula densa cells sense ↓ NaCl delivery to DCT → ↑ renin release → efferent arteriole vasoconstriction → ↑ GFR.	JGA maintains GFR via renin-angiotensin-aldosterone system. In addition to vasodilatory properties, β-blockers can decrease BP by inhibiting β_1-receptors of the JGA → ↓ renin release.

Kidney endocrine functions

Erythropoietin	Released by interstitial cells in peritubular capillary bed in response to hypoxia.	Stimulates RBC proliferation in bone marrow. Administered for anemia secondary to chronic kidney disease. ↑ risk of HTN.
Calciferol (vitamin D)	PCT cells convert 25-OH vitamin D_3 to 1,25-$(OH)_2$ vitamin D_3 (calcitriol, active form).	
Prostaglandins	Paracrine secretion vasodilates the afferent arterioles to ↑ RBF.	NSAIDs block renal-protective prostaglandin synthesis → constriction of afferent arteriole and ↓ GFR; this may result in acute kidney injury in low renal blood flow states.
Dopamine	Secreted by PCT cells, promotes natriuresis. At low doses; dilates interlobular arteries, afferent arterioles, efferent arterioles → ↑ RBF, little or no change in GFR. At higher doses; acts as vasoconstrictor.	

(c) ketabton.com: The Digital Library

Hormones acting on kidney

Atrial natriuretic peptide
Secreted in response to ↑ atrial pressure. Causes ↑ GFR and ↑ Na^+ filtration with no compensatory Na^+ reabsorption in distal nephron. Net effect: Na^+ loss and volume loss.



Angiotensin II
Synthesized in response to ↓ BP. Causes efferent arteriole constriction → ↑ GFR and ↑ FF but with compensatory Na^+ reabsorption in proximal and distal nephron. Net effect: preservation of renal function (↑ FF) in low-volume state with simultaneous Na^+ reabsorption (both proximal and distal) to maintain circulating volume.

Parathyroid hormone
Secreted in response to ↓ plasma [Ca^{2+}], ↑ plasma [PO_4^{3-}], or ↓ plasma 1,25-$(OH)_2$ D_3. Causes ↑ [Ca^{2+}] reabsorption (DCT), ↓ [PO_4^{3-}] reabsorption (PCT), and ↑ 1,25-$(OH)_2$ D_3 production (↑ Ca^{2+} and PO_4^{3-} absorption from gut via vitamin D).

Aldosterone
Secreted in response to ↓ blood volume (via AT II) and ↑ plasma [K^+]; causes ↑ Na^+ reabsorption, ↑ K^+ secretion, ↑ H^+ secretion.

ADH (vasopressin)
Secreted in response to ↑ plasma osmolarity and ↓ blood volume. Binds to receptors on principal cells, causing ↑ number of aquaporins and ↑ H_2O reabsorption.
↑ reabsorption of urea in collecting ducts to maximize corticopapillary osmotic gradient.

Potassium shifts

SHIFTS K^+ INTO CELL (CAUSING HYPOKALEMIA)	SHIFTS K^+ OUT OF CELL (CAUSING HYPERKALEMIA)
	Digitalis (blocks Na^+/K^+ ATPase)
Hypo-osmolarity	Hyper**O**smolarity
	Lysis of cells (eg, crush injury, rhabdomyolysis, tumor lysis syndrome)
Alkalosis	**A**cidosis
β-adrenergic agonist (↑ Na^+/K^+ ATPase)	β-blocker
Insulin (↑ Na^+/K^+ ATPase)	High blood **S**ugar (insulin deficiency)
Insulin shifts K^+ **in**to cells	**S**uccinylcholine (↑ risk in burns/muscle trauma)
	Hyperkalemia? **DO LAβSS**

(c) ketabton.com: The Digital Library

Electrolyte disturbances

ELECTROLYTE	LOW SERUM CONCENTRATION	HIGH SERUM CONCENTRATION
Sodium	Nausea, malaise, stupor, coma, seizures	Irritability, stupor, coma
Potassium	U waves and flattened T waves on ECG, arrhythmias, muscle cramps, spasm, weakness	Wide QRS and peaked T waves on ECG, arrhythmias, muscle weakness
Calcium	Tetany, seizures, QT prolongation, twitching (eg, Chvostek sign), spasm (eg, Trousseau sign)	**Stones** (renal), **bones** (pain), **groans** (abdominal pain), **thrones** (↑ urinary frequency), **psychiatric overtones** (anxiety, altered mental status)
Magnesium	Tetany, torsades de pointes, hypokalemia, hypocalcemia (when $[Mg^{2+}] < 1.0$ mEq/L)	↓ DTRs, lethargy, bradycardia, hypotension, cardiac arrest, hypocalcemia
Phosphate	Bone loss, osteomalacia (adults), rickets (children)	Renal stones, metastatic calcifications, hypocalcemia

Features of renal disorders

CONDITION	BLOOD PRESSURE	PLASMA RENIN	ALDOSTERONE	SERUM Mg^{2+}	URINE Ca^{2+}
SIADH	—/↑	↓	↓		
Primary hyperaldosteronism	↑	↓	↑		
Renin-secreting tumor	↑	↑	↑		
Bartter syndrome		↑	↑		↑
Gitelman syndrome		↑	↑	↓	↓
Liddle syndrome, syndrome of apparent mineralocorticoid excess	↑	↓	↓		

↑ ↓ = important differentiating feature.

(c) ketabton.com: The Digital Library

Acid-base physiology

	pH	P_{CO_2}	$[HCO_3^-]$	COMPENSATORY RESPONSE
Metabolic acidosis	↓	↓	↓	Hyperventilation (immediate)
Metabolic alkalosis	↑	↑	↑	Hypoventilation (immediate)
Respiratory acidosis	↓	↑	↑	↑ renal $[HCO_3^-]$ reabsorption (delayed)
Respiratory alkalosis	↑	↓	↓	↓ renal $[HCO_3^-]$ reabsorption (delayed)

Key: ↓ ↑ = compensatory response.

Henderson-Hasselbalch equation: $pH = 6.1 + \log \frac{[HCO_3^-]}{0.03\ P_{CO_2}}$

Predicted respiratory compensation for a simple metabolic acidosis can be calculated using the Winters formula. If measured P_{CO_2} > predicted P_{CO_2} → concomitant respiratory acidosis; if measured P_{CO_2} < predicted P_{CO_2} → concomitant respiratory alkalosis:

$$P_{CO_2} = 1.5\ [HCO_3^-] + 8 \pm 2$$

Acidosis and alkalosis



(c) ketabton.com: The Digital Library

Renal tubular acidosis

Disorder of the renal tubules that causes normal anion gap (hyperchloremic) metabolic acidosis.

RTA TYPE	DEFECT	URINE PH	SERUM K^+	CAUSES	ASSOCIATIONS
Distal renal tubular acidosis (type 1)	Inability of α-intercalated cells to secrete H^+ → no new HCO_3^- is generated → metabolic acidosis	> 5.5	↓	Amphotericin B toxicity, analgesic nephropathy, congenital anomalies (obstruction) of urinary tract, autoimmune diseases (eg, SLE)	↑ risk for calcium phosphate kidney stones (due to ↑ urine pH and ↑ bone turnover related to buffering)
Proximal renal tubular acidosis (type 2)	Defect in PCT HCO_3^- reabsorption → ↑ excretion of HCO_3^- in urine → metabolic acidosis Urine can be acidified by α-intercalated cells in collecting duct, but not enough to overcome ↑ HCO_3^- excretion	> 5.5 when resorptive threshold for serum HCO_3^- exceeded; < 5.5 when HCO_3^- depleted below resorptive threshold	↓	Fanconi syndrome, multiple myeloma, carbonic anhydrase inhibitors	↑ risk for hypophosphatemic rickets (in Fanconi syndrome)
Hyperkalemic tubular acidosis (type 4)	Hypoaldosteronism or aldosterone resistance; hyperkalemia → ↓ NH_3 synthesis in PCT → ↓ NH_4^+ excretion	< 5.5 (or variable)	↑	↓ aldosterone production (eg, diabetic hyporeninism, ACE inhibitors, ARBs, NSAIDs, heparin, cyclosporine, adrenal insufficiency) or aldosterone resistance (eg, K^+-sparing diuretics, nephropathy due to obstruction, TMP-SMX)	

(c) ketabton.com: The Digital Library

▸ RENAL—PATHOLOGY

Casts in urine	Presence of casts indicates that hematuria/pyuria is of glomerular or renal tubular origin. Bladder cancer, kidney stones → hematuria, no casts. Acute cystitis → pyuria, no casts.
RBC casts A	Glomerulonephritis, hypertensive emergency.
WBC casts B	Tubulointerstitial inflammation, acute pyelonephritis, transplant rejection.
Granular casts C	Acute tubular necrosis (ATN). Can be "muddy brown" in appearance.
Fatty casts ("oval fat bodies")	Nephrotic syndrome. Associated with "Maltese cross" sign D.
Waxy casts	End-stage renal disease/chronic kidney disease.
Hyaline casts E	Nonspecific, can be a normal finding. Form via solidification of Tamm–Horsfall mucoprotein (secreted by renal tubular cells).



Nomenclature of glomerular disorders

TYPE	CHARACTERISTICS	EXAMPLE
Focal	< 50% of glomeruli are involved	Focal segmental glomerulosclerosis
Diffuse	> 50% of glomeruli are involved	Diffuse proliferative glomerulonephritis
Proliferative	Hypercellular glomeruli	Membranoproliferative glomerulonephritis
Membranous	Thickening of glomerular basement membrane (GBM)	Membranous nephropathy
Primary glomerular disease	1° disease of the kidney specifically impacting the glomeruli	Minimal change disease
Secondary glomerular disease	Systemic disease or disease of another organ system that also impacts the glomeruli	SLE, diabetic nephropathy

(c) ketabton.com: The Digital Library

Glomerular diseases



TYPE	ETIOLOGY	CLINICAL PRESENTATION	EXAMPLES
Nephritic syndrome	Glomerular inflammation → GBM damage → loss of RBCs into urine → hematuria	Hematuria, RBC casts in urine ↓ GFR → oliguria, azotemia, ↑ renin release, HTN Proteinuria often in the subnephrotic range (< 3.5 g/day) but in severe cases may be in nephrotic range	▪ Acute poststreptococcal glomerulonephritis ▪ Rapidly progressive glomerulonephritis ▪ IgA nephropathy (Berger disease) ▪ Alport syndrome ▪ Membranoproliferative glomerulonephritis
Nephrotic syndrome	Podocyte damage → impaired charge barrier → proteinuria	Massive proteinuria (> 3.5 g/day) with hypoalbuminemia, edema Frothy urine with fatty casts Associated with hypercoagulable state due to antithrombin III loss in urine and ↑ risk of infection (loss of IgGs in urine and soft tissue compromise by edema)	May be 1° (eg, direct podocyte damage) or 2° (podocyte damage from systemic process): ▪ Focal segmental glomerulosclerosis (1° or 2°) ▪ Minimal change disease (1° or 2°) ▪ Membranous nephropathy (1° or 2°) ▪ Amyloidosis (2°) ▪ Diabetic glomerulonephropathy (2°)
Nephritic-nephrotic syndrome	Severe GBM damage → loss of RBCs into urine + impaired charge barrier → hematuria + proteinuria	Nephrotic-range proteinuria (> 3.5 g/day) and concomitant features of nephrotic syndrome	Can occur with any form of nephritic syndrome, but is most common with: ▪ Diffuse proliferative glomerulonephritis ▪ Membranoproliferative glomerulonephritis

(c) ketabton.com: The Digital Library

Nephritic syndrome

NephrItic syndrome = Inflammatory process.

Acute poststreptococcal glomerulonephritis

Most frequently seen in children. ~ 2–4 weeks after group A streptococcal infection of pharynx or skin. Resolves spontaneously in most children; may progress to renal insufficiency in adults. Type III hypersensitivity reaction. Presents with peripheral and periorbital edema, tea or cola-colored urine, HTN. ⊕ strep titers/serologies, ↓ complement levels (C3) due to consumption.

- LM—glomeruli enlarged and hypercellular A
- IF—("starry sky") granular appearance ("lumpy-bumpy") B due to IgG, IgM, and C3 deposition along GBM and mesangium
- EM—subepithelial IC humps

Rapidly progressive (crescentic) glomerulonephritis

Poor prognosis, rapidly deteriorating renal function (days to weeks).

- LM—crescent moon shape C. Crescents consist of fibrin and plasma proteins (eg, C3b) with glomerular parietal cells, monocytes, macrophages

Several disease processes may result in this pattern which may be delineated via IF pattern.

- Linear IF due to antibodies to GBM and alveolar basement membrane: Goodpasture syndrome—hematuria/hemoptysis; type II hypersensitivity reaction. Treatment: plasmapheresis
- Negative IF/Pauci-immune (no Ig/C3 deposition): granulomatosis with polyangiitis (Wegener)—PR3-ANCA/c-ANCA, eosinophilic granulomatosis with polyangiitis (Churg-Strauss) or Microscopic polyangiitis—MPO-ANCA/p-ANCA
- Granular IF—PSGN or DPGN

Diffuse proliferative glomerulonephritis

Often due to SLE (think "wire lupus"). DPGN and MPGN often present as nephrotic syndrome and nephritic syndrome concurrently.

- LM—"wire looping" of capillaries D
- IF—granular; EM—subendothelial, sometimes subepithelial or intramembranous IgG-based ICs often with C3 deposition

IgA nephropathy (Berger disease)

Episodic hematuria that usually occurs concurrently with respiratory or GI tract infections (IgA is secreted by mucosal linings). Renal pathology of IgA vasculitis (HSP).

- LM—mesangial proliferation
- IF—IgA-based IC deposits in mesangium; EM—mesangial IC deposition

Alport syndrome

Mutation in type IV collagen → thinning and splitting of glomerular basement membrane.

Most commonly X-linked dominant. Eye problems (eg, retinopathy, anterior lenticonus), glomerulonephritis, sensorineural deafness; "can't see, can't pee, can't hear a bee."

- EM—"basket-weave" appearance due to irregular thickening of GBM

Membrano-proliferative glomerulonephritis

MPGN is a nephritic syndrome that often co-presents with nephrotic syndrome.

Type I may be 2° to hepatitis B or C infection. May also be idiopathic.

- Subendothelial IC deposits with granular IF

Type II is associated with C3 nephritic factor (IgG autoantibody that stabilizes C3 convertase → persistent complement activation → ↓ C3 levels).

- Intramembranous deposits, also called dense deposit disease

Both types: mesangial ingrowth → GBM splitting → "tram-track" on H&E and PAS E stains.



FAS1_2019_14-Renal.indd 596 11/7/19 5:42 PM

(c) ketabton.com: The Digital Library

Nephrotic syndrome	NephrOtic syndrome—massive prOteinuria (> 3.5 g/day)
Minimal change disease	Also known as lipoid nephrosis. Most common cause of nephrotic syndrome in children. Often 1° (Idiopathic) and may be triggered by recent Infection, Immunization, Immune stimulus (4 I's of MCD). Rarely, may be 2° to lymphoma (eg, cytokine-mediated damage). 1° disease has excellent response to corticosteroids. ▪ LM—Normal glomeruli (lipid may be seen in PCT cells) ▪ IF—⊖ ▪ EM—effacement of podocyte foot processes A
Focal segmental glomerulosclerosis	Most common cause of nephrotic syndrome in African-Americans and Hispanics. Can be 1° (idiopathic) or 2° to other conditions (eg, HIV infection, sickle cell disease, heroin abuse, massive obesity, interferon treatment, or congenital malformations). 1° disease has inconsistent response to steroids. May progress to CKD. ▪ LM—segmental sclerosis and hyalinosis B ▪ IF—often ⊖ but may be ⊕ for nonspecific focal deposits of IgM, C3, C1 ▪ EM—effacement of foot processes similar to minimal change disease
Membranous nephropathy	Also known as membranous glomerulonephritis. Can be 1° (eg, antibodies to phospholipase A_2 receptor) or 2° to drugs (eg, NSAIDs, penicillamine, gold), infections (eg, HBV, HCV, syphilis), SLE, or solid tumors. 1° disease has poor response to steroids. May progress to CKD. ▪ LM—diffuse capillary and GBM thickening C ▪ IF—granular due to immune complex (IC) deposition ▪ EM—"Spike and dome" appearance of subepithelial deposits
Amyloidosis	Kidney is the most commonly involved organ (systemic amyloidosis). Associated with chronic conditions that predispose to amyloid deposition (eg, AL amyloid, AA amyloid). ▪ LM—Congo red stain shows apple-green birefringence under polarized light due to amyloid deposition in the mesangium
Diabetic glomerulo-nephropathy	Most common cause of ESRD in the United States. Hyperglycemia → nonenzymatic glycation of tissue proteins → mesangial expansion; GBM thickening and ↑ permeability. Hyperfiltration (glomerular HTN and ↑ GFR) → glomerular hypertrophy and glomerular scarring (glomerulosclerosis) → further progression of nephropathy. ▪ LM—Mesangial expansion, GBM thickening, eosinophilic nodular glomerulosclerosis (Kimmelstiel-Wilson lesions D)



(c) ketabton.com: The Digital Library

Kidney stones Can lead to severe complications such as hydronephrosis, pyelonephritis, and acute kidney injury. Obstructed stone presents with unilateral flank tenderness, colicky pain radiating to groin, hematuria. Treat and prevent by encouraging fluid intake.

CONTENT	PRECIPITATES WITH	X-RAY FINDINGS	CT FINDINGS	URINE CRYSTAL	NOTES
Calcium	Calcium oxalate: hypocitraturia	Radiopaque	Radiopaque	Shaped like envelope A or dumbbell	Calcium stones most common (80%); calcium oxalate more common than calcium phosphate stones. Can result from ethylene glycol (antifreeze) ingestion, vitamin C abuse, hypocitraturia (associated with ↓ urine pH), malabsorption (eg, Crohn disease). Treatment: thiazides, citrate, low-sodium diet.
	Calcium phosphate: ↑ pH	Radiopaque	Radiopaque	Wedge-shaped prism	Treatment: low-sodium diet, thiazides.
Ammonium magnesium phosphate (struvite)	↑ pH	Radiopaque	Radiopaque	Coffin lid B	Account for 15% of stones. Caused by infection with urease ⊕ bugs (eg, *Proteus mirabilis*, *Staphylococcus saprophyticus*, *Klebsiella*) that hydrolyze urea to ammonia → urine alkalinization. Commonly form staghorn calculi C. Treatment: eradication of underlying infection, surgical removal of stone.
Uric acid	↓ pH	RadiolUcent	Minimally visible	Rhomboid D or rosettes	About 5% of all stones. Risk factors: ↓ urine volume, arid climates, acidic pH. Strong association with hyperuricemia (eg, gout). Often seen in diseases with ↑ cell turnover (eg, leukemia). Treatment: alkalinization of urine, allopurinol.
Cystine	↓ pH	Faintly radiopaque	Moderately radiopaque	Hexagonal E	Hereditary (autosomal recessive) condition in which Cystine-reabsorbing PCT transporter loses function, causing cystinuria. Transporter defect also results in poor reabsorption of Ornithine, Lysine, Arginine (COLA). Cystine is poorly soluble, thus stones form in urine. Usually begins in childhood. Can form staghorn calculi. Sodium cyanide nitroprusside test ⊕. "SIXtine" stones have SIX sides. Treatment: low sodium diet, alkalinization of urine, chelating agents (eg, penicillamine) if refractory.



FAS1_2019_14-Renal.indd 598 11/7/19 5:42 PM

(c) ketabton.com: The Digital Library

Hydronephrosis



Distention/dilation of renal pelvis and calyces A. Usually caused by urinary tract obstruction (eg, renal stones, severe BPH, congenital obstructions, cervical cancer, injury to ureter); other causes include retroperitoneal fibrosis, vesicoureteral reflux. Dilation occurs proximal to site of pathology. Serum creatinine becomes elevated if obstruction is bilateral or if patient has an obstructed solitary kidney. Leads to compression and possible atrophy of renal cortex and medulla.

Urinary incontinence

Mixed incontinence has features of both stress and urgency incontinence.

	Stress incontinence	Urgency incontinence	Overflow incontinence
MECHANISM	Outlet incompetence (urethral hypermobility or intrinsic sphincter deficiency) → leak with ↑ intra-abdominal pressure (eg, sneezing, lifting) ⊕ bladder stress test (directly observed leakage from urethra upon coughing or Valsalva maneuver)	Detrusor overactivity → leak with urge to void immediately	Incomplete emptying (detrusor underactivity or outlet obstruction) → leak with overfilling, ↑ postvoid residual on catheterization or ultrasound
ASSOCIATIONS	Obesity, vaginal delivery, prostate surgery	UTI	Polyuria (eg, diabetes), bladder outlet obstruction (eg, BPH), neurogenic bladder (eg, MS)
TREATMENT	Pelvic floor muscle strengthening (Kegel) exercises, weight loss, pessaries	Kegel exercises, bladder training (timed voiding, distraction or relaxation techniques), antimuscarinics (eg, oxybutynin for overactive bladder), mirabegron	Catheterization, relieve obstruction (eg, α-blockers for BPH)





(c) ketabton.com: The Digital Library

Acute cystitis

Inflammation of urinary bladder. Presents as suprapubic pain, dysuria, urinary frequency, urgency. Systemic signs (eg, high fever, chills) are usually absent.

Risk factors include female sex (short urethra), sexual intercourse, indwelling catheter, diabetes mellitus, impaired bladder emptying.

Causes:

- *E coli* (most common)
- *Staphylococcus saprophyticus*—seen in sexually active young women (*E coli* is still more common in this group)
- *Klebsiella*
- *Proteus mirabilis*—urine has ammonia scent

Labs: ⊕ leukocyte esterase. ⊕ nitrites (indicate gram ⊖ organisms). Sterile pyuria (pyuria with ⊖ urine cultures) could suggest urethritis by *Neisseria gonorrhoeae* or *Chlamydia trachomatis*.

Treatment: antibiotics (eg, TMP-SMX, nitrofurantoin).

Pyelonephritis

Acute pyelonephritis

Neutrophils infiltrate renal interstitium A. Affects cortex with relative sparing of glomeruli/vessels. Presents with fevers, flank pain (costovertebral angle tenderness), nausea/vomiting, chills.

Causes include ascending UTI (*E coli* is most common), hematogenous spread to kidney. Presents with WBCs in urine +/– WBC casts. CT would show striated parenchymal enhancement B.

Risk factors include indwelling urinary catheter, urinary tract obstruction, vesicoureteral reflux, diabetes mellitus, pregnancy.

Complications include chronic pyelonephritis, renal papillary necrosis, perinephric abscess, urosepsis.

Treatment: antibiotics.

Chronic pyelonephritis

The result of recurrent or inadequately treated episodes of acute pyelonephritis. Typically requires predisposition to infection such as vesicoureteral reflux or chronically obstructing kidney stones.

Coarse, asymmetric corticomedullary scarring, blunted calyx. Tubules can contain eosinophilic casts resembling thyroid tissue C (thyroidization of kidney).

Xanthogranulomatous pyelonephritis—rare; grossly orange nodules that can mimic tumor nodules; characterized by widespread kidney damage due to granulomatous tissue containing foamy macrophages. Associated with *Proteus* infection.



(c) ketabton.com: The Digital Library

Acute kidney injury

	Prerenal azotemia	Intrinsic renal failure	Postrenal azotemia
ETIOLOGY	Hypovolemia ↓ cardiac output ↓ effective circulating volume (eg, HF, liver failure)	Tubules and interstitium: ▪ Acute tubular necrosis (ischemia, sepsis, infection, nephrotoxins) ▪ Acute interstitial nephritis Glomerulus: ▪ Acute glomerulonephritis Vascular: ▪ Vasculitis ▪ Malignant hypertension ▪ TTP-HUS	Stones BPH Neoplasm Congenital anomalies
PATHOPHYSIOLOGY	↓ RBF → ↓ GFR → ↑ reabsorption of Na^+/H_2O and urea	In ATN, patchy necrosis → debris obstructing tubules and fluid backflow → ↓ GFR In ATN, epithelial/granular casts	Outflow obstruction (bilateral)
URINE OSMOLALITY (mOsm/kg)	>500	<350	<350
URINE Na^+ (mEq/L)	<20	>40	Varies
FE_{Na}	<1%	>2%	Varies
SERUM BUN/Cr	>20	<15	Varies

Acute interstitial nephritis

Also called tubulointerstitial nephritis. Acute interstitial renal inflammation. Pyuria (classically eosinophils) and azotemia occurring after administration of drugs that act as haptens, inducing hypersensitivity (eg, diuretics, NSAIDs, penicillin derivatives, proton pump inhibitors, rifampin, quinolones, sulfonamides). Less commonly may be 2° to other processes such as systemic infections (eg, *Mycoplasma*) or autoimmune diseases (eg, Sjögren syndrome, SLE, sarcoidosis).

Associated with fever, rash, hematuria, pyuria, and costovertebral angle tenderness, but can be asymptomatic.

Remember these **5 P'S**:

- **P**ee (diuretics)
- **P**ain-free (NSAIDs)
- **P**enicillins and cephalosporins
- **P**roton pump inhibitors
- Rifam**P**in
- **S**ulfa drugs

(c) ketabton.com: The Digital Library

Acute tubular necrosis





Most common cause of acute kidney injury in hospitalized patients. Spontaneously resolves in many cases. Can be fatal, especially during initial oliguric phase. ↑ FE_{Na}.

Key finding: granular casts (often muddy brown in appearance) **A**.

3 stages:

1. Inciting event
2. Maintenance phase—oliguric; lasts 1–3 weeks; risk of hyperkalemia, metabolic acidosis, uremia
3. Recovery phase—polyuric; BUN and serum creatinine fall; risk of hypokalemia and renal wasting of other electrolytes and minerals

Can be caused by ischemic or nephrotoxic injury:

- Ischemic—2° to ↓ renal blood flow (eg, hypotension, shock, sepsis, hemorrhage, HF). Results in death of tubular cells that may slough into tubular lumen **B** (PCT and thick ascending limb are highly susceptible to injury).
- Nephrotoxic—2° to injury resulting from toxic substances (eg, aminoglycosides, radiocontrast agents, lead, cisplatin, ethylene glycol), crush injury (myoglobinuria), hemoglobinuria. Proximal tubules are particularly susceptible to injury.

Diffuse cortical necrosis

Acute generalized cortical infarction of both kidneys. Likely due to a combination of vasospasm and DIC.

Associated with obstetric catastrophes (eg, abruptio placentae), septic shock.

Renal papillary necrosis



Sloughing of necrotic renal papillae **A** → gross hematuria and proteinuria. May be triggered by recent infection or immune stimulus.

Associated with: **S**ickle cell disease or trait, **A**cute pyelonephritis, **A**nalgesics (NSAIDs), **D**iabetes mellitus (**SAAD papa** with **pap**illary necrosis).

(c) ketabton.com: The Digital Library

Consequences of renal failure

Decline in renal filtration can lead to excess retained nitrogenous waste products and electrolyte disturbances.

Consequences (**MAD HUNGER**):

- **M**etabolic **A**cidosis
- **D**yslipidemia (especially ↑ triglycerides)
- **H**igh potassium
- **U**remia—clinical syndrome marked by:
  - Nausea and anorexia
  - Pericarditis
  - Asterixis
  - Encephalopathy
  - Platelet dysfunction
- **N**a$^+$/H_2O retention (HF, pulmonary edema, hypertension)
- **G**rowth retardation and developmental delay
- **E**rythropoietin deficiency (anemia)
- **R**enal osteodystrophy

2 forms of renal failure: acute (eg, ATN) and chronic (eg, hypertension, diabetes mellitus, congenital anomalies).

Incremental reductions in GFR define the stages of chronic kidney disease.

Renal osteodystrophy

Hypocalcemia, hyperphosphatemia, and failure of vitamin D hydroxylation associated with chronic kidney disease → 2° hyperparathyroidism → 3° hyperparathyroidism (if 2° poorly managed). High serum phosphate can bind with Ca^{2+} → tissue deposits → ↓ serum Ca^{2+}. ↓ 1,25-$(OH)_2D_3$ → ↓ intestinal Ca^{2+} absorption. Causes subperiosteal thinning of bones.

(c) ketabton.com: The Digital Library

Renal cyst disorders

Autosomal dominant polycystic kidney disease

Numerous cysts in cortex and medulla A causing bilateral enlarged kidneys ultimately destroy kidney parenchyma. Presents with flank pain, hematuria, hypertension, urinary infection, progressive renal failure in ~ 50% of individuals.

Mutation in *PKD1* (85% of cases, chromosome 16) or *PKD2* (15% of cases, chromosome 4).

Complications include chronic kidney disease and hypertension (caused by ↑ renin production). Associated with berry aneurysms, mitral valve prolapse, benign hepatic cysts, diverticulosis.

Treatment: If hypertension or proteinuria develops, treat with ACE inhibitors or ARBs.

Autosomal recessive polycystic kidney disease

Cystic dilation of collecting ducts B. Often presents in infancy. Associated with congenital hepatic fibrosis. Significant oliguric renal failure in utero can lead to Potter sequence. Concerns beyond neonatal period include systemic hypertension, progressive renal insufficiency, and portal hypertension from congenital hepatic fibrosis.

Autosomal dominant tubulointerstitial kidney disease

Also called medullary cystic kidney disease. Causes tubulointerstitial fibrosis and progressive renal insufficiency with inability to concentrate urine. Medullary cysts usually not visualized; smaller kidneys on ultrasound. Poor prognosis.

Simple vs complex renal cysts

Simple cysts are filled with ultrafiltrate (anechoic on ultrasound C). Very common and account for majority of all renal masses. Found incidentally and typically asymptomatic.

Complex cysts, including those that are septated, enhanced, or have solid components on imaging require follow-up or removal due to risk of renal cell carcinoma.

Renovascular disease

Renal impairment due to ischemia from renal artery stenosis or microvascular disease.

↓ renal perfusion (one or both kidneys) → ↑ renin → ↑ angiotensin → HTN.

Main causes of renal artery stenosis:

- Atherosclerotic plaques—proximal 1/3 of renal artery, usually in older males, smokers.
- Fibromuscular dysplasia—distal 2/3 of renal artery or segmental branches, usually young or middle-aged females.

Clinically, patients can have refractory HTN with negative family history of HTN, asymmetric renal size, epigastric/flank bruits.

Most common cause of 2° HTN in adults. Other large vessels are often involved.

(c) ketabton.com: The Digital Library

Renal cell carcinoma

Polygonal clear cells A filled with accumulated lipids and carbohydrate. Often golden-yellow B due to ↑ lipid content.

Originates from PCT → invades renal vein (may develop varicocele if left sided) → IVC → hematogenous spread → metastasis to lung and bone.

Manifests with hematuria, palpable masses, 2° polycythemia, flank pain, fever, weight loss.

Treatment: surgery/ablation for localized disease. Immunotherapy (eg, aldesleukin) or targeted therapy for metastatic disease, rarely curative. Resistant to chemotherapy and radiation therapy.

Most common 1° renal malignancy C.

Most common in men 50–70 years old, ↑ incidence with smoking and obesity.

Associated with paraneoplastic syndromes, eg, **P**THrP, **E**ctopic EPO, **A**CTH, **R**enin ("**PEAR**"-aneoplastic).

Clear cell (most common subtype) associated with gene deletion on chromosome 3 (sporadic, or inherited as von Hippel-Lindau syndrome).

RCC = **3** letters = chromosome **3**.



Renal oncocytoma



Benign epithelial cell tumor arising from collecting ducts (arrows in A point to well-circumscribed mass with central scar).

Large eosinophilic cells with abundant mitochondria without perinuclear clearing B (vs chromophobe renal cell carcinoma).

Presents with painless hematuria, flank pain, abdominal mass.

Often resected to exclude malignancy (eg, renal cell carcinoma).



(c) ketabton.com: The Digital Library

Nephroblastoma



Also called Wilms tumor. Most common renal malignancy of early childhood (ages 2–4). Contains embryonic glomerular structures. Presents with large, palpable, unilateral flank mass A and/or hematuria and possible HTN.

"Loss of function" mutations of tumor suppressor genes *WT1* or *WT2* on chromosome 11.

May be a part of several syndromes:

- **WAGR** complex—**W**ilms tumor, **A**niridia (absence of iris), **G**enitourinary malformations, mental **R**etardation/intellectual disability (*WT1* deletion)
- **D**enys-**D**rash syndrome—Wilms tumor, **D**iffuse mesangial sclerosis (early-onset nephrotic syndrome), **D**ysgenesis of gonads (male pseudohermaphroditism), *WT1* mutation
- Beckwith-**W**iedemann syndrome—**W**ilms tumor, macroglossia, organomegaly, hemihyperplasia (*WT2* mutation), omphalocele

Urothelial carcinoma of the bladder



Also called transitional cell carcinoma. Most common tumor of urinary tract system (can occur in renal calyces, renal pelvis, ureters, and bladder) A B. Can be suggested by painless hematuria (no casts).

Associated with problems in your **P**ee **SAC**: **P**henacetin, **S**moking, **A**niline dyes, and **C**yclophosphamide.



Squamous cell carcinoma of the bladder

Chronic irritation of urinary bladder → squamous metaplasia → dysplasia and squamous cell carcinoma.

Risk factors include *Schistosoma haematobium* infection (Middle East), chronic cystitis, smoking, chronic nephrolithiasis. Presents with painless hematuria (no casts).

(c) ketabton.com: The Digital Library

▸ RENAL—PHARMACOLOGY

Diuretics site of action



Mannitol

MECHANISM	Osmotic diuretic. ↑ tubular fluid osmolarity → ↑ urine flow, ↓ intracranial/intraocular pressure.
CLINICAL USE	Drug overdose, elevated intracranial/intraocular pressure.
ADVERSE EFFECTS	Pulmonary edema, dehydration, hypo- or hypernatremia. Contraindicated in anuria, HF.

(c) ketabton.com: The Digital Library

Acetazolamide

MECHANISM	Carbonic anhydrase inhibitor. Causes self-limited $NaHCO_3$ diuresis and ↓ total body HCO_3^- stores. Alkalinizes urine.
CLINICAL USE	Glaucoma, metabolic alkalosis, altitude sickness, idiopathic intracranial hypertension.
ADVERSE EFFECTS	Proximal renal tubular acidosis, paresthesias, NH_3 toxicity, sulfa allergy, hypokalemia. Promotes calcium phosphate stone formation (insoluble at high pH).



"Acid"azolamide causes Acidosis.

Loop diuretics

Furosemide, bumetanide, torsemide

MECHANISM	Sulfonamide loop diuretics. Inhibit cotransport system ($Na^+/K^+/2Cl^-$) of thick ascending limb of loop of Henle. Abolish hypertonicity of medulla, preventing concentration of urine. Associated with ↑ PGE (vasodilatory effect on afferent arteriole); inhibited by NSAIDs. ↑ Ca^{2+} excretion. Loops Lose Ca^{2+}.
CLINICAL USE	Edematous states (HF, cirrhosis, nephrotic syndrome, pulmonary edema), hypertension, hypercalcemia.
ADVERSE EFFECTS	Ototoxicity, Hypokalemia, Hypomagnesemia, Dehydration, Allergy (sulfa), metabolic Alkalosis, Nephritis (interstitial), Gout.

OHH DAANG!

Ethacrynic acid

MECHANISM	Nonsulfonamide inhibitor of cotransport system ($Na^+/K^+/2Cl^-$) of thick ascending limb of loop of Henle.
CLINICAL USE	Diuresis in patients allergic to sulfa drugs.
ADVERSE EFFECTS	Similar to furosemide, but more ototoxic.



Loop earrings hurt your ears.

(c) ketabton.com: The Digital Library

Thiazide diuretics

Hydrochlorothiazide, chlorthalidone, metolazone.

MECHANISM

Inhibit NaCl reabsorption in early DCT → ↓ diluting capacity of nephron. ↓ Ca^{2+} excretion.

CLINICAL USE

Hypertension, HF, idiopathic hypercalciuria, nephrogenic diabetes insipidus, osteoporosis.

ADVERSE EFFECTS

Hypokalemic metabolic alkalosis, hyponatremia, hyperGlycemia, hyperLipidemia, hyperUricemia, hyperCalcemia. Sulfa allergy.

HyperGLUC.



Potassium-sparing diuretics

Spironolactone, Eplerenone, Amiloride, Triamterene.

Keep your SEAT

MECHANISM

Spironolact**one** and epleren**one** are competitive aldoster**one** receptor antagonists in cortical collecting tubule. Triamterene and amiloride block Na^+ channels at the same part of the tubule.

CLINICAL USE

Hyperaldosteronism, K^+ depletion, HF, hepatic ascites (spironolactone), nephrogenic DI (amiloride), antiandrogen.

ADVERSE EFFECTS

Hyperkalemia (can lead to arrhythmias), endocrine effects with spironolactone (eg, gynecomastia, antiandrogen effects).



Diuretics: electrolyte changes

Urine NaCl

↑ with all diuretics (strength varies based on potency of diuretic effect). Serum NaCl may decrease as a result.

Urine K^+

↑ especially with loop and thiazide diuretics. Serum K^+ may decrease as a result.

Blood pH

↓ (**acidemia**): carbonic anhydrase inhibitors: ↓ HCO_3^- reabsorption. K^+ sparing: aldosterone blockade prevents K^+ secretion and H^+ secretion. Additionally, hyperkalemia leads to K^+ entering all cells (via H^+/K^+ exchanger) in exchange for H^+ exiting cells.

↑ (**alkalemia**): loop diuretics and thiazides cause alkalemia through several mechanisms:

- Volume contraction → ↑ AT II → ↑ Na^+/H^+ exchange in PCT → ↑ HCO_3^- reabsorption ("contraction alkalosis")
- K^+ loss leads to K^+ exiting all cells (via H^+/K^+ exchanger) in exchange for H^+ entering cells
- In low K^+ state, H^+ (rather than K^+) is exchanged for Na^+ in cortical collecting tubule → alkalosis and "paradoxical aciduria"

Urine Ca^{2+}

↑ with loop diuretics: ↓ paracellular Ca^{2+} reabsorption → hypocalcemia.
↓ with thiazides: enhanced Ca^{2+} reabsorption.

(c) ketabton.com: The Digital Library

Angiotensin-converting enzyme inhibitors	Captopril, enalapril, lisinopril, ramipril.	
MECHANISM	Inhibit ACE → ↓ AT II → ↓ GFR by preventing constriction of efferent arterioles. ↑ renin due to loss of negative feedback. Inhibition of ACE also prevents inactivation of bradykinin, a potent vasodilator.	
CLINICAL USE	Hypertension, HF (↓ mortality), proteinuria, diabetic nephropathy. Prevent unfavorable heart remodeling as a result of chronic hypertension.	In chronic kidney disease (eg, diabetic nephropathy), ↓ intraglomerular pressure, slowing GBM thickening.
ADVERSE EFFECTS	Cough, Angioedema (both due to ↑ bradykinin; contraindicated in C1 esterase inhibitor deficiency), Teratogen (fetal renal malformations), ↑ Creatinine (↓ GFR), Hyperkalemia, and Hypotension. Used with caution in bilateral renal artery stenosis because ACE inhibitors will further ↓ GFR → renal failure.	Captopril's CATCHH.

Angiotensin II receptor blockers	Losartan, candesartan, valsartan.
MECHANISM	Selectively block binding of angiotensin II to AT_1 receptor. Effects similar to ACE inhibitors, but ARBs do not increase bradykinin.
CLINICAL USE	Hypertension, HF, proteinuria, or chronic kidney disease (eg, diabetic nephropathy) with intolerance to ACE inhibitors (eg, cough, angioedema).
ADVERSE EFFECTS	Hyperkalemia, ↓ GFR, hypotension; teratogen.

Aliskiren	
MECHANISM	Direct renin inhibitor, blocks conversion of angiotensinogen to angiotensin I. Aliskiren Kills Renin.
CLINICAL USE	Hypertension.
ADVERSE EFFECTS	Hyperkalemia, ↓ GFR, hypotension, angioedema. Relatively contraindicated in patients already taking ACE inhibitors or ARBs and contraindicated in pregnancy.

(c) ketabton.com: The Digital Library

Reproductive

"Artificial insemination is when the farmer does it to the cow instead of the bull."

—Student essay

Make no mistake about why these babies are here - they are here to replace us.

—Jerry Seinfeld

"Whoever called it necking was a poor judge of anatomy."

—Groucho Marx

"See, the problem is that God gives men a brain and a penis, and only enough blood to run one at a time."

—Robin Williams

Embryology 612

Anatomy 624

Physiology 629

Pathology 638

Pharmacology 655

The reproductive system can be intimidating at first but is manageable once you organize the concepts into the pregnancy, endocrinologic, embryologic, and oncologic aspects of reproduction. Study the endocrine and reproductive chapters together, because mastery of the hypothalamic-pituitary-gonadal axis is key to answering questions on ovulation, menstruation, disorders of sexual development, contraception, and many pathologies.

Embryology is a nuanced subject that covers multiple organ systems. Approaching it from a clinical perspective will allow for better understanding. For instance, make the connection between the presentation of DiGeorge syndrome and the 3rd/4th pharyngeal pouch, and between the Müllerian/Wolffian systems and disorders of sexual development.

As for oncology, don't worry about remembering screening or treatment guidelines. It is more important to know how these cancers present (eg, signs and symptoms) and their associated labs, histopathology, and risk factors. In addition, some of the testicular and ovarian cancers have distinct patterns of hCG, AFP, LH, or FSH derangements that serve as helpful clues in exam questions.

FAS1_2019_15-Repro.indd 611 11/7/19 5:52 PM

(c) ketabton.com: The Digital Library

▸ REPRODUCTIVE—EMBRYOLOGY

Important genes of embryogenesis

GENE	LOCATION	FUNCTION	NOTES
Sonic hedgehog (*SHH*) gene	Zone of polarizing activity at base of limb buds	Anterior-posterior axis patterning, CNS development	Mutations → holoprosencephaly
***Wnt*-7 gene**	Apical ectodermal ridge at distal end of each limb	Dorsal-ventral axis patterning, limb development	
Fibroblast growth factor (*FGF*) gene	Apical ectodermal ridge	Limb lengthening (via mitosis of mesoderm)	"Look at that Fetus, Growing Fingers"
Homeobox (*Hox*) genes	Multiple	Segmental organization in cranial-caudal direction, transcription factor coding	Mutations → appendages in wrong locations. Isotretinoin → ↑ *Hox* gene expression

Early fetal development

Early embryonic development



Within week 1	hCG secretion begins around the time of implantation of blastocyst.	Blastocyst "sticks" at day 6.
Within week 2	Bilaminar disc (epiblast, hypoblast).	2 weeks = 2 layers.
Within week 3	Gastrulation forms trilaminar embryonic disc. Cells from epiblast invaginate → primitive streak → endoderm, mesoderm, ectoderm. Notochord arises from midline mesoderm; overlying ectoderm becomes neural plate.	3 weeks = 3 layers.
Weeks 3–8 (embryonic period)	Neural tube formed by neuroectoderm and closes by week 4. Organogenesis.	Extremely susceptible to teratogens.
Week 4	Heart begins to beat. Upper and lower limb buds begin to form.	4 weeks = 4 limbs and 4 heart chambers.
Week 6	Fetal cardiac activity visible by transvaginal ultrasound.	
Week 8	Fetal movements start.	Gait at week 8.
Week 10	Genitalia have male/female characteristics.	Tenitalia.

(c) ketabton.com: The Digital Library

Embryologic derivatives

Ectoderm		External/outer layer
Surface ectoderm	Epidermis; adenohypophysis (from Rathke pouch); lens of eye; epithelial linings of oral cavity, sensory organs of ear, and olfactory epithelium; anal canal below the pectinate line; parotid, sweat, mammary glands.	Craniopharyngioma—benign Rathke pouch tumor with cholesterol crystals, calcifications.
Neural tube	Brain (neurohypophysis, CNS neurons, oligodendrocytes, astrocytes, ependymal cells, pineal gland), retina, spinal cord.	Neuroectoderm—think CNS.
Neural crest	Melanocytes, Odontoblasts, Tracheal cartilage, Enterochromaffin cells, Leptomeninges (arachnoid, pia), PNS ganglia (cranial, dorsal root, autonomic), Adrenal medulla, Schwann cells, Spiral membrane (aorticopulmonary septum), Endocardial cushions (also derived partially from mesoderm), Skull bones.	MOTEL PASSES Neural crest—think PNS and non-neural structures nearby.
Mesoderm	Muscle, bone, connective tissue, serous linings of body cavities (eg, peritoneum, pericardium, pleura), spleen (develops within foregut mesentery), cardiovascular structures, lymphatics, blood, wall of gut tube, upper vagina, kidneys, adrenal cortex, dermis, testes, ovaries, microglia. Notochord induces ectoderm to form neuroectoderm (neural plate); its only postnatal derivative is the nucleus pulposus of the intervertebral disc.	Middle/"meat" layer. Mesodermal defects = VACTERL: Vertebral defects Anal atresia Cardiac defects Tracheo-Esophageal fistula Renal defects Limb defects (bone and muscle)
Endoderm	Gut tube epithelium (including anal canal above the pectinate line), most of urethra and lower vagina (derived from urogenital sinus), luminal epithelial derivatives (eg, lungs, liver, gallbladder, pancreas, eustachian tube, thymus, parathyroid, thyroid follicular and parafollicular [C] cells).	"Enternal" layer.

Types of errors in morphogenesis

Agenesis	Absent organ due to absent primordial tissue.
Aplasia	Absent organ despite presence of primordial tissue.
Hypoplasia	Incomplete organ development; primordial tissue present.
Disruption	2° breakdown of previously normal tissue or structure (eg, amniotic band syndrome).
Deformation	Extrinsic disruption (eg, multiple gestations → crowding → foot deformities); occurs after embryonic period.
Malformation	Intrinsic disruption; occurs during embryonic period (weeks 3–8).
Sequence	Abnormalities result from a single 1° embryologic event (eg, oligohydramnios → Potter sequence).

(c) ketabton.com: The Digital Library

Teratogens Most susceptible in 3rd–8th weeks (embryonic period—organogenesis) of pregnancy. Before week 3, "all-or-none" effects. After week 8, growth and function affected.

TERATOGEN	EFFECTS ON FETUS	NOTES
Medications		
ACE inhibitors	Renal failure, oligohydramnios, hypocalvaria.	
Alkylating agents	Absence of digits, multiple anomalies.	
Aminoglycosides	Ototoxicity.	A mean guy hit the baby in the ear.
Antiepileptic drugs	Neural tube defects, cardiac defects, cleft palate, skeletal abnormalities (eg, phalanx/nail hypoplasia, facial dysmorphism).	High-dose folate supplementation recommended. Most commonly valproate, carbamazepine, phenytoin, phenobarbital.
Diethylstilbestrol (DES)	Vaginal clear cell adenocarcinoma, congenital Müllerian anomalies.	
Fluoroquinolones	Cartilage damage.	
Folate antagonists	Neural tube defects.	Antiepileptics, trimethoprim, methotrexate.
Isotretinoin	Multiple severe birth defects.	Contraception mandatory. IsoTERATinoin.
Lithium	Ebstein anomaly.	
Methimazole	Aplasia cutis congenita (congenital absence of skin, particularly on scalp).	
Tetracyclines	Discolored teeth, inhibited bone growth.	"Teethracyclines."
Thalidomide	Limb defects (phocomelia, micromelia—"flipper" limbs).	Limb defects with "tha-limb-domide."
Warfarin	Bone and cartilage deformities (stippled epiphyses, nasal and limb hypoplasia), optic nerve atrophy, fetal cerebral hemorrhage.	Do not wage warfare on the baby; keep it heppy with heparin (does not cross placenta).
Substance abuse		
Alcohol	Fetal alcohol syndrome.	
Cocaine	Low birth weight, preterm birth, IUGR, placental abruption.	Cocaine → vasoconstriction.
Smoking (nicotine, CO)	Low birth weight (leading cause in developed countries), preterm labor, placental problems, IUGR, SIDS, ADHD.	Nicotine → vasoconstriction. CO → impaired O_2 delivery.
Other		
Iodine (lack or excess)	Congenital goiter or hypothyroidism (cretinism).	
Maternal diabetes	Caudal regression syndrome, cardiac defects (eg, VSD), neural tube defects, macrosomia, neonatal hypoglycemia (due to islet cell hyperplasia), polycythemia.	
Methylmercury	Neurotoxicity.	Highest in swordfish, shark, tilefish, king mackerel.
Vitamin A excess	Extremely high risk for spontaneous abortions and birth defects (cleft palate, cardiac).	
X-rays	Microcephaly, intellectual disability.	Minimized by lead shielding.

(c) ketabton.com: The Digital Library

Fetal alcohol syndrome

One of the leading preventable causes of intellectual disability in the US. Newborns of mothers who consumed alcohol during any stage of pregnancy have ↑ incidence of congenital abnormalities, including pre- and postnatal developmental retardation, microcephaly, facial abnormalities A (eg, smooth philtrum, thin vermillion border, small palpebral fissures), limb dislocation, heart defects. Heart-lung fistulas and holoprosencephaly in most severe form. One mechanism is due to impaired migration of neuronal and glial cells.

Neonatal abstinence syndrome

Complex disorder involving CNS, ANS, and GI systems. Secondary to maternal substance use/abuse (most commonly opioids).

Universal screening for substance abuse is recommended in all pregnant patients.

Newborns may present with uncoordinated sucking reflexes, irritability, high-pitched crying, tremors, tachypnea, sneezing, diarrhea, and possibly seizures.

Treatment (for opiate abuse): methadone, morphine, buprenorphine.

(c) ketabton.com: The Digital Library

Twinning

Dizygotic ("fraternal") twins arise from 2 eggs that are separately fertilized by 2 different sperm (always 2 zygotes) and will have 2 separate amniotic sacs and 2 separate placentas (chorions).

Monozygotic ("identical") twins arise from 1 fertilized egg (1 egg + 1 sperm) that splits in early pregnancy. The timing of cleavage determines chorionicity (number of chorions) and amnionicity (number of amnions) (**SCAB**):

- Cleavage 0–4 days: **S**eparate chorion and amnion
- Cleavage 4–8 days: shared **C**horion
- Cleavage 8–12 days: shared **A**mnion
- Cleavage 13+ days: shared **B**ody (conjoined)



(c) ketabton.com: The Digital Library

Placenta	1° site of nutrient and gas exchange between mother and fetus.	
Fetal component		
Cytotrophoblast	Inner layer of chorionic villi.	Cytotrophoblast makes Cells.
Syncytiotrophoblast	Outer layer of chorionic villi; synthesizes and secretes hormones, eg, hCG (structurally similar to LH; stimulates corpus luteum to secrete progesterone during first trimester).	Syncytiotrophoblast synthesizes hormones. Lacks MHC-I expression → ↓ chance of attack by maternal immune system.
Maternal component		
Decidua basalis	Derived from endometrium. Maternal blood in lacunae.	



(c) ketabton.com: The Digital Library

Umbilical cord

Two umbilical arteries return deoxygenated blood from fetal internal iliac arteries to placenta A.

One umbilical vein supplies oxygenated blood from placenta to fetus; drains into IVC via liver or via ductus venosus.

Single umbilical artery (2-vessel cord) is associated with congenital and chromosomal anomalies.

Umbilical arteries and vein are derived from allantois.





Urachus

Allantois forms from hindgut and extends into urogenital sinus. Allantois becomes the urachus, a duct between fetal bladder and umbilicus. Failure of urachus to involute can lead to anomalies that may increase risk of infection and/or malignancy (eg, adenocarcinoma) if not treated.

Obliterated urachus is represented by the median umbilical ligament after birth, which is covered by median umbilical fold of the peritoneum.

Patent urachus	Total failure of urachus to obliterate → **u**rine discharge from **u**mbilicus.
Urachal cyst	Partial failure of urachus to obliterate; fluid-filled cavity lined with uroepithelium, between umbilicus and bladder. Cyst can become infected and present as painful mass below umbilicus.
Vesicourachal diverticulum	Slight failure of urachus to obliterate → outpouching of bladder.



Vitelline duct

7th week—obliteration of vitelline duct (omphalomesenteric duct), which connects yolk sac to midgut lumen.

Vitelline fistula	Vitelline duct fails to close → meconium discharge from umbilicus.
Meckel diverticulum	Partial closure of vitelline duct, with patent portion attached to ileum (true diverticulum, white arrow in B). May be asymptomatic. May have heterotopic gastric and/or pancreatic tissue → melena, hematochezia, abdominal pain.





(c) ketabton.com: The Digital Library

Aortic arch derivatives	Develop into arterial system.	
1st	Part of maxillary artery (branch of external carotid)	1st arch is maximal
2nd	Stapedial artery and hyoid artery	Second = Stapedial
3rd	Common Carotid artery and proximal part of internal Carotid artery	C is 3rd letter of alphabet
4th	On left, aortic arch; on right, proximal part of right subclavian artery	4th arch (4 limbs) = systemic
6th	Proximal part of pulmonary arteries and (on left only) ductus arteriosus	6th arch = pulmonary and the pulmonary-to-systemic shunt (ductus arteriosus)



6 months postnatal

Pharyngeal apparatus

Composed of pharyngeal clefts, arches, pouches.
Pharyngeal clefts—derived from ectoderm. Also called pharyngeal grooves.
Pharyngeal arches—derived from mesoderm (muscles, arteries) and neural crest (bones, cartilage).
Pharyngeal pouches—derived from endoderm.

CAP covers outside to inside:
Clefts = ectoderm
Arches = mesoderm + neural crest
Pouches = endoderm



Pharyngeal cleft derivatives

1st cleft develops into external auditory meatus.
2nd through 4th clefts form temporary cervical sinuses, which are obliterated by proliferation of 2nd arch mesenchyme.
Persistent cervical sinus → pharyngeal cleft cyst within lateral neck, anterior to sternocleidomastoid muscle (does not move with swallowing, vs thyroglossal duct cyst).

(c) ketabton.com: The Digital Library

Pharyngeal arch derivatives

ARCH	CARTILAGE	MUSCLES	NERVES[a]	NOTES
1st pharyngeal arch	**M**axillary process → **M**axilla, zygo**M**atic bone **M**andibular process → **M**eckel cartilage → **M**andible, **M**alleus and incus, spheno**M**andibular ligament	**M**uscles of **M**astication (temporalis, **M**asseter, lateral and **M**edial pterygoids), **M**ylohyoid, anterior belly of digastric, tensor tympani, anterior 2/3 of tongue, tensor veli palatini	CN V_3 **chew**	Pierre Robin sequence—micrognathia, glossoptosis, cleft palate, airway obstruction Treacher Collins syndrome—autosomal dominant neural crest dysfunction → craniofacial abnormalities (eg, zygomatic bone and mandibular hypoplasia), hearing loss, airway compromise
2nd pharyngeal arch	Reichert cartilage: **S**tapes, **S**tyloid process, le**SS**er horn of hyoid, **S**tylohyoid ligament	Muscles of facial expression, **S**tapedius, **S**tylohyoid, platy**S**ma, posterior belly of digastric	CN VII (facial expression) **smile**	
3rd pharyngeal arch	Greater horn of hyoid	Stylopharyngeus (think of stylo**pharyngeus** innervated by glosso**pharyngeal** nerve)	CN IX (**stylo**-pharyngeus) **swallow sty**lishly	
4th and 6th pharyngeal arches	**A**rytenoids, **C**ricoid, **C**orniculate, **C**uneiform, **T**hyroid (used to sing and **ACCCT**)	4th arch: most pharyngeal constrictors; cricothyroid, levator veli palatini 6th arch: all intrinsic muscles of larynx except cricothyroid	4th arch: CN X (superior laryngeal branch) **simply swallow** 6th arch: CN X (recurrent/inferior laryngeal branch) **speak**	Arches 3 and 4 form posterior 1/3 of tongue Arch 5 makes no major developmental contributions

[a] Sensory and motor nerves are not pharyngeal arch derivatives. They grow into the arches and are derived from neural crest (sensory) and neuroectoderm (motor).

When at the restaurant of the golden **arches**, children tend to first **chew** (1), then **smile** (2), then **swallow sty**lishly (3) or **simply swallow** (4), and then **speak** (6).

(c) ketabton.com: The Digital Library

Pharyngeal pouch derivatives

POUCH	DERIVATIVES	NOTES	MNEMONIC
1st pharyngeal pouch	Middle ear cavity, eustachian tube, mastoid air cells	1st pouch contributes to endoderm-lined structures of ear	**Ear**, **tonsils**, **bottom-to-top**: 1 (**ear**) 2 (**tonsils**) 3 dorsal (**bottom** for inferior parathyroids) 3 ventral (**t**o = **t**hymus) 4 (**top** = superior parathyroids)
2nd pharyngeal pouch	Epithelial lining of palatine tonsil		
3rd pharyngeal pouch	Dorsal wings → **inferior** parathyroids Ventral wings → thymus	3rd pouch contributes to 3 structures (thymus, left and right inferior parathyroids) 3rd-pouch structures end up **below** 4th-pouch structures	
4th pharyngeal pouch	Dorsal wings → **superior** parathyroids Ventral wings → ultimopharyngeal body → parafollicular (C) cells of thyroid		

Cleft lip and cleft palate

Distinct, multifactorial etiologies, but often occur together.

Cleft lip

Due to failure of fusion of the maxillary and merged medial nasal processes (formation of 1° palate).

Cleft palate

Due to failure of fusion of the two lateral palatine shelves or failure of fusion of lateral palatine shelf with the nasal septum and/or 1° palate (formation of 2° palate).



(c) ketabton.com: The Digital Library

Genital embryology

Female
Default development. Mesonephric duct degenerates and paramesonephric duct develops.

Male
SRY gene on Y chromosome—produces testis-determining factor → testes development.
Sertoli cells secrete Müllerian inhibitory factor (MIF, also called antimullerian hormone) that suppresses development of paramesonephric ducts.
Leydig cells secrete androgens that stimulate development of mesonephric ducts.

Paramesonephric (Müllerian) duct
Develops into female internal structures—fallopian tubes, uterus, upper portion of vagina (lower portion from urogenital sinus). Male remnant is appendix testis.
Müllerian agenesis (Mayer-Rokitansky-Küster-Hauser syndrome)—may present as 1° amenorrhea (due to a lack of uterine development) in females with fully developed 2° sexual characteristics (functional ovaries).

Mesonephric (Wolffian) duct
Develops into male internal structures (except prostate)—Seminal vesicles, Epididymis, Ejaculatory duct, Ductus deferens (SEED).
Female remnant is Gartner duct.



Sexual differentiation



1. Absence of Sertoli cells or lack of Müllerian inhibitory factor → develop both male and female internal genitalia and male external genitalia (streak gonads)
2. 5α-reductase deficiency—inability to convert testosterone into DHT → male internal genitalia, ambiguous external genitalia until puberty (when ↑ testosterone levels cause masculinization)

In the testes:
Leydig Leads to male (internal and external) sexual differentiation.
Sertoli Shuts down female (internal) sexual differentiation.

(c) ketabton.com: The Digital Library

Uterine (Müllerian duct) anomalies

Septate uterus	Common anomaly vs normal uterus **A**. Incomplete resorption of septum **B**. ↓ fertility and early miscarriage/pregnancy loss. Treat with septoplasty.
Bicornuate uterus	Incomplete fusion of Müllerian ducts **C**. ↑ risk of complicated pregnancy, early pregnancy loss, malpresentation, prematurity.
Uterus didelphys	Complete failure of fusion → double uterus, cervix, vagina **D**. Pregnancy possible.



Male/female genital homologs



Dihydrotestosterone		Estrogen
Glans penis	Genital tubercle	Glans clitoris
Corpus cavernosum and spongiosum	Genital tubercle	Vestibular bulbs
Bulbourethral glands (of Cowper)	Urogenital sinus	Greater vestibular glands (of Bartholin)
Prostate gland	Urogenital sinus	Urethral and paraurethral glands (of Skene)
Ventral shaft of penis (penile urethra)	Urogenital folds	Labia minora
Scrotum	Labioscrotal swelling	Labia majora

(c) ketabton.com: The Digital Library

Congenital penile abnormalities

Hypospadias



Abnormal opening of penile urethra on ventral surface of penis due to failure of urethral folds to fuse.

Hypospadias is more common than epispadias. Associated with inguinal hernia, cryptorchidism, chordee (downward or upward bending of penis).
Hypo is below.
Can be seen in 5α-reductase deficiency.

Epispadias



Abnormal opening of penile urethra on dorsal surface of penis due to faulty positioning of genital tubercle.

Exstrophy of the bladder is associated with Epispadias.
When you have Epispadias, you hit your Eye when you pEE.

Descent of testes and ovaries

	DESCRIPTION	MALE REMNANT	FEMALE REMNANT
Gubernaculum	Band of fibrous tissue	Anchors testes within scrotum	Ovarian ligament + round ligament of uterus
Processus vaginalis	Evagination of peritoneum	Forms tunica vaginalis Persistent patent processus vaginalis → hydrocele	Obliterated

▸ REPRODUCTIVE—ANATOMY

Gonadal drainage

Venous drainage

Left ovary/testis → left gonadal vein → left renal vein → IVC.
Right ovary/testis → right gonadal vein → IVC.
Because the left spermatic vein enters the left renal vein at a 90° angle, flow is less laminar on left than on right → left venous pressure > right venous pressure → varicocele more common on the left.

"Left gonadal vein takes the Longest way."



Lymphatic drainage

Ovaries/testes → para-aortic lymph nodes.
Body of uterus/cervix/superior part of bladder → external iliac nodes.
Prostate/cervix/corpus cavernosum/proximal vagina → internal iliac nodes.
Distal vagina/vulva/scrotum/distal anus → superficial inguinal nodes.
Glans penis → deep inguinal nodes.

(c) ketabton.com: The Digital Library

Female reproductive anatomy



LIGAMENT	CONNECTS	STRUCTURES CONTAINED	NOTES
Infundibulopelvic (suspensory) ligament	Ovaries to lateral pelvic wall	Ovarian vessels	Ligate vessels during oophorectomy to avoid bleeding Ureter courses retroperitoneally, close to gonadal vessels → at risk of injury during ligation of ovarian vessels
Cardinal (transverse cervical) ligament	Cervix to side wall of pelvis	Uterine vessels	Ureter at risk of injury during ligation of uterine vessels in hysterectomy
Round ligament of the uterus	Uterine horn to labia majora		Derivative of gubernaculum. Travels through round inguinal canal; above the artery of Sampson
Broad ligament	Uterus, fallopian tubes, and ovaries to pelvic side wall	Ovaries, fallopian tubes, round ligaments of uterus	Fold of peritoneum that comprises the mesosalpinx, mesometrium, and mesovarium
Ovarian ligament	Medial pole of ovary to uterine horn		Derivative of gubernaculum Ovarian ligament latches to lateral uterus

Adnexal torsion

Twisting of ovary and fallopian tube around infundibulopelvic ligament and ovarian ligament → compression of ovarian vessels in infundibulopelvic ligament → blockage of lymphatic and venous outflow. Continued arterial perfusion → ovarian edema → complete blockage of arterial inflow → necrosis, local hemorrhage.

Associated with ovarian masses. Presents with acute pelvic pain, adnexal mass, nausea/vomiting.

(c) ketabton.com: The Digital Library

Female reproductive epithelial histology



TISSUE	HISTOLOGY/NOTES
Vulva	Stratified squamous epithelium
Vagina	Stratified squamous epithelium, nonkeratinized
Ectocervix	Stratified squamous epithelium, nonkeratinized
Transformation zone	Squamocolumnar junction A (most common area for cervical cancer)
Endocervix	Simple columnar epithelium
Uterus	Simple columnar epithelium with long tubular glands in proliferative phase; coiled glands in secretory phase
Fallopian tube	Simple columnar epithelium, ciliated
Ovary, outer surface	Simple cuboidal epithelium (germinal epithelium covering surface of ovary)

Male reproductive anatomy





Pathway of sperm during ejaculation—
SEVEN UP:
Seminiferous tubules
Epididymis
Vas deferens
Ejaculatory ducts
(**N**othing)
Urethra
Penis

(c) ketabton.com: The Digital Library

Urethral injury

Occurs almost exclusively in men. Suspect if blood seen at urethral meatus. Urethral catheterization is relatively contraindicated.

	Anterior urethral injury	Posterior urethral injury
PART OF URETHRA	Bulbar (spongy) urethra	Membranous urethra
MECHANISM	Perineal straddle injury	Pelvic fracture
LOCATION OF URINE LEAK/BLOOD ACCUMULATION	Blood accumulates in scrotum If Buck fascia is torn, urine escapes into perineal space	Urine leaks into retropubic space
PRESENTATION	Blood at urethral meatus and scrotal hematoma	Blood at urethral meatus and high-riding prostate





Autonomic innervation of male sexual response

Erection—Parasympathetic nervous system (pelvic splanchnic nerves, S2-S4):
- NO → ↑ cGMP → smooth muscle relaxation → vasodilation → proerectile.
- Norepinephrine → ↑ $[Ca^{2+}]_{in}$ → smooth muscle contraction → vasoconstriction → antierectile.

Emission—Sympathetic nervous system (hypogastric nerve, T11-L2).

Expulsion—visceral and Somatic nerves (pudendal nerve).

Point, Squeeze, and Shoot.

S2, 3, 4 keep the penis off the floor.

PDE-5 inhibitors (eg, sildenafil) → ↓ cGMP breakdown.

(c) ketabton.com: The Digital Library

Seminiferous tubules

CELL	FUNCTION	LOCATION/NOTES
Spermatogonia	Maintain germ cell pool and produce 1° spermatocytes	Line seminiferous tubules A Germ cells
Sertoli cells	Secrete inhibin B → inhibit FSH Secrete androgen-binding protein → maintain local levels of testosterone Produce MIF Tight junctions between adjacent Sertoli cells form blood-testis barrier → isolate gametes from autoimmune attack Support and nourish developing spermatozoa Regulate spermatogenesis Temperature sensitive; ↓ sperm production and ↓ inhibin B with ↑ temperature	Line seminiferous tubules Non-germ cells Convert testosterone and androstenedione to estrogens via aromatase Sertoli cells are inSide Seminiferous tubules, Support Sperm Synthesis, and inhibit FSH Homolog of female granulosa cells ↑ temperature seen in varicocele, cryptorchidism
Leydig cells	Secrete **testosterone** in the presence of LH; testosterone production unaffected by temperature	Interstitium Endocrine cells Homolog of female theca interna cells **Ley**dies (ladies) **dig testosterone**





(c) ketabton.com: The Digital Library

▸ REPRODUCTIVE—PHYSIOLOGY

Spermatogenesis

Begins at puberty with spermatogonia. Full development takes 2 months. Occurs in seminiferous tubules. Produces spermatids that undergo spermiogenesis (loss of cytoplasmic contents, gain of acrosomal cap) to form mature spermatozoa.

"Gonium" is going to be a sperm; "Zoon" is "Zooming" to egg.

Tail mobility impaired in ciliary dyskinesia/Kartagener syndrome → infertility.

Tail mobility normal in cystic fibrosis (in CF, absent vas deferens → infertility).



(c) ketabton.com: The Digital Library

Estrogen

SOURCE	Ovary (17β-estradiol), placenta (estriol), adipose tissue (estrone via aromatization).	Potency: estradiol > estrone > estriol.
FUNCTION	Development of genitalia and breast, female fat distribution. Growth of follicle, endometrial proliferation, ↑ myometrial excitability. Upregulation of estrogen, LH, and progesterone receptors; feedback inhibition of FSH and LH, then LH surge; stimulation of prolactin secretion. ↑ transport proteins, SHBG; ↑ HDL; ↓ LDL.	Pregnancy: ▪ 50-fold ↑ in estradiol and estrone ▪ 1000-fold ↑ in estriol (indicator of fetal well-being) Estrogen receptors expressed in cytoplasm; translocate to nucleus when bound by estrogen.



Progesterone

SOURCE	Corpus luteum, placenta, adrenal cortex, testes.	Fall in progesterone after delivery disinhibits prolactin → lactation. ↑ progesterone is indicative of ovulation. **Progest**erone is **pro-gest**ation. **Prolact**in is **pro-lactat**ion.
FUNCTION	During luteal phase, prepares uterus for implantation of fertilized egg: ▪ Stimulation of endometrial glandular secretions and spiral artery development ▪ Production of thick cervical mucus → inhibits sperm entry into uterus ▪ Prevention of endometrial hyperplasia ▪ ↑ body temperature ▪ ↓ estrogen receptor expression ▪ ↓ gonadotropin (LH, FSH) secretion During pregnancy: ▪ Maintenance of pregnancy ▪ ↓ myometrial excitability → ↓ contraction frequency and intensity ▪ ↓ prolactin action on breasts	

(c) ketabton.com: The Digital Library

Oogenesis

1° oocytes begin meiosis I during fetal life and complete meiosis I just prior to ovulation.
Meiosis I is arrested in prOphase I for years until Ovulation (1° oocytes).
Meiosis II is arrested in metaphase II until fertilization (2° oocytes). "An egg met a sperm."
If fertilization does not occur within 1 day, the 2° oocyte degenerates.



Ovulation

↑ estrogen, ↑ GnRH receptors on anterior pituitary. Estrogen surge then stimulates LH release → ovulation (rupture of follicle).
↑ temperature (progesterone induced).

Mittelschmerz—transient mid-cycle ovulatory pain ("Middle hurts"); classically associated with peritoneal irritation (eg, follicular swelling/rupture, fallopian tube contraction). Can mimic appendicitis.

(c) ketabton.com: The Digital Library

Menstrual cycle

Follicular phase can vary in length. Luteal phase is 14 days. Ovulation day + 14 days = menstruation.
Follicular growth is fastest during 2nd week of the follicular phase.
Estrogen stimulates endometrial proliferation.
Progesterone maintains endometrium to support implantation.
↓ progesterone → ↓ fertility.



FAS1_2019_15-Repro.indd 632 11/7/19 5:52 PM

(c) ketabton.com: The Digital Library

Abnormal uterine bleeding

Characterized as either heavy menstrual bleeding (AUB/HMB) or intermenstrual bleeding (AUB/IMB).

These are further subcategorized by **PALM-COEIN**:

- Structural causes (**PALM**): **P**olyp, **A**denomyosis, **L**eiomyoma, or **M**alignancy/hyperplasia
- Non-structural causes (**COEIN**): **C**oagulopathy, **O**vulatory, **E**ndometrial, **I**atrogenic, **N**ot yet classified

Terms such as dysfunctional uterine bleeding, menorrhagia, oligomenorrhea are no longer recommended.

Pregnancy

Fertilization most commonly occurs in upper end of fallopian tube (the ampulla). Occurs within 1 day of ovulation.

Implantation within the wall of the uterus occurs 6 days after fertilization. Syncytiotrophoblasts secrete hCG, which is detectable in blood 1 week after conception and on home test in urine 2 weeks after conception.

Gestational age—calculated from date of last menstrual period.

Physiologic adaptations in pregnancy:

- ↑ GFR → ↓ BUN and creatinine, ↓ glucosuria threshold
- ↑ cardiac output (↑ preload, ↓ afterload, ↑ HR → ↑ placental and uterus perfusion)
- Anemia (↑↑ plasma, ↑ RBCs)
- Hypercoagulability (to ↓ blood loss at delivery)
- Hyperventilation (eliminate fetal CO_2)
- ↑ lipolysis and fat utilization (due to maternal hypoglycemia and insulin resistance) → preserves glucose and amino acids for utilization by the fetus



Placental hormone secretion generally increases over the course of pregnancy, but hCG peaks at 8–10 weeks.

Human chorionic gonadotropin

SOURCE	Syncytiotrophoblast of placenta.
FUNCTION	Maintains corpus luteum (and thus progesterone) for first 8–10 weeks of pregnancy by acting like LH (otherwise no luteal cell stimulation → abortion). After 8–10 weeks, placenta synthesizes its own estriol and progesterone and corpus luteum degenerates. Used to detect pregnancy because it appears early in urine (see above). Has identical α subunit as LH, FSH, TSH (states of ↑ hCG can cause hyperthyroidism). β subunit is unique (pregnancy tests detect β subunit). hCG is ↑ in multiple gestations, hydatidiform moles, choriocarcinomas, and Down syndrome; hCG is ↓ in ectopic/failing pregnancy, Edwards syndrome, and Patau syndrome.

(c) ketabton.com: The Digital Library

Human placental lactogen	Also called chorionic somatomammotropin.
SOURCE	Syncytiotrophoblast of placenta.
FUNCTION	Stimulates insulin production; overall ↑ insulin resistance. Gestational diabetes can occur if maternal pancreatic function cannot overcome the insulin resistance.

Apgar score

	Score 2	Score 1	Score 0
Appearance	Pink	Extremities blue	Pale or blue
Pulse	≥ 100 bpm	< 100 bpm	No pulse
Grimace	Cries and pulls away	Grimaces or weak cry	No response to stimulation
Activity	Active movement	Arms, legs flexed	No movement
Respiration	Strong cry	Slow, irregular	No breathing

Assessment of newborn vital signs following delivery via a 10-point scale evaluated at 1 minute and 5 minutes. **Apgar** score is based on **A**ppearance, **P**ulse, **G**rimace, **A**ctivity, and **R**espiration. Apgar scores < 7 may require further evaluation. If Apgar score remains low at later time points, there is ↑ risk the child will develop long-term neurologic damage.

(c) ketabton.com: The Digital Library

Infant and child development

Milestone dates are ranges that have been approximated and vary by source. Children not meeting milestones may need assessment for potential developmental delay.

AGE	MOTOR	SOCIAL	VERBAL/COGNITIVE
Infant	**Parents**	**Start**	**Observing,**
0–12 mo	Primitive reflexes disappear—Moro (by 3 mo), rooting (by 4 mo), palmar (by 6 mo), Babinski (by 12 mo) Posture—lifts head up prone (by 1 mo), rolls and sits (by 6 mo), crawls (by 8 mo), stands (by 10 mo), walks (by 12–18 mo) Picks—passes toys hand to hand (by 6 mo), Pincer grasp (by 10 mo) Points to objects (by 12 mo)	Social smile (by 2 mo) Stranger anxiety (by 6 mo) Separation anxiety (by 9 mo)	Orients—first to voice (by 4 mo), then to name and gestures (by 9 mo) Object permanence (by 9 mo) Oratory—says "mama" and "dada" (by 10 mo)
Toddler	**Child**	**Rearing**	**Working,**
12–36 mo	Cruises, takes first steps (by 12 mo) Climbs stairs (by 18 mo) Cubes stacked—number = age (yr) × 3 Cutlery—feeds self with fork and spoon (by 20 mo) Kicks ball (by 24 mo)	Recreation—parallel play (by 24–36 mo) Rapprochement—moves away from and returns to mother (by 24 mo) Realization—core gender identity formed (by 36 mo)	Words—uses 50-200 words by 2 yr, uses 300+ words by 3 yr.
Preschool	**Don't**	**Forget, they're still**	**Learning!**
3–5 yr	Drive—tricycle (3 wheels at 3 yr) Drawings—copies line or circle, stick figure (by 4 yr) Dexterity—hops on one foot (by 4 yr), uses buttons or zippers, grooms self (by 5 yr)	Freedom—comfortably spends part of day away from mother (by 3 yr) Friends—cooperative play, has imaginary friends (by 4 yr)	Language—understands 1000 words by 3 yr (3 zeros), uses complete sentences and prepositions (by 4 yr) Legends—can tell detailed stories (by 4 yr)

Low birth weight

Defined as < 2500 g. Caused by prematurity or intrauterine growth restriction (IUGR). Associated with ↑ risk of sudden infant death syndrome (SIDS) and with ↑ overall mortality.

(c) ketabton.com: The Digital Library

Lactation

After parturition and delivery of placenta, rapid ↓ in progesterone disinhibits prolactin → initiation of lactation. Suckling is required to maintain milk production and ejection, since ↑ nerve stimulation → ↑ oxytocin and prolactin.

Prolactin—induces and maintains lactation and ↓ reproductive function.

Oxytocin—assists in milk letdown; also promotes uterine contractions.

Breast milk is the ideal nutrition for infants < 6 months old. Contains maternal immunoglobulins (conferring passive immunity; mostly IgA), macrophages, lymphocytes. Breast milk reduces infant infections and is associated with ↓ risk for child to develop asthma, allergies, diabetes mellitus, and obesity. Guidelines recommend exclusively breastfed infants get vitamin D and possibly iron supplementation.

Breastfeeding ↓ maternal risk of breast and ovarian cancer and facilitates mother-child bonding.

Menopause

Diagnosed by amenorrhea for 12 months. ↓ estrogen production due to age-linked decline in number of ovarian follicles. Average age at onset is 51 years (earlier in smokers).

Usually preceded by 4–5 years of abnormal menstrual cycles. Source of estrogen (estrone) after menopause becomes peripheral conversion of androgens, ↑ androgens → hirsutism.

↑↑ FSH is specific for menopause (loss of negative feedback on FSH due to ↓ estrogen).

Hormonal changes: ↓ estrogen, ↑↑ FSH, ↑ LH (no surge), ↑ GnRH.

Causes **HAVOCS**: **H**ot flashes, **A**trophy of the **V**agina, **O**steoporosis, **C**oronary artery disease, **S**leep disturbances.

Menopause before age 40 suggests 1° ovarian insufficiency (premature ovarian failure); may occur in women who have received chemotherapy and/or radiation therapy.

Androgens

Testosterone, dihydrotestosterone (DHT), androstenedione.

SOURCE	DHT and testosterone (testis), **AnD**rostenedione (**AD**renal)	Potency: DHT > testosterone > androstenedione.
FUNCTION	Testosterone: ▪ Differentiation of epididymis, vas deferens, seminal vesicles (internal genitalia, except prostate) ▪ Growth spurt: penis, seminal vesicles, sperm, muscle, RBCs ▪ Deepening of voice ▪ Closing of epiphyseal plates (via estrogen converted from testosterone) ▪ Libido DHT: ▪ Early—differentiation of penis, scrotum, prostate ▪ Late—prostate growth, balding, sebaceous gland activity	Testosterone is converted to DHT by 5α-reductase, which is inhibited by finasteride. In the male, androgens are converted to estrogen by cytochrome P-450 aromatase (primarily in adipose tissue and testis). Aromatase is the key enzyme in conversion of androgens to estrogen. Androgenic steroid abuse—abuse of anabolic steroids to ↑ fat-free mass, muscle strength, and performance. Suspect in men who present with changes in behavior (eg, aggression), acne, gynecomastia, ↑ Hb and Hct, small testes (exogenous testosterone → hypothalamic-pituitary-gonadal axis inhibition → ↓ intratesticular testosterone → ↓ testicular size, ↓ sperm count, azoospermia). Women may present with virilization (eg, hirsutism, acne, breast atrophy, male pattern baldness).

(c) ketabton.com: The Digital Library

Tanner stages of sexual development

Tanner stage is assigned independently to genitalia, pubic hair, and breast (eg, a person can have Tanner stage 2 genitalia, Tanner stage 3 pubic hair). Earliest detectable secondary sexual characteristic is breast bud development in girls, testicular enlargement in boys.



Precocious puberty

Appearance of 2° sexual characteristics (eg, adrenarche, thelarche, menarche) before age 8 years in girls and 9 years in boys. ↑ sex hormone exposure or production → ↑ linear growth, somatic and skeletal maturation (eg, premature closure of epiphyseal plates → short stature). Types include:

- Central precocious puberty (↑ GnRH secretion): idiopathic (most common; early activation of hypothalamic-pituitary gonadal axis), CNS tumors.
- Peripheral precocious puberty (GnRH-independent; ↑ sex hormone production or exposure to exogenous sex steroids): congenital adrenal hyperplasia, estrogen-secreting ovarian tumor (eg, granulosa cell tumor), Leydig cell tumor, McCune-Albright syndrome.

(c) ketabton.com: The Digital Library

▸ REPRODUCTIVE—PATHOLOGY

Sex chromosome disorders

Aneuploidy most commonly due to meiotic nondisjunction.

Klinefelter syndrome

Male, 47,XXY.
Testicular atrophy, eunuchoid body shape, tall, long extremities, gynecomastia, female hair distribution A. May present with developmental delay. Presence of inactivated X chromosome (Barr body). Common cause of hypogonadism seen in infertility work-up.

Dysgenesis of seminiferous tubules → ↓ inhibin B → ↑ FSH.
Abnormal Leydig cell function → ↓ testosterone → ↑ LH → ↑ estrogen.

Turner syndrome

Female, 45,XO.
Short stature (associated with *SHOX* gene, preventable with growth hormone therapy), ovarian dysgenesis (streak ovary), shield chest B, bicuspid aortic valve, coarctation of the aorta (femoral < brachial pulse), lymphatic defects (result in webbed neck or cystic hygroma; lymphedema in feet, hands), horseshoe kidney, high-arched palate, shortened 4th metacarpals.
Most common cause of 1° amenorrhea. No Barr body.

Menopause before menarche.
↓ estrogen leads to ↑ LH, FSH.
Sex chromosome (X, or rarely Y) loss often due to nondisjunction during meiosis or mitosis.
Meiosis errors usually occur in paternal gametes → sperm missing the sex chromosome.
Mitosis errors occur after zygote formation → loss of sex chromosome in some but not all cells → mosaic karyotype (eg. 45,X/46XX).
(45,X/46,XY) mosaicism associated with increased risk for gonadoblastoma.
Pregnancy is possible in some cases (IVF, exogenous estradiol-17β and progesterone).

Double Y males

47, XYY.
Phenotypically normal (usually undiagnosed), very tall. Normal fertility. May be associated with severe acne, learning disability, autism spectrum disorders.

Ovotesticular disorder of sex development

46,XX > 46,XY.
Both ovarian and testicular tissue present (ovotestis); ambiguous genitalia. Previously called true hermaphroditism.

(c) ketabton.com: The Digital Library

Diagnosing disorders of sex hormones

Testosterone	LH	Diagnosis
↑	↑	Defective androgen receptor
↑	↓	Testosterone-secreting tumor, exogenous steroids
↓	↑	Hypergonadotropic hypogonadism (1°)
↓	↓	Hypogonadotropic hypogonadism (2°)

Other disorders of sex development

Disagreement between the phenotypic sex (external genitalia, influenced by hormonal levels) and the gonadal sex (testes vs ovaries, corresponds with Y chromosome). Includes the terms pseudohermaphrodite, hermaphrodite, and intersex.

46,XX DSD

Ovaries present, but external genitalia are virilized or ambiguous. Due to excessive and inappropriate exposure to androgenic steroids during early gestation (eg, congenital adrenal hyperplasia or exogenous administration of androgens during pregnancy).

46,XY DSD

Testes present, but external genitalia are female or ambiguous. Most common form is androgen insensitivity syndrome (testicular feminization).

Disorders by physical characteristics

UTERUS	BREASTS	DISORDERS
⊕	⊖	Hypergonadotropic hypogonadism (eg, Turner syndrome, genetic mosaicism, pure gonadal dysgenesis) Hypogonadotropic hypogonadism (eg, CNS lesions, Kallmann syndrome)
⊖	⊕	Uterovaginal agenesis in genotypic female or androgen insensitivity in genotypic male
⊖	⊖	Male genotype with insufficient production of testosterone

Placental aromatase deficiency

Inability to synthesize estrogens from androgens. Masculinization of female (46,XX DSD) infants (ambiguous genitalia), ↑ serum testosterone and androstenedione. Can present with maternal virilization during pregnancy (fetal androgens cross the placenta).

Androgen insensitivity syndrome

Defect in androgen receptor resulting in normal-appearing female (46,XY DSD); female external genitalia with scant axillary and pubic hair, rudimentary vagina; uterus and fallopian tubes absent due to persistence of anti-Müllerian hormone from testes. Patients develop normal functioning testes (often found in labia majora; surgically removed to prevent malignancy). ↑ testosterone, estrogen, LH (vs sex chromosome disorders).

5α-reductase deficiency

Autosomal recessive; sex limited to genetic males (46,XY DSD). Inability to convert testosterone to DHT. Ambiguous genitalia until puberty, when ↑ testosterone causes masculinization/↑ growth of external genitalia. Testosterone/estrogen levels are normal; LH is normal or ↑. Internal genitalia are normal.

Kallmann syndrome

Failure to complete puberty; a form of hypogonadotropic hypogonadism. Defective migration of neurons and subsequent failure of olfactory bulbs to develop → ↓ synthesis of GnRH in the hypothalamus; hyposmia/anosmia; ↓ GnRH, FSH, LH, testosterone. Infertility (low sperm count in males; amenorrhea in females).

(c) ketabton.com: The Digital Library

Pregnancy complications

Abruptio placentae

Premature separation (partial or complete) of placenta from uterine wall before delivery of infant. Risk factors: trauma (eg, motor vehicle accident), smoking, hypertension, preeclampsia, cocaine abuse.

Presentation: **abrupt**, painful bleeding (concealed or apparent) in third trimester; possible DIC (mediated by tissue factor activation), maternal shock, fetal distress. May be life threatening for mother and fetus.



Complete abruption with concealed hemorrhage

Partial abruption (blue arrow) with apparent hemorrhage (red arrow)

Morbidly adherent placenta

Defective decidual layer → abnormal attachment and separation after delivery. Risk factors: prior C-section or uterine surgery involving myometrium, inflammation, placenta previa, advanced maternal age, multiparity. Three types distinguishable by the depth of penetration:

Placenta **accreta**—placenta attaches to myometrium without penetrating it; most common type.

Placenta **increta**—placenta penetrates into myometrium.

Placenta **percreta**—placenta penetrates ("perforates") through myometrium and into uterine serosa (invades entire uterine wall); can result in placental attachment to rectum or bladder (can result in hematuria).

Presentation: often detected on ultrasound prior to delivery. No separation of placenta after delivery → postpartum bleeding (can cause Sheehan syndrome).



Placenta previa

Attachment of placenta over internal cervical os. Risk factors: multiparity, prior C-section. Associated with painless third-trimester bleeding. A "**previ**ew" of the **placenta** is visible through cervix.

Low-lying placenta (< 2 cm from internal cervical os, but not over it) is managed differently from placenta previa.



Partial placenta previa

Complete placenta previa

(c) ketabton.com: The Digital Library

Pregnancy complications *(continued)*

Vasa previa

Fetal vessels run over, or in close proximity to, cervical os. May result in vessel rupture, exsanguination, fetal death. Presents with triad of membrane rupture, painless vaginal bleeding, fetal bradycardia (< 110 beats/min). Emergency C-section usually indicated. Frequently associated with velamentous umbilical cord insertion (cord inserts in chorioamniotic membrane rather than placenta → fetal vessels travel to placenta unprotected by Wharton jelly).



Postpartum hemorrhage

Due to 4 T's: Tone (uterine atony; most common), Trauma (lacerations, incisions, uterine rupture), Thrombin (coagulopathy), Tissue (retained products of conception).

Treatment: uterine massage, oxytocin. If refractory, surgical ligation of uterine or internal iliac artery (will preserve fertility since ovarian arteries provide collateral circulation).

Ectopic pregnancy

Implantation of fertilized ovum in a site other than the uterus, most often in ampulla of fallopian tube A. Suspect with history of amenorrhea, lower-than-expected rise in hCG based on dates, and sudden lower abdominal pain; confirm with ultrasound, which may show extraovarian adnexal mass. Often clinically mistaken for appendicitis.

Pain +/– bleeding.

Risk factors:

- Prior ectopic pregnancy
- History of infertility
- Salpingitis (PID)
- Ruptured appendix
- Prior tubal surgery
- Smoking
- Advanced maternal age



Amniotic fluid abnormalities

Polyhydramnios

Too much amniotic fluid. Often idiopathic, but associated with fetal malformations (eg, esophageal/duodenal atresia, anencephaly; both result in inability to swallow amniotic fluid), maternal diabetes, fetal anemia, multiple gestations.

Oligohydramnios

Too little amniotic fluid. Associated with placental insufficiency, bilateral renal agenesis, posterior urethral valves (in males) and resultant inability to excrete urine. Any profound oligohydramnios can cause Potter sequence.

(c) ketabton.com: The Digital Library

Hydatidiform mole





Cystic swelling of chorionic villi and proliferation of chorionic epithelium (only trophoblast).
Presents with vaginal bleeding, emesis, uterine enlargement more than expected, pelvic pressure/pain. Associated with hCG-mediated sequelae: early preeclampsia (before 20 weeks), theca-lutein cysts, hyperemesis gravidarum, hyperthyroidism.
Treatment: dilation and curettage and methotrexate. Monitor hCG.

	Complete mole	Partial mole
KARYOTYPE	46,XX; 46,XY	69,XXX; 69,XXY; 69,XYY
COMPONENTS	Most commonly enucleated egg + single sperm (subsequently duplicates paternal DNA)	2 sperm + 1 egg
HISTOLOGY	Hydropic villi, circumferential and diffuse trophoblastic proliferation	Only some villi are hydropic, focal/minimal trophoblastic proliferation
FETAL PARTS	No	Yes (**part**ial = fetal **part**s)
STAINING FOR P57 PROTEIN	⊖ (paternally imprinted)	⊕ (maternally expressed)
UTERINE SIZE	↑	—
hCG	↑↑↑↑	↑
IMAGING	"Honeycombed" uterus or "clusters of grapes" A, "snowstorm" B on ultrasound	Fetal parts
RISK OF INVASIVE MOLE	15–20%	< 5%
RISK OF CHORIOCARCINOMA	2%	Rare

Choriocarcinoma



Rare; can develop during or after pregnancy in mother or baby. Malignancy of trophoblastic tissue A (cytotrophoblasts, syncytiotrophoblasts); **no** chorionic villi present. ↑ frequency of bilateral/multiple theca-lutein cysts. Presents with abnormal ↑ hCG, shortness of breath, hemoptysis. Hematogenous spread to lungs → "cannonball" metastases B.
Treatment: methotrexate.



(c) ketabton.com: The Digital Library

Hypertension in pregnancy

Gestational hypertension	BP > 140/90 mm Hg after 20th week of gestation. No pre-existing hypertension. No proteinuria or end-organ damage.	Treatment: antihypertensives (**H**ydralazine, α-**M**ethyldopa, **L**abetalol, **N**ifedipine), deliver at 37–39 weeks. **H**ypertensive **M**oms **L**ove **N**ifedipine.
Preeclampsia	New-onset hypertension with either proteinuria or end-organ dysfunction after 20th week of gestation (< 20 weeks suggests molar pregnancy). Caused by abnormal placental spiral arteries → endothelial dysfunction, vasoconstriction, ischemia. Incidence ↑ in patients with pre-existing hypertension, diabetes, chronic kidney disease, autoimmune disorders (eg, antiphospholipid antibody syndrome), age > 40 years. Complications: placental abruption, coagulopathy, renal failure, pulmonary edema, uteroplacental insufficiency; may lead to eclampsia (+ seizures) and/or HELLP syndrome.	Treatment: antihypertensives, IV magnesium sulfate (to prevent seizure); definitive is delivery of fetus. **P**roteinuria, **R**ising BP (new-onset HTN), **E**nd-organ dysfunction (eg, pulmonary edema).
Eclampsia	Preeclampsia + maternal seizures. Maternal death due to stroke, intracranial hemorrhage, or ARDS.	Treatment: IV magnesium sulfate, antihypertensives, immediate delivery.
HELLP syndrome	**H**emolysis, **E**levated **L**iver enzymes, **L**ow **P**latelets. A manifestation of severe preeclampsia. Blood smear shows schistocytes. Can lead to DIC (due to release of tissue factor from injured placenta) and hepatic subcapsular hematomas → rupture → severe hypotension.	Treatment: immediate delivery.

Gynecologic tumor epidemiology	Incidence (US)—endometrial > ovarian > cervical; cervical cancer is more common worldwide due to lack of screening or HPV vaccination. Prognosis: **C**ervical (**best** prognosis, diagnosed < 45 years old) > **E**ndometrial (middle-aged, about 55 years old) > **O**varian (**worst** prognosis, > 65 years).	**CEO**s often go from **best** to **worst** as they get **older**.

(c) ketabton.com: The Digital Library

Vulvar pathology

Non-neoplastic	
Bartholin cyst and abscess	Due to blockage of Bartholin gland duct causing accumulation of gland fluid. May lead to abscess 2° to obstruction and inflammation A. Usually in reproductive-age females.
Lichen sclerosus	Thinning of epidermis with fibrosis/sclerosis of dermis. Presents with porcelain-white plaques with a red or violet border. Skin fragility with erosions can be observed B. Most common in postmenopausal women. Benign, but slightly increased risk for SCC.
Lichen simplex chronicus	Hyperplasia of vulvar squamous epithelium. Presents with leathery, thick vulvar skin with enhanced skin markings due to chronic rubbing or scratching. Benign, no risk of SCC.
Neoplastic	
Vulvar carcinoma	Carcinoma from squamous epithelial lining of vulva C. Rare. Presents with leukoplakia, biopsy often required to distinguish carcinoma from other causes. HPV-related vulvar carcinoma—associated with high-risk HPV types 16, 18. Risk factors: multiple partners, early coitarche. Usually in reproductive-age females. Non-HPV vulvar carcinoma—usually from long-standing lichen sclerosus. Females > 70 years old.
Extramammary Paget disease	Intraepithelial adenocarcinoma. Carcinoma in situ, low risk of underlying carcinoma (vs Paget disease of the breast, which is always associated with underlying carcinoma). Presents with pruritus, erythema, crusting, ulcers D.



Imperforate hymen — Incomplete degeneration of the central portion of the hymen. Accumulation of vaginal mucus at birth → self-resolving bulge in introitus. If untreated, leads to 1° amenorrhea, cyclic abdominal pain, hematocolpos (accumulation of menstrual blood in vagina → bulging and bluish hymenal membrane).

Vaginal tumors

Vaginal squamous cell carcinoma	Usually 2° to cervical SCC; 1° vaginal carcinoma rare.
Clear cell adenocarcinoma	Affects women who had exposure to DES in utero.
Sarcoma botryoides	Embryonal rhabdomyosarcoma variant. Affects girls < 4 years old; spindle-shaped cells; desmin ⊕. Presents with clear, grape-like, polypoid mass emerging from vagina.

(c) ketabton.com: The Digital Library

Cervical pathology

Dysplasia and carcinoma in situ



Disordered epithelial growth; begins at basal layer of squamocolumnar junction (transformation zone) and extends outward. Classified as CIN 1, CIN 2, or CIN 3 (severe, irreversible dysplasia or carcinoma in situ), depending on extent of dysplasia. Associated with HPV-16 and HPV-18, which produce both the E6 gene product (inhibits *TP*53) and E7 gene product (inhibits *pRb*) (6 before 7; P before R). Koilocytes A are pathognomonic of HPV infection. May progress slowly to invasive carcinoma if left untreated. Typically asymptomatic (detected with Pap smear) or presents as abnormal vaginal bleeding (often postcoital).

Risk factors: multiple sexual partners, HPV, smoking, early coitarche, DES exposure, immunocompromise (eg, HIV, transplant).

Invasive carcinoma

Often squamous cell carcinoma. Pap smear can detect cervical dysplasia before it progresses to invasive carcinoma. Diagnose via colposcopy and biopsy. Lateral invasion can block ureters → hydronephrosis → renal failure.

Primary ovarian insufficiency

Also called premature ovarian failure.

Premature atresia of ovarian follicles in women of reproductive age. Most often idiopathic; associated with chromosomal abnormalities (especially in females < 30 years), autoimmunity. Need karyotype screening. Patients present with signs of menopause after puberty but before age 40. ↓ estrogen, ↑ LH, ↑ FSH.

Most common causes of anovulation

Pregnancy, polycystic ovarian syndrome, obesity, HPO axis abnormalities/immaturity, premature ovarian failure, hyperprolactinemia, thyroid disorders, eating disorders, competitive athletics, Cushing syndrome, adrenal insufficiency, chromosomal abnormalities (eg, Turner syndrome).

Functional hypothalamic amenorrhea

Also called exercise-induced amenorrhea. Severe caloric restriction, ↑ energy expenditure, and/or stress → functional disruption of pulsatile GnRH secretion → ↓ LH, FSH, estrogen. Pathogenesis includes ↓ leptin (due to ↓ fat) and ↑ cortisol (stress, excessive exercise).

Associated with eating disorders and "female athlete triad" (↓ calorie availability/excessive exercise, ↓ bone mineral density, menstrual dysfunction).

Polycystic ovarian syndrome



Hyperinsulinemia and/or insulin resistance hypothesized to alter hypothalamic hormonal feedback response → ↑ LH:FSH, ↑ androgens (eg, testosterone) from theca interna cells, ↓ rate of follicular maturation → unruptured follicles (cysts) + anovulation. Common cause of ↓ fertility in women.

Enlarged, bilateral cystic ovaries A; presents with amenorrhea/oligomenorrhea, hirsutism, acne, ↓ fertility. Associated with obesity, acanthosis nigricans. ↑ risk of endometrial cancer 2° to unopposed estrogen from repeated anovulatory cycles.

Treatment: cycle regulation via weight reduction (↓ peripheral estrone formation), OCPs (prevent endometrial hyperplasia due to unopposed estrogen); clomiphene (ovulation induction); spironolactone, finasteride, flutamide to treat hirsutism.

Primary dysmenorrhea

Painful menses, caused by uterine contractions to ↓ blood loss → ischemic pain. Mediated by prostaglandins. Treatment: NSAIDs.

(c) ketabton.com: The Digital Library

Ovarian cysts

Follicular cyst	Distention of unruptured Graafian follicle. May be associated with hyperestrogenism, endometrial hyperplasia. Most common ovarian mass in young women.
Theca-lutein cyst	Often bilateral/multiple. Due to gonadotropin stimulation. Associated with choriocarcinoma and hydatidiform moles.

Ovarian neoplasms

Most common adnexal mass in women >55 years old. Present with abdominal distention, bowel obstruction, pleural effusion.

Risk ↑ with advanced age, infertility, endometriosis, PCOS, genetic predisposition (eg, *BRCA1* or *BRCA2* mutations, Lynch syndrome, strong family history).

Risk ↓ with previous pregnancy, history of breastfeeding, OCPs, tubal ligation.

Epithelial tumors are typically serous (lined by serous epithelium natively found in fallopian tubes, and often bilateral) or mucinous (lined by mucinous epithelium natively found in cervix). Monitor response to therapy/relapse by measuring CA 125 levels (not good for screening).

Germ cell tumors can differentiate into somatic structures (eg, teratomas), or extra-embryonic structures (eg, yolk sac tumors), or can remain undifferentiated (eg, dysgerminoma).

Sex cord stromal tumors develop from embryonic sex cord (develops into theca and granulosa cells of follicle, Sertoli and Leydig cells of seminiferous tubules) and stromal (ovarian cortex) derivatives.



TYPE	MALIGNANT?	CHARACTERISTICS
Epithelial tumors		
Serous cystadenoma	Benign	Most common ovarian neoplasm.
Serous cystadenocarcinoma	Malignant	Most common malignant ovarian neoplasm. Psammoma bodies.
Mucinous cystadenoma	Benign	Multiloculated, large. Lined by mucus-secreting epithelium A.
Mucinous cystadenocarcinoma	Malignant	Rare. May be metastatic from appendiceal or GI tumors. Can result in pseudomyxoma peritonei (intraperitoneal accumulation of mucinous material).
Brenner tumor	Usually benign	Solid, pale yellow-tan tumor that appears encapsulated. "Coffee bean" nuclei on H&E stain.

(c) ketabton.com: The Digital Library

Ovarian neoplasms *(continued)*

Germ cell tumors		
Dysgerminoma	Malignant	Most common in adolescents. Equivalent to male seminoma but rarer. Sheets of uniform "fried egg" cells B. Tumor markers: ↑ hCG, LDH.
Mature cystic teratoma	Benign	Also called dermoid cyst. Most common ovarian tumor in young females. Cystic mass with elements from all 3 germ layers (eg, teeth, hair, sebum) C. May be painful 2° to ovarian enlargement or torsion. Monodermal form with thyroid tissue (struma ovarii D) may present with hyperthyroidism. Malignant transformation rare (usually to squamous cell carcinoma).
Immature teratoma	Malignant, aggressive	Contains fetal tissue, neuroectoderm. Commonly diagnosed before age 20. Typically represented by immature/embryonic-like neural tissue.
Yolk sac (endodermal sinus) tumor	Malignant, aggressive	Occur in ovaries and sacrococcygeal area in children. Yellow, friable (hemorrhagic) mass. 50% have Schiller-Duval bodies (resemble glomeruli, arrow in E). Tumor marker: ↑ AFP.
Sex cord stromal tumors		
Thecoma	Benign	May produce estrogen. Usually presents as abnormal uterine bleeding in a postmenopausal woman.
Granulosa cell tumor	Malignant	Most common malignant sex cord stromal tumor. Predominantly women in their 50s. Often produces estrogen and/or progesterone and presents with postmenopausal bleeding, endometrial hyperplasia, sexual precocity (in pre-adolescents), breast tenderness. Histology shows Call-Exner bodies (granulosa cells arranged haphazardly around collections of eosinophilic fluid, resembling primordial follicles; arrow in F). "Give Granny a Call!"
Sertoli-Leydig cell tumor	Benign	Small, grey to yellow-brown mass. Resembles testicular histology with tubules/cords lined by pink Sertoli cells. May produce androgens → virilization (eg, hirsutism, male pattern baldness, clitoral enlargement).
Fibromas	Benign	Bundles of spindle-shaped fibroblasts. Meigs syndrome—triad of ovarian fibroma, ascites, pleural effusion. "Pulling" sensation in groin.



(c) ketabton.com: The Digital Library

Uterine conditions

Non-neoplastic uterine conditions	
Adenomyosis	Extension of endometrial tissue (glandular) into uterine myometrium. Caused by hyperplasia of basal layer of endometrium. Presents with dysmenorrhea, AUB/HMB, and uniformly enlarged, soft, globular uterus. Treatment: GnRH agonists, hysterectomy, excision of an organized adenomyoma.
Asherman syndrome	Adhesions and/or fibrosis of the endometrium. Presents with ↓ fertility, recurrent pregnancy loss, AUB, pelvic pain. Often associated with dilation and curettage of intrauterine cavity.
Endometrial hyperplasia	Abnormal endometrial gland proliferation usually stimulated by excess estrogen. ↑ risk for endometrial carcinoma (especially with nuclear atypia). Presents as postmenopausal vaginal bleeding. ↑ risk with anovulatory cycles, hormone replacement therapy, PCOS, granulosa cell tumors.
Endometriosis	Endometrium-like glands/stroma outside endometrial cavity, most commonly in the ovary (frequently bilateral), pelvis, peritoneum (yellow-brown "powder burn" lesions). In ovary, appears as endometrioma (blood-filled "chocolate cysts" [oval structures above and below asterisks in **A**]). May be due to retrograde flow, metaplastic transformation of multipotent cells, transportation of endometrial tissue via lymphatic system. Characterized by cyclic pelvic pain, bleeding, dysmenorrhea, dyspareunia, dyschezia (pain with defecation), infertility; normal-sized uterus. Treatment: NSAIDs, OCPs, progestins, GnRH agonists, danazol, laparoscopic removal.
Endometritis	Inflammation of endometrium **B** associated with retained products of conception following delivery, miscarriage, abortion, or with foreign body (eg, IUD). Retained material is nidus for bacteria from vagina or GI tract. Chronic endometritis shows plasma cells on histology. Treatment: gentamicin + clindamycin +/– ampicillin.
Uterine neoplasms	
Endometrial carcinoma	Most common gynecologic malignancy **C**. Presents with irregular vaginal bleeding. Two types: Endometrioid—most cases caused by unopposed estrogen exposure due to obesity, but also associated with early menarche, late menopause, nulliparity. Histology shows abnormally arranged endometrial glands. Early pathogenic events include loss of PTEN or mismatch repair proteins. Serous—associated with endometrial atrophy in postmenopausal women. Aggressive. Psammoma bodies often seen on histology. Characterized by formation of papillae and tufts.
Leiomyoma (fibroid)	Most common tumor in females. Often presents with multiple discrete tumors **D**. ↑ incidence in African Americans. Benign smooth muscle tumor; malignant transformation to leiomyosarcoma is rare. Estrogen sensitive; tumor size ↑ with pregnancy and ↓ with menopause. Peak occurrence at 20-40 years of age. May be asymptomatic, cause AUB, or result in miscarriage. Severe bleeding may lead to iron deficiency anemia. Whorled pattern of smooth muscle bundles with well-demarcated borders on histology **E**.
Leiomyosarcoma	Malignant proliferation of smooth muscle arising from myometrium; arises de novo (not from leiomyomas), usually in postmenopausal women. Exam shows single lesion with areas of necrosis.



(c) ketabton.com: The Digital Library

Breast pathology



Benign breast diseases

Fibrocystic changes

Most common in premenopausal women 20-50 years old. Present with premenstrual breast pain or lumps; often bilateral and multifocal. Nonproliferative lesions include simple cysts (fluid-filled duct dilation, blue dome), papillary apocrine change/metaplasia, stromal fibrosis. Risk of cancer is usually not increased. Subtypes include:

- Sclerosing adenosis—acini and stromal fibrosis, associated with calcifications. Slight ↑ risk for cancer.
- Epithelial hyperplasia—cells in terminal ductal or lobular epithelium. ↑ risk of carcinoma with atypical cells.

Inflammatory processes

Fat necrosis—benign, usually painless, lump due to injury to breast tissue. Calcified oil cyst on mammography; necrotic fat and giant cells on biopsy. Up to 50% of patients may not report trauma.

Lactational mastitis—occurs during breastfeeding, ↑ risk of bacterial infection through cracks in nipple. *S aureus* is most common pathogen. Treat with antibiotics and continue breastfeeding.

Benign tumors

Fibroadenoma—most common in women < 35 years old. Small, well-defined, mobile mass A. Tumor composed of fibrous tissue and glands. ↑ size and tenderness with ↑ estrogen (eg, pregnancy, prior to menstruation). Risk of cancer is usually not increased.

Intraductal papilloma—small fibroepithelial tumor within lactiferous ducts, typically beneath areola. Most common cause of nipple discharge (serous or bloody). Slight ↑ risk for cancer.

Phyllodes tumor—large mass B of connective tissue and cysts with "leaf-like" lobulations C. Most common in 5th decade. Some may become malignant.

Gynecomastia

Breast enlargement in males due to ↑ estrogen compared with androgen activity. Physiologic in newborn, pubertal, and elderly males, but may persist after puberty. Other causes include cirrhosis, hypogonadism (eg, Klinefelter syndrome), testicular tumors, and drugs (Spironolactone, Hormones, Cimetidine, Finasteride, Ketoconazole: "Some Hormones Create Funny Knockers").



(c) ketabton.com: The Digital Library

Breast cancer



Commonly postmenopausal. Often presents as a palpable hard mass A most often in the upper outer quadrant. Invasive cancer can become fixed to pectoral muscles, deep fascia, Cooper ligaments, and overlying skin → nipple retraction/skin dimpling.

Usually arises from terminal duct lobular unit. Amplification/overexpression of estrogen/progesterone receptors or *c-erbB2* (HER2, an EGF receptor) is common; triple negative (ER ⊖, PR ⊖, and HER2/neu ⊖) form more aggressive.

Risk factors in women: ↑ age; history of atypical hyperplasia; family history of breast cancer; race (Caucasians at highest risk, African Americans at ↑ risk for triple ⊖ breast cancer); *BRCA1/BRCA2* mutations; ↑ estrogen exposure (eg, nulliparity); postmenopausal obesity (adipose tissue converts androstenedione to estrone); ↑ total number of menstrual cycles; absence of breastfeeding; later age of first pregnancy; alcohol intake. In men: *BRCA2* mutation, Klinefelter syndrome.

Axillary lymph node metastasis most important prognostic factor in early-stage disease.

TYPE	CHARACTERISTICS	NOTES
Noninvasive carcinomas		
Ductal carcinoma in situ	Fills ductal lumen (black arrow in B indicates neoplastic cells in duct; blue arrow shows engorged blood vessel). Arises from ductal atypia. Often seen early as microcalcifications on mammography.	Early malignancy without basement membrane penetration. Usually does not produce a mass. Comedocarcinoma—Subtype of DCIS. Cells have high-grade nuclei with extensive central necrosis C and dystrophic calcification.
Paget disease	Extension of underlying DCIS/invasive breast cancer up the lactiferous ducts and into the contiguous skin of nipple → eczematous patches over nipple and areolar skin D.	Paget cells = intraepithelial adenocarcinoma cells.
Lobular carcinoma in situ	↓ E-cadherin expression. No mass or calcifications → incidental biopsy finding.	↑ risk of cancer in either breast (vs DCIS, same breast and quadrant).
Invasive carcinomas[a]		
Invasive ductal	Firm, fibrous, "rock-hard" mass with sharp margins and small, glandular, duct-like cells in desmoplastic stroma.	
Invasive lobular	↓ E-cadherin expression → orderly row of cells ("single file" E) and no duct formation. Often lacks desmoplastic response.	Often bilateral with multiple lesions in the same location. Lines of cells = Lobular.
Medullary	Large, anaplastic cells growing in sheets with associated lymphocytes and plasma cells.	Well-circumscribed tumor can mimic fibroadenoma.
Inflammatory	Dermal lymphatic space invasion → breast pain with warm, swollen, erythematous skin around exaggerated hair follicles, peau d'orange F.	Poor prognosis (50% survival at 5 years). Often mistaken for mastitis or Paget disease. Usually lacks a palpable mass.

[a]All types of invasive breast carcinoma can be either of tubular subtype (well-differentiated tubules that lack myoepithelium) or mucinous subtype (abundant extracellular mucin, seen in older women).

(c) ketabton.com: The Digital Library

Penile pathology

Peyronie disease

Abnormal curvature of penis A due to fibrous plaque within tunica albuginea. Associated with erectile dysfunction. Can cause pain, anxiety. Consider surgical repair or treatment with collagenase injections once curvature stabilizes. Distinct from penile fracture (rupture of corpora cavernosa due to forced bending).

Ischemic priapism

Painful sustained erection lasting > 4 hours. Associated with sickle cell disease (sickled RBCs block venous drainage of corpus cavernosum vascular channels), medications (eg, sildenafil, trazodone). Treat immediately with corporal aspiration, intracavernosal phenylephrine, or surgical decompression to prevent ischemia.

Squamous cell carcinoma

Seen in the US, but more common in Asia, Africa, South America. Precursor in situ lesions: Bowen disease (in penile shaft, presents as leukoplakia "white plaque"), erythroplasia of Queyrat (carcinoma in situ of the glans B, presents as erythroplakia "red plaque"). Bowenoid papulosis (carcinoma in situ of unclear malignant potential, presenting as reddish papules). Associated with uncircumcised males and HPV.

Cryptorchidism



Descent failure of one A or both testes; impaired spermatogenesis (since sperm develop best at temperatures < 37°C); can have normal testosterone levels (Leydig cells are mostly unaffected by temperature); associated with ↑ risk of germ cell tumors. Prematurity ↑ risk of cryptorchidism. ↓ inhibin B, ↑ FSH, ↑ LH; testosterone ↓ in bilateral cryptorchidism, normal in unilateral. Most cases resolve spontaneously; otherwise, orchiopexy performed before 2 years of age.

Testicular torsion

Rotation of testicle around spermatic cord and vascular pedicle. Commonly presents in males 12–18 years old. May occur after an inciting event (eg, trauma) or spontaneously. Characterized by acute, severe pain, high-riding testis, and absent cremasteric reflex.

Treatment: surgical correction (orchiopexy) within 6 hours, manual detorsion if surgical option unavailable in timeframe. If testis is not viable, orchiectomy. Orchiopexy, when performed, should be bilateral because the contralateral testis is at risk for subsequent torsion.

Varicocele



Dilated veins in pampiniform plexus due to ↑ venous pressure; most common cause of scrotal enlargement in adult males; most often on left side because of ↑ resistance to flow from left gonadal vein drainage into left renal vein; can cause infertility because of ↑ temperature; diagnosed by standing clinical exam/Valsalva maneuver (distension on inspection and "bag of worms" on palpation; augmented by Valsalva) or ultrasound A; does not transilluminate.

Treatment: consider surgical ligation or embolization if associated with pain or infertility.

(c) ketabton.com: The Digital Library

Extragonadal germ cell tumors Arise in midline locations. In adults, most commonly in retroperitoneum, mediastinum, pineal, and suprasellar regions. In infants and young children, sacrococcygeal teratomas are most common.

Scrotal masses

Benign scrotal lesions present as testicular masses that can be transilluminated (vs solid testicular tumors).

Congenital hydrocele

Common cause of scrotal swelling A in infants, due to incomplete obliteration of processus vaginalis. Most spontaneously resolve within 1 year.

Transilluminating swelling.

Acquired hydrocele Scrotal fluid collection usually 2° to infection, trauma, tumor. If bloody → hematocele.

Spermatocele Cyst due to dilated epididymal duct or rete testis.

Paratesticular fluctuant nodule.



Testicular tumors



Germ cell tumors account for ~ 95% of all testicular tumors. Arise from germ cells that produce sperm. Most often occur in young men. Risk factors: cryptorchidism, Klinefelter syndrome. Can present as a mixed germ cell tumor. Do not transilluminate. Usually not biopsied (risk of seeding scrotum), removed via radical orchiectomy.

Sex cord stromal tumors develop from embryonic sex cord (develops into Sertoli and Leydig cells of seminiferous tubules, theca and granulosa cells of follicle) derivatives. 5% of all testicular tumors. Mostly benign.

(c) ketabton.com: The Digital Library

Testicular tumors *(continued)*

Germ cell tumors		
Seminoma	Malignant	Painless, homogenous testicular enlargement. Most common testicular tumor. Analogous to ovarian dysgerminoma. Does not occur in infancy. Large cells in lobules with watery cytoplasm and "fried egg" appearance on histology, ↑ placental ALP (PALP). Highly radiosensitive. Late metastasis, excellent prognosis.
Teratoma	May be malignant	Unlike in females, **M**ature teratoma in adult **M**ales may be **M**alignant. Benign in children.
Embryonal carcinoma	Malignant	Painful, hemorrhagic mass with necrosis. Often glandular/papillary morphology. "Pure" embryonal carcinoma is rare; most commonly mixed with other tumor types. May present with metastases. May be associated with ↑ hCG and normal AFP levels when pure (↑ AFP when mixed). Worse prognosis than seminoma.
Yolk sac (endodermal sinus) tumor	Malignant, aggressive	Yellow, mucinous. Analogous to ovarian yolk sac tumor. Schiller-Duval bodies resemble primitive glomeruli. ↑ AFP is highly characteristic. Most common testicular tumor in boys < 3 years old.
Choriocarcinoma	Malignant	Disordered syncytiotrophoblastic and cytotrophoblastic elements. Hematogenous metastases to lungs and brain. ↑ h**C**G, may produce gynecomastia, symptoms of hyperthyroidism (α-subunit of hCG is identical to LH, FSH, TSH).
Non-germ cell tumors		
Sertoli cell tumor	Mostly benign	Androblastoma from sex cord stroma.
Leydig cell tumor	Mostly benign	Golden brown color; contains Reinke crystals (eosinophilic cytoplasmic inclusions). Produces androgens or estrogens → gynecomastia in men, precocious puberty in boys.
Testicular lymphoma	Malignant, aggressive	Most common testicular cancer in older men. Not a 1° cancer; arises from metastatic lymphoma to testes.

Hormone levels in germ cell tumors

	SEMINOMA	YOLK SAC TUMOR	CHORIOCARCINOMA	TERATOMA	EMBRYONAL CARCINOMA
PALP	↑	–	–	–	–
AFP	–	↑↑	–	–	–/↑ (when mixed)
β-hCG	–/↑	–/↑	↑↑	–	↑

(c) ketabton.com: The Digital Library

Epididymitis and orchitis

Most common causes:
- *C trachomatis* and *N gonorrhoeae* (young men)
- *E coli* and *Pseudomonas* (elderly, associated with UTI and BPH)
- Autoimmune (eg, granulomas involving seminiferous tubules)

Epididymitis

Inflammation of epididymis. Presents with localized pain and tenderness over posterior testis. ⊕ Prehn sign (pain relief with scrotal elevation). May progress to involve testis.

Orchitis

Inflammation of testis. Presents with testicular pain and swelling. Mumps orchitis ↑ infertility risk. Rare in boys < 10 years old.

Benign prostatic hyperplasia

Common in men > 50 years old. Characterized by smooth, elastic, firm nodular enlargement (hyperplasia not hypertrophy) of periurethral (lateral and middle) lobes, which compress the urethra into a vertical slit. Not premalignant.

Often presents with ↑ frequency of urination, nocturia, difficulty starting and stopping urine stream, dysuria. May lead to distention and hypertrophy of bladder, hydronephrosis, UTIs. ↑ free prostate-specific antigen (PSA).

Treatment: α_1-antagonists (terazosin, tamsulosin), which cause relaxation of smooth muscle; 5α-reductase inhibitors (eg, finasteride); PDE-5 inhibitors (eg, tadalafil); surgical resection (eg, TURP, ablation).



Prostatitis

Characterized by dysuria, frequency, urgency, low back pain. Warm, tender, enlarged prostate.

Acute bacterial prostatitis—in older men most common bacterium is *E coli*; in young men consider *C trachomatis*, *N gonorrhoeae*.

Chronic prostatitis—either bacterial or nonbacterial (eg, 2° to previous infection, nerve problems, chemical irritation).

Prostatic adenocarcinoma

Common in men > 50 years old. Arises most often from posterior lobe (peripheral zone) of prostate gland and is most frequently diagnosed by ↑ PSA and subsequent needle core biopsies. Prostatic acid phosphatase (PAP) and PSA are useful tumor markers (↑ total PSA, with ↓ fraction of free PSA). Osteoblastic metastases in bone may develop in late stages, as indicated by lower back pain and ↑ serum ALP and PSA. Metastasis to the spine often occurs via Batson (vertebral) venous plexus.

(c) ketabton.com: The Digital Library

▸ REPRODUCTIVE—PHARMACOLOGY

Control of reproductive hormones



(c) ketabton.com: The Digital Library

Goserelin, leuprolide

MECHANISM	GnRH analogs. When used in pulsatile fashion act as GnRH agonists. When used in continuous fashion first transiently act as GnRH agonists (tumor flare), but subsequently act as GnRH antagonists (downregulate GnRH receptor in pituitary → ↓ FSH and ↓ LH).	**Leu**prolide can be used in **lieu** of GnRH.
CLINICAL USE	Uterine fibroids, endometriosis, precocious puberty, prostate cancer, infertility.	
ADVERSE EFFECTS	Hypogonadism, ↓ libido, erectile dysfunction, nausea, vomiting.	

Degarelix

MECHANISM	GnRH antagonist. No start-up flare.
CLINICAL USE	Prostate cancer.
ADVERSE EFFECTS	Hot flashes, liver toxicity.

Estrogens

Ethinyl estradiol, DES, mestranol.

MECHANISM	Bind estrogen receptors.
CLINICAL USE	Hypogonadism or ovarian failure, menstrual abnormalities (combined OCPs), hormone replacement therapy in postmenopausal women.
ADVERSE EFFECTS	↑ risk of endometrial cancer (when given without progesterone), bleeding in postmenopausal women, clear cell adenocarcinoma of vagina in females exposed to DES in utero, ↑ risk of thrombi. Contraindications—ER ⊕ breast cancer, history of DVTs, tobacco use in women > 35 years old.

Selective estrogen receptor modulators

Clomiphene	Antagonist at estrogen receptors in hypothalamus. Prevents normal feedback inhibition and ↑ release of LH and FSH from pituitary, which stimulates ovulation. Used to treat infertility due to anovulation (eg, PCOS). May cause hot flashes, ovarian enlargement, multiple simultaneous pregnancies, visual disturbances.
Tamoxifen	Antagonist at breast; agonist at bone, uterus; ↑ risk of **t**hromboembolic events (especially with smoking) and **en**dometrial cancer. Used to treat and prevent recurrence of ER/PR ⊕ breast cancer.
Raloxifene	Antagonist at breast, uterus; agonist at bone; ↑ risk of thromboembolic events (especially with smoking) but no increased risk of endometrial cancer (vs tamoxifen); used primarily to treat osteoporosis.

Aromatase inhibitors

Anastrozole, letrozole, exemestane.

MECHANISM	Inhibit peripheral conversion of androgens to estrogen.
CLINICAL USE	ER ⊕ breast cancer in postmenopausal women.

(c) ketabton.com: The Digital Library

Hormone replacement therapy

Used for relief or prevention of menopausal symptoms (eg, hot flashes, vaginal atrophy), osteoporosis (↑ estrogen, ↓ osteoclast activity).
Unopposed estrogen replacement therapy ↑ risk of endometrial cancer, progesterone/progestin is added. Possible increased cardiovascular risk.

Progestins

Levonorgestrel, medroxyprogesterone, etonogestrel, norethindrone, megestrol.

MECHANISM	Bind progesterone receptors, ↓ growth and ↑ vascularization of endometrium, thicken cervical mucus.
CLINICAL USE	Contraception (forms include pill, intrauterine device, implant, depot injection), endometrial cancer, abnormal uterine bleeding. Progestin challenge: presence of withdrawal bleeding excludes anatomic defects (eg, Asherman syndrome) and chronic anovulation without estrogen.

Antiprogestins

Mifepristone, ulipristal.

MECHANISM	Competitive inhibitors of progestins at progesterone receptors.
CLINICAL USE	Termination of pregnancy (mifepristone with misoprostol); emergency contraception (ulipristal).

Combined contraception

Progestins and ethinyl estradiol; forms include pill, patch, vaginal ring.
Estrogen and progestins inhibit LH/FSH and thus prevent estrogen surge. No estrogen surge → no LH surge → no ovulation.
Progestins cause thickening of cervical mucus, thereby limiting access of sperm to uterus. Progestins also inhibit endometrial proliferation → endometrium is less suitable to the implantation of an embryo.
Adverse effects: breakthrough menstrual bleeding, breast tenderness, VTE, hepatic adenomas.
Contraindications: smokers > 35 years old (↑ risk of cardiovascular events), patients with ↑ risk of cardiovascular disease (including history of venous thromboembolism, coronary artery disease, stroke), migraine (especially with aura), breast cancer, liver disease.

Copper intrauterine device

MECHANISM	Produces local inflammatory reaction toxic to sperm and ova, preventing fertilization and implantation; hormone free.
CLINICAL USE	Long-acting reversible contraception. Most effective emergency contraception.
ADVERSE EFFECTS	Heavier or longer menses, dysmenorrhea. Risk of PID with insertion (contraindicated in active pelvic infection).

Tocolytics

Medications that relax the uterus; include terbutaline (β_2-agonist action), nifedipine (Ca^{2+} channel blocker), indomethacin (NSAID). Used to ↓ contraction frequency in preterm labor and allow time for administration of steroids (to promote fetal lung maturity) or transfer to appropriate medical center with obstetrical care.

(c) ketabton.com: The Digital Library

Danazol

MECHANISM	Synthetic androgen that acts as partial agonist at androgen receptors.
CLINICAL USE	Endometriosis, hereditary angioedema.
ADVERSE EFFECTS	Weight gain, edema, acne, hirsutism, masculinization, ↓ HDL levels, hepatotoxicity, idiopathic intracranial hypertension.

Testosterone, methyltestosterone

MECHANISM	Agonists at androgen receptors.
CLINICAL USE	Treat hypogonadism and promote development of 2° sex characteristics; stimulate anabolism to promote recovery after burn or injury.
ADVERSE EFFECTS	Masculinization in females; ↓ intratesticular testosterone in males by inhibiting release of LH (via negative feedback) → gonadal atrophy. Premature closure of epiphyseal plates. ↑ LDL, ↓ HDL.

Antiandrogens

Finasteride	5α-reductase inhibitor (↓ conversion of testosterone to DHT). Used for BPH and male-pattern baldness. Adverse effects: gynecomastia and sexual dysfunction.	Testosterone $\xrightarrow{\text{5α-reductase}}$ DHT (more potent).
Flutamide, bicalutamide, apalutamide, enzalutamide	Nonsteroidal competitive inhibitors at androgen receptors. Used for prostate carcinoma.	
Ketoconazole	Inhibits steroid synthesis (inhibits 17,20 desmolase/17α-hydroxylase).	Used in PCOS to reduce androgenic symptoms. Both can cause gynecomastia and amenorrhea.
Spironolactone	Inhibits steroid binding, 17,20 desmolase/17α-hydroxylase.	

Tamsulosin — α_1-antagonist used to treat BPH by inhibiting smooth muscle contraction. Selective for $\alpha_{1A/D}$ receptors (found on prostate) vs vascular α_{1B} receptors.

Minoxidil

MECHANISM	Direct arteriolar vasodilator.
CLINICAL USE	Androgenetic alopecia (pattern baldness), severe refractory hypertension.

(c) ketabton.com: The Digital Library

Respiratory

"There's so much pollution in the air now that if it weren't for our lungs, there'd be no place to put it all."

—Robert Orben

"Freedom is the oxygen of the soul."

—Moshe Dayan

"Whenever I feel blue, I start breathing again."

—L. Frank Baum

"Life is not the amount of breaths you take; it's the moments that take your breath away."

—Will Smith, *Hitch*

Group key respiratory, cardiovascular, and renal concepts together for study whenever possible. Know obstructive vs restrictive lung disorders, $\dot{V}/\dot{Q}$ mismatch, lung volumes, mechanics of respiration, and hemoglobin physiology. Lung cancers and other causes of lung masses are high yield. Be comfortable reading basic chest x-rays, CT scans, and PFTs.

- Embryology 660
- Anatomy 662
- Physiology 664
- Pathology 671
- Pharmacology 686

FAS1_2019_16-Respiratory.indd 659 11/8/19 7:33 AM

(c) ketabton.com: The Digital Library

▸ RESPIRATORY—EMBRYOLOGY

Lung development

Occurs in five stages. Initial development includes development of lung bud from distal end of respiratory diverticulum during week 4. **E**very **P**ulmonologist **C**an **S**ee **A**lveoli.

STAGE	STRUCTURAL DEVELOPMENT	NOTES
Embryonic (weeks 4–7)	Lung bud → trachea → bronchial buds → mainstem bronchi → secondary (lobar) bronchi → tertiary (segmental) bronchi.	Errors at this stage can lead to tracheoesophageal fistula.
Pseudoglandular (weeks 5–17)	Endodermal tubules → terminal bronchioles. Surrounded by modest capillary network.	Respiration impossible, incompatible with life.
Canalicular (weeks 16–25)	Terminal bronchioles → respiratory bronchioles → alveolar ducts. Surrounded by prominent capillary network.	Airways increase in diameter. Respiration capable at 25 weeks. Pneumocytes develop starting at 20 weeks.
Saccular (week 26–birth)	Alveolar ducts → terminal sacs. Terminal sacs separated by 1° septae.	
Alveolar (week 36–8 years)	Terminal sacs → adult alveoli (due to 2° septation). In utero, "breathing" occurs via aspiration and expulsion of amniotic fluid → ↑ vascular resistance through gestation. At birth, fluid gets replaced with air → ↓ in pulmonary vascular resistance.	At birth: 20–70 million alveoli. By 8 years: 300–400 million alveoli.



Congenital lung malformations

Pulmonary hypoplasia	Poorly developed bronchial tree with abnormal histology. Associated with congenital diaphragmatic hernia (usually left-sided), bilateral renal agenesis (Potter sequence).
Bronchogenic cysts	Caused by abnormal budding of the foregut and dilation of terminal or large bronchi. Discrete, round, sharply defined, fluid-filled densities on CXR (air-filled if infected). Generally asymptomatic but can drain poorly, causing airway compression and/or recurrent respiratory infections.

(c) ketabton.com: The Digital Library

Club cells — Nonciliated; low columnar/cuboidal with secretory granules. Located in bronchioles. Degrade toxins; secrete component of surfactant; act as reserve cells.

Alveolar cell types

Type I pneumocytes — Squamous. 97% of alveolar surfaces. Thinly line the alveoli (two black arrows in A) for optimal gas exchange.

Type II pneumocytes





Cuboidal and clustered A.

2 functions:

1. Serve as stem cell precursors for 2 cell types (type I and type II cells); proliferate during lung damage.
2. Secrete surfactant from lamellar bodies (arrowheads in B)

Surfactant— ↓ alveolar surface tension, ↓ alveolar collapse, ↓ lung recoil, and ↑ compliance.

Composed of multiple lecithins, mainly dipalmitoylphosphatidylcholine (DPPC). Synthesis begins ~week 20 of gestation and achieves mature levels ~week 35. Corticosteroids important for fetal surfactant synthesis and lung development.

$$\text{Collapsing pressure } (P) = \frac{2\,(\text{surface tension})}{\text{radius}}$$

Law of Laplace—Alveoli have ↑ tendency to collapse on expiration as radius ↓.

Alveolar macrophages — Phagocytose foreign materials; release cytokines and alveolar proteases. Hemosiderin-laden macrophages ("HF cells") may be found in the setting of pulmonary edema or alveolar hemorrhage.

Neonatal respiratory distress syndrome



Surfactant deficiency → ↑ surface tension → alveolar collapse ("ground-glass" appearance of lung fields) A.

Risk factors: prematurity, maternal diabetes (due to ↑ fetal insulin), C-section delivery (↓ release of fetal glucocorticoids; less stressful than vaginal delivery).

Treatment: maternal steroids before birth; exogenous surfactant for infant.

Therapeutic supplemental O_2 can result in Retinopathy of prematurity, Intraventricular hemorrhage, Bronchopulmonary dysplasia (RIB).

Screening tests for fetal lung maturity: lecithin-sphingomyelin (L/S) ratio in amniotic fluid (≥ 2 is healthy; < 1.5 predictive of NRDS), foam stability index, surfactant-albumin ratio.

Persistently low O_2 tension → risk of PDA.



(c) ketabton.com: The Digital Library

▸ RESPIRATORY—ANATOMY

Respiratory tree

Conducting zone

Large airways consist of nose, pharynx, larynx, trachea, and bronchi. Airway resistance highest in the large- to medium-sized bronchi. Small airways consist of bronchioles that further divide into terminal bronchioles (large numbers in parallel → least airway resistance).

Warms, humidifies, and filters air but does not participate in gas exchange → "anatomic dead space."

Cartilage and goblet cells extend to the end of bronchi.

Pseudostratified ciliated columnar cells primarily make up epithelium of bronchus and extend to beginning of terminal bronchioles, then transition to cuboidal cells. Clear mucus and debris from lungs (mucociliary escalator).

Airway smooth muscle cells extend to end of terminal bronchioles (sparse beyond this point).

Respiratory zone

Lung parenchyma; consists of respiratory bronchioles, alveolar ducts, and alveoli. Participates in gas exchange.

Mostly cuboidal cells in respiratory bronchioles, then simple squamous cells up to alveoli. Cilia terminate in respiratory bronchioles. Alveolar macrophages clear debris and participate in immune response.



(c) ketabton.com: The Digital Library

Lung anatomy



Right lung has 3 lobes; **L**eft has **L**ess **L**obes (2) and **L**ingula (homolog of right middle lobe). Instead of a middle lobe, left lung has a space occupied by the heart **A**.

Relation of the pulmonary artery to the bronchus at each lung hilum is described by **RALS**—**R**ight **A**nterior; **L**eft **S**uperior. Carina is posterior to ascending aorta and anteromedial to descending aorta **B**.

Right lung is a more common site for inhaled foreign bodies because right main stem bronchus is wider, more vertical, and shorter than the left. If you aspirate a peanut:

- While supine—usually enters superior segment of right lower lobe.
- While lying on right side—usually enters right upper lobe.
- While upright—usually enters right lower lobe.







Diaphragm structures



Structures perforating diaphragm:

- At T8: IVC, right phrenic nerve
- At T10: esophagus, vagus (CN 10; 2 trunks)
- At T12: aorta (red), thoracic duct (white), azygos vein (blue) ("At T-1-2 it's the red, white, and blue")

Diaphragm is innervated by C3, 4, and 5 (phrenic nerve). Pain from diaphragm irritation (eg, air, blood, or pus in peritoneal cavity) can be referred to shoulder (C5) and trapezius ridge (C3, 4).

Number of letters = T level:
T8: vena cava (IVC)
T10: (O)esophagus
T12: aortic hiatus

I ate (8) ten eggs at twelve.

C3, 4, 5 keeps the diaphragm alive.

Other bifurcations:

- The common carotid bifourcates at C4.
- The trachea bifourcates at T4.
- The abdominal aorta bifourcates at L4.

(c) ketabton.com: The Digital Library

▸ RESPIRATORY—PHYSIOLOGY

Lung volumes

Note: a **capacity** is a sum of ≥ 2 physiologic **volumes.**

Inspiratory reserve volume	Air that can still be breathed in after normal inspiration
Tidal volume	Air that moves into lung with each quiet inspiration, typically 500 mL
Expiratory reserve volume	Air that can still be breathed out after normal expiration
Residual volume	Air in lung after maximal expiration; RV and any lung capacity that includes RV cannot be measured by spirometry
Inspiratory capacity	IRV + TV Air that can be breathed in after normal exhalation
Functional residual capacity	RV + ERV Volume of gas in lungs after normal expiration
Vital capacity	TV + IRV + ERV Maximum volume of gas that can be expired after a maximal inspiration
Total lung capacity	IRV + TV + ERV + RV Volume of gas present in lungs after a maximal inspiration



Determination of physiologic dead space

$$V_D = V_T \times \frac{Paco_2 - Peco_2}{Paco_2}$$

V_D = physiologic dead space = anatomic dead space of conducting airways plus alveolar dead space; apex of healthy lung is largest contributor of alveolar dead space. Volume of inspired air that does not take part in gas exchange.

V_T = tidal volume.

$Paco_2$ = arterial Pco_2.

$Peco_2$ = expired air Pco_2.

Taco, Paco, Peco, Paco (refers to order of variables in equation)

Physiologic dead space—approximately equivalent to anatomic dead space in normal lungs. May be greater than anatomic dead space in lung diseases with $\dot{V}/\dot{Q}$ defects.

Ventilation

Minute ventilation	Total volume of gas entering lungs per minute $V_E = V_T \times RR$	Normal values: Respiratory rate (RR) = 12–20 breaths/min V_T = 500 mL/breath V_D = 150 mL/breath
Alveolar ventilation	Volume of gas that reaches alveoli each minute $V_A = (V_T - V_D) \times RR$	

(c) ketabton.com: The Digital Library

Lung and chest wall

Elastic recoil

Tendency for lungs to collapse inward and chest wall to spring outward.

At FRC, airway and alveolar pressures equal atmospheric pressure (called zero), and intrapleural pressure is negative (preventing atelectasis). The inward pull of the lung is balanced by the outward pull of the chest wall. System pressure is atmospheric. Pulmonary vascular resistance (PVR) is at a minimum.



Compliance

Change in lung volume for a change in pressure ($\Delta V/\Delta P$). Inversely proportional to wall stiffness and increased by surfactant.

- ↑ compliance = lung easier to fill (eg, emphysema, aging)
- ↓ compliance = lung harder to fill (eg, pulmonary fibrosis, pneumonia, ARDS, pulmonary edema)



Hysteresis

Lung inflation follows a different pressure-volume curve than lung deflation due to need to overcome surface tension forces in inflation.

Respiratory system changes in the elderly

Aging is associated with progressive ↓ in lung function. TLC remains the same.

INCREASED	DECREASED
Lung compliance (loss of elastic recoil)	Chest wall compliance (↑ chest wall stiffness)
RV	FVC and FEV_1
$\dot{V}/\dot{Q}$ mismatch	Respiratory muscle strength (can impair cough)
A-a gradient	Ventilatory response to hypoxia/hypercapnia

Hemoglobin



Normal adult hemoglobin (Hb) is composed of 4 polypeptide subunits (2 α and 2 β) and exists in 2 forms:

- Deoxygenated form has low affinity for O_2, thus promoting release/unloading of O_2.
- Oxygenated form has high affinity for O_2 (300×). Hb exhibits positive cooperativity and positive allostery.

Hemoglobin acts as buffer for H^+ ions.

Myoglobin is composed of a single polypeptide chain associated with one heme moiety. Higher affinity for oxygen than Hb.

(c) ketabton.com: The Digital Library

Oxygen content of blood

O_2 content = (1.34 × Hb × Sao_2) + (0.003 × Pao_2)
Hb = hemoglobin concentration; Sao_2 = arterial O_2 saturation
Pao_2 = partial pressure of O_2 in arterial blood

Normally 1 g Hb can bind 1.34 mL O_2; normal Hb amount in blood is 15 g/dL.
O_2 binding capacity ≈ 20 mL O_2/dL of blood.
With ↓ Hb there is ↓ O_2 content of arterial blood, but no change in O_2 saturation and Pao_2.
O_2 delivery to tissues = cardiac output × O_2 content of blood.

	Hb CONCENTRATION	% O_2 SAT OF Hb	DISSOLVED O_2 (Pao_2)	TOTAL O_2 CONTENT
CO poisoning	Normal	↓ (CO competes with O_2)	Normal	↓
Anemia	↓	Normal	Normal	↓
Polycythemia	↑	Normal	Normal	↑

Methemoglobin

Iron in Hb is normally in a reduced state (ferrous Fe^{2+}; "just the 2 of us").
Oxidized form of Hb (ferric, Fe^{3+}) does not bind O_2 as readily as Fe^{2+}, but has ↑ affinity for cyanide → tissue hypoxia from ↓ O_2 saturation and ↓ O_2 content.
Methemoglobinemia may present with cyanosis and chocolate-colored blood.
Nitrites (eg, from dietary intake or polluted/high-altitude water sources) and benzocaine cause poisoning by oxidizing Fe^{2+} to Fe^{3+}.
Methemoglobinemia can be treated with methylene blue and vitamin C.

Oxygen-hemoglobin dissociation curve

ODC has sigmoidal shape due to positive cooperativity (ie, tetrameric Hb molecule can bind 4 O_2 molecules and has higher affinity for each subsequent O_2 molecule bound). Myoglobin is monomeric and thus does not show positive cooperativity; curve lacks sigmoidal appearance.
Shifting ODC to the right → ↓ Hb affinity for O_2 (facilitates unloading of O_2 to tissue) → ↑ P_{50} (higher Po_2 required to maintain 50% saturation).
Shifting ODC to the left → ↓ O_2 unloading → renal hypoxia → ↑ EPO synthesis → compensatory erythrocytosis.
Fetal Hb (2 α and 2 γ subunits) has higher affinity for O_2 than adult Hb (due to ↓ affinity for 2,3-BPG) → dissociation curve is shifted left, driving diffusion of O_2 across the placenta from mother to fetus.



Left shift	Right shift
(↓ O_2 unloading to tissue)	(↑ O_2 unloading to tissues)
Left = Lower	ACE BATs right handed
↓ H^+ (↑ pH, base)	↑ H^+ (↓ pH, Acid)
↓ Pco_2	↑ Pco_2
↓ 2,3–BPG	Exercise
↓ Temperature	↑ 2,3–BPG
↑ CO	High Altitude
↑ MetHb	↑ Temperature
↑ HbF	

(c) ketabton.com: The Digital Library

Cyanide vs carbon monoxide poisoning

Both inhibit aerobic metabolism via inhibition of complex IV (cytochrome c oxidase) → hypoxia that does not fully correct with supplemental O_2 and ↑ anaerobic metabolism.
Both can lead to pink or cherry red skin (usually postmortem finding), seizures, and coma.

	Cyanide	Carbon monoxide
SOURCE	Byproduct of synthetic product combustion, ingestion of amygdalin (cyanogenic glucoside found in apricot seeds) or cyanide.	Odorless gas from fires, car exhaust, or gas heaters.
TREATMENT	Hydroxocobalamin (binds cyanide → cyanocobalamin → renal excretion). Nitrites (oxidize Hb → methemoglobin → binds cyanide → cyanomethemoglobin → less toxicity). Sodium thiosulfate (↑ cyanide conversion to thiocyanate → renal excretion).	100% O_2, hyperbaric O_2.
SIGNS/SYMPTOMS 	Breath has bitter almond odor; cardiovascular collapse.	Headache, dizziness. Multiple individuals may be involved (eg, family with similar symptoms in winter). Classically associated with bilateral globus pallidus lesions on MRI A, although rarely seen with cyanide toxicity as well.
EFFECT ON OXYGEN-HEMOGLOBIN DISSOCIATION CURVE	Curve normal; oxygen saturation may appear normal initially.	Left shift in curve → ↑ affinity for O_2 → ↓ O_2 unloading in tissues. Binds competitively to Hb with 200× greater affinity than O_2 to form carboxyhemoglobin → ↓ %O_2 saturation of Hb.



(c) ketabton.com: The Digital Library

Pulmonary circulation

Normally a low-resistance, high-compliance system. A ↓ in P_{AO_2} causes a hypoxic vasoconstriction that shifts blood away from poorly ventilated regions of lung to well-ventilated regions of lung.

Perfusion limited—O_2 (normal health), CO_2, N_2O. Gas equilibrates early along the length of the capillary. Exchange can be ↑ only if blood flow ↑.

Diffusion limited—O_2 (emphysema, fibrosis, exercise), CO. Gas does not equilibrate by the time blood reaches the end of the capillary.

A consequence of pulmonary hypertension is cor pulmonale and subsequent right ventricular failure.

Diffusion: $\dot{V}_{gas} = A \times D_k \times \frac{P_1 - P_2}{\Delta_x}$ where

A = area, Δ_x = alveolar wall thickness, D_k = diffusion coefficient of gas, $P_1 - P_2$ = difference in partial pressures.

- A ↓ in emphysema.
- Δ_x ↑ in pulmonary fibrosis.

DLCO is the extent to which CO passes from air sacs of lungs into blood.



Pa = partial pressure of gas in pulmonary capillary blood
PA = partial pressure of gas in alveolar air

Pulmonary vascular resistance

$$PVR = \frac{P_{pulm\ artery} - P_{L\ atrium}}{Q}$$

Remember: $\Delta P = Q \times R$, so $R = \Delta P / Q$

$$R = \frac{8\eta l}{\pi r^4}$$

$P_{pulm\ artery}$ = pressure in pulmonary artery
$P_{L\ atrium}$ ≈ pulmonary capillary wedge pressure
Q = cardiac output (flow)
R = resistance
η = viscosity of blood
l = vessel length
r = vessel radius

Alveolar gas equation

$$P_{AO_2} = PI_{O_2} - \frac{P_{aCO_2}}{R}$$

$$\approx 150\ mm\ Hg^a - \frac{P_{aCO_2}}{0.8}$$

[a]At sea level breathing room air

P_{AO_2} = alveolar P_{O_2} (mm Hg)
PI_{O_2} = P_{O_2} in inspired air (mm Hg)
P_{aCO_2} = arterial P_{CO_2} (mm Hg)
R = respiratory quotient = CO_2 produced/O_2 consumed
A-a gradient = $P_{AO_2} - P_{aO_2}$. Normal A-a gradient estimated as (age/4) + 4 (eg, for a person <40 years old, gradient should be <14).

(c) ketabton.com: The Digital Library

Oxygen deprivation

Hypoxia (↓ O_2 delivery to tissue)	Hypoxemia (↓ Pao_2)	Ischemia (loss of blood flow)
↓ cardiac output Hypoxemia Ischemia Anemia CO poisoning	Normal A-a gradient ▪ High altitude ▪ Hypoventilation (eg, opioid use, obesity hypoventilation syndrome) ↑ A-a gradient ▪ $\dot{V}/\dot{Q}$ mismatch ▪ Diffusion limitation (eg, fibrosis) ▪ Right-to-left shunt	Impeded arterial flow ↓ venous drainage

Ventilation/perfusion mismatch

Ideally, ventilation is matched to perfusion (ie, $\dot{V}/\dot{Q} = 1$) for adequate gas exchange.

Lung zones:
- $\dot{V}/\dot{Q}$ at apex of lung = 3 (wasted ventilation)
- $\dot{V}/\dot{Q}$ at base of lung = 0.6 (wasted perfusion)

Both ventilation and perfusion are greater at the base of the lung than at the apex of the lung.

With exercise (↑ cardiac output), there is vasodilation of apical capillaries → $\dot{V}/\dot{Q}$ ratio approaches 1.

Certain organisms that thrive in high O_2 (eg, TB) flourish in the apex.

$\dot{V}/\dot{Q} = 0$ = "oirway" obstruction (shunt). In shunt, 100% O_2 does not improve Pao_2 (eg, foreign body aspiration).

$\dot{V}/\dot{Q} = \infty$ = blood flow obstruction (physiologic dead space). Assuming < 100% dead space, 100% O_2 improves Pao_2 (eg, pulmonary embolus).



(c) ketabton.com: The Digital Library

Carbon dioxide transport

CO_2 is transported from tissues to lungs in 3 forms:

1. HCO_3^- (70%).
2. Carbaminohemoglobin or $HbCO_2$ (21–25%). CO_2 bound to Hb at N-terminus of globin (not heme). CO_2 favors deoxygenated form (O_2 unloaded).
3. Dissolved CO_2 (5–9%).

In lungs, oxygenation of Hb promotes dissociation of H^+ from Hb. This shifts equilibrium toward CO_2 formation; therefore, CO_2 is released from RBCs (Haldane effect).

In peripheral tissue, ↑ H^+ from tissue metabolism shifts curve to right, unloading O_2 (Bohr effect).

Majority of blood CO_2 is carried as HCO_3^- in the plasma.



Response to high altitude

↓ atmospheric oxygen (PiO_2) → ↓ Pao_2 → ↑ ventilation → ↓ $Paco_2$ → respiratory alkalosis → altitude sickness.

Chronic ↑ in ventilation.

↑ erythropoietin → ↑ Hct and Hb (due to chronic hypoxia).

↑ 2,3-BPG (binds to Hb causing rightward shift of the ODC so that Hb releases more O_2).

Cellular changes (↑ mitochondria).

↑ renal excretion of HCO_3^- to compensate for respiratory alkalosis (can augment with acetazolamide).

Chronic hypoxic pulmonary vasoconstriction results in pulmonary hypertension and RVH.

Response to exercise

↑ CO_2 production.

↑ O_2 consumption.

Right shift of ODC.

↑ ventilation rate to meet O_2 demand.

$\dot{V}/\dot{Q}$ ratio from apex to base becomes more uniform.

↑ pulmonary blood flow due to ↑ cardiac output.

↓ pH during strenuous exercise (2° to lactic acidosis).

No change in Pao_2 and $Paco_2$, but ↑ in venous CO_2 content and ↓ in venous O_2 content.

(c) ketabton.com: The Digital Library

▸ RESPIRATORY—PATHOLOGY

Rhinosinusitis



Obstruction of sinus drainage into nasal cavity → inflammation and pain over affected area. Typically affects maxillary sinuses, which drain against gravity due to ostia located superomedially (red arrow points to fluid-filled right maxillary sinus in A).

Superior meatus—drains sphenoid, posterior ethmoid; middle meatus—drains frontal, maxillary, and anterior ethmoid; inferior meatus—drains nasolacrimal duct.

Most common acute cause is viral URI; may lead to superimposed bacterial infection, most commonly *H influenzae*, *S pneumoniae*, *M catarrhalis*.

Paranasal sinus infections may extend to the orbits, cavernous sinus, and brain, causing complications (eg, orbital cellulitis, cavernous sinus syndrome, meningitis).

Epistaxis

Nose bleed. Most commonly occurs in anterior segment of nostril (**Kiesselbach** plexus). Life-threatening hemorrhages occur in posterior segment (sphenopalatine artery, a branch of maxillary artery). Common causes include foreign body, trauma, allergic rhinitis, and nasal angiofibromas (common in adolescent males).

Kiesselbach drives his **L**exus with his **LEGS**: superior **L**abial artery, anterior and posterior **E**thmoidal arteries, **G**reater palatine artery, **S**phenopalatine artery.

Head and neck cancer

Mostly squamous cell carcinoma. Risk factors include tobacco, alcohol, HPV-16 (oropharyngeal), EBV (nasopharyngeal). Field cancerization: carcinogen damages wide mucosal area → multiple tumors that develop independently after exposure.

Deep venous thrombosis

Blood clot within a deep vein → swelling, redness A, warmth, pain. Predisposed by Virchow triad (**SHE**):

- **S**tasis (eg, post-op, long drive/flight)
- **H**ypercoagulability (eg, defect in coagulation cascade proteins, such as factor V Leiden; oral contraceptive use; pregnancy)
- **E**ndothelial damage (exposed collagen triggers clotting cascade)

Most pulmonary emboli arise from proximal deep veins of lower extremity.

D-dimer lab test used clinically to rule out DVT in low-to-moderate risk patients (high sensitivity, low specificity).

Imaging test of choice is compression ultrasound with Doppler.

Use unfractionated heparin or low-molecular weight heparins (eg, enoxaparin) for prophylaxis and acute management.

Use oral anticoagulants (eg, rivaroxaban, apixaban) for treatment and long-term prevention.

(c) ketabton.com: The Digital Library

Pulmonary emboli

V̇/Q̇ mismatch, hypoxemia, respiratory alkalosis. Sudden-onset dyspnea, pleuritic chest pain, tachypnea, tachycardia. Large emboli or saddle embolus A may cause sudden death due to electromechanical dissociation (pulseless electrical activity). CT pulmonary angiography is imaging test of choice for PE (look for filling defects) B. May have S1Q3T3 abnormality on ECG.

Lines of Zahn C are interdigitating areas of pink (platelets, fibrin) and red (RBCs) found only in thrombi formed before death; help distinguish pre- and postmortem thrombi.

Types: **F**at, **A**ir, **T**hrombus, **B**acteria, **A**mniotic fluid, **T**umor. An embolus moves like a **FAT BAT**.

Fat emboli—associated with long bone fractures and liposuction; classic triad of hypoxemia, neurologic abnormalities, petechial rash.

Air emboli—nitrogen bubbles precipitate in ascending divers (caisson disease/decompression sickness); treat with hyperbaric O_2; or, can be iatrogenic 2° to invasive procedures (eg, central line placement).

Amniotic fluid emboli—typically occurs during labor or postpartum, but can be due to uterine trauma. Can lead to DIC. Rare, but high mortality.



Mediastinal pathology

Normal mediastinum contains heart, thymus, lymph nodes, esophagus, and aorta.

Mediastinal masses

Some pathologies (eg, lymphoma, lung cancer, abscess) can occur in any compartment, but there are common associations:

- Anterior—**4T**'s: **T**hyroid (substernal goiter), **T**hymic neoplasm, **T**eratoma, "**T**errible" lymphoma.
- Middle—esophageal carcinoma, metastases, hiatal hernia, bronchogenic cysts.
- Posterior—neurogenic tumor (eg, neurofibroma), multiple myeloma.

Mediastinitis

Inflammation of mediastinal tissues. Commonly due to postoperative complications of cardiothoracic procedures (≤ 14 days), esophageal perforation, or contiguous spread of odontogenic/retropharyngeal infection.

Chronic mediastinitis—also known as fibrosing mediastinitis; due to ↑ proliferation of connective tissue in mediastinum. *Histoplasma capsulatum* is common cause.

Clinical features: fever, tachycardia, leukocytosis, chest pain, and sternal wound drainage.

Pneumomediastinum



Presence of gas (usually air) in the mediastinum (black arrows show air around the aorta, red arrow shows air dissecting into the neck A). Can either be spontaneous (due to rupture of pulmonary bleb) or 2° (eg, trauma, iatrogenic, Boerhaave syndrome).

Ruptured alveoli allow tracking of air into the mediastinum via peribronchial and perivascular sheaths.

Clinical features: chest pain, dyspnea, voice change, subcutaneous emphysema, ⊕ Hamman sign (crepitus on cardiac auscultation).

(c) ketabton.com: The Digital Library

Flow-volume loops

FLOW-VOLUME PARAMETER	Obstructive lung disease	Restrictive lung disease
RV	↑	↓
FRC	↑	↓
TLC	↑	↓
FEV_1	↓↓	↓
FVC	↓	↓
FEV_1/FVC	↓ FEV_1 decreased more than FVC	Normal or ↑ FEV_1 decreased proportionately to FVC



(c) ketabton.com: The Digital Library

Obstructive lung diseases

Obstruction of air flow → air trapping in lungs. Airways close prematurely at high lung volumes → ↑ **FRC**, ↑ **RV**, ↑ **TLC**. PFTs: ↓↓ FEV_1, ↓ FVC → ↓ FEV_1/FVC ratio (hallmark), $\dot{V}/\dot{Q}$ mismatch. Chronic hypoxic pulmonary vasoconstriction can lead to cor pulmonale. Chronic obstructive pulmonary disease (**COPD**) includes chronic bronchitis and emphysema. "**FR**i**C**kin' **RV** needs some increased **TLC**, but it's hard with **COPD**!"

TYPE	PRESENTATION	PATHOLOGY	OTHER
Chronic bronchitis ("blue bloater")	Findings: wheezing, crackles, cyanosis (hypoxemia due to shunting), dyspnea, CO_2 retention, 2° polycythemia.	Hypertrophy and hyperplasia of mucus-secreting glands in bronchi → Reid index (thickness of mucosal gland layer to thickness of wall between epithelium and cartilage) > 50%. DLCO usually normal.	Diagnostic criteria: productive cough for ≥ 3 months in a year for > 2 consecutive years.
Emphysema ("pink puffer") 	Findings: barrel-shaped chest D, exhalation through pursed lips (increases airway pressure and prevents airway collapse).	Centriacinar—affects respiratory bronchioles while sparing distal alveoli, associated with **smoking** A B. Frequently in **up**per lobes (**smoke** rises **up**). Panacinar—affects respiratory bronchioles and alveoli, associated with α_1-antitrypsin deficiency. Frequently in lower lobes. Enlargement of air spaces ↓ recoil, ↑ compliance, ↓ DLCO from destruction of alveolar walls (arrow in C) and ↓ blood volume in pulmonary capillaries. Imbalance of proteases and antiproteases → ↑ elastase activity → ↑ loss of elastic fibers → ↑ lung compliance.	CXR: ↑ AP diameter, flattened diaphragm, ↑ lung field lucency.
Asthma	Findings: cough, wheezing, tachypnea, dyspnea, hypoxemia, ↓ inspiratory/expiratory ratio, pulsus paradoxus, mucus plugging E. Triggers: viral URIs, allergens, stress.	Hyperresponsive bronchi → reversible bronchoconstriction. Smooth muscle hypertrophy and hyperplasia, Curschmann spirals F (shed epithelium forms whorled mucous plugs), and Charcot-Leyden crystals G (eosinophilic, hexagonal, double-pointed crystals formed from breakdown of eosinophils in sputum). DLCO normal or ↑.	Type I hypersensitivity reaction. Diagnosis supported by spirometry and methacholine challenge. NSAID-exacerbated respiratory disease is a combination of COX inhibition (leukotriene overproduction → airway constriction), chronic sinusitis with nasal polyps, and asthma symptoms.

(c) ketabton.com: The Digital Library

Obstructive lung diseases *(continued)*

TYPE	PRESENTATION	PATHOLOGY	OTHER
Bronchiectasis	Findings: purulent sputum, recurrent infections (most often *P aeruginosa*), hemoptysis, digital clubbing.	Chronic necrotizing infection of bronchi or obstruction → permanently dilated airways.	Associated with bronchial obstruction, poor ciliary motility (eg, smoking, Kartagener syndrome), cystic fibrosis H, allergic bronchopulmonary aspergillosis.



Restrictive lung diseases



Restricted lung expansion causes ↓ lung volumes (↓ FVC and TLC). PFTs: ↑ FEV_1/FVC ratio. Patient presents with short, shallow breaths.

Types:

- Poor breathing mechanics (extrapulmonary, normal D_{LCO}, normal A-a gradient):
  - Poor muscular effort—polio, myasthenia gravis, Guillain-Barré syndrome
  - Poor structural apparatus—scoliosis, morbid obesity
- Interstitial lung diseases (pulmonary, ↓ D_{LCO}, ↑ A-a gradient):
  - Pneumoconioses (eg, coal workers' pneumoconiosis, silicosis, asbestosis)
  - Sarcoidosis: bilateral hilar lymphadenopathy, noncaseating granulomas; ↑ ACE and Ca^{2+}
  - Idiopathic pulmonary fibrosis (repeated cycles of lung injury and wound healing with ↑ collagen deposition, "honeycomb" lung appearance [red arrows in A], traction bronchiectasis [blue arrow in A] and digital clubbing).
  - Granulomatosis with polyangiitis (Wegener)
  - Pulmonary Langerhans cell histiocytosis (eosinophilic granuloma)
  - Hypersensitivity pneumonitis
  - Drug toxicity (eg, bleomycin, busulfan, amiodarone, methotrexate)

Hypersensitivity pneumonitis—mixed type III/IV hypersensitivity reaction to environmental antigen. Causes dyspnea, cough, chest tightness, fever, headache. Often seen in farmers and those exposed to birds. Reversible in early stages if stimulus is avoided.

(c) ketabton.com: The Digital Library

Sarcoidosis

Characterized by immune-mediated, widespread noncaseating granulomas A, elevated serum ACE levels, and elevated CD4/CD8 ratio in bronchoalveolar lavage fluid. More common in African-American females. Often asymptomatic except for enlarged lymph nodes. CXR shows bilateral adenopathy and coarse reticular opacities B; CT of the chest better demonstrates the extensive hilar and mediastinal adenopathy C.

Associated with **Bell palsy**, **U**veitis, **G**ranulomas (noncaseating epithelioid, containing microscopic Schaumann and asteroid bodies), **L**upus pernio (skin lesions on face resembling lupus), **I**nterstitial fibrosis (restrictive lung disease), **E**rythema nodosum, **R**heumatoid arthritis-like arthropathy, hypercalcemia (due to ↑ 1α-hydroxylase–mediated vitamin D activation in macrophages). A **facial droop** is **UGLIER**.

Treatment: steroids (if symptomatic).

Inhalation injury and sequelae

Complication of inhalation of noxious stimuli (eg, smoke). Caused by heat, particulates (< 1 μm diameter), or irritants (eg, NH_3) → chemical tracheobronchitis, edema, pneumonia, ARDS. Many patients present 2° to burns, CO inhalation, cyanide poisoning, or arsenic poisoning. Singed nasal hairs or soot in oropharynx common on exam.

Bronchoscopy shows severe edema, congestion of bronchus, and soot deposition (A, 18 hours after inhalation injury; B, resolution at 11 days after injury).



(c) ketabton.com: The Digital Library

Pneumoconioses

Asbestos is from the roof (was common in insulation), but affects the base (lower lobes).
Silica and coal are from the base (earth), but affect the roof (upper lobes).

Asbestosis

Associated with shipbuilding, roofing, plumbing. "Ivory white," calcified, supradiaphragmatic A and pleural B plaques are pathognomonic of asbestosis.
Risk of bronchogenic carcinoma > risk of mesothelioma. ↑ risk of Caplan syndrome (rheumatoid arthritis and pneumoconioses with intrapulmonary nodules).

Affects lower lobes.
Asbestos (ferruginous) bodies are golden-brown fusiform rods resembling dumbbells C, found in alveolar sputum sample, visualized using Prussian blue stain, often obtained by bronchoalveolar lavage.
↑ risk of pleural effusions.

Berylliosis

Associated with exposure to beryllium in aerospace and manufacturing industries. Granulomatous (noncaseating) D on histology and therefore occasionally responsive to steroids. ↑ risk of cancer and cor pulmonale.

Affects upper lobes.

Coal workers' pneumoconiosis

Prolonged coal dust exposure → macrophages laden with carbon → inflammation and fibrosis.
Also known as black lung disease. ↑ risk of Caplan syndrome.

Affects upper lobes.
Small, rounded nodular opacities seen on imaging.
Anthracosis—asymptomatic condition found in many urban dwellers exposed to sooty air.

Silicosis

Associated with sandblasting, foundries, mines. Macrophages respond to silica and release fibrogenic factors, leading to fibrosis. It is thought that silica may disrupt phagolysosomes and impair macrophages, increasing susceptibility to TB. ↑ risk of cancer, cor pulmonale, and Caplan syndrome.

Affects upper lobes.
"Eggshell" calcification of hilar lymph nodes on CXR.
The silly egg sandwich I found is mine!



(c) ketabton.com: The Digital Library

Mesothelioma



Malignancy of the pleura associated with asbestosis. May result in hemorrhagic pleural effusion (exudative), pleural thickening A.

Psammoma bodies seen on histology.
Calretinin and cytokeratin 5/6 ⊕ in almost all mesotheliomas, ⊖ in most carcinomas.
Smoking not a risk factor.

Acute respiratory distress syndrome

PATHOPHYSIOLOGY	Alveolar insult → release of pro-inflammatory cytokines → neutrophil recruitment, activation, and release of toxic mediators (eg, reactive oxygen species, proteases, etc) → capillary endothelial damage and ↑ vessel permeability → leakage of protein-rich fluid into alveoli → formation of intra-alveolar hyaline membranes (arrows in A) and noncardiogenic pulmonary edema (normal PCWP). Loss of surfactant also contributes to alveolar collapse.
CAUSES	Sepsis (most common), aspiration, pneumonia, trauma, pancreatitis.
DIAGNOSIS	Diagnosis of exclusion with the following criteria (**ARDS**): ▪ **A**bnormal chest X-ray (bilateral lung opacities) B ▪ **R**espiratory failure within 1 week of alveolar insult ▪ **D**ecreased Pao_2/Fio_2 (ratio < 300, hypoxemia due to ↑ intrapulmonary shunting and diffusion abnormalities) ▪ **S**ymptoms of respiratory failure are not due to HF/fluid overload
CONSEQUENCES	Impaired gas exchange, ↓ lung compliance; pulmonary hypertension.
MANAGEMENT	Treat the underlying cause. Mechanical ventilation: ↓ tidal volume, ↑ PEEP.





(c) ketabton.com: The Digital Library

Sleep apnea	Repeated cessation of breathing > 10 seconds during sleep → disrupted sleep → daytime somnolence. Diagnosis confirmed by sleep study. Nocturnal hypoxia → systemic/pulmonary hypertension, arrhythmias (atrial fibrillation/flutter), sudden death. Hypoxia → ↑ EPO release → ↑ erythropoiesis.
Obstructive sleep apnea	Respiratory effort against airway obstruction. Normal Pao_2 during the day. Associated with obesity, loud snoring, daytime sleepiness. Caused by excess parapharyngeal tissue in adults, adenotonsillar hypertrophy in children. Treatment: weight loss, CPAP, dental devices.
Central sleep apnea	Impaired respiratory effort due to CNS injury/toxicity, HF, opioids. May be associated with Cheyne-Stokes respirations (oscillations between apnea and hyperpnea). Think 3 C's: Congestive HF, CNS toxicity, Cheyne-Stokes respirations. Treat with positive airway pressure.
Obesity hypoventilation syndrome	Obesity (BMI ≥ 30 kg/m^2) → hypoventilation → ↑ $Paco_2$ during waking hours (retention); ↓ Pao_2 and ↑ $Paco_2$ during sleep. Also known as Pickwickian syndrome.

Pulmonary hypertension	Normal mean pulmonary artery pressure = 10–14 mm Hg; pulmonary hypertension ≥ 25 mm Hg at rest. Results in arteriosclerosis, medial hypertrophy, intimal fibrosis of pulmonary arteries, plexiform lesions. Course: severe respiratory distress → cyanosis and RVH → death from decompensated cor pulmonale.
ETIOLOGIES	
Pulmonary arterial hypertension	Often idiopathic. Heritable PAH can be due to an inactivating mutation in *BMPR2* gene (normally inhibits vascular smooth muscle proliferation); poor prognosis. Pulmonary vasculature endothelial dysfunction results in ↑ vasoconstrictors (eg, endothelin) and ↓ vasodilators (eg, NO and prostacyclins). Other causes include drugs (eg, amphetamines, cocaine), connective tissue disease, HIV infection, portal hypertension, congenital heart disease, schistosomiasis.
Left heart disease	Causes include systolic/diastolic dysfunction and valvular disease.
Lung diseases or hypoxia	Destruction of lung parenchyma (eg, COPD), lung inflammation/fibrosis (eg, interstitial lung diseases), hypoxemic vasoconstriction (eg, obstructive sleep apnea, living in high altitude).
Chronic thromboembolic	Recurrent microthrombi → ↓ cross-sectional area of pulmonary vascular bed.
Multifactorial	Causes include hematologic, systemic, and metabolic disorders, along with compression of the pulmonary vasculature by a tumor.

(c) ketabton.com: The Digital Library

Physical findings in select lung diseases

ABNORMALITY	BREATH SOUNDS	PERCUSSION	FREMITUS	TRACHEAL DEVIATION
Pleural effusion	↓	Dull	↓	None if small Away from side of lesion if large
Atelectasis	↓	Dull	↓	Toward side of lesion
Simple pneumothorax	↓	Hyperresonant	↓	None
Tension pneumothorax	↓	Hyperresonant	↓	Away from side of lesion
Consolidation (lobar pneumonia, pulmonary edema)	Bronchial breath sounds; late inspiratory crackles, egophony, whispered pectoriloquy	Dull	↑	None

Atelectasis



Alveolar collapse (right upper lobe collapse against mediastinum in A). Multiple causes:

- Obstructive—airway obstruction prevents new air from reaching distal airways, old air is resorbed (eg, foreign body, mucous plug, tumor)
- Compressive—external compression on lung decreases lung volumes (eg, space-occupying lesion, pleural effusion)
- Contraction (cicatrization)—scarring of lung parenchyma that distorts alveoli (eg, sarcoidosis)
- Adhesive—due to lack of surfactant (eg, NRDS in premature babies)



(c) ketabton.com: The Digital Library

Pleural effusions

Excess accumulation of fluid A between pleural layers → restricted lung expansion during inspiration. Can be treated with thoracentesis to remove/reduce fluid B.

Lymphatic

Also known as chylothorax. Due to thoracic duct injury from trauma or malignancy. Milky-appearing fluid; ↑ triglycerides.

Exudate

↑ protein content (> 2.9 g/dL), cloudy (cellular). Due to malignancy, inflammation/infection (eg, pneumonia, collagen vascular disease), trauma (occurs in states of ↑ vascular permeability). Must be drained due to risk of infection.

Transudate

↓ protein content (< 2.5 g/dL), clear (hypocellular). Due to ↑ hydrostatic pressure (eg, HF, Na^+ retention) or ↓ oncotic pressure (eg, nephrotic syndrome, cirrhosis).





(c) ketabton.com: The Digital Library

Pneumothorax	Accumulation of air in pleural space A. Dyspnea, uneven chest expansion. Chest pain, ↓ tactile fremitus, hyperresonance, and diminished breath sounds, all on the affected side.
Primary spontaneous pneumothorax	Due to rupture of apical subpleural bleb or cysts. Occurs most frequently in tall, thin, young males and smokers.
Secondary spontaneous pneumothorax	Due to diseased lung (eg, bullae in emphysema, infections), mechanical ventilation with use of high pressures → barotrauma.
Traumatic pneumothorax	Caused by blunt (eg, rib fracture), penetrating (eg, gunshot), or iatrogenic (eg, central line placement, lung biopsy, barotrauma due to mechanical ventilation) trauma.
Tension pneumothorax	Can be from any of the above. Air enters pleural space but cannot exit. Increasing trapped air → tension pneumothorax. Trachea deviates away from affected lung B. May lead to increased intrathoracic pressure → mediastinal displacement → kinking of IVC → ↓ venous return → ↓ cardiac output. Needs immediate needle decompression and chest tube placement.



(c) ketabton.com: The Digital Library

Pneumonia

TYPE	TYPICAL ORGANISMS	CHARACTERISTICS
Lobar pneumonia	*S pneumoniae* most frequently, also *Legionella*, *Klebsiella*	Intra-alveolar exudate → consolidation A; may involve entire lobe B or the whole lung.
Bronchopneumonia	*S pneumoniae*, *S aureus*, *H influenzae*, *Klebsiella*	Acute inflammatory infiltrates C from bronchioles into adjacent alveoli; patchy distribution involving ≥ 1 lobe D.
Interstitial (atypical) pneumonia	*Mycoplasma*, *Chlamydophila pneumoniae*, *Chlamydophila psittaci*, *Legionella*, viruses (RSV, CMV, influenza, adenovirus)	Diffuse patchy inflammation localized to interstitial areas at alveolar walls; CXR shows bilateral multifocal opacities E. Generally follows a more indolent course ("walking" pneumonia).
Cryptogenic organizing pneumonia	Etiology unknown. Secondary organizing pneumonia is caused by chronic inflammatory diseases (eg, rheumatoid arthritis) or medication side effects (eg, amiodarone). ⊖ sputum and blood cultures, often responds to steroids but not to antibiotics.	Formerly known as bronchiolitis obliterans organizing pneumonia (BOOP). Noninfectious pneumonia characterized by inflammation of bronchioles and surrounding structure.



Natural history of lobar pneumonia

	Congestion	Red hepatization	Gray hepatization	Resolution
DAYS	1–2	3–4	5–7	8+
FINDINGS	Red-purple, partial consolidation of parenchyma Exudate with mostly bacteria	Red-brown consolidation Exudate with fibrin, bacteria, RBCs, WBCs Reversible	Uniformly gray Exudate full of WBCs, lysed RBCs, and fibrin	Enzymatic digestion of exudate by macrophages

(c) ketabton.com: The Digital Library

Lung cancer

Leading cause of cancer death.

Presentation: cough, hemoptysis, bronchial obstruction, wheezing, pneumonic "coin" lesion on CXR or noncalcified nodule on CT.

Sites of metastases from lung cancer: **L**iver (jaundice, hepatomegaly), **A**drenals, **Bone** (pathologic fracture), **Brain**; "Lung 'mets' **L**ove **A**ffective **Bone**heads and **Brain**iacs."

In the lung, metastases (usually multiple lesions) are more common than 1° neoplasms. Most often from breast, colon, prostate, and bladder cancer.

SPHERE of complications:
- **S**uperior vena cava/thoracic outlet syndromes
- **P**ancoast tumor
- **H**orner syndrome
- **E**ndocrine (paraneoplastic)
- **R**ecurrent laryngeal nerve compression (hoarseness)
- **E**ffusions (pleural or pericardial)

Risk factors include smoking, secondhand smoke, radon, asbestos, family history.

Squamous and **S**mall cell carcinomas are **S**entral (central) and often caused by **S**moking.

TYPE	LOCATION	CHARACTERISTICS	HISTOLOGY
Small cell			
Small cell (oat cell) carcinoma	Central	Undifferentiated → very aggressive. May produce **A**CTH (Cushing syndrome), **A**DH (SIADH), or **A**ntibodies against presynaptic Ca^{2+} channels (Lambert-Eaton myasthenic syndrome) or neurons (paraneoplastic myelitis, encephalitis, subacute cerebellar degeneration). **A**mplification of *myc* oncogenes common. Managed with chemotherapy +/– radiation.	Neoplasm of neuroendocrine Kulchitsky cells → small dark blue cells **A**. Chromogranin **A** ⊕, neuron-specific enolase ⊕, synaptophysin ⊕.
Non-small cell			
Adenocarcinoma	Peripheral	Most common 1° lung cancer. More common in women than men, most likely to arise in nonsmokers. Activating mutations include *KRAS*, *EGFR*, and *ALK*. Associated with hypertrophic osteoarthropathy (clubbing). Bronchioloalveolar subtype (adenocarcinoma in situ): CXR often shows hazy infiltrates similar to pneumonia; better prognosis.	Glandular pattern on histology, often stains mucin ⊕ **B**. Bronchioloalveolar subtype: grows along alveolar septa → apparent "thickening" of alveolar walls. Tall, columnar cells containing mucus.
Squamous cell carcinoma	**C**entral	Hilar mass **C** arising from bronchus; **C**avitation; **C**igarettes; hyper**C**alcemia (produces PTHrP).	Keratin pearls **D** and intercellular bridges.
Large cell carcinoma	Peripheral	Highly anaplastic undifferentiated tumor; poor prognosis. Less responsive to chemotherapy; removed surgically. Strong association with smoking.	Pleomorphic giant cells **E**.
Bronchial carcinoid tumor	Central or peripheral	Excellent prognosis; metastasis rare. Symptoms due to mass effect or carcinoid syndrome (flushing, diarrhea, wheezing).	Nests of neuroendocrine cells; chromogranin A ⊕.



(c) ketabton.com: The Digital Library

Lung abscess





Localized collection of pus within parenchyma A. Caused by aspiration of oropharyngeal contents (especially in patients predisposed to loss of consciousness [eg, alcoholics, epileptics]) or bronchial obstruction (eg, cancer).

Air-fluid levels B often seen on CXR; presence suggests cavitation. Due to anaerobes (eg, *Bacteroides*, *Fusobacterium*, *Peptostreptococcus*) or *S aureus*.

Treatment: antibiotics, drainage, or surgery.

Lung abscess 2° to aspiration is most often found in right lung. Location depends on patient's position during aspiration: RLL if upright, RUL or RML if recumbent.

Pancoast tumor



Also known as superior sulcus tumor. Carcinoma that occurs in the apex of lung A may cause Pancoast syndrome by invading/compressing local structures.

Compression of locoregional structures may cause array of findings:

- Recurrent laryngeal nerve → hoarseness
- Stellate ganglion → Horner syndrome (ipsilateral ptosis, miosis, anhidrosis)
- Superior vena cava → SVC syndrome
- Brachiocephalic vein → brachiocephalic syndrome (unilateral symptoms)
- Brachial plexus → sensorimotor deficits
- Phrenic nerve → hemidiaphragm paralysis (hemidiaphragm elevation on CXR)

Superior vena cava syndrome



An obstruction of the SVC that impairs blood drainage from the head ("facial plethora"; note blanching after fingertip pressure in A), neck (jugular venous distention), and upper extremities (edema). Commonly caused by malignancy (eg, mediastinal mass, Pancoast tumor) and thrombosis from indwelling catheters B. Medical emergency. Can raise intracranial pressure (if obstruction is severe) → headaches, dizziness, ↑ risk of aneurysm/rupture of intracranial arteries.



(c) ketabton.com: The Digital Library

▸ RESPIRATORY—PHARMACOLOGY

Histamine-1 blockers

Reversible inhibitors of H_1 histamine receptors.

First generation	Diphenhydramine, dimenhydrinate, chlorpheniramine, doxylamine.	Names usually contain "-en/-ine" or "-en/-ate."
CLINICAL USE	Allergy, motion sickness, sleep aid.	
ADVERSE EFFECTS	Sedation, antimuscarinic, anti-α-adrenergic.	
Second generation	Loratadine, fexofenadine, desloratadine, cetirizine.	Names usually end in "-adine."
CLINICAL USE	Allergy.	
ADVERSE EFFECTS	Far less sedating than 1st generation because of ↓ entry into CNS.	

Guaifenesin

Expectorant—thins respiratory secretions; does not suppress cough reflex.

N-acetylcysteine

Mucolytic—liquifies mucus in chronic bronchopulmonary diseases (eg, COPD, CF) by disrupting disulfide bonds. Also used as an antidote for acetaminophen overdose.

Dextromethorphan

Antitussive (antagonizes NMDA glutamate receptors). Synthetic codeine analog. Has mild opioid effect when used in excess. Naloxone can be given for overdose. Mild abuse potential. May cause serotonin syndrome if combined with other serotonergic agents.

Pseudoephedrine, phenylephrine

MECHANISM	α-adrenergic agonists.
CLINICAL USE	Reduce hyperemia, edema (used as nasal decongestants); open obstructed eustachian tubes.
ADVERSE EFFECTS	Hypertension. Rebound congestion if used more than 4–6 days. Can also cause CNS stimulation/anxiety (pseudoephedrine).

Pulmonary hypertension drugs

DRUG	MECHANISM	CLINICAL NOTES
Endothelin receptor antagonists	Competitively antagonizes endothelin-1 receptors → ↓ pulmonary vascular resistance.	Hepatotoxic (monitor LFTs). Example: bosentan.
PDE-5 inhibitors	Inhibits PDE-5 → ↑ cGMP → prolonged vasodilatory effect of NO.	Also used to treat erectile dysfunction. Contraindicated when taking nitroglycerin or other nitrates (due to risk of severe hypotension). Example: sildenafil.
Prostacyclin analogs	PGI_2 (prostacyclin) with direct vasodilatory effects on pulmonary and systemic arterial vascular beds. Inhibits platelet aggregation.	Side effects: flushing, jaw pain. Examples: epoprostenol, iloprost.

(c) ketabton.com: The Digital Library

Asthma drugs

Bronchoconstriction is mediated by (1) inflammatory processes and (2) parasympathetic tone; therapy is directed at these 2 pathways.

β_2-agonists

Albuterol—relaxes bronchial smooth muscle (short acting β_2-agonist). For acute exacerbations. Can cause tremor, arrhythmia.

Salmeterol, formoterol—long-acting agents for prophylaxis. Can cause tremor, arrhythmia.

Inhaled corticosteroids

Fluticasone, budesonide—inhibit the synthesis of virtually all cytokines. Inactivate NF-κB, the transcription factor that induces production of TNF-α and other inflammatory agents. 1st-line therapy for chronic asthma. Use a spacer or rinse mouth after use to prevent oral thrush.

Muscarinic antagonists

Tiotropium, ipratropium—competitively block muscarinic receptors, preventing bronchoconstriction. Also used for COPD. Tiotropium is long acting.

Antileukotrienes

Montelukast, zafirlukast—block leukotriene receptors (CysLT1). Especially good for aspirin-induced and exercise-induced asthma.

Zileuton—5-lipoxygenase pathway inhibitor. Blocks conversion of arachidonic acid to leukotrienes. Hepatotoxic.

Anti-IgE monoclonal therapy

Omalizumab—binds mostly unbound serum IgE and blocks binding to FcεRI. Used in allergic asthma with ↑ IgE levels resistant to inhaled steroids and long-acting β_2-agonists.

Methylxanthines

Theophylline—likely causes bronchodilation by inhibiting phosphodiesterase → ↑ cAMP levels due to ↓ cAMP hydrolysis. Limited use due to narrow therapeutic index (cardiotoxicity, neurotoxicity); metabolized by cytochrome P-450. Blocks actions of adenosine.

Chromones

Cromolyn—prevents mast cell degranulation. Prevents acute asthma symptoms. Rarely used.

Anti-IL-5 monoclonal therapy

Prevents eosinophil differentiation, maturation, activation, and survival mediated by IL-5 stimulation. For maintenance therapy in severe eosinophilic asthma.

Mepolizumab, reslizumab—against IL-5.

Benralizumab—against IL-5 receptor α.





(c) ketabton.com: The Digital Library

▸ NOTES

(c) ketabton.com: The Digital Library

Rapid Review

"Study without thought is vain: thought without study is dangerous."
—Confucius

"It is better, of course, to know useless things than to know nothing."
—Lucius Annaeus Seneca

"For every complex problem there is an answer that is clear, simple, and wrong."
—H. L. Mencken

The following tables represent a collection of high-yield associations between diseases and their clinical findings, treatments, and key associations. They can be quickly reviewed in the days before the exam.

- Classic Presentations 690
- Classic Labs/ Findings 695
- Classic/Relevant Treatments 699
- Key Associations 702
- Equation Review 707
- Easily Confused Medications 709

FAS1_2019_17_Rapid Rev.indd 689 11/7/19 6:09 PM

(c) ketabton.com: The Digital Library

▸ CLASSIC PRESENTATIONS

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Gout, intellectual disability, self-mutilating behavior in a boy	Lesch-Nyhan syndrome (HGPRT deficiency, X-linked recessive)	37
Situs inversus, chronic sinusitis, bronchiectasis, infertility	Kartagener syndrome (dynein arm defect affecting cilia)	49
Blue sclera	Osteogenesis imperfecta (type I collagen defect)	51
Elastic skin, hypermobility of joints, ↑ bleeding tendency	Ehlers-Danlos syndrome (type V collagen defect, type III collagen defect seen in vascular subtype of ED)	51
Arachnodactyly, lens dislocation (upward and temporal), aortic dissection, hyperflexible joints	Marfan syndrome (fibrillin defect)	52
Café-au-lait spots (unilateral), polyostotic fibrous dysplasia, precocious puberty, multiple endocrine abnormalities	McCune-Albright syndrome (G_s-protein activating mutation)	57
Calf pseudohypertrophy	Muscular dystrophy (most commonly Duchenne, due to X-linked recessive frameshift mutation of dystrophin gene)	61
Child uses arms to stand up from squat	Duchenne muscular dystrophy (Gowers sign)	61
Slow, progressive muscle weakness in boys	Becker muscular dystrophy (X-linked non-frameshift deletions in dystrophin; less severe than Duchenne)	61
Infant with cleft lip/palate, microcephaly or holoprosencephaly, polydactyly, cutis aplasia	Patau syndrome (trisomy 13)	63
Infant with microcephaly, rocker-bottom feet, clenched hands, and structural heart defect	Edwards syndrome (trisomy 18)	63
Single palmar crease	Down syndrome	63
Dilated cardiomyopathy, edema, alcoholism or malnutrition	Wet beriberi (thiamine [vitamin B_1] deficiency)	66
Dermatitis, dementia, diarrhea	Pellagra (niacin [vitamin B_3] deficiency)	67
Swollen gums, mucosal bleeding, poor wound healing, petechiae	Scurvy (vitamin C deficiency: can't hydroxylate proline/lysine for collagen synthesis)	69
Chronic exercise intolerance with myalgia, fatigue, painful cramps, myoglobinuria	McArdle disease (skeletal muscle glycogen phosphorylase deficiency)	87
Infant with hypoglycemia, hepatomegaly	Cori disease (debranching enzyme deficiency) or Von Gierke disease (glucose-6-phosphatase deficiency, more severe)	87
Myopathy (infantile hypertrophic cardiomyopathy), exercise intolerance	Pompe disease (lysosomal α-1,4-glucosidase deficiency)	87
"Cherry-red spots" on macula	Tay-Sachs (ganglioside accumulation) or Niemann-Pick (sphingomyelin accumulation), central retinal artery occlusion	88
Hepatosplenomegaly, pancytopenia, osteoporosis, avascular necrosis of femoral head, bone crises	Gaucher disease (glucocerebrosidase [β-glucosidase] deficiency)	88
Achilles tendon xanthoma	Familial hypercholesterolemia (↓ LDL receptor signaling)	94
Anaphylaxis following blood transfusion	IgA deficiency	116
Male child, recurrent infections, no mature B cells	Bruton disease (X-linked agammaglobulinemia)	116

(c) ketabton.com: The Digital Library

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Recurrent cold (noninflamed) abscesses, eczema, high serum IgE, ↑ eosinophils	Hyper-IgE syndrome (Job syndrome: neutrophil chemotaxis abnormality)	116
"Strawberry tongue"	Scarlet fever Kawasaki disease	136, 314
Abdominal pain, diarrhea, leukocytosis, recent antibiotic use	*Clostridium difficile* infection	138
Back pain, fever, night sweats	Pott disease (vertebral TB)	140
Adrenal hemorrhage, hypotension, DIC	Waterhouse-Friderichsen syndrome (meningococcemia)	142, 349
Red "currant jelly" sputum in alcoholic or diabetic patients	*Klebsiella pneumoniae* pneumonia	145
Large rash with bull's-eye appearance	Erythema migrans from *Ixodes* tick bite (Lyme disease: *Borrelia*)	146
Ulcerated genital lesion	Nonpainful, indurated: chancre (1° syphilis, *Treponema pallidum*) Painful, with exudate: chancroid (*Haemophilus ducreyi*)	147, 184
Pupil accommodates but doesn't react	Neurosyphilis (Argyll Robertson pupil)	147
Smooth, moist, painless, wart-like white lesions on genitals	Condylomata lata (2° syphilis)	147
Fever, chills, headache, myalgia following antibiotic treatment for syphilis	Jarisch-Herxheimer reaction (rapid lysis of spirochetes results in endotoxin-like release)	148
Dog or cat bite resulting in infection	*Pasteurella multocida* (cellulitis at inoculation site)	149
Rash on palms and soles	Coxsackie A, 2° syphilis, Rocky Mountain spotted fever	150
Black eschar on face of patient with diabetic ketoacidosis	*Mucor* or *Rhizopus* fungal infection	153
Chorioretinitis, hydrocephalus, intracranial calcifications	Congenital toxoplasmosis	156
Child with fever later develops red rash on face that spreads to body	Erythema infectiosum/fifth disease ("slapped cheeks" appearance, caused by parvovirus B19)	164
Fever, cough, conjunctivitis, coryza, diffuse rash	Measles	170
Small, irregular red spots on buccal/lingual mucosa with blue-white centers	Koplik spots (measles [rubeola] virus)	170
Bounding pulses, wide pulse pressure, diastolic heart murmur, head bobbing	Aortic regurgitation	291
Systolic ejection murmur (crescendo-decrescendo)	Aortic stenosis	291
Continuous "machine-like" heart murmur	PDA (close with indomethacin; keep open with PGE analogs)	291
Chest pain on exertion	Angina (stable: with moderate exertion; unstable: with minimal exertion or at rest)	304
Chest pain with ST depressions on ECG	Angina (⊖ troponins) or NSTEMI (⊕ troponins)	304
Chest pain, pericardial effusion/friction rub, persistent fever following MI	Dressler syndrome (autoimmune-mediated post-MI fibrinous pericarditis, 2 weeks to several months after acute episode)	307
Painful, raised red lesions on pads of fingers/toes	Osler nodes (infective endocarditis, immune complex deposition)	311

(c) ketabton.com: The Digital Library

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Painless erythematous lesions on palms and soles	Janeway lesions (infective endocarditis, septic emboli/microabscesses)	311
Splinter hemorrhages in fingernails	Bacterial endocarditis	311
Retinal hemorrhages with pale centers	Roth spots (bacterial endocarditis)	311
Distant heart sounds, distended neck veins, hypotension	Beck triad of cardiac tamponade	310
Cervical lymphadenopathy, desquamating rash, coronary aneurysms, red conjunctivae and tongue, hand-foot changes	Kawasaki disease (mucocutaneous lymph node syndrome, treat with IVIG and aspirin)	314
Palpable purpura on buttocks/legs, joint pain, abdominal pain (child), hematuria	Immunoglobulin A vasculitis (Henoch-Schönlein purpura, affects skin and kidneys)	315
Telangiectasias, recurrent epistaxis, skin discoloration, arteriovenous malformations, GI bleeding, hematuria	Hereditary hemorrhagic telangiectasia (Osler-Weber-Rendu syndrome)	316
Skin hyperpigmentation, hypotension, fatigue	1° adrenocortical insufficiency → ↑ ACTH, ↑ α-MSH (eg, Addison disease)	349
Cutaneous flushing, diarrhea, bronchospasm	Carcinoid syndrome (right-sided cardiac valvular lesions, ↑ 5-HIAA)	352
Cold intolerance, weight gain, brittle hair	Hypothyroidism	341
Cutaneous/dermal edema due to deposition of mucopolysaccharides in connective tissue	Myxedema (caused by hypothyroidism, Graves disease [pretibial])	340
Facial muscle spasm upon tapping	Chvostek sign (hypocalcemia)	344
No lactation postpartum, absent menstruation, cold intolerance	Sheehan syndrome (postpartum hemorrhage leading to pituitary infarction)	339
Deep, labored breathing/hyperventilation	Diabetic ketoacidosis (Kussmaul respirations)	347
Pancreatic, pituitary, parathyroid tumors	MEN 1 (autosomal dominant)	351
Thyroid tumors, pheochromocytoma, ganglioneuromatosis, Marfanoid habitus	MEN 2B (autosomal dominant *RET* mutation)	351
Thyroid and parathyroid tumors, pheochromocytoma	MEN 2A (autosomal dominant *RET* mutation)	351
Jaundice, palpable distended non-tender gallbladder	Courvoisier sign (distal malignant obstruction of biliary tree)	398
Vomiting blood following gastroesophageal lacerations	Mallory-Weiss syndrome (alcoholic and bulimic patients)	377
Dysphagia (esophageal webs), glossitis, iron deficiency anemia	Plummer-Vinson syndrome (may progress to esophageal squamous cell carcinoma)	377
Enlarged, hard left supraclavicular node	Virchow node (abdominal metastasis)	379
Arthralgias, adenopathy, cardiac and neurological symptoms, diarrhea	Whipple disease (*Tropheryma whipplei*)	381
Severe RLQ pain with palpation of LLQ	Rovsing sign (acute appendicitis)	383
Severe RLQ pain with deep tenderness	McBurney sign (acute appendicitis)	383
Hamartomatous GI polyps, hyperpigmented macules on mouth, feet, hands, genitalia	Peutz-Jeghers syndrome (inherited, benign polyposis can cause bowel obstruction; ↑ cancer risk, mainly GI)	387
Multiple colon polyps, osteomas/soft tissue tumors, impacted/supernumerary teeth	Gardner syndrome (subtype of FAP)	387
Abdominal pain, ascites, hepatomegaly	Budd-Chiari syndrome (posthepatic venous thrombosis)	392

(c) ketabton.com: The Digital Library

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Severe jaundice in neonate	Crigler-Najjar syndrome (congenital unconjugated hyperbilirubinemia)	394
Golden brown rings around peripheral cornea	Wilson disease (Kayser-Fleischer rings due to copper accumulation)	395
Fat, female, forty, fertile	Cholelithiasis (gallstones)	396
Painless jaundice	Cancer of the pancreatic head obstructing bile duct	398
Bluish line on gingiva	Burton line (lead poisoning)	419
Short stature, café-au-lait spots, thumb/radial defects, ↑ incidence of tumors/leukemia, aplastic anemia	Fanconi anemia (genetic loss of DNA crosslink repair; often progresses to AML)	421
Red/pink urine, fragile RBCs	Paroxysmal nocturnal hemoglobinuria	422
Painful blue fingers/toes, hemolytic anemia	Cold agglutinin disease (autoimmune hemolytic anemia caused by *Mycoplasma pneumoniae*, infectious mononucleosis, CLL)	423
Petechiae, mucosal bleeding, prolonged bleeding time	Platelet disorders (eg, Glanzmann thrombasthenia, Bernard Soulier, HUS, TTP, ITP)	427
Fever, night sweats, weight loss	B symptoms of malignancy	429
Skin patches/plaques, Pautrier microabscesses, atypical T cells	Mycosis fungoides (cutaneous T-cell lymphoma) or Sézary syndrome (mycosis fungoides + malignant T cells in blood)	430
WBCs that look "smudged"	CLL	432
Neonate with arm paralysis following difficult birth, arm in "waiter's tip" position	Erb-Duchenne palsy (superior trunk [C5–C6] brachial plexus injury)	448
Anterior drawer sign ⊕	Anterior cruciate ligament injury	454
Bone pain, bone enlargement, arthritis	Osteitis deformans (Paget disease of bone, ↑ osteoblastic and osteoclastic activity)	463
Swollen, hard, painful finger joints in an elderly individual, pain worse with activity	Osteoarthritis (osteophytes on PIP [Bouchard nodes], DIP [Heberden nodes])	466
Sudden swollen/painful big toe joint, tophi	Gout/podagra (hyperuricemia)	467
Dry eyes, dry mouth, arthritis	Sjögren syndrome (autoimmune destruction of exocrine glands)	468
Urethritis, conjunctivitis, arthritis in a male	Reactive arthritis associated with HLA-B27	469
"Butterfly" facial rash and Raynaud phenomenon in a young female	Systemic lupus erythematosus	470
Painful fingers/toes changing color from white to blue to red with cold or stress	Raynaud phenomenon (vasospasm in extremities)	472
Anticentromere antibodies	Scleroderma (CREST)	473
Dark purple skin/mouth nodules in a patient with AIDS	Kaposi sarcoma, associated with HHV-8	478
Anti-desmoglein (anti-desmosome) antibodies	Pemphigus vulgaris (blistering)	480
Pruritic, purple, polygonal planar papules and plaques (6 P's)	Lichen planus	482
↑ AFP in amniotic fluid/maternal serum	Dating error, anencephaly, spina bifida (open neural tube defects)	491
Ataxia, nystagmus, vertigo, dysarthria	Cerebellar lesion	499

(c) ketabton.com: The Digital Library

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Toe extension/fanning upon plantar scrape	Babinski sign (UMN lesion)	510
Hyperphagia, hypersexuality, hyperorality	Klüver-Bucy syndrome (bilateral amygdala lesion)	511
Resting tremor, athetosis, chorea	Basal ganglia lesion	511
Lucid interval after traumatic brain injury	Epidural hematoma (middle meningeal artery rupture)	513
"Worst headache of my life"	Subarachnoid hemorrhage	513
Resting tremor, rigidity, akinesia, postural instability, shuffling gait	Parkinson disease (loss of dopaminergic neurons in substantia nigra pars compacta)	520
Chorea, dementia, caudate degeneration	Huntington disease (autosomal dominant CAG repeat expansion)	520
Nystagmus, intention tremor, scanning speech, bilateral internuclear ophthalmoplegia	Multiple sclerosis	523
Rapidly progressive limb weakness that ascends following GI/upper respiratory infection	Guillain-Barré syndrome (acute inflammatory demyelinating polyradiculopathy subtype)	524
Café-au-lait spots, Lisch nodules (iris hamartoma), cutaneous neurofibromas, pheochromocytomas, optic gliomas	Neurofibromatosis type I	525
Vascular birthmark (port-wine stain) of the face	Nevus flammeus (benign, but associated with Sturge-Weber syndrome)	525
Renal cell carcinoma (bilateral), hemangioblastomas, angiomatosis, pheochromocytoma	von Hippel-Lindau disease (dominant tumor suppressor gene mutation)	525
Bilateral vestibular schwannomas	Neurofibromatosis type 2	525
Hyperreflexia, hypertonia, Babinski sign present	UMN damage	529
Hyporeflexia, hypotonia, atrophy, fasciculations	LMN damage	529
Spastic weakness, sensory loss, bowel/bladder dysfunction	Spinal cord lesion	530
Unilateral facial drooping involving forehead	LMN facial nerve (CN VII) palsy; UMN lesions spare the forehead	532
Episodic vertigo, tinnitus, hearing loss	Ménière disease	534
Ptosis, miosis, anhidrosis	Horner syndrome (sympathetic chain lesion)	540
Conjugate horizontal gaze palsy, horizontal diplopia	Internuclear ophthalmoplegia (damage to MLF; may be unilateral or bilateral)	543
Polyuria, renal tubular acidosis type II, growth failure, electrolyte imbalances, hypophosphatemic rickets	Fanconi syndrome (multiple combined dysfunction of the proximal convoluted tubule)	586
Athlete with polycythemia	2° to erythropoietin injection	589
Periorbital and/or peripheral edema, proteinuria (> 3.5g/day), hypoalbuminemia, hypercholesterolemia	Nephrotic syndrome	597
Hereditary nephritis, sensorineural hearing loss, retinopathy, lens dislocation	Alport syndrome (mutation in collagen IV)	596
Streak ovaries, congenital heart disease, horseshoe kidney, cystic hygroma at birth, short stature, webbed neck, lymphedema	Turner syndrome (45,XO)	638
Red, itchy, swollen rash of nipple/areola	Paget disease of the breast (sign of underlying neoplasm)	650

(c) ketabton.com: The Digital Library

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Fibrous plaques in tunica albuginea of penis with abnormal curvature	Peyronie disease (connective tissue disorder)	651
Hypoxemia, polycythemia, hypercapnia	Chronic bronchitis (hyperplasia of mucous cells, "blue bloater")	674
Pink complexion, dyspnea, hyperventilation	Emphysema ("pink puffer," centriacinar [smoking] or panacinar [α_1-antitrypsin deficiency])	674
Bilateral hilar adenopathy, uveitis	Sarcoidosis (noncaseating granulomas)	676

▸ CLASSIC LABS/FINDINGS

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
↓ AFP in amniotic fluid/maternal serum	Down syndrome, Edwards syndrome	63
Large granules in phagocytes, immunodeficiency	Chédiak-Higashi disease (congenital failure of phagolysosome formation)	117
Recurrent infections, eczema, thrombocytopenia	Wiskott-Aldrich syndrome	117
Optochin sensitivity	Sensitive: *S pneumoniae*; resistant: viridans streptococci (*S mutans*, *S sanguis*)	134
Novobiocin response	Sensitive: *S epidermidis*; resistant: *S saprophyticus*	134
Bacitracin response	Sensitive: *S pyogenes* (group A); resistant: *S agalactiae* (group B)	134
Streptococcus bovis bacteremia	Colon cancer	137
Branching gram ⊕ rods with sulfur granules	*Actinomyces israelii*	139
Hilar lymphadenopathy, peripheral granulomatous lesion in middle or lower lung lobes (can calcify)	Ghon complex (1° TB: *Mycobacterium* bacilli)	140
"Thumb sign" on lateral neck x-ray	Epiglottitis (*Haemophilus influenzae*)	142
Bacteria-covered vaginal epithelial cells	"Clue cells" (*Gardnerella vaginalis*)	148
Cardiomegaly with apical atrophy	Chagas disease (*Trypanosoma cruzi*)	158
Atypical lymphocytes	EBV	165
Enlarged cells with intranuclear inclusion bodies	"Owl eye" appearance of CMV	165
Heterophile antibodies	Infectious mononucleosis (EBV)	165
Intranuclear eosinophilic droplet-like bodies	Cowdry type A bodies (HSV or VZV)	166
Eosinophilic globule in liver	Councilman body (viral hepatitis, yellow fever), represents hepatocyte undergoing apoptosis	168
"Steeple" sign on frontal CXR	Croup (parainfluenza virus)	170
Eosinophilic inclusion bodies in cytoplasm of hippocampal and cerebellar neurons	Negri bodies of rabies	171
Ring-enhancing brain lesion on CT/MRI in AIDS	*Toxoplasma gondii*, CNS lymphoma	177
Psammoma bodies	Meningiomas, papillary thyroid carcinoma, mesothelioma, papillary serous carcinoma of the endometrium and ovary	211

(c) ketabton.com: The Digital Library

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
"Delta wave" on ECG, short PR interval, supraventricular tachycardia	Wolff-Parkinson-White syndrome (Bundle of Kent bypasses AV node)	294
"Boot-shaped" heart on x-ray	Tetralogy of Fallot (due to RVH)	298
Rib notching (inferior surface, on x-ray)	Coarctation of the aorta	299
Heart nodules (granulomatous)	Aschoff bodies (rheumatic fever)	312
Electrical alternans (alternating amplitude on ECG)	Cardiac tamponade	310
Antineutrophil cytoplasmic antibodies (ANCAs)	Microscopic polyangiitis and eosinophilic granulomatosis with polyangiitis (MPO-ANCA/p-ANCA); granulomatosis with polyangiitis (Wegener; PR3-ANCA/c-ANCA); primary sclerosing cholangitis (MPO-ANCA/p-ANCA)	315
Hypertension, hypokalemia, metabolic alkalosis	1° hyperaldosteronism (Conn syndrome)	349
Enlarged thyroid cells with ground-glass nuclei with central clearing	"Orphan Annie" eyes nuclei (papillary carcinoma of the thyroid)	343
Mucin-filled cell with peripheral nucleus	"Signet ring" (gastric carcinoma)	379
Anti-transglutaminase/anti-gliadin/anti-endomysial antibodies	Celiac disease (diarrhea, weight loss)	381
Narrowing of bowel lumen on barium x-ray	"String sign" (Crohn disease)	382
"Lead pipe" appearance of colon on abdominal imaging	Ulcerative colitis (loss of haustra)	382
Thousands of polyps on colonoscopy	Familial adenomatous polyposis (autosomal dominant, mutation of APC gene)	387
"Apple core" lesion on barium enema x-ray	Colorectal cancer (usually left-sided)	388
Eosinophilic cytoplasmic inclusion in liver cell	Mallory body (alcoholic liver disease)	391
Triglyceride accumulation in liver cell vacuoles	Fatty liver disease (alcoholic or metabolic syndrome)	391
"Nutmeg" appearance of liver	Chronic passive congestion of liver due to right heart failure or Budd-Chiari syndrome	392
Antimitochondrial antibodies (AMAs)	1° biliary cholangitis (female, cholestasis, portal hypertension)	395
Low serum ceruloplasmin	Wilson disease (hepatolenticular degeneration; Kayser-Fleischer rings due to copper accumulation)	395
Migratory thrombophlebitis (leading to migrating DVTs and vasculitis)	Trousseau syndrome (adenocarcinoma of pancreas or lung)	398
Basophilic nuclear remnants in RBCs	Howell-Jolly bodies (due to splenectomy or nonfunctional spleen)	416
Basophilic stippling of RBCs	Lead poisoning or sideroblastic anemia	416
Hypochromic, microcytic anemia	Iron deficiency anemia, lead poisoning, thalassemia (fetal hemoglobin sometimes present)	418, 419
"Hair on end" ("Crew-cut") appearance on x-ray	β-thalassemia, sickle cell disease (marrow expansion)	422
Hypersegmented neutrophils	Megaloblastic anemia (B_{12} deficiency: neurologic symptoms; folate deficiency: no neurologic symptoms)	420
Antiplatelet antibodies	Idiopathic thrombocytopenic purpura	427
High level of D-dimers	DVT, PE, DIC	428
Giant B cells with bilobed nuclei with prominent inclusions ("owl's eye")	Reed-Sternberg cells (Hodgkin lymphoma)	429

(c) ketabton.com: The Digital Library

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
Sheets of medium-sized lymphoid cells with scattered pale, tingible body–laden macrophages ("starry sky" histology)	Burkitt lymphoma (t[8:14] c-*myc* activation, associated with EBV; "starry sky" made up of malignant cells)	430
Lytic ("punched-out") bone lesions on x-ray	Multiple myeloma	431
Monoclonal antibody spike	▪ Multiple myeloma (usually IgG or IgA) ▪ Monoclonal gammopathy of undetermined significance (MGUS consequence of aging) ▪ Waldenström (M protein = IgM) macroglobulinemia ▪ Primary amyloidosis	431
Stacks of RBCs	Rouleaux formation (high ESR, multiple myeloma)	423
Azurophilic peroxidase ⊕ granular inclusions in granulocytes and myeloblasts	Auer rods (AML, especially the promyelocytic [M3] type)	432
WBCs that look "smudged"	CLL (almost always B cell)	432
"Tennis racket"-shaped cytoplasmic organelles (EM) in Langerhans cells	Birbeck granules (Langerhans cell histiocytosis)	434
"Brown" tumor of bone	Hyperparathyroidism or osteitis fibrosa cystica (deposited hemosiderin from hemorrhage gives brown color)	464
"Soap bubble" in femur or tibia on x-ray	Giant cell tumor of bone (generally benign)	464
Raised periosteum (creating a "Codman triangle")	Aggressive bone lesion (eg, osteosarcoma, Ewing sarcoma, osteomyelitis)	465
"Onion skin" periosteal reaction	Ewing sarcoma (malignant small blue cell tumor)	465
Anti-IgG antibodies	Rheumatoid arthritis (systemic inflammation, joint pannus, boutonniere and swan neck deformities)	466
Rhomboid crystals, ⊕ birefringent	Pseudogout (calcium pyrophosphate dihydrate crystals)	467
Needle-shaped, ⊖ birefringent crystals	Gout (monosodium urate crystals)	467
↑ uric acid levels	Gout, Lesch-Nyhan syndrome, tumor lysis syndrome, loop and thiazide diuretics	467
"Bamboo spine" on x-ray	Ankylosing spondylitis (chronic inflammatory arthritis: HLA-B27)	469
Antinuclear antibodies (ANAs: anti-Smith and anti-dsDNA)	SLE (type III hypersensitivity)	470
Anti-histone antibodies	Drug-induced SLE (eg, hydralazine, isoniazid, phenytoin, procainamide)	250
Anti-topoisomerase antibodies	Diffuse scleroderma	473
Keratin pearls on a skin biopsy	Squamous cell carcinoma	484
Bloody or yellow tap on lumbar puncture	Xanthochromia (due to subarachnoid hemorrhage)	513
Eosinophilic cytoplasmic inclusion in neuron	Lewy body (Parkinson disease and Lewy body dementia)	520
Extracellular amyloid deposition in gray matter of brain	Senile plaques (Alzheimer disease)	520
Depigmentation of neurons in substantia nigra	Parkinson disease (basal ganglia disorder: rigidity, resting tremor, bradykinesia)	520
Protein aggregates in neurons from hyperphosphorylation of tau protein	Neurofibrillary tangles (Alzheimer disease) and Pick bodies (Pick disease)	520
Silver-staining spherical aggregation of tau proteins in neurons	Pick bodies (Pick disease: progressive dementia, changes in personality)	520

(c) ketabton.com: The Digital Library

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
Pseudopalisading tumor cells on brain biopsy	Glioblastoma multiforme	526
Circular grouping of dark tumor cells surrounding pale neurofibrils	Homer-Wright rosettes (neuroblastoma, medulloblastoma)	528
"Waxy" casts with very low urine flow	Chronic end-stage renal disease	594
Nodular hyaline deposits in glomeruli	Kimmelstiel-Wilson nodules (diabetic nephropathy)	597
Podocyte fusion or "effacement" on electron microscopy	Minimal change disease (child with nephrotic syndrome)	597
"Spikes" on basement membrane, "dome-like" subepithelial deposits	Membranous nephropathy (nephrotic syndrome)	597
RBC casts in urine	Glomerulonephritis	594
"Tram-track" appearance of capillary loops of glomerular basement membranes on light microscopy	Membranoproliferative glomerulonephritis	596
Anti–glomerular basement membrane antibodies	Goodpasture syndrome (glomerulonephritis and hemoptysis)	596
Cellular crescents in Bowman capsule	Rapidly progressive (crescentic) glomerulonephritis	596
"Wire loop" glomerular capillary appearance on light microscopy	Diffuse proliferative glomerulonephritis (usually seen with lupus)	596
Linear appearance of IgG deposition on glomerular and alveolar basement membranes	Goodpasture syndrome	596
"Lumpy bumpy" appearance of glomeruli on immunofluorescence	Poststreptococcal glomerulonephritis (due to deposition of IgG, IgM, and C3)	596
Necrotizing vasculitis (lungs) and necrotizing glomerulonephritis	Granulomatosis with polyangiitis (Wegener; PR3-ANCA/c-ANCA) and Goodpasture syndrome (anti–basement membrane antibodies)	596
Thyroid-like appearance of kidney	Chronic pyelonephritis (usually due to recurrent infections)	600
WBC casts in urine	Acute pyelonephritis	600
Renal epithelial casts in urine	Intrinsic renal failure (eg, ischemia or toxic injury)	601
hCG elevated	Choriocarcinoma, hydatidiform mole (occurs with and without embryo, and multiple pregnancy)	633
Dysplastic squamous cervical cells with "raisinoid" nuclei and hyperchromasia	Koilocytes (HPV: predisposes to cervical cancer)	645
Disarrayed granulosa cells arranged around collections of eosinophilic fluid	Call-Exner bodies (granulosa cell tumor of the ovary)	647
"Chocolate cyst" of ovary	Endometriosis (frequently involves both ovaries)	648
Mammary gland ("blue domed") cyst	Fibrocystic change of the breast	649
Glomerulus-like structure surrounding vessel in germ cells	Schiller-Duval bodies (yolk sac tumor)	647
Rectangular, crystal-like, cytoplasmic inclusions in Leydig cells	Reinke crystals (Leydig cell tumor)	653
Thrombi made of white/red layers	Lines of Zahn (arterial thrombus, layers of platelets/RBCs)	672
Hexagonal, double-pointed, needle-like crystals in bronchial secretions	Bronchial asthma (Charcot-Leyden crystals: eosinophilic granules)	674

(c) ketabton.com: The Digital Library

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
Desquamated epithelium casts in sputum	Curschmann spirals (bronchial asthma; can result in whorled mucous plugs)	674
"Honeycomb lung" on x-ray or CT	Idiopathic pulmonary fibrosis	675
Colonies of mucoid *Pseudomonas* in lungs	Cystic fibrosis (autosomal recessive mutation in *CFTR* gene → fat-soluble vitamin deficiency and mucous plugs)	675
Iron-containing nodules in alveolar septum	Ferruginous bodies (asbestosis: ↑ chance of lung cancer)	677
Bronchogenic apical lung tumor on imaging	Pancoast tumor (can compress cervical sympathetic chain and cause Horner syndrome)	685

▸ CLASSIC/RELEVANT TREATMENTS

CONDITION	COMMON TREATMENT(S)	PAGE
Ethylene glycol/methanol intoxication	Fomepizole (alcohol dehydrogenase inhibitor)	72
Chronic hepatitis B or C	IFN-α (HBV and HCV); ribavirin, simeprevir, sofosbuvir (HCV)	121
Streptococcus bovis	Penicillin prophylaxis; evaluation for colon cancer if linked to endocarditis	137
Clostridium botulinum	Antitoxin	138
Clostridium tetani	Antitoxin	138
Haemophilus influenzae (B)	Amoxicillin ± clavulanate (mucosal infections), ceftriaxone (meningitis), rifampin (prophylaxis)	142
Neisseria gonorrhoeae	Ceftriaxone (add doxycycline to cover likely concurrent *C trachomatis*)	142
Neisseria meningitidis	Penicillin/ceftriaxone, rifampin (prophylaxis)	142
Legionella pneumophila	Macrolides (eg, azithromycin)	143
Pseudomonas aeruginosa	Piperacillin/tazobactam, aminoglycosides, carbapenems	143
Treponema pallidum	Penicillin G	147
Chlamydia trachomatis	Doxycycline (+ ceftriaxone for gonorrhea coinfection), oral erythromycin to treat chlamydial conjunctivitis in infants	148
Candida albicans	Topical azoles (vaginitis); nystatin, fluconazole, caspofungin (oral/esophageal); fluconazole, caspofungin, amphotericin B (systemic)	153
Cryptococcus neoformans	Induction with amphotericin B and flucytosine, maintenance with fluconazole (in AIDS patients)	153
Sporothrix schenckii	Itraconazole, oral potassium iodide	154
Pneumocystis jirovecii	TMP-SMX (prophylaxis and treatment in immunosuppressed patients, CD4 < 200/mm^3)	154
Toxoplasma gondii	Sulfadiazine + pyrimethamine	156
Malaria	Chloroquine, mefloquine, atovaquone/proguanil (for blood schizont), primaquine (for liver hypnozoite)	157

(c) ketabton.com: The Digital Library

CONDITION	COMMON TREATMENT(S)	PAGE
Trichomonas vaginalis	Metronidazole (patient and partner)	158
Streptococcus pyogenes	Penicillin prophylaxis	187
Streptococcus pneumoniae	Penicillin/cephalosporin (systemic infection, pneumonia), vancomycin (meningitis)	187, 190
Staphylococcus aureus	MSSA: nafcillin, oxacillin, dicloxacillin (antistaphylococcal penicillins); MRSA: vancomycin, daptomycin, linezolid, ceftaroline	188, 190, 195
Enterococci	Vancomycin, aminopenicillins/cephalosporins	189, 190
Rickettsia rickettsii	Doxycycline, chloramphenicol	192
Clostridium difficile	Oral metronidazole; if refractory, oral vancomycin	190, 195
Mycobacterium tuberculosis	RIPE (rifampin, isoniazid, pyrazinamide, ethambutol)	196
UTI prophylaxis	TMP-SMX	198
Influenza	Oseltamivir, zanamivir	201
CMV	Ganciclovir, foscarnet, cidofovir	202
Patent ductus arteriosus	Close with indomethacin; keep open with PGE analogs	282
Stable angina	Sublingual nitroglycerin	304
Buerger disease	Smoking cessation	314
Kawasaki disease	IVIG, aspirin	314
Temporal arteritis	High-dose steroids	314
Granulomatosis with polyangiitis (Wegener)	Cyclophosphamide, corticosteroids	315
Hypercholesterolemia	Statin (first-line)	320
Hypertriglyceridemia	Fibrate	320
Arrhythmia in damaged cardiac tissue	Class IB antiarrhythmic (lidocaine, mexiletine)	322
Prolactinoma	Cabergoline/bromocriptine (dopamine agonists)	330
Diabetes insipidus	Desmopressin (central); hydrochlorothiazide, indomethacin, amiloride (nephrogenic)	338
SIADH	Fluid restriction, IV hypertonic saline, conivaptan/tolvaptan, demeclocycline	338
Diabetic ketoacidosis	Fluids, insulin, K^+	347
Diabetes mellitus type 1	Dietary intervention (low carbohydrate) + insulin replacement	347
Diabetes mellitus type 2	Dietary intervention, oral hypoglycemics, and insulin (if refractory)	347
Pheochromocytoma	α-antagonists (eg, phenoxybenzamine)	350
Carcinoid syndrome	Octreotide	352
Crohn disease	Corticosteroids, infliximab, azathioprine	382
Ulcerative colitis	5-ASA preparations (eg, mesalamine), 6-mercaptopurine, infliximab, colectomy	382
Sickle cell disease	Hydroxyurea (↑ fetal hemoglobin)	422

(c) ketabton.com: The Digital Library

CONDITION	COMMON TREATMENT(S)	PAGE
Chronic myelogenous leukemia	Imatinib	433
Acute promyelocytic leukemia (M3)	All-*trans* retinoic acid, arsenic trioxide	432
Drug of choice for anticoagulation in pregnancy or renal failure	Low-molecular-weight heparin	436
Heparin reversal	Protamine sulfate	436
Immediate anticoagulation	Heparin	436
Long-term anticoagulation	Warfarin, dabigatran, rivaroxaban and apixaban	436, 437
Warfarin reversal	Fresh frozen plasma (acute), vitamin K (non-acute)	436
Cyclophosphamide-induced hemorrhagic cystitis	Mesna	441
HER2/neu ⊕ breast cancer	Trastuzumab	443
Osteoporosis	Calcium/vitamin D supplementation (prophylaxis); bisphosphonates, PTH analogs, SERMs, calcitonin, denosumab (treatment)	462
Osteomalacia/rickets	Vitamin D supplementation	463
Chronic gout	Xanthine oxidase inhibitors (eg, allopurinol, febuxostat); pegloticase; probenecid	467
Acute gout attack	NSAIDs, colchicine, glucocorticoids	467
Neural tube defect prevention	Prenatal folic acid	491
Migraine	Abortive therapies (eg, sumatriptan, NSAIDs); prophylaxis (eg, propranolol, topiramate, CCBs, amitriptyline)	518
Multiple sclerosis	Disease-modifying therapies (eg, β-interferon, natalizumab); for acute flares, use IV steroids	523
Tonic-clonic seizures	Levetiracetam, phenytoin, valproate, carbamazepine	544
Absence seizures	Ethosuximide	544
Trigeminal neuralgia (tic douloureux)	Carbamazepine	544
Malignant hyperthermia	Dantrolene	551
Anorexia	Nutrition, psychotherapy, SSRIs	567
Bulimia nervosa	SSRIs	567
Alcoholism	Disulfiram, acamprosate, naltrexone, supportive care	571
ADHD	Methylphenidate, amphetamines, CBT, atomoxetine, guanfacine, clonidine	572
Alcohol withdrawal	Long-acting benzodiazepines	572
Bipolar disorder	Mood stabilizers (eg, lithium, valproic acid, carbamazepine), atypical antipsychotics	572
Depression	SSRIs (first-line)	572
Generalized anxiety disorder	SSRIs, SNRIs (first line); buspirone (second line)	572
Schizophrenia (positive symptoms)	Typical and atypical antipsychotics	573
Schizophrenia (negative symptoms)	Atypical antipsychotics	573

(c) ketabton.com: The Digital Library

CONDITION	COMMON TREATMENT(S)	PAGE
Hyperaldosteronism	Spironolactone	609
Benign prostatic hyperplasia	α_1-antagonists, 5α-reductase inhibitors, PDE-5 inhibitors	654
Infertility	Leuprolide, GnRH (pulsatile), clomiphene	656
Breast cancer in postmenopausal woman	Aromatase inhibitor (anastrozole)	656
ER ⊕ breast cancer	Tamoxifen	656
Prostate adenocarcinoma/uterine fibroids	Leuprolide, GnRH (continuous)	656
Medical abortion	Mifepristone	657
Prostate adenocarcinoma	Flutamide	658
Erectile dysfunction	Sildenafil, tadalafil, vardenafil	686
Pulmonary arterial hypertension (idiopathic)	Sildenafil, bosentan, epoprostenol	686

▸ KEY ASSOCIATIONS

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Mitochondrial inheritance	Disease occurs in both males and females, inherited through females only	59
Intellectual disability	Down syndrome, fragile X syndrome	62, 63
Vitamin deficiency (USA)	Folate (pregnant women are at high risk; body stores only 3- to 4-month supply; prevents neural tube defects)	68
Lysosomal storage disease	Gaucher disease	88
Bacterial meningitis (adults and elderly)	*S pneumoniae*	180
Bacterial meningitis (newborns and kids)	Group B streptococcus/*E coli*/*Listeria monocytogenes* (newborns), *S pneumoniae*/*N meningitidis* (kids/teens)	180
HLA-DR3	Diabetes mellitus type 1, SLE, Graves disease, Hashimoto thyroiditis (also associated with HLA-DR5), Addison disease	100
HLA-DR4	Diabetes mellitus type 1, rheumatoid arthritis, Addison disease	100
Bacteria associated with gastritis, peptic ulcer disease, and gastric malignancies (eg, adenocarcinoma, MALToma)	*H pylori*	146
Opportunistic infection in AIDS	*Pneumocystis jirovecii* pneumonia	154
Helminth infection (US)	*Enterobius vermicularis*	159
Viral encephalitis affecting temporal lobe	HSV-1	164
Infection 2° to blood transfusion	Hepatitis C	172
Food poisoning (exotoxin mediated)	*S aureus*, *B cereus*	178
Osteomyelitis	*S aureus* (most common overall)	180
Osteomyelitis in sickle cell disease	*Salmonella*	180
Osteomyelitis with IV drug use	*Pseudomonas*, *Candida*, *S aureus*	180

(c) ketabton.com: The Digital Library

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
UTI	*E coli*, *Staphylococcus saprophyticus* (young women)	181
Sexually transmitted disease	*C trachomatis* (usually coinfected with *N gonorrhoeae*)	184
Nosocomial pneumonia	*S aureus*, *Pseudomonas*, other enteric gram ⊖ rods	185
Pelvic inflammatory disease	*C trachomatis*, *N gonorrhoeae*	185
Infections in chronic granulomatous disease	*S aureus*, *E coli*, *Aspergillus* (catalase ⊕)	186
Metastases to bone	Prostate, breast > kidney, thyroid, lung	223
Metastases to brain	Lung > breast > melanoma, colon, kidney	223
Metastases to liver	Colon >> stomach > pancreas	223
S3 heart sound	↑ ventricular filling pressure (eg, mitral regurgitation, HF), common in dilated ventricles	287
S4 heart sound	Stiff/hypertrophic ventricle (aortic stenosis, restrictive cardiomyopathy)	287
Constrictive pericarditis	TB (developing world); idiopathic, viral illness (developed world)	287
Holosystolic murmur	VSD, tricuspid regurgitation, mitral regurgitation	291
Ejection click	Aortic stenosis	291
Mitral valve stenosis	Rheumatic heart disease	291
Opening snap	Mitral stenosis	291
Heart murmur, congenital	Mitral valve prolapse	291
Chronic arrhythmia	Atrial fibrillation (associated with high risk of emboli)	295
Cyanosis (early; less common)	Tetralogy of Fallot, transposition of great vessels, truncus arteriosus, total anomalous pulmonary venous return, tricuspid atresia	298
Late cyanotic shunt (uncorrected left to right becomes right to left)	Eisenmenger syndrome (caused by ASD, VSD, PDA; results in pulmonary hypertension/polycythemia)	299
Congenital cardiac anomaly	VSD	299
Hypertension, 2°	Renal artery stenosis, chronic kidney disease (eg, polycystic kidney disease, diabetic nephropathy), hyperaldosteronism	300
Aortic aneurysm, thoracic	Marfan syndrome (idiopathic cystic medial degeneration)	302
Aortic aneurysm, abdominal	Atherosclerosis, smoking is major risk factor	302
Aortic aneurysm, ascending or arch	3° syphilis (syphilitic aortitis), vasa vasorum destruction	303
Sites of atherosclerosis	Abdominal aorta > coronary artery > popliteal artery > carotid artery	302
Aortic dissection	Hypertension	303
Right heart failure due to a pulmonary cause	Cor pulmonale	309
Heart valve in bacterial endocarditis	Mitral > aortic (rheumatic fever), tricuspid (IV drug abuse)	310
Endocarditis presentation associated with bacterium	*S aureus* (acute, IVDA, tricuspid valve), viridans streptococci (subacute, dental procedure), *S bovis* (colon cancer), culture negative (*Coxiella*, *Bartonella*, HACEK)	310
Temporal arteritis	Risk of ipsilateral blindness due to occlusion of ophthalmic artery; polymyalgia rheumatica	314

(c) ketabton.com: The Digital Library

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Recurrent inflammation/thrombosis of small/medium vessels in extremities	Buerger disease (strongly associated with tobacco)	314
Cardiac 1° tumor (kids)	Rhabdomyoma, often seen in tuberous sclerosis	316
Cardiac tumor (adults)	Metastasis, myxoma (90% in left atrium; "ball valve")	316
Congenital adrenal hyperplasia, hypotension	21-hydroxylase deficiency	335
Hypopituitarism	Pituitary adenoma (usually benign tumor)	339
Cretinism	Iodine deficit/congenital hypothyroidism	341
Thyroid cancer	Papillary carcinoma (childhood irradiation)	343
Hypoparathyroidism	Accidental excision during thyroidectomy	344
1° hyperparathyroidism	Adenomas, hyperplasia, carcinoma	345
2° hyperparathyroidism	Hypocalcemia of chronic kidney disease	345
Cushing syndrome	▪ Iatrogenic (from corticosteroid therapy) ▪ Adrenocortical adenoma (secretes excess cortisol) ▪ ACTH-secreting pituitary adenoma (Cushing disease) ▪ Paraneoplastic (due to ACTH secretion by tumors)	348
1° hyperaldosteronism	Adrenal hyperplasia or adenoma	349
Tumor of the adrenal medulla (kids)	Neuroblastoma (malignant)	350
Tumor of the adrenal medulla (adults)	Pheochromocytoma (usually benign)	350
Refractory peptic ulcers and high gastrin levels	Zollinger-Ellison syndrome (gastrinoma of duodenum or pancreas), associated with MEN1	351, 352
Esophageal cancer	Squamous cell carcinoma (worldwide); adenocarcinoma (US)	378
Acute gastric ulcer associated with CNS injury	Cushing ulcer (↑ intracranial pressure stimulates vagal gastric H^+ secretion)	379
Acute gastric ulcer associated with severe burns	Curling ulcer (greatly reduced plasma volume results in sloughing of gastric mucosa)	379
Bilateral ovarian metastases from gastric carcinoma	Krukenberg tumor (mucin-secreting signet ring cells)	379
Chronic atrophic gastritis (autoimmune)	Predisposition to gastric carcinoma (can also cause pernicious anemia)	379
Gastric cancer	Adenocarcinoma	379
Alternating areas of transmural inflammation and normal colon	Skip lesions (Crohn disease)	382
Site of diverticula	Sigmoid colon	383
Diverticulum in pharynx	Zenker diverticulum (diagnosed by barium swallow)	384
Hepatocellular carcinoma	Cirrhotic liver (associated with hepatitis B and C, alcoholism, and hemochromatosis)	392
Liver disease	Alcoholic cirrhosis	391
1° liver cancer	Hepatocellular carcinoma (chronic hepatitis, cirrhosis, hemochromatosis, α_1-antitrypsin deficiency, Wilson disease)	392
Congenital conjugated hyperbilirubinemia (black liver)	Dubin-Johnson syndrome (inability of hepatocytes to secrete conjugated bilirubin into bile)	394

(c) ketabton.com: The Digital Library

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Hereditary harmless jaundice	Gilbert syndrome (benign congenital unconjugated hyperbilirubinemia)	394
Hemochromatosis	Multiple blood transfusions or hereditary *HFE* mutation (can result in heart failure, "bronze diabetes," and ↑ risk of hepatocellular carcinoma)	395
Pancreatitis (acute)	Gallstones, alcohol	397
Pancreatitis (chronic)	Alcohol (adults), cystic fibrosis (kids)	397
Microcytic anemia	Iron deficiency	418
Autosplenectomy (fibrosis and shrinkage)	Sickle cell disease (hemoglobin S)	422
Bleeding disorder with GpIb deficiency	Bernard-Soulier syndrome (defect in platelet adhesion to von Willebrand factor)	427
Hereditary bleeding disorder	von Willebrand disease	428
DIC	Severe sepsis, obstetric complications, cancer, burns, trauma, major surgery, acute pancreatitis, APL	428
Malignancy associated with noninfectious fever	Hodgkin lymphoma	429
Type of Hodgkin lymphoma	Nodular sclerosis (vs mixed cellularity, lymphocytic predominance, lymphocytic depletion)	429
t(14;18)	Follicular lymphomas (*BCL-2* activation, anti-apoptotic oncogene)	430
t(8;14)	Burkitt lymphoma (c-*myc* fusion, transcription factor oncogene)	430
Type of non-Hodgkin lymphoma	Diffuse large B-cell lymphoma	430
1° bone tumor (adults)	Multiple myeloma	431
Age ranges for patient with ALL/CLL/AML/CML	ALL: child, CLL: adult > 60, AML: adult ~ 65, CML: adult 45–85	432, 433
Malignancy (kids)	Leukemia, brain tumors	432, 526
Death in CML	Blast crisis	433
t(9;22)	Philadelphia chromosome, CML (*BCR-ABL* oncogene, tyrosine kinase activation), more rarely associated with ALL	434
Vertebral compression fracture	Osteoporosis (type I: postmenopausal woman; type II: elderly man or woman)	462
HLA-B27	Psoriatic arthritis, ankylosing spondylitis, IBD-associated arthritis, reactive arthritis (formerly Reiter syndrome)	469
Death in SLE	Lupus nephropathy	470
Tumor of infancy	Strawberry hemangioma (grows rapidly and regresses spontaneously by childhood)	478
Actinic (solar) keratosis	Precursor to squamous cell carcinoma	482
Cerebellar tonsillar herniation	Chiari I malformation	492
Atrophy of the mammillary bodies	Wernicke encephalopathy (thiamine deficiency causing ataxia, ophthalmoplegia, and confusion)	511

(c) ketabton.com: The Digital Library

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Epidural hematoma	Rupture of middle meningeal artery (trauma; lentiform shaped)	513
Subdural hematoma	Rupture of bridging veins (crescent shaped)	513
Dementia	Alzheimer disease, multiple infarcts (vascular dementia)	520, 521
Demyelinating disease in young women	Multiple sclerosis	523
Brain tumor (adults)	Supratentorial: metastasis, astrocytoma (including glioblastoma multiforme), meningioma, schwannoma	526
Pituitary tumor	Prolactinoma, somatotropic adenoma	527
Brain tumor (kids)	Infratentorial: medulloblastoma (cerebellum) or supratentorial: craniopharyngioma	528
Mixed (UMN and LMN) motor neuron disease	Amyotrophic lateral sclerosis	530
Degeneration of dorsal column fibers	Tabes dorsalis (3° syphilis), subacute combined degeneration (dorsal columns, lateral corticospinal, spinocerebellar tracts affected)	530
Nephrotic syndrome (adults)	Membranous nephropathy	597
Nephrotic syndrome (kids)	Minimal change disease	597
Glomerulonephritis (adults)	Berger disease (IgA nephropathy)	596
Kidney stones	▪ Calcium = radiopaque ▪ Struvite (ammonium) = radiopaque (formed by urease ⊕ organisms such as *Klebsiella*, *Proteus* species, and *S saprophyticus*) ▪ Uric acid = radiolucent ▪ Cystine = faintly radiopaque	598
Renal tumor	Renal cell carcinoma: associated with von Hippel-Lindau and cigarette smoking; paraneoplastic syndromes (EPO, renin, PTHrP, ACTH)	605
Obstruction of male urinary tract	BPH	654
1° amenorrhea	Turner syndrome (45,XO or 45,XO/46,XX mosaic)	638
Neuron migration failure	Kallmann syndrome (hypogonadotropic hypogonadism and anosmia)	639
Clear cell adenocarcinoma of the vagina	DES exposure in utero	644
Ovarian tumor (benign, bilateral)	Serous cystadenoma	646
Ovarian tumor (malignant)	Serous cystadenocarcinoma	646
Tumor in women	Leiomyoma (estrogen dependent, not precancerous)	648
Gynecologic malignancy	Endometrial carcinoma (most common in US); cervical carcinoma (most common worldwide)	648
Breast mass	Fibrocystic change, carcinoma (in postmenopausal women)	649
Breast tumor (benign, young woman)	Fibroadenoma	649
Breast cancer	Invasive ductal carcinoma	650
Testicular tumor	Seminoma (malignant, radiosensitive), ↑ placental ALP	652, 653

(c) ketabton.com: The Digital Library

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Pulmonary hypertension	Idiopathic, heritable, left heart disease (eg, HF), lung disease (eg, COPD), hypoxemic vasoconstriction (eg, OSA), thromboembolic (eg, PE)	679
Hypercoagulability, endothelial damage, blood stasis	Virchow triad (↑ risk of thrombosis)	671
SIADH	Small cell carcinoma of the lung	684

▸ EQUATION REVIEW

TOPIC	EQUATION	PAGE
Volume of distribution	$V_d = \frac{\text{amount of drug in the body}}{\text{plasma drug concentration}}$	231
Half-life	$t_{½} = \frac{0.7 \times V_d}{CL}$	231
Drug clearance	$CL = \frac{\text{rate of elimination of drug}}{\text{plasma drug concentration}} = V_d \times K_e$ (elimination constant)	231
Loading dose	$LD = \frac{C_p \times V_d}{F}$	231
Maintenance dose	$D = \frac{C_p \times CL \times \tau}{F}$	231
Sensitivity	Sensitivity = TP / (TP + FN)	257
Specificity	Specificity = TN / (TN + FP)	257
Positive predictive value	PPV = TP / (TP + FP)	257
Negative predictive value	NPV = TN / (FN + TN)	257
Odds ratio (for case-control studies)	$OR = \frac{a/c}{b/d} = \frac{ad}{bc}$	258
Relative risk	$RR = \frac{a/(a+b)}{c/(c+d)}$	258
Attributable risk	$AR = \frac{a}{a+b} - \frac{c}{c+d}$	258
Relative risk reduction	RRR = 1 − RR	258
Absolute risk reduction	$ARR = \frac{c}{c+d} - \frac{a}{a+b}$	258
Number needed to treat	NNT = 1/ARR	258
Number needed to harm	NNH = 1/AR	258
Cardiac output	$CO = \frac{\text{rate of } O_2 \text{ consumption}}{(\text{arterial } O_2 \text{ content} - \text{venous } O_2 \text{ content})}$ CO = stroke volume × heart rate	285 285

(c) ketabton.com: The Digital Library

TOPIC	EQUATION	PAGE
Mean arterial pressure	MAP = cardiac output × total peripheral resistance	285
	MAP = ⅔ diastolic + ⅓ systolic	285
Stroke volume	SV = EDV – ESV	285
Ejection fraction	$EF = \frac{SV}{EDV} = \frac{EDV - ESV}{EDV}$	285
Resistance	$\text{Resistance} = \frac{\text{driving pressure } (\Delta P)}{\text{flow } (Q)} = \frac{8\eta \text{ (viscosity)} \times \text{length}}{\pi r^4}$	286
Capillary fluid exchange	J_v = net fluid flow = $K_f[(P_c - P_i) - \sigma(\pi_c - \pi_i)]$	297
Renal clearance	$C_x = (U_x V)/P_x$	582
Glomerular filtration rate	$C_{inulin} = GFR = U_{inulin} \times V/P_{inulin}$	582
	$= K_f [(P_{GC} - P_{BS}) - (\pi_{GC} - \pi_{BS})]$	
Effective renal plasma flow	$eRPF = U_{PAH} \times \frac{V}{P_{PAH}} = C_{PAH}$	582
Renal blood flow	$RBF = \frac{RPF}{1 - Hct}$	582
Filtration fraction	$FF = \frac{GFR}{RPF}$	583
Henderson-Hasselbalch equation (for extracellular pH)	$pH = 6.1 + \log \frac{[HCO_3^-]}{0.03\ P_{CO_2}}$	592
Winters formula	$P_{CO_2} = 1.5\ [HCO_3^-] + 8 \pm 2$	592
Anion gap	$Na^+ - (Cl^- + HCO_3^-)$	592
Physiologic dead space	$V_D = V_T \times \frac{PaCO_2 - PECO_2}{PaCO_2}$	664
Pulmonary vascular resistance	$PVR = \frac{P_{\text{pulm artery}} - P_{\text{L atrium}}}{\text{cardiac output}}$	668
Alveolar gas equation	$P_{AO_2} = P_{IO_2} - \frac{PaCO_2}{R}$	668

(c) ketabton.com: The Digital Library

▸ EASILY CONFUSED MEDICATIONS

DRUG	CLINICAL USE/MECHANISM OF ACTION
Amiloride	K^+-sparing diuretic
Amiodarone	Class III antiarrhythmic
Amlodipine	Dihydropyridine Ca^{2+} channel blocker
Benztropine	Cholinergic antagonist
Bromocriptine	Dopamine agonist
Buspirone	Generalized anxiety disorder (5-HT_{1A}-receptor agonist)
Bupropion	Depression, smoking cessation (NE-DA reuptake inhibitor)
Cimetidine	H_2-receptor antagonist
Cetirizine	2nd-generation antihistamine
Chloramphenicol	Antibiotic (blocks 50S subunit)
Chlordiazepoxide	Long-acting benzodiazepine
Chlorpromazine	Typical antipsychotic
Chlorpropamide	1st-generation sulfonylurea
Chlorpheniramine	1st-generation antihistamine
Chlorthalidone	Thiazide diuretic
Clozapine	5-HT_{2A}-agonist
Clomipramine	Tricyclic antidepressant
Clomiphene	Selective estrogen receptor modulator
Clonidine	α_2-agonist
Doxepin	Tricyclic antidepressant
Doxazosin	α_1-antagonist
Eplerenone	K^+-sparing diuretic
Propafenone	Class IC antiarrhythmic
Fluoxetine	Selective serotonin reuptake inhibitor
Fluphenazine	Typical antipsychotic
Duloxetine	Serotonin-norepinephrine reuptake inhibitor
Guaifenesin	Expectorant (thins respiratory secretions)
Guanfacine	α_2-agonist
Mifepristone	Progesterone receptor antagonist
Misoprostol	PGE_1 synthetic analog
Naloxone	Opioid receptor antagonist (treats toxicity)
Naltrexone	Opioid receptor antagonist (prevents relapse)
Nitroprusside	Hypertensive emergency (↑ cGMP/NO)
Nitroglycerin	Antianginal (↑ cGMP/NO)
Omeprazole	Proton pump inhibitor
Ketoconazole	Antifungal (inhibits fungal sterol synthesis)

(c) ketabton.com: The Digital Library

DRUG	CLINICAL USE/MECHANISM OF ACTION
Aripiprazole	Atypical antipsychotic
Anastrozole	Aromatase inhibitor
Rifaximin	Hepatic encephalopathy (↓ ammoniagenic bacteria)
Rifampin	Antimicrobial (inhibits DNA-dependent RNA polymerase)
Sertraline	Selective serotonin reuptake inhibitor
Selegiline	MAO-B inhibitor
Trazodone	Insomnia (blocks 5-HT_2, α_1-adrenergic, and H_1 receptors)
Tramadol	Chronic pain (weak opioid agonist)
Varenicline	Smoking cessation (nicotinic ACh receptor partial agonist)
Venlafaxine	Serotonin-norepinephrine reuptake inhibitor

(c) ketabton.com: The Digital Library

SECTION IV

Top-Rated Review Resources

"Some books are to be tasted, others to be swallowed, and some few to be chewed and digested."

—Sir Francis Bacon

"Always read something that will make you look good if you die in the middle of it."

—P.J. O'Rourke

"So many books, so little time."

—Frank Zappa

"If one cannot enjoy reading a book over and over again, there is no use in reading it at all."

—Oscar Wilde

- How to Use the Database 712
- Question Banks and Books 714
- Web and Mobile Apps 714
- Comprehensive 715
- Anatomy, Embryology, and Neuroscience 715
- Behavioral Science 716
- Biochemistry 716
- Cell Biology and Histology 716
- Microbiology and Immunology 717
- Pathology 717
- Pharmacology 718
- Physiology 718

FAS1_2019_18_PrintEditionBookRev.indd 711 11/8/19 8:09 AM

(c) ketabton.com: The Digital Library

HOW TO USE THE DATABASE

This section is a database of top-rated basic science review books, sample examination books, software, websites, and apps that have been marketed to medical students studying for the USMLE Step 1. For each recommended resource, we list (where applicable) the **Title**, the **First Author** (or editor), the **Current Publisher**, the **Copyright Year**, the **Number of Pages**, the **Approximate List Price**, the **Format** of the resource, and the **Number of Test Questions**. Finally, each recommended resource receives a **Rating**. Within each section, resources are arranged first by Rating and then alphabetically by the first author within each Rating group.

For a complete list of resources, including summaries that describe their overall style and utility, go to **www.firstaidteam.com/bonus**.

A letter rating scale with six different grades reflects the detailed student evaluations for **Rated Resources**. Each rated resource receives a rating as follows:

A+	Excellent for boards review.
A A–	Very good for boards review; choose among the group.
B+ B	Good, but use only after exhausting better resources.
B–	Fair, but there are many better resources in the discipline; or low-yield subject material.

The Rating is meant to reflect the overall usefulness of the resource in helping medical students prepare for the USMLE Step 1. This is based on a number of factors, including:

- The cost
- The readability of the text or usability of the app
- The appropriateness and accuracy of the material
- The quality and number of sample questions
- The quality of written answers to sample questions
- The quality and appropriateness of the illustrations (eg, graphs, diagrams, photographs)
- The length of the text (longer is not necessarily better)
- The quality and number of other resources available in the same discipline
- The importance of the discipline for the USMLE Step 1

Please note that ratings do not reflect the quality of the resources for purposes other than reviewing for the USMLE Step 1. Many books with lower ratings are well written and informative but are not ideal for boards

(c) ketabton.com: The Digital Library

preparation. We have not listed or commented on general textbooks available in the basic sciences.

Evaluations are based on the cumulative results of formal and informal surveys of thousands of medical students at many medical schools across the country. The ratings represent a consensus opinion, but there may have been a broad range of opinion or limited student feedback on any particular resource.

Please note that the data listed are subject to change in that:

- Publishers' prices change frequently.
- Bookstores often charge an additional markup.
- New editions come out frequently, and the quality of updating varies.
- The same book may be reissued through another publisher.

We actively encourage medical students and faculty to submit their opinions and ratings of these basic science review materials so that we may update our database. (See p. xvii, How to Contribute.) In addition, we ask that publishers and authors submit for evaluation review copies of basic science review books, including new editions and books not included in our database. We also solicit reviews of new books or suggestions for alternate modes of study that may be useful in preparing for the examination, such as flash cards, computer software, commercial review courses, apps, and websites.

Disclaimer/Conflict of Interest Statement

No material in this book, including the ratings, reflects the opinion or influence of the publisher. All errors and omissions will gladly be corrected if brought to the attention of the authors through our blog at www.firstaidteam.com. Please note that USMLE-Rx and the entire *First Aid for the USMLE* series are publications by certain authors of this book; the following ratings are based solely on recommendations from the student authors of this book as well as data from the student survey and feedback forms.

(c) ketabton.com: The Digital Library

▸ TOP-RATED REVIEW RESOURCES

Question Banks and Books

		AUTHOR	PUBLISHER	TYPE	PRICE
A+	*UWorld Qbank*	UWorld	www.uworld.com	Test/2400 q	$249–$749
A	*NBME Practice Exams*	National Board of Medical Examiners	www.nbme.org/students/sas/Comprehensive.html	Test/200 q	$60
A−	*AMBOSS*	Amboss	www.amboss.com	Test/3500 q	$9–$365
A−	*USMLE-Rx Qmax*	USMLE-Rx	www.usmle-rx.com	Test/2300 q	$89–$339
B+	*Kaplan Qbank*	Kaplan	www.kaptest.com	Test/2100 q	$99–$349
B	*BoardVitals*		www.boardvitals.com	Test/1750 q	$59–$179
B	*Kaplan USMLE Step 1 Qbook*	Kaplan	Kaplan, 2017, 468 pages	Test/850 q	$50
B	*Pastest*		www.pastest.com	Test/2100 q	$79–$249
B	*TrueLearn Review*		www.truelearn.com	Test/2200 q	$159–$399

Web and Mobile Apps

		AUTHOR	PUBLISHER	TYPE	PRICE
A	*Anki*		www.ankisrs.net	Flash cards	Free
A	*Boards and Beyond*		www.boardsbeyond.com	Review/Test/1300 q	$19–$249
A	*Physeo*		www.physeo.com	Review	$30–$150
A	*SketchyMedical*		www.sketchymedical.com	Review	$99–$369
A−	*Cram Fighter*		www.cramfighter.com	Study plan	$29–$159
A−	*First Aid Step 1 Express*		www.usmle-rx.com	Review/Test	$69–$299
B+	*First Aid Step 1 Flash Facts*		www.usmle-rx.com	Flash cards	$29–$149
B+	*Medbullets*		www.medbullets.com	Review/Test/1000 q	Free
B+	*Medical School Pathology*		www.medicalschoolpathology.com	Review	Free
B+	*OnlineMedEd*		www.onlinemeded.org	Review	Free
B+	*Osmosis*		www.osmosis.org	Test	$179–$279
B+	*USMLE Step 1 Mastery*		builtbyhlt.com/medical/usmle-step-1-mastery	Test/1400 q	$2–$10
B+	*WebPath: The Internet Pathology Laboratory*		webpath.med.utah.edu	Review/Test/1300 q	Free
B	*Blue Histology*		www.lab.anhb.uwa.edu.au/mb140	Review/Test	Free
B	*Digital Anatomist Project: Interactive Atlases*	University of Washington	da.si.washington.edu/da.html	Review	Free
B	*Dr. Najeeb Lectures*		www.drnajeeblectures.com	Review	$99

(c) ketabton.com: The Digital Library

B	*Firecracker*	Firecracker Inc.	firecracker.lww.com	Review/ Test/2800 q	$39–$660
B	*KISSPrep*		www.kissprep.com	Review	$99–$135
B	*Lecturio*		www.lecturio.com	Review/ Test/2150 q	$50–$300
B	*Memorang*	Memorang Inc.	www.memorangapp.com	Flash cards	$19–$239
B	*Picmonic*		www.picmonic.com	Review	$25–$480
B⁻	*Radiopaedia.org*		www.radiopaedia.org	Cases/Test	Free
B⁻	*The Pathology Guy*	Friedlander	www.pathguy.com	Review	Free

Comprehensive

		AUTHOR	PUBLISHER	TYPE	PRICE
A	*First Aid for the Basic Sciences: General Principles*	Le	McGraw-Hill, 2017, 528 pages	Review	$55
A	*First Aid Cases for the USMLE Step 1*	Le	McGraw-Hill, 2018, 496 pages	Cases	$50
A⁻	*First Aid for the Basic Sciences: Organ Systems*	Le	McGraw-Hill, 2017, 912 pages	Review	$72
A⁻	*Crush Step 1: The Ultimate USMLE Step 1 Review*	O'Connell	Elsevier, 2017, 704 pages	Review	$45
A⁻	*Cracking the USMLE Step 1*	Princeton Review	Princeton Review, 2013, 832 pages	Review	$45
B⁺	*USMLE Step 1 Secrets in Color*	Brown	Elsevier, 2016, 800 pages, ISBN 9780323396790	Review	$43
B⁺	*Step-Up to USMLE Step 1 2015*	Jenkins	Lippincott Williams & Wilkins, 2014, 528 pages	Review	$50
B⁺	*USMLE Step 1 Lecture Notes 2018*	Kaplan	Kaplan Medical, 2018, ~2700 pages	Review	$330
B⁺	*USMLE Images for the Boards: A Comprehensive Image-Based Review*	Tully	Elsevier, 2012, 296 pages	Review	$42
B	*USMLE Step 1 Made Ridiculously Simple*	Carl	MedMaster, 2017, 416 pages,	Review/Test 1000 q	$30
B	*medEssentials for the USMLE Step 1*	Manley	Kaplan, 2012, 588 pages	Review	$55

Anatomy, Embryology, and Neuroscience

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁻	*High-Yield Gross Anatomy*	Dudek	Lippincott Williams & Wilkins, 2014, 320 pages	Review	$43
A⁻	*Clinical Anatomy Made Ridiculously Simple*	Goldberg	MedMaster, 2016, 175 pages	Review	$30
B⁺	*High-Yield Embryology*	Dudek	Lippincott Williams & Wilkins, 2013, 176 pages	Review	$56
B⁺	*High-Yield Neuroanatomy*	Fix	Lippincott Williams & Wilkins, 2015, 208 pages	Review/ Test/50 q	$40

(c) ketabton.com: The Digital Library

Anatomy, Embryology, and Neuroscience *(continued)*

		AUTHOR	PUBLISHER	TYPE	PRICE
B+	*Anatomy—An Essential Textbook*	Gilroy	Thieme, 2017, 528 pages	Text/ Test/400 q	$48
B+	*Netter's Anatomy Flash Cards*	Hansen	Saunders, 2018, 688 flash cards	Flash cards	$40
B+	*Crash Course: Anatomy*	Stenhouse	Elsevier, 2015, 288 pages	Review	$45
B	*BRS Embryology*	Dudek	Lippincott Williams & Wilkins, 2014, 336 pages	Review/ Test/220 q	$56
B	*Anatomy Flash Cards: Anatomy on the Go*	Gilroy	Thieme, 2013, 752 flash cards	Flash cards	$60
B	*Clinical Neuroanatomy Made Ridiculously Simple*	Goldberg	MedMaster, 2014, 90 pages + CD-ROM	Review/Test/ Few q	$26
B	*Netter's Anatomy Coloring Book*	Hansen	Elsevier, 2018, 392 pages	Review	$20
B	*Case Files: Anatomy*	Toy	McGraw-Hill, 2014, 416 pages	Cases	$35
B−	*Case Files: Neuroscience*	Toy	McGraw-Hill, 2014, 432 pages	Cases	$35

Behavioral Science

		AUTHOR	PUBLISHER	TYPE	PRICE
A−	*BRS Behavioral Science*	Fadem	Lippincott Williams & Wilkins, 2016, 384 pages	Review/ Test/700 q	$52
B+	*High-Yield Biostatistics, Epidemiology, and Public Health*	Glaser	Lippincott Williams & Wilkins, 2013, 168 pages	Review	$43

Biochemistry

		AUTHOR	PUBLISHER	TYPE	PRICE
A−	*Pixorize*		www.pixorize.com	Review	$100–$130
B+	*Medical Biochemistry—An Illustrated Review*	Panini	Thieme, 2013, 441 pages	Review/ Test/400 q	$40
B	*Lange Flash Cards Biochemistry and Genetics*	Baron	McGraw-Hill, 2017, 196 flash cards	Flash cards	$40
B	*Lippincott Illustrated Reviews: Biochemistry*	Ferrier	Lippincott Williams & Wilkins, 2017, 560 pages	Review/ Test/200 q	$78
B	*BRS Biochemistry, Molecular Biology, and Genetics*	Lieberman	Lippincott Williams & Wilkins, 2013, 432 pages	Review/Test	$54
B	*Case Files: Biochemistry*	Toy	McGraw-Hill, 2014, 480 pages	Cases	$35
B	*PreTest Biochemistry and Genetics*	Wilson	McGraw-Hill, 2017, 592 pages	Test/500 q	$38

Cell Biology and Histology

		AUTHOR	PUBLISHER	TYPE	PRICE
B+	*BRS Cell Biology and Histology*	Gartner	Lippincott Williams & Wilkins, 2018, 448 pages	Review/ Test/320 q	$54
B+	*Crash Course: Cell Biology and Genetics*	Stubbs	Elsevier, 2015, 216 pages	Review/Print + online	$47
B	*Wheater's Functional Histology*	Young	Elsevier, 2013, 464 pages	Text	$83

(c) ketabton.com: The Digital Library

Microbiology and Immunology

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁻	*Basic Immunology*	Abbas	Elsevier, 2019, 336 pages	Review	$70
A⁻	*Clinical Microbiology Made Ridiculously Simple*	Gladwin	MedMaster, 2019, 418 pages	Review	$38
A⁻	*Medical Microbiology and Immunology Flash Cards*	Rosenthal	Elsevier, 2016, 192 flash cards	Flash cards	$40
B⁺	*Lippincott Illustrated Reviews: Immunology*	Doan	Lippincott Williams & Wilkins, 2012, 384 pages	Reference/ Test/Few q	$75
B⁺	*Microcards: Microbiology Flash Cards*	Harpavat	Lippincott Williams & Wilkins, 2015, 312 flash cards	Flash cards	$53
B⁺	*Review of Medical Microbiology and Immunology*	Levinson	McGraw-Hill, 2018, 832 pages	Review/ Test/654 q	$63
B⁺	*How the Immune System Works*	Sompayrac	Wiley-Blackwell, 2019, 168 pages	Review	$50
B	*Case Studies in Immunology: Clinical Companion*	Geha	W. W. Norton & Company, 2016, 384 pages	Cases	$62
B	*Pretest: Microbiology*	Kettering	McGraw-Hill, 2013, 480 pages	Test/500 q	$38
B	*Case Files: Microbiology*	Toy	McGraw-Hill, 2014, 416 pages	Cases	$36
B	*Lange Microbiology and Infectious Diseases Flash Cards, 3e*	Somers	McGraw-Hill Education, 2017, 358 pages	Flash cards	$46
B⁻	*Lippincott Illustrated Reviews: Microbiology*	Cornelissen	Lippincott Williams & Wilkins, 2019, 448 pages	Review/Test/ Few q	$73

Pathology

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁺	*Pathoma: Fundamentals of Pathology*	Sattar	Pathoma, 2019, 218 pages	Review/ Lecture	$85–$120
A⁻	*Rapid Review: Pathology*	Goljan	Elsevier, 2018, 864 pages	Review/ Test/500 q	$65
A⁻	*Robbins and Cotran Review of Pathology*	Klatt	Elsevier, 2014, 504 pages	Test/1100 q	$55
A⁻	*Crash Course: Pathology*	Xiu	Elsevier, 2019, 438 pages	Review	$40
B	*High-Yield Histopathology*	Dudek	Lippincott Williams & Wilkins, 2017, 320 pages	Review	$36
B	*Pathophysiology of Disease: Introduction to Clinical Medicine*	Hammer	McGraw-Hill, 2018, 832 pages	Text	$90
B	*Haematology at a Glance*	Mehta	Blackwell Science, 2014, 136 pages	Review	$49
B	*Pocket Companion to Robbins and Cotran Pathologic Basis of Disease*	Mitchell	Elsevier, 2016, 896 pages	Review	$40
B	*BRS Pathology*	Schneider	Lippincott Williams & Wilkins, 2013, 480 pages	Review/ Test/450 q	$54

(c) ketabton.com: The Digital Library

Pharmacology

		AUTHOR	PUBLISHER	TYPE	PRICE
B+	*Crash Course: Pharmacology*	Battista	Elsevier, 2019, 336 pages	Review	$40
B+	*Master the Boards USMLE Step 1 Pharmacology Flashcards*	Fischer	Kaplan, 2015, 200 flash cards	Flash cards	$55
B+	*BRS Pharmacology*	Rosenfeld	Lippincott Williams & Wilkins, 2019, 384 pages	Review/ Test/200 q	$55
B	*Lange Pharmacology Flash Cards*	Baron	McGraw-Hill, 2017, 266 flash cards	Flash cards	$39
B	*Pharmacology Flash Cards*	Brenner	Elsevier, 2017, 230 flash cards	Flash cards	$45
B	*Katzung & Trevor's Pharmacology: Examination and Board Review*	Trevor	McGraw-Hill, 2018, 592 pages	Review/ Test/800 q	$54
B	*Lippincott Illustrated Reviews: Pharmacology*	Whalen	Lippincott Williams & Wilkins, 2018, 576 pages	Review/ Test/380 q	$75

Physiology

		AUTHOR	PUBLISHER	TYPE	PRICE
A−	*BRS Physiology*	Costanzo	Lippincott Williams & Wilkins, 2018, 304 pages	Review/ Test/350 q	$54
A−	*Pathophysiology of Heart Disease*	Lilly	Lippincott Williams & Williams, 2015, 480 pages	Review	$57
A−	*PreTest Physiology*	Metting	McGraw-Hill, 2013, 528 pages	Test/500 q	$38
A−	*Color Atlas of Physiology*	Silbernagl	Thieme, 2015, 472 pages	Review	$50
B+	*BRS Physiology Cases and Problems*	Costanzo	Lippincott Williams & Wilkins, 2012, 368 pages	Cases	$58
B+	*Physiology*	Costanzo	Saunders, 2017, 528 pages	Text	$60
B+	*Vander's Renal Physiology*	Eaton	McGraw-Hill, 2018, 224 pages	Text	$49
B+	*Acid-Base, Fluids, and Electrolytes Made Ridiculously Simple*	Preston	MedMaster, 2017, 166 pages	Review	$24
B+	*Pulmonary Pathophysiology: The Essentials*	West	Lippincott Williams & Wilkins, 2017, 264 pages	Review/ Test/75 q	$57
B	*Rapid Review: Physiology*	Brown	Elsevier, 2011, 384 pages	Test/350 q	$39
B	*Endocrine Physiology*	Molina	McGraw-Hill, 2018, 320 pages	Review	$59
B−	*Netter's Physiology Flash Cards*	Mulroney	Saunders, 2015, 450 flash cards	Flash cards	$40

(c) ketabton.com: The Digital Library

SECTION IV

Abbreviations and Symbols

ABBREVIATION	MEANING
1st MC*	1st metacarpal
A-a	alveolar-arterial [gradient]
AA	Alcoholics Anonymous, amyloid A
AAMC	Association of American Medical Colleges
AAo*	ascending aorta
Ab	antibody
ABPA	allergic bronchopulmonary aspergillosis
AC	adenylyl cyclase
ACA	anterior cerebral artery
Acetyl-CoA	acetyl coenzyme A
ACD	anemia of chronic disease
ACE	angiotensin-converting enzyme
ACh	acetylcholine
AChE	acetylcholinesterase
ACL	anterior cruciate ligament
ACom	anterior communicating [artery]
ACTH	adrenocorticotropic hormone
AD	Alzheimer disease, autosomal dominant
ADA	adenosine deaminase, Americans with Disabilities Act
ADH	antidiuretic hormone
ADHD	attention-deficit hyperactivity disorder
ADP	adenosine diphosphate
ADPKD	autosomal-dominant polycystic kidney disease
AFP	α-fetoprotein
Ag	antigen, silver
AICA	anterior inferior cerebellar artery
AIDS	acquired immunodeficiency syndrome
AIHA	autoimmune hemolytic anemia
AKI	acute kidney injury
AKT	protein kinase B
AL	amyloid light [chain]
ALA	aminolevulinate
ALI	acute lung injury
ALL	acute lymphoblastic (lymphocytic) leukemia
ALP	alkaline phosphatase
ALS	amyotrophic lateral sclerosis
ALT	alanine transaminase
AMA	American Medical Association, antimitochondrial antibody
AML	acute myelogenous (myeloid) leukemia
AMP	adenosine monophosphate
ANA	antinuclear antibody
ANCA	antineutrophil cytoplasmic antibody
ANOVA	analysis of variance

ABBREVIATION	MEANING
ANP	atrial natriuretic peptide
ANS	autonomic nervous system
Ant*	anterior
anti-CCP	anti-cyclic citrullinated peptide
Ao*	aorta
AOA	American Osteopathic Association
AP	action potential, A & P [ribosomal binding sites]
APC	antigen-presenting cell, activated protein C
Apo	apolipoprotein
APP	amyloid precursor protein
APRT	adenine phosphoribosyltransferase
aPTT	activated partial thromboplastin time
APUD	amine precursor uptake decarboxylase
AR	attributable risk, autosomal recessive, aortic regurgitation
ARB	angiotensin receptor blocker
ARDS	acute respiratory distress syndrome
Arg	arginine
ARPKD	autosomal-recessive polycystic kidney disease
ART	antiretroviral therapy
AS	aortic stenosis
ASA	anterior spinal artery
ASD	atrial septal defect
ASO	anti–streptolysin O
AST	aspartate transaminase
AT	angiotensin, antithrombin
ATN	acute tubular necrosis
ATP	adenosine triphosphate
ATPase	adenosine triphosphatase
ATTR	transthyretin-mediated amyloidosis
AUB	abnormal uterine bleeding
AV	atrioventricular
AZT	azidothymidine
BAL	British anti-Lewisite [dimercaprol]
BBB	blood-brain barrier
BCG	bacille Calmette-Guérin
BH_4	tetrahydrobiopterin
BM	basement membrane
BOOP	bronchiolitis obliterans organizing pneumonia
BP	bisphosphate, blood pressure
BPG	bisphosphoglycerate
BPH	benign prostatic hyperplasia
BT	bleeding time
BUN	blood urea nitrogen
Ca*	capillary

*Image abbreviation only

(c) ketabton.com: The Digital Library

ABBREVIATION	MEANING
Ca^{2+}	calcium ion
CAD	coronary artery disease
CAF	common application form
cAMP	cyclic adenosine monophosphate
CBG	corticosteroid-binding globulin
Cbm*	cerebellum
CBSE	Comprehensive Basic Science Examination
CBSSA	Comprehensive Basic Science Self-Assessment
CBT	computer-based test, cognitive behavioral therapy
CC*	corpus callosum
CCA*	common carotid artery
CCK	cholecystokinin
CCS	computer-based case simulation
CD	cluster of differentiation
CDK	cyclin-dependent kinase
cDNA	complementary deoxyribonucleic acid
CEA	carcinoembryonic antigen
CETP	cholesteryl-ester transfer protein
CF	cystic fibrosis
CFTR	cystic fibrosis transmembrane conductance regulator
CGD	chronic granulomatous disease
cGMP	cyclic guanosine monophosphate
CGRP	calcitonin gene–related peptide
C_H1–C_H3	constant regions, heavy chain [antibody]
ChAT	choline acetyltransferase
CHD*	common hepatic duct
χ^2	chi-squared
CI	confidence interval
CIN	candidate identification number, carcinoma in situ, cervical intraepithelial neoplasia
CIS	Communication and Interpersonal Skills
CK	clinical knowledge, creatine kinase
CKD	chronic kidney disease
CK-MB	creatine kinase, MB fraction
C_L	constant region, light chain [antibody]
CL	clearance
Cl^-	chloride ion
CLL	chronic lymphocytic leukemia
CMC	carpometacarpal (joint)
CML	chronic myelogenous (myeloid) leukemia
CMV	cytomegalovirus
CN	cranial nerve
CN^-	cyanide ion
CNS	central nervous system
CNV	copy number variation
CO	carbon monoxide, cardiac output
CO_2	carbon dioxide
CoA	coenzyme A
COL1A1	collagen, type I, alpha 1
COL1A2	collagen, type I, alpha 2
COMT	catechol-O-methyltransferase
COP	coat protein
COPD	chronic obstructive pulmonary disease
CoQ	coenzyme Q

ABBREVIATION	MEANING
COX	cyclooxygenase
C_p	plasma concentration
CPAP	continuous positive airway pressure
CPR	cardiopulmonary resuscitation
Cr	creatinine
CRC	colorectal cancer
CREST	calcinosis, Raynaud phenomenon, esophageal dysfunction, sclerosis, and telangiectasias [syndrome]
CRH	corticotropin-releasing hormone
CRP	C-reactive protein
CS	clinical skills
C-section	cesarean section
CSF	cerebrospinal fluid
CT	computed tomography
CTP	cytidine triphosphate
CXR	chest x-ray
DA	dopamine
DAF	decay-accelerating factor
DAG	diacylglycerol
dATP	deoxyadenosine triphosphate
DCIS	ductal carcinoma in situ
DCT	distal convoluted tubule
ddI	didanosine
DES	diethylstilbestrol
DH	dehydrogenase
DHAP	dihydroxyacetone phosphate
DHEA	dehydroepiandrosterone
DHF	dihydrofolic acid
DHT	dihydrotestosterone
DI	diabetes insipidus
DIC	disseminated intravascular coagulation
DIP	distal interphalangeal [joint]
DKA	diabetic ketoacidosis
DLCO	diffusing capacity for carbon monoxide
DM	diabetes mellitus
DNA	deoxyribonucleic acid
DNR	do not resuscitate
dNTP	deoxynucleotide triphosphate
DO	doctor of osteopathy
DPGN	diffuse proliferative glomerulonephritis
DPM	doctor of podiatric medicine
DPP-4	dipeptidyl peptidase-4
DPPC	dipalmitoylphosphatidylcholine
DS	double stranded
dsDNA	double-stranded deoxyribonucleic acid
dsRNA	double-stranded ribonucleic acid
DRG	dorsal root ganglion
d4T	didehydrodeoxythymidine [stavudine]
dTMP	deoxythymidine monophosphate
DTR	deep tendon reflex
DTs	delirium tremens
dUDP	deoxyuridine diphosphate
dUMP	deoxyuridine monophosphate
DVT	deep venous thrombosis
E*	euthromatin, esophagus

*Image abbreviation only

(c) ketabton.com: The Digital Library

ABBREVIATION	MEANING
EBV	Epstein-Barr virus
ECA*	external carotid artery
ECF	extracellular fluid
ECFMG	Educational Commission for Foreign Medical Graduates
ECG	electrocardiogram
ECL	enterochromaffin-like [cell]
ECM	extracellular matrix
ECT	electroconvulsive therapy
ED_{50}	median effective dose
EDRF	endothelium-derived relaxing factor
EDTA	ethylenediamine tetra-acetic acid
EDV	end-diastolic volume
EEG	electroencephalogram
EF	ejection fraction
EGF	epidermal growth factor
EHEC	enterohemorrhagic *E coli*
EIEC	enteroinvasive *E coli*
ELISA	enzyme-linked immunosorbent assay
EM	electron micrograph/microscopy
EMB	eosin–methylene blue
EPEC	eneteropathogenic *E coli*
Epi	epinephrine
EPO	erythropoietin
EPS	extrapyramidal system
ER	endoplasmic reticulum, estrogen receptor
ERAS	Electronic Residency Application Service
ERCP	endoscopic retrograde cholangiopancreatography
ERP	effective refractory period
eRPF	effective renal plasma flow
ERT	estrogen replacement therapy
ERV	expiratory reserve volume
ESR	erythrocyte sedimentation rate
ESRD	end-stage renal disease
ESV	end-systolic volume
ETEC	enterotoxigenic *E coli*
EtOH	ethyl alcohol
EV	esophageal vein
F	bioavailability
FA	fatty acid
Fab	fragment, antigen-binding
FAD	flavin adenine dinucleotide
$FADH_2$	reduced flavin adenine dinucleotide
FAP	familial adenomatous polyposis
F1,6BP	fructose-1,6-bisphosphate
F2,6BP	fructose-2,6-bisphosphate
FBPase	fructose bisphosphatase
FBPase-2	fructose bisphosphatase-2
Fc	fragment, crystallizable
FcR	Fc receptor
5f-dUMP	5-fluorodeoxyuridine monophosphate
Fe^{2+}	ferrous ion
Fe^{3+}	ferric ion
Fem*	femur
FENa	excreted fraction of filtered sodium

ABBREVIATION	MEANING
FEV_1	forced expiratory volume in 1 second
FF	filtration fraction
FFA	free fatty acid
FGF	fibroblast growth factor
FGFR	fibroblast growth factor receptor
FISH	fluorescence in situ hybridization
FIT	fecal immunochemical testing
FKBP	FK506 binding protein
fMet	formylmethionine
FMG	foreign medical graduate
FMN	flavin mononucleotide
FN	false negative
FP, FP*	false positive, foot process
FRC	functional residual capacity
FSH	follicle-stimulating hormone
FSMB	Federation of State Medical Boards
FTA-ABS	fluorescent treponemal antibody—absorbed
FTD*	frontotemporal dementia
5-FU	5-fluorouracil
FVC	forced vital capacity
GABA	γ-aminobutyric acid
GAG	glycosaminoglycan
Gal	galactose
GBM	glomerular basement membrane
GC	glomerular capillary
G-CSF	granulocyte colony-stimulating factor
GERD	gastroesophageal reflux disease
GFAP	glial fibrillary acid protein
GFR	glomerular filtration rate
GGT	γ-glutamyl transpeptidase
GH	growth hormone
GHB	γ-hydroxybutyrate
GHRH	growth hormone–releasing hormone
G_I	G protein, I polypeptide
GI	gastrointestinal
GIP	gastric inhibitory peptide
GIST	gastrointestinal stromal tumor
GLUT	glucose transporter
GM	granulocyte macrophage
GM-CSF	granulocyte-macrophage colony stimulating factor
GMP	guanosine monophosphate
GnRH	gonadotropin-releasing hormone
GP	glycoprotein
G6P	glucose-6-phosphate
G6PD	glucose-6-phosphate dehydrogenase
GPe	globus pallidus externa
GPi	globus pallidus interna
GPI	glycosyl phosphatidylinositol
GRP	gastrin-releasing peptide
G_S	G protein, S polypeptide
GSH	reduced glutathione
GSSG	oxidized glutathione
GTP	guanosine triphosphate
GTPase	guanosine triphosphatase

*Image abbreviation only

(c) ketabton.com: The Digital Library

ABBREVIATION	MEANING
GU	genitourinary
H*	heterochromatin
H^+	hydrogen ion
H_1, H_2	histamine receptors
H_2S	hydrogen sulfide
HAV	hepatitis A virus
HAVAb	hepatitis A antibody
Hb	hemoglobin
HBcAb/HBcAg	hepatitis B core antibody/antigen
HBeAb/HBeAg	hepatitis B early antibody/antigen
HBsAb/HBsAg	hepatitis B surface antibody/antigen
$HbCO_2$	carbaminohemoglobin
HBV	hepatitis B virus
HCC	hepatocellular carcinoma
hCG	human chorionic gonadotropin
HCO_3^-	bicarbonate
Hct	hematocrit
HCTZ	hydrochlorothiazide
HCV	hepatitis C virus
HDL	high-density lipoprotein
HDN	hemolytic disease of the newborn
HDV	hepatitis D virus
H&E	hematoxylin and eosin
HEV	hepatitis E virus
HF	heart failure
Hfr	high-frequency recombination [cell]
HFpEF	heart failure with preserved ejection fraction
HFrEF	heart failure with reduced ejection fraction
HGPRT	hypoxanthine-guanine phosphoribosyltransferase
HHb	deoxygenated hemoglobin
HHS	hyperosmolar hyperglycemic state
HHV	human herpesvirus
5-HIAA	5-hydroxyindoleacetic acid
HIT	heparin-induced thrombocytopenia
HIV	human immunodeficiency virus
HL	hepatic lipase
HLA	human leukocyte antigen
HMG-CoA	hydroxymethylglutaryl-coenzyme A
HMP	hexose monophosphate
HMWK	high-molecular-weight kininogen
HNPCC	hereditary nonpolyposis colorectal cancer
hnRNA	heterogeneous nuclear ribonucleic acid
H_2O_2	hydrogen peroxide
HOCM	hypertrophic obstructive cardiomyopathy
HPA	hypothalamic-pituitary-adrenal [axis]
HPL	human placental lactogen
HPO	hypothalamic-pituitary-ovarian [axis]
HPV	human papillomavirus
HR	heart rate
HSP	Henoch-Schönlein purpura
HSV	herpes simplex virus
5-HT	5-hydroxytryptamine (serotonin)
HTLV	human T-cell leukemia virus
HTN	hypertension

ABBREVIATION	MEANING
HUS	hemolytic-uremic syndrome
HVA	homovanillic acid
IBD	inflammatory bowel disease
IBS	irritable bowel syndrome
IC	inspiratory capacity, immune complex
I_{Ca}	calcium current [heart]
I_f	funny current [heart]
ICA	internal carotid artery
ICAM	intercellular adhesion molecule
ICD	implantable cardioverter defibrillator
ICE	Integrated Clinical Encounter
ICF	intracellular fluid
ICP	intracranial pressure
ID	identification
ID_{50}	median infective dose
IDL	intermediate-density lipoprotein
IF	immunofluorescence, initiation factor
IFN	interferon
Ig	immunoglobulin
IGF	insulin-like growth factor
I_K	potassium current [heart]
IL	interleukin
IM	intramuscular
IMA	inferior mesenteric artery
IMG	international medical graduate
IMP	inosine monophosphate
IMV	inferior mesenteric vein
I_{Na}	sodium current [heart]
INH	isoniazid
INO	internuclear ophthalmoplegia
INR	International Normalized Ratio
IO	inferior oblique [muscle]
IOP	intraocular pressure
IP_3	inositol triphosphate
IPV	inactivated polio vaccine
IR	current × resistance [Ohm's law], inferior rectus [muscle]
IRV	inspiratory reserve volume
ITP	idiopathic thrombocytopenic purpura
IUD	intrauterine device
IUGR	intrauterine growth restriction
IV	intravenous
IVC	inferior vena cava
IVDU	intravenous drug use
IVIG	intravenous immunoglobulin
JAK/STAT	Janus kinase/signal transducer and activator of transcription [pathway]
JGA	juxtaglomerular apparatus
JVD	jugular venous distention
JVP	jugular venous pulse
K^+	potassium ion
KatG	catalase-peroxidase produced by *M tuberculosis*
K_e	elimination constant
K_f	filtration constant
KG	ketoglutarate

*Image abbreviation only

(c) ketabton.com: The Digital Library

ABBREVIATION	MEANING
K_m	Michaelis-Menten constant
KOH	potassium hydroxide
L	left, liver
LA	left atrial, left atrium
LAD	left anterior descending coronary artery
LAP	leukocyte alkaline phosphatase
Lat cond*	lateral condyle
Lb*	lamellar body
LCA	left coronary artery
LCAT	lecithin-cholesterol acyltransferase
LCC*	left common carotid artery
LCFA	long-chain fatty acid
LCL	lateral collateral ligament
LCME	Liaison Committee on Medical Education
LCMV	lymphocytic choriomeningitis virus
LCX	left circumflex coronary artery
LD	loading dose
LD_{50}	median lethal dose
LDH	lactate dehydrogenase
LDL	low-density lipoprotein
LES	lower esophageal sphincter
LFA	leukocyte function–associated antigen
LFT	liver function test
LH	luteinizing hormone
LLL*	left lower lobe (of lung)
LLQ	left lower quadrant
LM	lateral meniscus, left main coronary artery, light microscopy
LMN	lower motor neuron
LOS	lipooligosaccharide
LPA*	left pulmonary artery
LPL	lipoprotein lipase
LPS	lipopolysaccharide
LR	lateral rectus [muscle]
LT	labile toxin, leukotriene
LUL*	left upper lobe (of lung)
LV	left ventricle, left ventricular
M_1-M_5	muscarinic (parasympathetic) ACh receptors
MAC	membrane attack complex, minimum alveolar concentration
MALT	mucosa-associated lymphoid tissue
MAO	monoamine oxidase
MAOI	monoamine oxidase inhibitor
MAP	mean arterial pressure, mitogen-activated protein
Max*	maxillary sinus
MC	midsystolic click
MCA	middle cerebral artery
MCAT	Medical College Admissions Test
MCHC	mean corpuscular hemoglobin concentration
MCL	medial collateral ligament
MCP	metacarpophalangeal [joint]
MCV	mean corpuscular volume
MD	maintenance dose
MDD	major depressive disorder
Med cond*	medial condyle

ABBREVIATION	MEANING
MELAS syndrome	mitochondrial encephalopathy, lactic acidosis, and stroke-like episodes
MEN	multiple endocrine neoplasia
Mg^{2+}	magnesium ion
$MgSO_4$	magnesium sulfate
MGUS	monoclonal gammopathy of undetermined significance
MHC	major histocompatibility complex
MI	myocardial infarction
MIF	müllerian inhibiting factor
MIRL	membrane inhibitor of reactive lysis
MLCK	myosin light-chain kinase
MLF	medial longitudinal fasciculus
MMC	migrating motor complex
MMR	measles, mumps, rubella [vaccine]
6-MP	6-mercaptopurine
MPGN	membranoproliferative glomerulonephritis
MPO	myeloperoxidase
MPO-ANCA/ p-ANCA	myeloperoxidase/perinuclear antineutrophil cytoplasmic antibody
MR	medial rectus [muscle], mitral regurgitation
MRI	magnetic resonance imaging
miRNA	microribonucleic acid
mRNA	messenger ribonucleic acid
MRSA	methicillin-resistant *S aureus*
MS	mitral stenosis, multiple sclerosis
MSH	melanocyte-stimulating hormone
mtDNA	mitochondrial DNA
mTOR	mammalian target of rapamycin
MTP	metatarsophalangeal [joint]
MTX	methotrexate
MVO_2	myocardial oxygen consumption
MVP	mitral valve prolapse
N*	nucleus
Na^+	sodium ion
NAT	nucleic acid testing
NAD	nicotinamide adenine dinucleotide
NAD^+	oxidized nicotinamide adenine dinucleotide
NADH	reduced nicotinamide adenine dinucleotide
$NADP^+$	oxidized nicotinamide adenine dinucleotide phosphate
NADPH	reduced nicotinamide adenine dinucleotide phosphate
NBME	National Board of Medical Examiners
NBOME	National Board of Osteopathic Medical Examiners
NBPME	National Board of Podiatric Medical Examiners
NE	norepinephrine
NF	neurofibromatosis
NFAT	nuclear factor of activated T-cell
NH_3	ammonia
NH_4^+	ammonium
NK	natural killer [cells]
N_M	muscarinic ACh receptor in neuromuscular junction
NMDA	*N*-methyl-d-aspartate
NMJ	neuromuscular junction
NMS	neuroleptic malignant syndrome
N_N	nicotinic ACh receptor in autonomic ganglia

*Image abbreviation only

(c) ketabton.com: The Digital Library

ABBREVIATION	MEANING
NRMP	National Residency Matching Program
NNRTI	non-nucleoside reverse transcriptase inhibitor
NO	nitric oxide
N_2O	nitrous oxide
NPH	neutral protamine Hagedorn, normal pressure hydrocephalus
NPV	negative predictive value
NRTI	nucleoside reverse transcriptase inhibitor
NSAID	nonsteroidal anti-inflammatory drug
NSE	neuron-specific enolase
NSTEMI	non–ST-segment elevation myocardial infarction
Nu*	nucleolus
OAA	oxaloacetic acid
OCD	obsessive-compulsive disorder
OCP	oral contraceptive pill
ODC	oxygen-hemoglobin dissociation curve
OH	hydroxy
1,25-OH D_3	calcitriol (active form of vitamin D)
25-OH D_3	storage form of vitamin D
OPV	oral polio vaccine
OR	odds ratio
OS	opening snap
OSA	obstructive sleep apnea
OVLT	organum vasculosum of the lamina terminalis
P-body	processing body (cytoplasmic)
P-450	cytochrome P-450 family of enzymes
PA	posteroanterior, pulmonary artery
PABA	*para*-aminobenzoic acid
$Paco_2$	arterial Pco_2
$PAco_2$	alveolar Pco_2
PAH	*para*-aminohippuric acid
PAN	polyarteritis nodosa
Pao_2	partial pressure of oxygen in arterial blood
PAo_2	partial pressure of oxygen in alveolar blood
PAP	Papanicolaou [smear], prostatic acid phosphatase
PAPPA	pregnancy-associated plasma protein A
PAS	periodic acid–Schiff
Pat*	patella
PBP	penicillin-binding protein
PC	platelet count, pyruvate carboxylase
PCA	posterior cerebral artery
PCC	prothrombin complex concentrate
PCL	posterior cruciate ligament
Pco_2	partial pressure of carbon dioxide
PCom	posterior communicating [artery]
PCOS	polycystic ovarian syndrome
PCP	phencyclidine hydrochloride, *Pneumocystis jirovecii* pneumonia
PCR	polymerase chain reaction
PCT	proximal convoluted tubule
PCV13	pneumococcal conjugate vaccine
PCWP	pulmonary capillary wedge pressure
PDA	patent ductus arteriosus, posterior descending artery
PDE	phosphodiesterase

ABBREVIATION	MEANING
PDGF	platelet-derived growth factor
PDH	pyruvate dehydrogenase
PE	pulmonary embolism
PECAM	platelet–endothelial cell adhesion molecule
$PEco_2$	expired air Pco_2
PEP	phosphoenolpyruvate
PF	platelet factor
PFK	phosphofructokinase
PFK-2	phosphofructokinase-2
PFT	pulmonary function test
PG	phosphoglycerate
P_i	plasma interstitial osmotic pressure, inorganic phosphate
PICA	posterior inferior cerebellar artery
PID	pelvic inflammatory disease
PIo_2	Po_2 in inspired air
PIP	proximal interphalangeal [joint]
PIP_2	phosphatidylinositol 4,5-bisphosphate
PIP_3	phosphatidylinositol 3,4,5-bisphosphate
PKD	polycystic kidney disease
PKR	interferon-α–induced protein kinase
PKU	phenylketonuria
PLP	pyridoxal phosphate
PML	progressive multifocal leukoencephalopathy
PMN	polymorphonuclear [leukocyte]
P_{net}	net filtration pressure
PNET	primitive neuroectodermal tumor
PNS	peripheral nervous system
Po_2	partial pressure of oxygen
PO_4^{3-}	phosphate
Pop*	popliteal artery
Pop a*	popliteal artery
Post*	posterior
PPAR	peroxisome proliferator-activated receptor
PPD	purified protein derivative
PPI	proton pump inhibitor
PPM	parts per million
PPSV23	pneumococcal polysaccharide vaccine
PPV	positive predictive value
PR3-ANCA/ c-ANCA	cytoplasmic antineutrophil cytoplasmic antibody
PrP	prion protein
PRPP	phosphoribosylpyrophosphate
PSA	prostate-specific antigen
PSS	progressive systemic sclerosis
PT	prothrombin time
PTEN	phosphatase and tensin homolog
PTH	parathyroid hormone
PTHrP	parathyroid hormone–related protein
PTSD	post-traumatic stress disorder
PTT	partial thromboplastin time
PV	plasma volume, venous pressure
Pv*	pulmonary vein
PVC	polyvinyl chloride
PVR	pulmonary vascular resistance

*Image abbreviation only

(c) ketabton.com: The Digital Library

ABBREVIATION	MEANING
R	correlation coefficient, right, R variable [group]
R_3	Registration, Ranking, & Results [system]
RA	right atrium
RAAS	renin-angiotensin-aldosterone system
RANK-L	receptor activator of nuclear factor-κ B ligand
RAS	reticular activating system
RBF	renal blood flow
RCA	right coronary artery
REM	rapid eye movement
RER	rough endoplasmic reticulum
Rh	*rhesus* antigen
RLL*	right lower lobe (of lungs)
RLQ	right lower quadrant
RML*	right middle lobe (of lung)
RNA	ribonucleic acid
RNP	ribonucleoprotein
ROS	reactive oxygen species
RPF	renal plasma flow
RPGN	rapidly progressive glomerulonephritis
RPR	rapid plasma reagin
RR	relative risk, respiratory rate
rRNA	ribosomal ribonucleic acid
RS	Reed-Sternberg [cells]
RSC*	right subclavian artery
RSV	respiratory syncytial virus
RTA	renal tubular acidosis
RUL*	right upper lobe (of lung)
RUQ	right upper quadrant
RV	residual volume, right ventricle, right ventricular
RVH	right ventricular hypertrophy
[S]	substrate concentration
SA	sinoatrial
SAA	serum amyloid–associated [protein]
SAM	*S*-adenosylmethionine
SARS	severe acute respiratory syndrome
SCC	squamous cell carcinoma
SCD	sudden cardiac death
SCID	severe combined immunodeficiency disease
SCJ	squamocolumnar junction
SCM	sternocleidomastoid muscle
SCN	suprachiasmatic nucleus
SD	standard deviation
SE	standard error [of the mean]
SEP	Spoken English Proficiency
SER	smooth endoplasmic reticulum
SERM	selective estrogen receptor modulator
SGLT	sodium-glucose transporter
SHBG	sex hormone–binding globulin
SIADH	syndrome of inappropriate [secretion of] antidiuretic hormone
SIDS	sudden infant death syndrome
SJS	Stevens-Johnson syndrome
SLE	systemic lupus erythematosus
SLL	small lymphocytic lymphoma
SLT	Shiga-like toxin
SMA	superior mesenteric artery
SMX	sulfamethoxazole
SNARE	soluble NSF attachment protein receptor
SNc	substantia nigra pars compacta
SNP	single nucleotide polymorphism
SNr	substantia nigra pars reticulata
SNRI	serotonin and norepinephrine receptor inhibitor
snRNA	small nuclear RNA
snRNP	small nuclear ribonucleoprotein
SO	superior oblique [muscle]
SOAP	Supplemental Offer and Acceptance Program
Sp*	spleen
spp	species
SR	superior rectus [muscle]
SS	single stranded
ssDNA	single-stranded deoxyribonucleic acid
SSPE	subacute sclerosing panencephalitis
SSRI	selective serotonin reuptake inhibitor
ssRNA	single-stranded ribonucleic acid
St*	stomach
ST	Shiga toxin
StAR	steroidogenic acute regulatory protein
STEMI	ST-segment elevation myocardial infarction
STI	sexually transmitted infection
STN	subthalamic nucleus
SV	splenic vein, stroke volume
SVC	superior vena cava
SVR	systemic vascular resistance
SVT	supraventricular tachycardia
T*	trachea
$t_{1/2}$	half-life
T_3	triiodothyronine
T_4	thyroxine
TAPVR	total anomalous pulmonary venous return
TB	tuberculosis
TBG	thyroxine-binding globulin
TBV	total blood volume
3TC	dideoxythiacytidine [lamivudine]
TCA	tricarboxylic acid [cycle], tricyclic antidepressant
Tc cell	cytotoxic T cell
TCR	T-cell receptor
TDF	tenofovir disoproxil fumarate
TdT	terminal deoxynucleotidyl transferase
TE	tracheoesophageal
TFT	thyroid function test
TG	triglyceride
TGF	transforming growth factor
Th cell	helper T cell
THF	tetrahydrofolic acid
TI	therapeutic index
TIA	transient ischemic attack
Tib*	tibia
TIBC	total iron-binding capacity
TIPS	transjugular intrahepatic portosystemic shunt

*Image abbreviation only

(c) ketabton.com: The Digital Library

ABBREVIATION	MEANING
TLC	total lung capacity
T_m	maximum rate of transport
TMP	trimethoprim
TN	true negative
TNF	tumor necrosis factor
TNM	tumor, node, metastases [staging]
TOP	topoisomerase
ToRCHeS	*Toxoplasma gondii*, rubella, CMV, HIV, HSV-2, syphilis
TP	true positive
tPA	tissue plasminogen activator
TPO	thyroid peroxidase, thrombopoietin
TPP	thiamine pyrophosphate
TPPA	*Treponema pallidum* particle agglutination assay
TPR	total peripheral resistance
TR	tricuspid regurgitation
TRAP	tartrate-resistant acid phosphatase
TRECs	T-cell receptor excision circles
TRH	thyrotropin-releasing hormone
tRNA	transfer ribonucleic acid
TSH	thyroid-stimulating hormone
TSI	triple sugar iron
TSS	toxic shock syndrome
TSST	toxic shock syndrome toxin
TTP	thrombotic thrombocytopenic purpura
TTR	transthyretin
TV	tidal volume
TXA_2	thromboxane A_2
UDP	uridine diphosphate
UMN	upper motor neuron
UMP	uridine monophosphate
UPD	uniparental disomy
URI	upper respiratory infection
USMLE	United States Medical Licensing Examination
UTI	urinary tract infection
UTP	uridine triphosphate

ABBREVIATION	MEANING
UV	ultraviolet
V_1, V_2	vasopressin receptors
VC	vital capacity
V_d	volume of distribution
VD	physiologic dead space
V(D)J	variable, (diversity), joining gene segments rearranged to form Ig genes
VDRL	Venereal Disease Research Laboratory
VEGF	vascular endothelial growth factor
V_H	variable region, heavy chain [antibody]
VHL	von Hippel-Lindau [disease]
VIP	vasoactive intestinal peptide
VIPoma	vasoactive intestinal polypeptide-secreting tumor
VJ	light-chain hypervariable region [antibody]
V_L	variable region, light chain [antibody]
VLCFA	very-long-chain fatty acids
VLDL	very low density lipoprotein
VMA	vanillylmandelic acid
VMAT	vesicular monoamine transporter
V_{max}	maximum velocity
VPL	ventral posterior nucleus, lateral
VPM	ventral posterior nucleus, medial
VPN	vancomycin, polymyxin, nystatin [media]
$\dot{V}/\dot{Q}$	ventilation/perfusion [ratio]
VRE	vancomycin-resistant enterococcus
VSD	ventricular septal defect
V_T	tidal volume
VTE	venous thromboembolism
vWF	von Willebrand factor
VZV	varicella-zoster virus
VMAT	vesicular monoamine transporter
XR	X-linked recessive
XX/XY	normal complement of sex chromosomes for female/male
ZDV	zidovudine [formerly AZT]

*Image abbreviation only

(c) ketabton.com: The Digital Library

SECTION IV

Image Acknowledgments

In this edition, in collaboration with MedIQ Learning, LLC, and a variety of other partners, we are pleased to include the following clinical images and diagrams for the benefit of integrative student learning.

Portions of this book identified with the symbol Rx are copyright © USMLE-Rx.com (MedIQ Learning, LLC).

Portions of this book identified with the symbol RU are copyright © Dr. Richard Usatine and are provided under license through MedIQ Learning, LLC.

Portions of this book identified with the symbol * are listed below by page number.

This symbol refers to material that is available in the public domain.

This symbol refers to the Creative Commons Attribution license, full text at http://creativecommons.org/licenses/by/4.0/legalcode.

This symbol refers to the Creative Commons Attribution-Share Alike license, full text at: http://creativecommons.org/licenses/by-sa/4.0/legalcode.

Biochemistry

34 **Chromatin structure.** Electron micrograph showing heterochromatin, euchromatin, and nucleolus. This image is a derivative work, adapted from the following source, available under : Roller RA, Rickett JD, Stickle WB. The hypobranchial gland of the estuarine snail *Stramonita haemastoma canaliculata* (Gray) (Prosobranchia: Muricidae): a light and electron microscopical study. *Am Malac Bull.* 1995;11(2):177-190. Available at https://archive.org/details/americanm101119931994amer.

49 **Cilia structure: Image A.** Nine doublet + 2 singlet arrangement of microtubule. Courtesy of Louisa Howard and Michael Binder. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

49 **Cilia structure: Image B.** Cilia structure of basal body. This image is a derivative work, adapted from the following source, available under : Riparbelli MG, Cabrera OA, Callaini G, et al. Unique properties of *Drosophila* spermatocyte primary cilia. *Biol Open.* 2013 Nov 15; 2(11): 1137–1147. DOI: 10.1242/bio.20135355.

49 **Cilia structure: Image C.** Dextrocardia. This image is a derivative work, adapted from the following source, available under : Oluwadare O, Ayoka AO, Akomolafe RO, et al. The role of electrocardiogram in the diagnosis of dextrocardia with mirror image atrial arrangement and ventricular position in a young adult Nigerian in Ile-Ife: a case report. *J Med Case Rep.* 2015;9:222. DOI: 10.1186/s13256-015-0695-4.

51 **Osteogenesis imperfecta: Image A.** Skeletal deformities in upper extremity of child. This image is a derivative work, adapted from the following source, available under : Vanakker OM, Hemelsoet D, De Paepe. Hereditary connective tissue diseases in young adult stroke: a comprehensive synthesis. *Stroke Res Treat.* 2011;712903. DOI: 10.4061/2011/712903. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

51 **Osteogenesis imperfecta: Image B.** Blue sclera. This image is a derivative work, adapted from the following source, available under : Wheatley K, Heng EL, Sheppard M, et al. A case of spontaneous intestinal perforation in osteogenesis imperfects. *J Clin Med Res.* 2010;2(4):198–200. DOI: 10.4021/jocmr369w.

51 **Ehlers-Danlos syndrome: Images A and B.** Hyperextensibility of skin and DIP joint. This image is a derivative work, adapted from the following source, available under : Whitaker JK, Alexander, P, Chau DYS, et al. Severe conjunctivochalasis in association with classic type Ehlers-Danlos syndrome. *BMC Ophthalmol.* 2012;2:47. DOI: 10.1186/1471-2415-12-47.

52 **Elastin: Image A.** Pes escavatum. This image is a derivative work, adapted from the following source, available under : De Maio F, Fichera A, De Luna V, et al. Orthopaedic aspects of Marfan syndrome: the experience of a referral center for diagnosis of rare diseases. *Adv Orthop.* 2016; 2016: 8275391. DOI 10.1155/2016/8275391.

55 **Karyotyping.** Paar C, Herber G, Voskova, et al. This image is a derivative work, adapted from the following source, available under : A case of acute myeloid leukemia (AML) with an unreported combination of chromosomal abnormalities: gain of isochromosome 5p, tetrasomy 8 and unbalanced translocation der(19)t(17;19)(q23;p13). *Mol Cytogenet.* 2013;6:40. DOI: 10.1186/1755-8166-6-40.

55 **Fluorescence in situ hybridization.** This image is a derivative work, adapted from the following source, available under : Paar C, Herber G, Voskova, et al. A case of acute myeloid leukemia (AML) with an unreported combination of chromosomal abnormalities: gain of isochromosome 5p, tetrasomy 8 and unbalanced translocation der(19)t(17;19)(q23;p13). *Mol Cytogenet.* 2013;6:40. DOI: 10.1186/1755-8166-6-40.

57 **Genetic terms.** Café-au-lait spots. This image is a derivative work, adapted from the following source, available under : Dumitrescu CE and Collins MT. *Orphanet J Rare Dis.* 2008;3:12. DOI: 10.1186/1750-1172-3-12.

61 **Muscular dystrophies: Image A.** Fibrofatty replacement of muscle. Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

64 **Williams syndrome.** This image is a derivative work, adapted from the following source, available under : Mazumdar J, Sarkar R, Badveli A, et al. Double chamber right ventricle in Williams syndrome: a rare cardiac anomaly reported. *Springerplus.* 2016; 5: 275. DOI: 10.1186/s40064-016-1897-y.

66 **Vitamin A.** Bitot sponts on conjunctiva. This image is a derivative work, adapted from the following source, available under : Baiyeroju A, Bowman R, Gilbert C, et al. Managing eye health in young children. *Comm Eye Health.* 2010;23(72):4-11. Available at https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2873666.

(c) ketabton.com: The Digital Library

67 **Vitamin B_3.** Pellagra. This image is a derivative work, adapted from the following source, available under CC BY: van Dijk HA, Fred H. Images of memorable cases: case 2. Connexions Web site. Dec 4, 2008. Available at: http://cnx.org/contents/3d3dcb2e-8e98-496f-91c2-fe94e93428a1@3@3/.

70 **Vitamin D.** X-ray of lower extremity in child with rickets. This image is a derivative work, adapted from the following source, available under CC BY-NC-SA: Dr. Michael L. Richardson. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-NC-SA.

71 **Protein-energy malnutrition: Image A.** Child with kwashiorkor. Courtesy of the Department of Health and Human Services and Dr. Lyle Conrad.

71 **Protein-energy malnutrition: Image B.** Child with marasmus. Courtesy of the Department of Health and Human Services.

84 **Alkaptonuria.** Pigment granules on dorsum of hand. This image is a derivative work, adapted from the following source, available under CC BY: Vasudevan B, Sawhney MPS, Radhakrishnan S. Alkaptonuria associated with degenerative collagenous palmar plaques. *Indian J Dermatol.* 2009;54:299-301. DOI: 10.4103/0019-5154.55650.

85 **Cystinuria.** Hexagonal cystine stones in urine. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of Cayla Devine.

88 **Lysosomal storage diseases: Image A.** "Cherry-red" spot on macula in Tay-Sachs disease. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of Dr. Jonathan Trobe.

88 **Lysosomal storage diseases: Image B.** Angiokeratomas. This image is a derivative work, adapted from the following source, available under CC BY: Burlina AP, Sims KB, Politei JM, et al. Early diagnosis of peripheral nervous system involvement in Fabry disease and treatment of neuropathic pain: the report of an expert panel. *BMC Neurol.* 2011;11:61. DOI: 10.1186/1471-2377-11-61. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

88 **Lysosomal storage diseases: Image C.** Gaucher cells in Gaucher disease. This image is a derivative work, adapted from the following source, available under CC BY: Sokołowska B, Skomra D, Czartoryska B. et al. Gaucher disease diagnosed after bone marrow trephine biopsy—a report of two cases. *Folia Histochem Cytobiol.* 2011;49:352-356. DOI: 10.5603/FHC.2011.0048. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

88 **Lysosomal storage diseases: Image D.** Foam cells in Niemann-Pick disease. This image is a derivative work, adapted from the following source, available under CC BY: Prieto-Potin I, Roman-Blas JA, Martinez-Calatrava MJ, et al. Hypercholesterolemia boosts joint destruction in chronic arthritis. An experimental model aggravated by foam macrophage infiltration. *Arthritis Res Ther.* 2013;15:R81. DOI: 10.1186/ar4261.

Immunology

96 **Lymph node: Images A and B.** Lymph node histology. This image is a derivative work, adapted from the following source, available under CC BY: Navid Golpur.

98 **Spleen.** Red and white pulp. This image is a derivative work, adapted from the following source, available under CC BY: Heinrichs S, Conover LF, Bueso-Ramos CE, et al. MYBL2 is a sub-haploinsufficient tumor suppressor gene in myeloid malignancy. *eLife.* 2013;2:e00825. DOI: 10.7554/eLife.00825. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

98 **Thymus: Image A.** Hassall corpuscles. This image is a derivative work, adapted from the following source, available under CC BY: Minato H, Kinoshita E, Nakada S, et al. Thymic lymphoid hyperplasia with multilocular thymic cysts diagnosed before the Sjögren syndrome diagnosis. *Diagn Pathol.* 2015;10:103. DOI: 10.1186/s13000-015-0332-y.

98 **Thymus: Image B.** "Sail sign" on x-ray of normal thymus in neonate. This image is a derivative work, adapted from the following source, available under CC BY: Di Serafino M, Esposito F, Severino R, et al. Think thymus, think well: the chest x-ray thymic signs. *J Pediatr Moth Care.* 2016;1(2):108-109. DOI: 10.19104/japm.2016.108.

107 **Complement disorders.** Paroxysmal nocturnal hemoglobinuria. This image is a derivative work, adapted from the following source, available under CC BY: Nakamura N, Sugawara T, Shirato K, et al. *J Med Case Reports.* 2011;5:550. doi: 10.1186/1752-1947-5-550

117 **Immunodeficiencies: Image A.** Spider angioma (telangiectasia). This image is a derivative work, adapted from the following source, available under CC BY: Liapakis IE, Englander M, Sinani R, et al. Management of facial telangiectasias with hand cautery. *World J Plast Surg.* 2015 Jul;4(2):127-133.

117 **Immunodeficiencies: Image B.** Giant granules in granulocytes in Chédiak-Higashi syndrome. This image is a derivative work, adapted from the following source, available under CC BY: Bharti S, Bhatia P, Bansal D, et al. The accelerated phase of Chediak-Higashi syndrome: the importance of hematological evaluation. *Turk J Haematol.* 2013;30:85-87. DOI: 10.4274/tjh.2012.0027. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

Microbiology

125 **Stains: Image A.** *Trypanosoma lewisi* on Giemsa stain. Courtesy of the Department of Health and Human Services and Dr. Mae Melvin.

125 **Stains: Image B.** Periodic acid–Schiff stain reveals *Tropheryma whipplei* infection. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of Dr. Ed Uthman.

125 **Stains: Image C.** *Mycobacterium tuberculosis* on Ziehl-Neelsen stain. Courtesy of the Department of Health and Human Services and Dr. George P. Kubica.

125 **Stains: Image D.** *Cryptococcus neoformans* on India ink stain. Courtesy of the Department of Health and Human Services.

125 **Stains: Image E.** *Coccidioides immitis* on silver stain. Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr.

127 **Encapsulated bacteria.** Capsular swelling of *Streptococcus pneumoniae* using the Neufeld-Quellung test. Courtesy of the Department of Health and Human Services.

128 **Catalase-positive organisms.** Oxygen bubbles released during catalase reaction. This image is a derivative work, adapted from the following source, available under CC BY-NC-SA: Stefano Nase. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-NC-SA.

129 **Spore-forming bacteria.** This image is a derivative work, adapted from the following source, available under CC BY: Jones SW, Paredes CJ, Tracy B. The transcriptional program underlying the physiology of clostridial sporulation. *Genome Biol.* 2008;9:R114. DOI: 10.1186/gb-2008-9-7-r114.

135 **α-hemolytic bacteria.** α-hemolysis. This image is a derivative work, adapted from the following source, available under CC BY-NC-SA: Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-NC-SA.

(c) ketabton.com: The Digital Library

135 **β-hemolytic bacteria.** β-hemolysis. This image is a derivative work, adapted from the following source, available under CC BY: Wikimedia Commons.

135 ***Staphylococcus aureus*.** Courtesy of the Department of Health and Human Services and Dr. Richard Facklam.

136 ***Streptococcus pneumoniae*.** Courtesy of the Department of Health and Human Services and Dr. Mike Miller.

136 ***Streptococcus pyogenes:* (group A streptococci).** This image is a derivative work, adapted from the following source, available under CC BY: Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

137 ***Bacillus anthracis*.** Ulcer with black eschar. Courtesy of the Department of Health and Human Services and James H. Steele.

138 **Clostridia: Image A.** Gas gangrene due to *Clostridium perfringens*. This image is a derivative work, adapted from the following source, available under CC BY: Schröpfer E, Rauthe S, Meyer T. Diagnosis and misdiagnosis of necrotizing soft tissue infections: three case reports. *Cases J.* 2008;1:252. DOI: 10.1186/1757-1626-1-252.

138 **Clostridia: Image B.** Pseudomembranous enterocolitis on colonoscopy. This image is a derivative work, adapted from the following source, available under CC BY: Klinikum Dritter Orden für die Überlassung des Bildes zur Veröffentlichu. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

139 ***Corynebacterium diphtheriae*.** Pseudomembranous pharyngitis. This image is a derivative work, adapted from the following source, available under CC BY: Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

139 ***Listeria monocytogenes*.** Actin rockets. This image is a derivative work, adapted from the following source, available under CC BY: Schuppler M, Loessner MJ. The opportunistic pathogen *Listeria monocytogenes*: pathogenicity and interaction with the mucosal immune system. *Int J Inflamm.* 2010;2010:704321. DOI: 10.4061/2010/704321. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

139 ***Nocardia* vs *Actinomyces*: Image A.** *Nocardia* on acid-fast stain. This image is a derivative work, adapted from the following source, available under CC BY: Venkataramana K. Human *Nocardia* infections: a review of pulmonary nocardiosis. *Cereus.* 2015;7(8):e304. DOI: 10.7759/cureus.304.

139 ***Nocardia* vs *Actinomyces*: Image B.** *Actinomyces israelii* on Gram stain. Courtesy of the Department of Health and Human Services.

140 **Mycobacteria.** Acid-fast stain. Courtesy of the Department of Health and Human Services and Dr. George P. Kubica

140 **Tuberculosis.** Langhans giant cell in caseating granuloma. Courtesy of J. Hayman.

141 **Leprosy: Image A.** "Glove and stocking" distribution. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of Bruno Jehle.

142 ***Neisseria*: Image A.** Intracellular *N gonorrhoeae*. Courtesy of the Department of Health and Human Services and Dr. Mike Miller.

142 ***Haemophilus influenzae*: Image A.** Epiglottitis. This image is a derivative work, adapted from the following source, available under CC BY: Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

143 ***Legionella pneumophila*.** Lung findings of unilateral and lobar infiltrate. This image is a derivative work, adapted from the following source, available under CC BY: Robbins NM, Kumar A, Blair BM. *Legionella pneumophila* infection presenting as headache, confusion and dysarthria in a human immunodeficiency virus-1 (HIV-1) positive patient: case report. *BMC Infect Dis.* 2012;12:225. DOI: 10.1186/1471-2334-12-225.

143 ***Pseudomonas aeruginosa*: Image A.** Blue-green pigment on centrimide agar. This image is a derivative work, adapted from the following source, available under CC BY: Hansen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

143 ***Pseudomonas aeruginosa*: Image B.** Ecthyma gangrenosum. This image is a derivative work, adapted from the following source, available under CC BY: Uludokumaci S, Balkan II, Mete B, et al. Ecthyma gangrenosum-like lesions in a febrile neutropenic patient with simultaneous *Pseudomonas* sepsis and disseminated fusariosis. *Turk J Haematol.* 2013 Sep;30(3):321-4. DOI: 10.4274/Tjh.2012.0030.

145 ***Klebsiella*.** Courtesy of the Department of Health and Human Services.

145 ***Campylobacter jejuni*.** Courtesy of the Department of Health and Human Services.

146 ***Vibrio cholerae*.** This image is a derivative work, adapted from the following source, available under CC BY: Phetsouvanh R, Nakatsu M, Arakawa E, et al. Fatal bacteremia due to immotile *Vibrio cholerae* serogroup O21 in Vientiane, Laos—a case report. *Ann Clin Microbiol Antimicrob.* 2008;7:10. DOI: 10.1186/1476-0711-7-10.

146 ***Helicobacter pylori*.** Courtesy of the Department of Health and Human Services, Dr. Patricia Fields, and Dr. Collette Fitzgerald.

146 **Spirochetes.** Appearance on darkfield microscopy. Courtesy of the Department of Health and Human Services.

146 **Lyme disease: Image A.** *Ixodes* tick. Courtesy of the Department of Health and Human Services and Dr. Michael L. Levin.

146 **Lyme disease: Image B.** Erythema migrans. Courtesy of the Department of Health and Human Services and James Gathany.

147 **Syphilis: Image A.** Painless chancre in primary syphilis. Courtesy of the Department of Health and Human Services and M. Rein.

147 **Syphilis: Image B.** Treponeme on darkfield microscopy. Courtesy of the Department of Health and Human Services and Renelle Woodall.

147 **Syphilis: Image D.** Rash on palms. This image is a derivative work, adapted from the following source, available under CC BY: Drahansky M, Dolezel M, Urbanek J, et al. Influence of skin diseases on fingerprint recognition. *J Biomed Biotechnol.* 2012;626148. DOI: 10.1155/2012/626148.

147 **Syphilis: Image E.** Condyloma lata. Courtesy of the Department of Health and Human Services and Susan Lindsley.

147 **Syphilis: Image F.** Gumma. This image is a derivative work, adapted from the following source, available under CC BY: Chakir K, Benchikhi H. Granulome centro-facial révélant une syphilis tertiaire. *Pan Afr Med J.* 2013;15:82. DOI: 10.11604/pamj.2013.15.82.3011.

147 **Syphilis: Image G.** Congenital syphilis. Courtesy of the Department of Health and Human Services and Dr. Norman Cole.

147 **Syphilis: Image H.** Hutchinson teeth. Courtesy of the Department of Health and Human Services and Susan Lindsley.

148 ***Gardnerella vaginalis*.** Courtesy of the Department of Health and Human Services and M. Rein.

150 **Rickettsial diseases and vector-borne illnesses: Image A.** Rash of Rocky Mountain spotted fever. Courtesy of the Department of Health and Human Services.

(c) ketabton.com: The Digital Library

150 **Rickettsial diseases and vector-borne illnesses: Image B.** *Ehrlichia* morulae. This image is a derivative work, adapted from the following source, available under CC BY: Dantas-Torres F. Canine vector-borne diseases in Brazil. *Parasit Vectors.* 2008;1:25. DOI: 10.1186/1756-3305-1-25. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

150 **Rickettsial diseases and vector-borne illnesses: Image C.** *Anaplasma phagocytophilium* in neutrophil. Courtesy of the Department of Health and Human Services and Dumler JS, Choi K, Garcia-Garcia JC, et al. Human granulocytic anaplasmosis. *Emerg Infect Dis.* 2005. DOI 10.3201/eid1112.050898.

150 ***Mycoplasma pneumoniae*.** This image is a derivative work, adapted from the following source, available under CC BY: Rottem S, Kosower ND, Kornspan JD. Contamination of tissue cultures by *Mycoplasma*. In: Ceccherini-Nelli L, ed: *Biomedical tissue culture.* 2016. DOI: 10.5772/51518.

151 **Systemic mycoses: Image A.** *Histoplasma.* Courtesy of the Department of Health and Human Services and Dr. D.T. McClenan.

151 **Systemic mycoses: Image B.** *Blastomyces dermatitidis* undergoing broad-base budding. Courtesy of the Department of Health and Human Services and Dr. Libero Ajello.

151 **Systemic mycoses: Image C.** Coccidiomycosis with endospheres. Courtesy of the Department of Health and Human Services.

151 **Systemic mycoses: Image D.** "Captain's wheel" shape of *Paracoccidioides*.. Courtesy of the Department of Health and Human Services and Dr. Lucille K. Georg.

152 **Cutaneous mycoses: Image G.** Tinea versicolor. This image is a derivative work, adapted from the following source, available under CC BY-SA: Sarah (Rosenau) Korf. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

153 **Opportunistic fungal infections: Image A.** Budding yeast of *Candida albicans.* This image is a derivative work, adapted from the following source, available under CC BY-SA: Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

153 **Opportunistic fungal infections: Image B.** Germ tubes of *Candida albicans*.. This image is a derivative work, adapted from the following source, available under CC BY-SA: Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

153 **Opportunistic fungal infections: Image C.** Oral thrush. Courtesy of the Department of Health and Human Services and Dr. Sol Silverman, Jr.

153 **Opportunistic fungal infections: Image E.** Conidiophores of *Aspergillus fumigatus*. Courtesy of the Department of Health and Human Services.

153 **Opportunistic fungal infections: Image F.** Aspergilloma in left lung. This image is a derivative work, adapted from the following source, available under CC BY: Souilamas R, Souilamas JI, Alkhamees K, et al. Extra corporal membrane oxygenation in general thoracic surgery: a new single veno-venous cannulation. *J Cardiothorac Surg.* 2011;6:52. DOI: 10.1186/1749-8090-6-52.

153 **Opportunistic fungal infections: Image G.** *Cryptococcus neoformans*. Courtesy of the Department of Health and Human Services and Dr. Leanor Haley.

153 **Opportunistic fungal infections: Image H.** *Cryptococcus neoformans* on mucicarmine stain. Courtesy of the Department of Health and Human Services and Dr. Leanor Haley.

153 **Opportunistic fungal infections: Image I.** Mucor. Courtesy of the Department of Health and Human Services and Dr. Lucille K. Georg.

153 **Opportunistic fungal infections: Image J.** Mucormycosis. This image is a derivative work, adapted from the following source, available under CC BY: Jiang N, Zhao G, Yang S, et al. A retrospective analysis of eleven cases of invasive rhino-orbito-cerebral mucormycosis presented with orbital apex syndrome initially. *BMC Ophthalmol.* 2016; 16: 10. DOI: 10.1186/s12886-016-0189-1.

154 ***Pneumocystis jirovecii*: Image A.** Interstitial opacities in lung. This image is a derivative work, adapted from the following source, available under CC BY: Chuang C, Zhanhong X, Yinyin G, et al. Unsuspected *Pneumocystis* pneumonia in an HIV-seronegative patient with untreated lung cancer: circa case report. *J Med Case Rep.* 2007;1:15. DOI: 10.1186/1752-1947-1-115.

154 ***Pneumocystis jirovecii*: Image B.** CT of lung. This image is a derivative work, adapted from the following source, available under CC BY: Allen CM, Al-Jahdali HH, Irion KL, et al. Imaging lung manifestations of HIV/AIDS. *Ann Thorac Med.* 2010 Oct-Dec; 5(4): 201–216. DOI: 10.4103/1817-1737.69106.

154 ***Pneumocystis jiroveci*: Image C.** Disc-shaped yeast. This image is a derivative work, adapted from the following source, available under CC BY: Kirby S, Satoskar A, Brodsky S, et al. Histological spectrum of pulmonary manifestations in kidney transplant recipients on sirolimus inclusive immunosuppressive regimens. *Diagn Pathol.* 2012;7:25. DOI: 10.1186/1746-1596-7-25.

154 ***Sporothrix schenckii*.** Subcutaneous mycosis. This image is a derivative work, adapted from the following source, available under CC BY: Govender NP, Maphanga TG, Zulu TG, et al. An outbreak of lymphocutaneous sporotrichosis among mine-workers in South Africa. *PLoS Negl Trop Dis.* 2015 Sep; 9(9): e0004096. DOI: 10.1371/journal.pntd.0004096.

155 **Protozoa—GI infections: Image A.** *Giardia lamblia* trophozoite. This image is a derivative work, adapted from the following source, available under CC BY: Lipoldová M. Giardia and Vilém Dušan Lambl. *PLoS Negl Trop Dis.* 2014;8:e2686. DOI: 10.1371/journal.pntd.0002686.

155 **Protozoa—GI infections: Image B.** *Giardia lamblia* cyst. Courtesy of the Department of Health and Human Services.

155 **Protozoa—GI infections: Image C.** *Entamoeba histolytica* trophozoites. Courtesy of the Department of Health and Human Services.

155 **Protozoa—GI infections: Image D.** *Entamoeba histolytica* cyst. Courtesy of the Department of Health and Human Services.

155 **Protozoa—GI infections: Image E.** *Cryptosporidium* oocysts. Courtesy of the Department of Health and Human Services.

156 **Protozoa—CNS infections: Image A.** *Toxoplasma gondii*. This image is a derivative work, adapted from the following source, available under CC BY: Agrawal A, Bhake A, Sangole VM, et al. Multiple-ring enhancing lesions in an immunocompetent adult. *J Glob Infect Dis.* 2010 Sep-Dec;2(3):313-4. DOI: 10.4103/0974-777X.68545.

156 **Protozoa—CNS infections: Image B.** *Toxoplasma gondii* tachyzoite. Courtesy of the Department of Health and Human Services and Dr. L.L. Moore, Jr.

156 **Protozoa—CNS infections: Image C.** *Naegleria fowleri* amoebas. Courtesy of the Department of Health and Human Services.

156 **Protozoa—CNS infections: Image D.** *Trypanosoma brucei gambiense*. Courtesy of the Department of Health and Human Services and Dr. Mae Melvin.

(c) ketabton.com: The Digital Library

157 **Protozoa—hematologic infections: Image A.** *Plasmodium* trophozoite ring form. Courtesy of the Department of Health and Human Services.

157 **Protozoa—hematologic infections: Image B.** *Plasmodium* schizont containing merozoites. Courtesy of the Department of Health and Human Services and Steven Glenn.

157 **Protozoa—hematologic infections: Image C.** *Babesia* with ring form and with "Maltese cross" form. Courtesy of the Department of Health and Human Services.

158 **Protozoa—others: Image A.** *Trypanosoma cruzi.* Courtesy of the Department of Health and Human Services and Dr. Mae Melvin.

158 **Protozoa—others: Image B.** Cutaneous leishmaniasis. This image is a derivative work, adapted from the following source, available under : Sharara SL, Kanj SS. War and infectious diseases: challenges of the Syrian civil war. *PLoS Pathog.* 2014 Nov;10(11):e1004438. DOI: 10.1371/journal.ppat.1004438.

158 **Protozoa—others: Image C.** *Leishmania* spp. Courtesy of the Department of Health and Human Services and Dr. Francis W. Chandler. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

158 **Protozoa—others: Image D.** *Trichomonas vaginalis*. Courtesy of the Department of Health and Human Services.

159 **Nematodes (roundworms): Image A.** *Enterobius vermicularis* eggs. Courtesy of the Department of Health and Human Services, BG Partin, and Dr. Moore.

159 **Nematodes (roundworms): Image B.** *Ascaris lumbricoides* egg. Courtesy of the Department of Health and Human Services.

159 **Nematodes (roundworms): Image C.** *Ancylostoma* spp rash This image is a derivative work, adapted from the following source, available under : Archer M. Late presentation of cutaneous larva migrans: a case report. *Cases J.* 2009; 2: 7553. doi:10.4076/1757-1626-2-7553.

159 **Nematodes (roundworms): Image D.** *Trichinella spiralis* cysts in muscle. This image is a derivative work, adapted from the following source, available under : Franssen FFJ, Fonville M, Takumi K, et al. *Vet Res.* 2011; 42(1): 113. DOI: 10.1186/1297-9716-42-113.

159 **Nematodes (roundworms): Image E.** *Wuchereria bancrofti* Elephantiasis. Courtesy of the Department of Health and Human Services.

160 **Cestodes (tapeworms): Image A.** *Taenia solium*. Courtesy of the Department of Health and Human Services Robert J. Galindo. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

160 **Cestodes (tapeworms): Image B.** Neurocysticercosis. This image is a derivative work, adapted from the following source, available under : Coyle CM, Tanowitz HB. Diagnosis and treatment of neurocysticercosis. *Interdiscip Perspect Infect Dis.* 2009;2009:180742. DOI: 10.1155/2009/180742. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

160 **Cestodes (tapeworms): Image C.** *Echinococcus granulosus.* Courtesy of the Department of Health and Human Services.

160 **Cestodes (tapeworms): Image D.** Hyatid cyst of *Echinococcus granulosus.* Courtesy of the Department of Health and Human Services and Dr. I. Kagan.

160 **Cestodes (tapeworms): Image E.** *Echinococcus granulosus* cyst in liver. This image is a derivative work, adapted from the following source, available under : Ma Z, Yang W, Yao Y, et al. The adventitia resection in treatment of liver hydatid cyst: a case report of a 15-year-old boy. *Case Rep Surg.* 2014;2014:123149. DOI: 10.1155/2014/123149.

160 **Trematodes (flukes): Image A.** *Schistosoma mansoni* egg with lateral spine. Courtesy of the Department of Health and Human Services.

160 **Trematodes (flukes): Image B.** *Schistosoma haematobium* egg with terminal spine. Courtesy of the Department of Health and Human Services.

161 **Ectoparasites: Image A.** Scabies. This image is a derivative work, adapted from the following source, available under : Siegfried EC, Hebert AA. Diagnosis of atopic dermatitis: mimics, overlaps, and complications. *Clin Med.* 2015 May; 4(5): 884–917. DOI: 10.3390/jcm4050884.

161 **Ectoparasites: Image B.** Nit of a louse. Courtesy of the Department of Health and Human Services and Joe Miller.

164 **DNA viruses: Image A.** Febrile pharyngitis. Balfour HH Jr, Dunmire SK, Hogquist KA. *Clin Transl Immunology.*2015 Feb 27. DOI: 10.1038/cti.2015.1.

165 **Herpesviruses: Image A.** Keratoconjunctivitis in HSV-1 infection. This image is a derivative work, adapted from the following source, available under : Yang HK, Han YK, Wee WR, et al. Bilateral herpetic keratitis presenting with unilateral neurotrophic keratitis in pemphigus foliaceus: a case report. *J Med Case Rep.* 2011;5:328. DOI: 10.1186/1752-1947-5-328.

165 **Herpesviruses: Image B.** Herpes labialis. Courtesy of the Department of Health and Human Services and Dr. Herrmann.

165 **Herpesviruses: Image E.** Shingles (varicella-zoster virus infection). This image is a derivative work, adapted from the following source, available under : Fisle. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

165 **Herpesviruses: Image F.** Hepatosplenomegaly due to EBV infection. This image is a derivative work, adapted from the following source, available under : Gow NJ, Davidson RN, Ticehurst R, et al. Case report: no response to liposomal daunorubicin in a patient with drug-resistant HIV-associated visceral leishmaniasis. *PLoS Negl Trop Dis.* 2015 Aug; 9(8):e0003983. DOI: 10.1371/journal.pntd.0003983.

165 **Herpesviruses: Image G.** Atypical lymphocytes in Epstein-Barr virus infection. This image is a derivative work, adapted from the following source, available under : Coutesy of Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

165 **Herpesviruses: Image I.** Roseola. Courtesy of Emiliano Burzagli.

165 **Herpesviruses: Image J.** Kaposi sarcoma. Courtesy of the Department of Health and Human Services.

166 **HSV identification.** Positive Tzanck smear in HSV-2 infection. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

168 **Rotavirus.** Courtesy of the Department of Health and Human Services and Erskine Palmer.

169 **Rubella virus.** Rubella rash. Courtesy of the Department of Health and Human Services.

170 **Acute laryngotracheobronchitis.** Steeple sign. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.

170 **Measles (rubeola) virus: Image A.** Koplik spots. Courtesy of the Department of Health and Human Services. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

(c) ketabton.com: The Digital Library

170 **Measles (rubeola) virus: Image B.** Rash of measles. Courtesy of the Department of Health and Human Services.

170 **Mumps virus.** Swollen neck and parotid glands. Courtesy of the Department of Health and Human Services.

171 **Rabies virus: Image A.** Transmission electron micrograph. Courtesy of the Department of Health and Human Services Dr. Fred Murphy, and Sylvia Whitfield.

171 **Rabies virus: Image B.** Negri bodies. Courtesy of the Department of Health and Human Services and Dr. Daniel P. Perl.

171 **Ebola virus.** Courtesy of the Department of Health and Human Services and Cynthia Goldsmith.

180 **Osteomyelitis.** X-ray (left) and MRI (right) views. This image is a derivative work, adapted from the following source, available under : Huang P-Y, Wu P-K, Chen C-F, et al. Osteomyelitis of the femur mimicking bone tumors: a review of 10 cases. *World J Surg Oncol.* 2013;11:283. DOI: 10.1186/1477-7819-11-283.

181 **Common vaginal infections: Image B.** Motile trichomonads. Courtesy of Joe Miller.

181 **Common vaginal infections: Image C.** *Candida* vulvovaginitis. Courtesy of Mikael Häggström.

182 **TORCH infections: Image A.** "Blueberry muffin" rash. This image is a derivative work, adapted from the following source, available under : Benmiloud S, Elhaddou G, Belghiti ZA, et al. Blueberry muffin syndrome. *Pan Afr Med J.* 2012;13:23.

182 **TORCH infections: Image B.** Cataract in infant with contenital rubella. Courtesy of the Department of Health and Human Services .

182 **TORCH infections: Image C.** Periventricular calcifications in congenital cytomegalovirus infection. This image is a derivative work, adapted from the following source, available under : Bonthius D, Perlman S. Congenital viral infections of the brain: lessons learned from lymphocytic choriomeningitis virus in the neonatal rat. *PLoS Pathog.* 2007;3:e149. DOI: 10.1371/journal.ppat.0030149. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

183 **Red rashes of childhood: Image C.** Child with scarlet fever. This image is a derivative work, adapted from the following source, available under : www.badobadop.co.uk.

183 **Red rashes of childhood: Image D.** Chicken pox. Courtesy of the Department of Health and Human Services.

184 **Sexually transmitted infections: Image A.** Chancroid. Courtesy of the Department of Health and Human Services and Susan Lindsley.

184 **Sexually transmitted infections: Image B.** Donovanosis. Courtesy of the Department of Health and Human Services and Dr. Pinozzi.

185 **Pelvic inflammatory disease: Image A.** Purulent cervical discharge. This image is a derivative work, adapted from the following source, available under : SOS-AIDS Amsterdam The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

185 **Pelvic inflammatory disease: Image B.** Adhesions in Fitz-Hugh–Curtis syndrome. Courtesy of Hic et nunc.

190 **Vancomycin.** Red man syndrome. This image is a derivative work, adapted from the following source, available under : O'Meara P, Borici-Mazi R, Morton R, et al. DRESS with delayed onset acute interstitial nephritis and profound refractory eosinophilia secondary to vancomycin. *Allergy Asthma Clin Immunol.* 2011;7:16. DOI: 10.1186/1710-1492-7-16.

Pathology

209 **Necrosis: Image A.** Coagulative necrosis. Courtesy of the Department of Health and Human Services and Dr. Steven Rosenberg.

209 **Necrosis: Image B.** Liquefactive necrosis. Courtesy of Daftblogger.

209 **Necrosis: Image C.** Caseous necrosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

209 **Necrosis: Image D.** Fat necrosis. This image is a derivative work, adapted from the following source, available under : Patho. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

209 **Necrosis: Image E.** Fibrinoid necrosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

209 **Necrosis: Image F.** Acral gangrene. Courtesy of the Department of Health and Human Services and William Archibald.

210 **Ischemia.** This image is a derivative work, adapted from the following source, available under : Van Assche LM, Kim HW, Jensen CJ, et al. A new CMR protocol for non-destructive, high resolution, ex-vivo assessment of the area at risk simultaneous with infarction: validation with histopathology. *J Cardiovasc Magn Reson.* 2012; 14(Suppl 1): O7. DOI: 10.1186/1532-429X-14-S1-O7.

210 **Types of infarcts: Image B.** Pale infarct. Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

211 **Types of calcification: Image A.** Dystrophic calcification. This image is a derivative work, adapted from the following source, available under : Chun J-S, Hong R, Kim J-A. Osseous metaplasia with mature bone formation of the thyroid gland: three case reports. *Oncol Lett.* 2013;6:977-979. DOI: 10.3892/ol.2013.1475. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

211 **Lipofuscin.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

212 **Amyloidosis: Image A.** Amyloid deposits on Congo red stain. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman.

212 **Amyloidosis: Image B.** Apple green birefringence under polarized light. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman.

214 **Acute inflammation: Image A.** Pericardium with severe inflammation, neutrophilic infiltration and fibrin with entrapped clusters of bacteria. This image is a derivative work, adapted from the following source, available under : Faida Ajili, et al. Coexistence of pyoderma gangrenosum and sweet's syndrome in a patient with ulcerative colitis. *Pan Afr Med J.* 2015 Jun 24. DOI: 10.11604/pamj.2015.21.151.6364.

217 **Granulomatous diseases.** Granuloma. Courtesy of Sanjay Mukhopadhyay.

218 **Scar formation: Image A.** Hypertrophic scar. This image is a derivative work, adapted from the following source, available under : Baker R, Urso-Baiarda F, Linge C, et al. Cutaneous scarring: a clinical review. *Dermatol Res Pract.* 2009;2009:625376. DOI: 10.1155/2009/625376.

(c) ketabton.com: The Digital Library

218 **Scar formation: Image B.** Keloid scar. This image is a derivative work, adapted from the following source, available under CC BY: Dr. Andreas Settje. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

219 **Neoplasia and neoplastic progression: Image A.** Cervical tissue. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

223 **Common metastases: Image A.** Brain metastases from breast cancer. This image is a derivative work, adapted from the following source, available under CC BY: Jmarchn. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

223 **Common metastases: Image B.** Brain metastasis. Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

223 **Common metastases: Image C.** Liver metastasis. This image is a derivative work, adapted from the following source, available under CC BY: Dr. James Heilman The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

223 **Common metastases: Image D.** Liver metastasis. Courtesy of J. Hayman.

223 **Common metastases: Image E.** Bone metastasis. This image is a derivative work, adapted from the following source, available under CC BY: Dr. Paul Hellerhoff.

223 **Common metastases: Image F.** Bone metastasis. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of M Emmanuel.

227 **Psammoma bodies.** Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

Cardiovascular

283 **Anatomy of the heart: Image A.** MRI showing normal cardiac anatomy. This image is a derivative work, adapted from the following source, available under CC BY: Zhang J, Chen L, Wang X, et al. Compounding local invariant features and global deformable geometry for medical image registration. *PLoS One.* 2014;9(8):e105815. DOI: 10.1371/journal.pone.0105815.

298 **Congenital heart diseases: Image A.** Tetralogy of Fallot. This image is a derivative work, adapted from the following source, available under CC BY: Rashid AKM: Heart diseases in Down syndrome. In: Dey S, ed: Down syndrome. DOI: 10.5772/46009. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

299 **Congenital heart diseases: Image B.** Ventricular septal defect. This image is a derivative work, adapted from the following source, available under CC BY: Bardo DME, Brown P. Cardiac multidetector computed tomography: basic physics of image acquisition and clinical applications. *Curr Cardiol Rev.* 2008 Aug;4(3):231–243. DOI: 10.2174/157340308785160615.

299 **Congenital heart diseases: Image C.** Atrial septal defect. This image is a derivative work, adapted from the following source, available under CC BY: Teo KSL, Dundon BK, Molaee P, et al. Percutaneous closure of atrial septal defects leads to normalisation of atrial and ventricular volumes. *J Cardiovasc Magn Reson.* 2008;10(1):55. DOI: 10.1186/1532-429X-10-55.

299 **Congenital heart diseases: Image D.** Patent ductus arteriosus. This image is a derivative work, adapted from the following source, available under CC BY: Henjes CR, Nolte I, Wesfaedt P. Multidetector-row computed tomography of thoracic aortic anomalies in dogs and cats: patent ductus arteriosus and vascular rings. *BMC Vet Res.* 2011;7:57. DOI: 10.1186/1746-6148-7-57.

299 **Congenital heart diseases: Image E.** Clubbing of fingers. Courtesy of Ann McGrath.

299 **Congenital heart diseases: Image F.** MRI showing coarctation of the aorta. This image is a derivative work, adapted from the following source, available under CC BY: Vergales JE, Gangemi JJ, Rhueban KS, Lim DS. Coarctation of the aorta — the current state of surgical and transcatheter therapies. *Curr Cardiol Rev.* 2013 Aug; 9(3): 211–219. DOI: 10.2174/1573403X113099990032

300 **Hypertension: Image A.** "String of beads" appearance in fibromuscular dysplasia. This image is a derivative work, adapted from the following source, available under CC BY: Plouin PF, Perdu J, LaBatide-Alanore A, et al. Fibromuscular dysplasia. *Orphanet J Rare Dis.* 2007;7:28. DOI: 10.1186/1750-1172-2-28. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

301 **Hyperlipidemia signs: Image C.** Tendinous xanthoma. This image is a derivative work, adapted from the following source, available under CC BY: Raffa W, Hassam B. Xanthomes tendineux et tubéreux révélant une hypercholestérolémie familiale. *Pan Afr Med J.* 2013; 15: 49. DOI: 10.11604/pamj.2013.15.49.2636.

301 **Arteriosclerosis: Image A.** Hyaline type. This image is a derivative work, adapted from the following source, available under CC BY: Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

301 **Arteriosclerosis: Image B.** Hyperplastic type. This image is a derivative work, adapted from the following source, available under CC BY: Paco Larosa. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

301 **Arteriosclerosis: Image C.** Monckeberg sclerosis (medial calcific sclerosis). This image is a derivative work, adapted from the following source, available under CC BY: Couri CE, da Silva GA, Martinez JA, et al. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

303 **Aortic dissection.** This image is a derivative work, adapted from the following source, available under CC BY: Qi Y, Ma X, Li G, et al. Three-dimensional visualization and imaging of the entry tear and intimal flap of aortic dissection using CT virtual intravascular endoscopy. *PLoS One.* 2016; 11(10): e0164750. DOI: 10.1371/journal.pone.0164750.

307 **Myocardial infarction complications: Image A.** Papillary muscle rupture. This image is a derivative work, adapted from the following source, available under CC BY: Routy B, Huynh T, Fraser R, et al. Vascular endothelial cell function in catastrophic antiphospholipid syndrome: a case report and review of the literature. *Case Rep Hematol.* 2013;2013:710365. DOI: 10.1155/2013/710365.

307 **Myocardial infarction complications: Image B.** Drawing of pseudoaneurysm. This image is a derivative work, adapted from the following source, available under CC BY: Patrick J. Lynch and Dr. C. Carl Jaffe.

307 **Myocardial infarction complications: Image C.** Free wall rupture of left ventricle. This image is a derivative work, adapted from the following source, available under CC BY: Zacarias ML, da Trindade H, Tsutsu J, et al. Left ventricular free wall impeding rupture in post-myocardial infarction period diagnosed by myocardial contrast echocardiography: case report. *Cardiovasc Ultrasound.* 2006;4:7. DOI: 10.1186/1476-7120-4-7.

308 **Cardiomyopathies: Image A.** Dilated cardiomyopathy. This image is a derivative work, adapted from the following source, available under

(c) ketabton.com: The Digital Library

: Gho JMIH, van Es R, Stathonikos N, et al. High resolution systematic digital histological quantification of cardiac fibrosis and adipose tissue in phospholamban p.Arg14del mutation associated cardiomyopathy. *PLoS One.* 2014;9:e94820. DOI: 10.1371/journal.pone.0094820.

308 **Cardiomyopathies: Image B.** Hypertrophic obstructive cardiomyopathy. This image is a derivative work, adapted from the following source, available under : Benetti MA, Belo Nunes RA, Benvenuti LA. Case 2/2016 - 76-year-old male with hypertensive heart disease, renal tumor and shock. *Arq Bras Cardiol.* 2016 May; 106(5): 439–446. DOI: 10.5935/abc.20160067.

309 **Heart failure.** Pedal edema. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

310 **Cardiac tamponade: Image A.** This image is a derivative work, adapted from the following source, available under : Yousuf T, Kramer J, Kopiec A, et al. A rare case of cardiac tamponade induced by chronic rheumatoid arthritis. *J Clin Med Res.* 2015 Sep;7(9):720–723. DOI: 10.14740/jocmr2226w.

310 **Cardiac tamponade: Image B.** This image is a derivative work, adapted from the following source, available under : Maharaj SS, Chang SM. Cardiac tamponade as the initial presentation of systemic lupus erythematosus: a case report and review of the literature. *Pediatr Rheumatol Online J.* 2015; 13: 9. DOI: 10.1186/s12969-015-0005-0.

311 **Bacterial endocarditis: Image A.** Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr.

311 **Bacterial endocarditis: Image C.** Osler nodes. This image is a derivative work, adapted from the following source, available under : Yang ML, Chen YH, Lin WR, et al. Case report: infective endocarditis caused by Brevundimonas vesicularis. *BMC Infect Dis.* 2006;6:179. DOI: 10.1186/1471-2334-6-179.

311 **Bacterial endocarditis: Image D.** Janeway lesions on sole. This image is a derivative work, adapted from the following source, available under : Courtesy of DeNanneke.

312 **Rheumatic fever.** Aschoff body and Anitschkow cells. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

313 **Acute pericarditis.** This image is a derivative work, adapted from the following source, available under : Bogaert J, Francone M. Cardiovascular magnetic resonance in pericardial diseases. *J Cardiovasc Magn Reson.* 2009;11:14. DOI: 10.1186/1532-429X-11-14. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

315 **Vasculitides: Image A.** Temporal arteritis histology. This image is a derivative work, adapted from the following source, available under : Marvin. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

315 **Vasculitides: Image B.** Angiogram in patient with Takayasu arteritis. Courtesy of the Department of Health and Human Services and Justin Ly.

315 **Vasculitides: Image C.** Gangrene as a consequence of Buerger disease. This image is a derivative work, adapted from the following source, available under : Afsjarfard A, Mozaffar M, Malekpour F, et al. The wound healing effects of iloprost in patients with Buerger's disease: claudication and prevention of major amputations. *Iran Red Crescent Med J.* 2011;13:420-423.

315 **Vasculitides: Image D.** Strawberry tongue in patient with Kawasaki disease. This image is a derivative work, adapted from the following source, available under : Courtesy of Natr.

315 **Vasculitides: Image E.** Coronary artery aneurysm in Kawasaki disease. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

315 **Vasculitides.** Polyarteritis nodosa. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.

315 **Vasculitides: Image G.** Churg-Strauss syndrome histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

315 **Vasculitides: Image H.** Granulomatosis with polyangiitis (formerly Wegener) and PR3-ANCA/c-ANCA. Courtesy of M.A. Little.

315 **Vasculitides: Image I.** Henoch-Schönlein purpura. Courtesy of Okwikikim.

315 **Vasculitides: Image J.** MPO-ANCA/p-ANCA in microscopic polyangiitis. Courtesy of and M.A. Little.

Endocrine

326 **Thyroid development.** Thyroglossal duct cyst. This image is a derivative work, adapted from the following source, available under : Adelchi C, Mara P, Melissa L, et al. Ectopic thyroid tissue in the head and neck: a case series. *BMC Res Notes.* 2014;7:790. DOI: 10.1186/1756-0500-7-790.

340 **Hypothyroidism vs hyperthyroidism.** Onycholysis. This image is a derivative work, adapted from the following source, available under : Rajebi MR, Shahrokni A, Chaisson M. Uncommon osseous involvement in multisystemic sarcoidosis. *Ann Saudi Med.* 2009 Nov-Dec;29(6):485–486.

341 **Hypothyroidism: Image B.** Before and after treatment of congenital hypothyroidism. Courtesy of the Department of Health and Human Services.

341 **Hypothyroidism: Image C.** Congenital hypothyroidism. This image is a derivative work, adapted from the following source, available under : Sadasiv Swain. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

342 **Thyroid adenoma: Image A.** This image is a derivative work, adapted from the following source, available under : Terada T. Brain metastasis from thyroid adenomatous nodules or an encapsulated thyroid follicular tumor without capsular and vascular invasion: a case report. *Cases J.* 2009; 2: 7180. DOI: 10.4076/1757-1626-2-7180.

344 **Hypoparathyroidism.** Shortened 4th and 5th digits. This image is a derivative work, adapted from the following source, available under : Ferrario C, Gastaldi G, Portmann L, et al. Bariatric surgery in an obese patient with Albright hereditary osteodystrophy: a case report. *J Med Case Rep.* 2013; 7: 111. DOI: 10.1186/1752-1947-7-111.

345 **Hyperparathyroidism.** Multiple lytic lesions. This image is a derivative work, adapted from the following source, available under : Khaoula BA, Kaouther BA, Ines C, et al. An unusual presentation of primary hyperparathyroidism: pathological fracture. *Case Rep Orthop.* 2011;2011:521578. DOI: 10.1155/2011/521578. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

349 **Adrenal insufficiency: Image A.** Mucosal hyperpigmentation in primary adrenal insufficiency. Courtesy of FlatOut. The image

(c) ketabton.com: The Digital Library

may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

350 **Pheochromocytoma.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Feldman.

351 **Multiple endocrine neoplasias.** Mucosal neuroma. This image is a derivative work, adapted from the following source, available under : Martucciello G, Lerone M, Bricco L, et al. Multiple endocrine neoplasias type 2B and RET proto-oncogene. *Ital J Pediatr.* 2012;38:9. DOI: 10.1186/1824-7288-38-9.

352 **Carcinoid syndrome.** Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

Gastrointestinal

358 **Ventral wall defects.** Gastroschisis. This image is a derivative work, adapted from the following source, available under : Zvizdic Z. Gastroschisis with concomitant jejuno-ileal atresia complicated by jejunal perforation. *J Neonatal Surg.* 2016 Apr-Jun; 5(2): 25.

358 **Ventral wall defects.** Omphalocele. This image is a derivative work, adapted from the following source, available under : Khan YA, Qureshi MA, Akhtar J. Omphalomesenteric duct cyst in an omphalocele: a rare association. *Pak J Med Sci.* 2013 May-Jun; 29(3): 866–868.

358 **Ventral wall defects.** Drawings of gastroschisis (left) and omphalocele (right). Courtesy of the Department of Health and Human Services.

359 **Intestinal atresia.** This image is a derivative work, adapted from the following source, available under : Saha M. Alimentary tract atresias associated with anorectal malformations: 10 years' experience. *J Neonatal Surg.* 2016 Oct-Dec; 5(4): 43. DOI: 10.21699/jns.v5i4.449.

359 **Hypertrophic pyloric stenosis.** This image is a derivative work, adapted from the following source, available under : Hassan RAA, Choo YU, Noraida R, et al. Infantile hypertrophic pyloric stenosis in postoperative esophageal atresia with tracheoesophageal fistula. *J Neonatal Surg.* 2015 Jul-Sep;4(3):32.

360 **Pancreas and spleen embryology.** Annular pancreas. This image is a derivative work, adapted from the following source, available under : Mahdi B, Selim S, Hassen T, et al. A rare cause of proximal intestinal obstruction in adults—annular pancreas: a case report. *Pan Afr Med J.* 2011;10:56. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

360 **Retroperitoneal structures.** This image is a derivative work, adapted from the following source, available under : Sammut J, Ahiaku E, Williams DT. Complete regression of renal tumour following ligation of an accessory renal artery during repair of an abdominal aortic aneurysm. *Ann R Coll Surg Engl.* 2012 Sep; 94(6): e198–e200. DOI: 10.1308/003588412X13373405384972.

362 **Digestive tract anatomy.** Histology of stomach wall. This image is a derivative work, adapted from the following source, available under : Alexander Klepnev.

362 **Digestive tract histology: Image A.** Courtesy of Dr. Michale Bonert.

362 **Digestive tract histology: Image B.** Courtesy of W. Ben Smith.

362 **Digestive tract histology: Images C, D, E.** This image is a derivative work, adapted from the following source, available under : Wikimedia Commons.

367 **Liver tissue architecture: Image A.** Portal triad. This image is a derivative work, adapted from the following source, available under : Liver development. In: Zorn AM. Stem book. Cambridge: Harvard Stem Cell Institute, 2008.

367 **Liver tissue architecture: Image B.** Kupffer cells. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

368 **Biliary structures.** Gallstones. This image is a derivative work, adapted from the following source, available under : J. Guntau. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

370 **Hernias: Image A.** Congenital diaphragmatic hernia. This image is a derivative work, adapted from the following source, available under : Tovar J. Congenital diaphragmatic hernia. *Orphanet J Rare Dis.* 2012;7:1. DOI: 10.1186/1750-1172-7-1.

372 **Gastrointestinal secretory products.** Histology of gastric pit. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

374 **Peyer patches.** This image is a derivative work, adapted from the following source, available under : Plainpaper. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

376 **Sialolithiasis.** This image is a derivative work, adapted from the following source, available under : Pastor-Ramos V, Cuervo-Diaz A, Aracil-Kessler L. Sialolithiasis. Proposal for a new minimally invasive procedure: piezoelectric surgery. *J Clin Exp Dent.* 2014 Jul;6(3):e295–e298. DOI: 10.4317/jced.51253.

376 **Salivary gland tumors.** Pleomorphic adenoma histology. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

376 **Achalasia.** This image is a derivative work, adapted from the following source, available under : Farnoosh Farrokhi and Michael F. Vaezi. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

377 **Esophageal pathologies: Image A.** Pneumomediastinum in Boerhaave syndrome. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

377 **Esophageal pathologies: Image B.** Esophageal varices on endoscopy. This image is a derivative work, adapted from the following source, available under : Costaguta A, Alvarez F. Etiology and management of hemorrhagic complications of portal hypertension in children. *Int J Hepatol.* 2012;2012:879163. DOI: 10.1155/2012/879163.

377 **Esophageal pathologies: Image C.** Esophageal varices on CT. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

377 **Esophageal pathologies: Image D.** Esophagitis. This image is a derivative work, adapted from the following source, available under : Takahashi Y, Nagata N, Shimbo T. Long-term trends in esophageal candidiasis prevalence and associated risk factors with or without HIV infection: lessons from an endoscopic study of 80,219 patients. *PLoS One.* 2015; 10(7): e0133589. DOI: 10.1371/journal.pone.0133589.

378 **Barrett esophagus: Image A.** Endoscopy. This image is a derivative work, adapted from the following source, available under : Coda S, Thillainayagam AV. State of the art in advanced endoscopic

(c) ketabton.com: The Digital Library

imaging for the detection and evaluation of dysplasia and early cancer of the gastrointestinal tract. *Clin Exp Gastroenterol.* 2014;7:133-150. DOI: 10.2147/CEG.S58157. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

378 **Barrett esophagus: Image B.** Goblet cells. This image is a derivative work, adapted from the following source, available under CC BY: Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

379 **Ménétriere disease.** This image is a derivative work, adapted from the following source, available under CC BY: Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

379 **Gastric cancer Tan Y, Fu J, Li X.** This image is a derivative work, adapted from the following source, available under CC BY: A minor (<50%) signet-ring cell component associated with poor prognosis in colorectal cancer patients: a 26-year retrospective study in China. *PLoS One.* 2015; 10(3): e0121944. DOI: 10.1371/journal.pone.0121944.

380 **Ulcer complications.** Free air under diaphragm in perforated ulcer. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.

381 **Malabsorption syndromes: Image A.** This image is a derivative work, adapted from the following source, available under CC BY: Celiac disease. Sedda S, Caruso R, Marafini I, et al. Pyoderma gangrenosum in refractory celiac disease: a case report. *BMC Gastroenterol.* 2013; 13: 162. DOI: 10.1186/1471-230X-13-162.

381 **Malabsorption syndromes: Image B.** *Tropheryma whippeli* on PAS stain. This image is a derivative work, adapted from the following source, available under CC BY: Tran HA. Reversible hypothyroidism and Whipple's disease. *BMC Endocr Disord.* 2006;6:3. DOI: 10.1186/1472-6823-6-3.

382 **Inflammatory bowel diseases: Image A.** "String sign" on barium swallow in Crohn disease. This image is a derivative work, adapted from the following source, available under CC BY: Al-Mofarreh MA, Al Mofleh IA, Al-Teimi IN, et al. Crohn's disease in a Saudi outpatient population: is it still rare? *Saudi J Gastroenterol.* 2009;15:111-116. DOI: 10.4103/1319-3767.45357. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

382 **Inflammatory bowel diseases: Images B (normal mucosa) and C (punched-out ulcers) in ulcerative colitis.** This image is a derivative work, adapted from the following source, available under CC BY: Ishikawa D, Ando T, Watanabe O, et al. Images of colonic real-time tissue sonoelastography correlate with those of colonoscopy and may predict response to therapy in patients with ulcerative colitis. *BMC Gastroenterol.* 2011;11:29. DOI: 10.1186/1471-230X-11-29.

383 **Appendicitis.** Fecalith. This image is a derivative work, adapted from the following source, available under CC BY: Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY.

383 **Diverticula of the GI tract: Image B.** Diverticulosis. This image is a derivative work, adapted from the following source, available under CC BY: Sartelli M, Moore FA, Ansaloni L, et al. A proposal for a CT driven classification of left colon acute diverticulitis. *World J Emerg Surg.* 2015;10:3. DOI: 10.1186/1749-7922-10-3.

383 **Diverticula of the GI tract: Image C.** This image is a derivative work, adapted from the following source, available under CC BY: Hupfeld L, Burcharth J, Pommergaard HC, Rosenberg J. The best choice of treatment for acute colonic diverticulitis with purulent peritonitis is uncertain. *Biomed Res Int.* 2014; 2014: 380607. DOI: 10.1155/2014/380607.

384 **Zenker diverticulum.** This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of Bernd Brägelmann.

385 **Maltotation.** This image is a derivative work, adapted from the following source, available under CC BY: Mathews R, Thenabadu S, Jaiganesh T. Abdominal pain with a twist. *Int J Emerg Med.* 2011;4:21. DOI: 10.1186/1865-1380-4-21.

385 **Intussusception: Image A.** Interoperative image of intussusception. This image is a derivative work, adapted from the following source, available under CC BY: Vasiliadis K, Kogopoulos E, Katsamakas M, et al. Ileoileal intussusception induced by a gastrointestinal stromal tumor. *World J Surg Oncol.* 2008;6:133. DOI: 10.1186/1477-7819-6-133.

385 **Intussusception: Image B.** Ultrasound showing target sign. This image is a derivative work, adapted from the following source, available under CC BY: Abbo O, Pinnagoda K, Micol LA. Osteosarcoma metastasis causing ileo-ileal intussusception. *World J Surg Oncol.* 2013 Aug 12;11(1):188. DOI: 10.1186/1477-7819-11-188.

386 **Volvulus.** Coffee bean sign. This image is a derivative work, adapted from the following source, available under CC BY: Yigit M, Turkdogan KA. Coffee bean sign, whirl sign and bird's beak sign in the diagnosis of sigmoid volvulus. *Pan Afr Med J.* 1014;19:56. DOI: 10.11604/pamj.2014.19.56.5142.

386 **Other intestinal disorders: Image A.** Necrosis due to occlusion of SMA. This image is a derivative work, adapted from the following source, available under CC BY: Van De Winkel N, Cheragwandi A, Nieboer K, et al. Superior mesenteric arterial branch occlusion causing partial jejunal ischemia: a case report. *J Med Case Rep.* 2012;6:48. DOI: 10.1186/1752-1947-6-48.

386 **Other intestinal disorders: Image B.** Loops of dilated bowel suggestive of small bowel obstruction. This image is a derivative work, adapted from the following source, available under CC BY: Welte FJ, Crosso M. Left-sided appendicitis in a patient with congenital gastrointestinal malrotation: a case report. *J Med Case Rep.* 2007;1:92. DOI: 10.1186/1752-1947-1-92.

386 **Other intestinal disorders: Image C.** Endoscopy showing dilated vessels. This image is a derivative work, adapted from the following source, available under CC BY: Gunjan D, Sharma V, Rana SS, et al. Small bowel bleeding: a comprehensive review. *Gastroenterol Rep.* 2014 Nov;2(4):262-75. DOI: 10.1093/gastro/gou025.

386 **Other intestinal disorders: Image D.** Pneumatosis intestinalis. This image is a derivative work, adapted from the following source, available under CC BY: Pelizzo G, Nakib G, Goruppi I, et al. Isolated colon ischemia with norovirus infection in preterm babies: a case series. *J Med Case Rep.* 2013;7:108. DOI: 10.1186/1752-1947-7-108.

387 **Colonic polyps: Image A.** This image is a derivative work, adapted from the following source, available under CC BY: M. Emannuel.

387 **Colonic polyps: Image B.** Adenomatous polyps. This image is a derivative work, adapted from the following source, available under CC BY: Shussman N, Wexner SD. Colorectal polyps and polyposis syndromes. *Gastroenterol Rep (Oxf).* 2014 Feb;2(1):1-15. DOI: 10.1093/gastro/got041.

387 **Colonic polyps: Image C.** This image is a derivative work, adapted from the following source, available under CC BY: Rehani B, Chasen RM, Dowdy Y, et al. Advanced adenoma diagnosis with FDG PET in a visibly normal mucosa: a case report. *J Med Case Reports.* 2007; 1: 99. DOI: 10.1186/1752-1947-1-99.

388 **Colorectal cancer: Image A.** Polyp. This image is a derivative work, adapted from the following source, available under CC BY: Takiyama A, Nozawa H, Ishihara S, et al. Secondary metastasis in the lymph node of the bowel invaded by colon cancer: a report of three cases. *World J Surg Oncol.* 2016; 14: 273. DOI: 10.1186/s12957-016-1026-y.

389 **Cirrhosis and portal hypertension: Image A.** Splenomegaly and liver nodularity in cirrhosis. This image is a derivative work, adapted from

(c) ketabton.com: The Digital Library

the following source, available under : Inversitus. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

389 **Cirrhosis and portal hypertension: Image B.** This image is a derivative work, adapted from the following source, available under : Blackburn PR, Hickey RD, Nace RA, et al. Silent tyrosinemia type I without elevated tyrosine or succinylacetone associated with liver cirrhosis and hepatocellular carcinoma. *Hum Mutat.* 2016 Oct; 37(10): 1097–1105.DOI: 10.1002/humu.23047.

391 **Alcoholic liver disease: Image B.** Mallory bodies. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

391 **Alcoholic liver disease: Image C.** Sclerosis in alcoholic cirrhosis. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

391 **Non-alcoholic fatty liver disease.** This image is a derivative work, adapted from the following source, available under : El-Karaksy HM, El-Koofy NM, Anwar GM, et al. Predictors of non-alcoholic fatty liver disease in obese and overweight Egyptian children: single center study. *Saudi J Gastroenterol.* 2011;17:40-46. DOI: 10.4103/1319-3767.74476.

392 **Hepatocellular carcinoma/hepatoma: Image A.** Gross specimen. Reproduced, with permission, from Jean-Christophe Fournet and Humpath.

392 **Other liver tumors.** Cavernous liver hemangioma. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

392 **α_1-antitrypsin deficiency.** Liver histology. This image is a derivative work, adapted from the following source, available under : Dr. Jerad M. Gardner. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

393 **Jaundice.** Yellow sclera. Courtesy of the Department of Health and Human Services and Dr. Thomas F. Sellers.

395 **Wilson disease.** This image is a derivative work, adapted from the following source, available under : Kodama H, Fujisawa C, Bhadhprasit W. Inherited copper transport disorders: biochemical mechanisms, diagnosis, and treatment. *Curr Drug Metab.* 2012 Mar; 13(3): 237–250. DOI: 10.2174/138920012799320455.

395 **Hemochromatosis.** Hemosiderin deposits. This image is a derivative work, adapted from the following source, available under : Mathew J, Leong MY, Morley N, et al. A liver fibrosis cocktail? Psoriasis, methotrexate and genetic hemochromatosis. *BMC Dermatol.* 2005;5:12. DOI: 10.1186/1471-5945-5-12.

396 **Cholelithiasis and related pathologies: Image A.** Gross specimen of gallstones. This image is a derivative work, adapted from the following source, available under : Courtesy of M. Emmanuel.

396 **Cholelithiasis and related pathologies: Image B.** Large gallstone. This image is a derivative work, adapted from the following source, available under : Spangler R, Van Pham T, Khoujah D, et al. Abdominal emergencies in the geriatric patient. *Int J Emerg Med.* 2014; 7: 43. DOI: 10.1186/s12245-014-0043-2.

397 **Cholelithiasis and related pathologies: Image C.** Porcelain gallbladder. This image is a derivative work, adapted from the following source, available under : Fred H, van Dijk H. Images of memorable cases: case 19. Connexions Web site. December 4, 2008. Available at: http://cnx.org/content/m14939/1.3/. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

397 **Acute pancreatitis: Image A.** Acute exudative pancreatitis. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

397 **Acute pancreatitis: Image B.** Pancreatic pseudocyst. This image is a derivative work, adapted from the following source, available under : Thomas Zimmerman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

397 **Chronic pancreatitis.** This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

398 **Pancreatic adenocarcinoma: Image A.** Histology. This image is a derivative work, adapted from the following source, available under : KGH. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

398 **Pancreatic adenocarcinoma: Image B.** CT scan. Courtesy of MBq. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

Hematology and Oncology

406 **Neutrophils.** Courtesy of B. Lennert.

407 **Erythrocytes.** Courtesy of the Department of Health and Human Services and Drs. Noguchi, Rodgers, and Schechter.

407 **Thrombocytes (platelets).** This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

407 **Monocytes.** This image is a derivative work, adapted from the following source, available under : Dr. Graham Beards. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

407 **Macrophages.** This image is a derivative work, adapted from the following source, available under : De Tommasi AS, Otranto D, Furlanello T, et al. Evaluation of blood and bone marrow in selected canine vector-borne diseases. *Parasit Vectors.* 2014;7:534. DOI: 10.1186/s13071-014-0534-2.

408 **Eosinophils.** This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman.

408 **Basophils.** This image is a derivative work, adapted from the following source, available under : Dr. Erhabor Osaro. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

408 **Mast cells.** Courtesy of Wikimedia Commons.

408 **Dendritic cells.** This image is a derivative work, adapted from the following source, available under : Cheng J-H, Lee S-Y, Lien Y-Y, et al. Immunomodulating activity of *Nymphaea rubra* roxb. extracts: activation of rat dendritic cells and improvement of the TH1 immune response. *Int J Mol Sci.* 2012;13:10722-10735. DOI: 10.3390/ijms130910722.

409 **Lymphocytes.** This image is a derivative work, adapted from the following source, available under : Fickleandfreckled.

409 **Plasma cells.** Courtesy of the Department of Health and Human Services and Dr. Francis W. Chandler. The image may have been

(c) ketabton.com: The Digital Library

modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

415 **RBC morphology: Image J.** Sickle cell. Courtesy of the Department of Health and Human Services and the Sickle Cell Foundation of Georgia, Jackie George, and Beverly Sinclair.

416 **RBC inclusions: Image A.** Ringed sideroblast. This image is a derivative work, adapted from the following source, available under : Paulo Henrique Orlandi Mourao. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

416 **RBC inclusions: Image B.** Howell-Jolly bodies. This image is a derivative work, adapted from the following source, available under : Serio B, Pezzullo L, Giudice V, et al. OPSI threat in hematological patients. *Transl Med UniSa.* 2013 May-Aug;6:2-10.

416 **RBC inclusions: Image C.** Bsaophilic stippling. This image is a derivative work, adapted from the following source, available under : Dr. Erhabor Osaro.

416 **RBC inclusions: Image D.** Pappenheimer bodies. This image is a derivative work, adapted from the following source, available under : Paulo Henrique Orlandi Mourao. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

419 **Microcytic, hypochromic anemias: Image A.** This image is a derivative work, adapted from the following source, available under : Bock F, Borucki K, Vorwerk P, et al. A two-and-a-half-year-old breastfed toddler presenting with anemia: a case report. *BMC Res Notes.* 2014; 7: 917. DOI: 10.1186/1756-0500-7-917.

419 **Microcytic, hypochromic anemia: Image D.** Lead lines in lead poisoning. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.

419 **Microcytic, hypochromic anemia: Image E.** Sideroblastic anemia. This image is a derivative work, adapted from the following source, available under : Paulo Henrique Orlandi Moura. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

420 **Macrocytic anemias.** Megaloblastic anemia. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Ed Uthman.

422 **Intrinsic hemolytic anemias.** This image is a derivative work, adapted from the following source, available under : El Ariss AB, Younes M, Matar J. Prevalence of sickle cell trait in the southern suburb of Beirut, Lebanon. *Mediterr J Hematol Infect Dis.* 2016; 8(1): e2016015. DOI: 10.4084/MJHID.2016.015.

425 **Heme synthesis, porphyrias, and lead poisoning: Image A.** Basophilic stippling in lead poisoning. This image is a derivative work, adapted from the following source, available under : van Dijk HA, Fred HL. Images of memorable cases: case 81. Connexions Web site. December 3, 2008. Available at http://cnx.org/contents/3196bf3e-1e1e-4c4d-a1ac-d4fc9ab65443@4@4.

425 **Heme synthesis, porphyrias, and lead poisoning: Image B.** Porphyria cutanea tarda. This image is a derivative work, adapted from the following source, available under : Bovenschen HJ, Vissers WHPM. Primary hemochromatosis presented by porphyria cutanea tarda: a case report. *Cases J.* 2009;2:7246. DOI: 10.4076/1757-1626-2-7246.

426 **Coagulation disorders.** This image is a derivative work, adapted from the following source, available under : Lakjiri S, Mernissi FZ. Tabetic arthropathy revealing neurosyphilis: a new observation. *Pan Afr Med J.* 2014; 18: 198. DOI: 10.11604/pamj.2014.18.198.4893.

429 **Hodgkin lymphoma.** This image is a derivative work, adapted from the following source, available under : Knecht H, Righolt C, Mai S. Genomic instability: the driving force behind refractory/relapsing Hodgkin's lymphoma. *Cancers (Basel).* 2013 Jun; 5(2): 714–725. DOI: 10.3390/cancers5020714.

430 **Non-Hodgkin lymphoma: Image B.** This image is a derivative work, adapted from the following source, available under : Bi CF, Tang Y, Zhang WY, et al. Sporadic Burkitt lymphomas of children and adolescents in Chinese: a clinicopathological study of 43 cases. *Diagn Pathol.* 2012;7:72. DOI:10.1186/1746-1596-7-72.

430 **Non-Hodgkin lymphoma: Image C.** This image is a derivative work, adapted from the following source, available under : Mansour A, Qandeel M, Abdel-Razeq H, et al. MR imaging features of intracranial primary CNS lymphoma in immune competent patients. *Cancer Imaging.* 2014;14(1):22. DOI: 10.1186/1470-7330-14-22.

430 **Non-Hodgkin lymphoma: Image D.** This image is a derivative work, adapted from the following source, available under : Chaudhary S, Bansal C, Ranga U, et al. Erythrodermic mycosis fungoides with hypereosinophilic syndrome: a rare presentation. *Ecancermedicalscience.* 2013;7:337. DOI:10.3332/ecancer.2013.337

431 **Plasma cell dyscrasias: Image C.** This image is a derivative work, adapted from the following source, available under : Mehrotra R, Singh M, Singh PA, et al. Should fine needle aspiration biopsy be the first pathological investigation in the diagnosis of a bone lesion? An algorithmic approach with review of literature. *Cytojournal.* 2007; 4: 9. DOI: 10.1186/1742-6413-4-9.

432 **Myelodysplastic syndromes.** This image is a derivative work, adapted from the following source, available under : Lukaszewska J, Allison RW, Stepkowska J. Congenital Pelger-Huët anomaly in a Danish/Swedish farmdog: case report. *Acta Vet Scand.* 2011; 53(1): 14. DOI: 10.1186/1751-0147-53-14.

433 **Leukemias: Image A.** This image is a derivative work, adapted from the following source, available under : Chiaretti S, Zini G, Bassan R. Diagnosis and subclassification of acute lymphoblastic leukemia. *Mediterr J Hematol Infect Dis.* 2014; 6(1): e2014073. DOI: 10.4084/MJHID.2014.073.

433 **Leukemias: Image C.** Hairy cell leukemia. This image is a derivative work, adapted from the following source, available under : Chan SM, George T, Cherry AM, et al. Complete remission of primary plasma cell leukemia with bortezomib, doxorubicin, and dexamethasone: a case report. *Cases J.* 2009;2:121. DOI: 10.1186/1757-1626-2-121.

433 **Chronic myeloproliferative disorders: Image A.** Erythromelalgia in polycythemia vera. This image is a derivative work, adapted from the following source, available under : Fred H, van Dijk H. Images of memorable cases: case 151. Connexions Web site. December 4, 2008. Available at http://cnx.org/content/m14932/1.3/.

433 **Chronic myeloproliferative disorders: Image C.** Myelofibrosis. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Ed Uthman.

434 **Langerhans cell histiocytosis: Image A.** Lytic bone lesion. This image is a derivative work, adapted from the following source, available under : Dehkordi NR, Rajabi P, Naimi A, et al. Langerhans cell histiocytosis following Hodgkin lymphoma: a case report from Iran. *J Res Med Sci.* 2010;15:58-61. PMCID PMC3082786.

434 **Langerhans cell histiocytosis: Image B.** Birbeck granules. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

435 **Hemophagocytic lymphohistiocytosis.** This image is a derivative work, adapted from the following source, available under : Kashif M, Tariq H, Ijaz M. Disseminated histoplasmosis and secondary hemophagocytic syndrome in a non-HIV patient. *Case Rep Crit Care.* 2015; 2015: 295735. DOI: 10.1155/2015/295735.

(c) ketabton.com: The Digital Library

436 **Warfarin.** This image is a derivative work, adapted from the following source, available under : Bakoyiannis C, Karaolanis G, Patelis N. Dabigatran in the treatment of warfarin-induced skin necrosis: A new hope. *Case Rep Dermatol Med.* 2016; 2016: 3121469. DOI: 10.1155/2016/3121469.

Musculoskeletal, Skin, and Connective Tissue

446 **Rotator cuff muscles.** Glenohumeral instability. This image is a derivative work, adapted from the following source, available under : Koike Y, Sano H, Imamura I, et al. Changes with time in skin temperature of the shoulders in healthy controls and a patient with shoulder-hand syndrome. *Ups J Med Sci* 2010;115:260-265. DOI: 10.3109/03009734.2010.503354. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

448 **Brachial plexus lesions: Image A.** Cervical rib. This image is a derivative work, adapted from the following source, available under : Dahlin LB, Backman C, Duppe H, et al. Compression of the lower trunk of the brachial plexus by a cervical rib in two adolescent girls: case reports and surgical treatment. *J Brachial Plex Peripher Nerve Inj.* 2009;4:14. DOI: 10.1186/1749-7221-4-14.

448 **Brachial plexus lesions: Image B.** Winged scapula. This image is a derivative work, adapted from the following source, available under : Boukhris J, Boussouga M, Jaafar A, et al. Stabilisation dynamique d'un winging scapula (à propos d'un cas avec revue de la littérature). *Pan Afr Med J.* 2014; 19: 331. DOI: 10.11604/pamj.2014.19.331.3429.

449 **Wrist region: Image B.** Anatomic snuff box. This image is a derivative work, adapted from the following source, available under : Rhemrev SJ, Ootes D, Beeres FJP, et al. Current methods ofdiagnosis and treatment of scaphoid fractures. *Int J Emerg Med.* 2011;4:4. DOI: 10.1186/1865-1380-4-4.

456 **Motoneuron action potential to muscle contraction: Image A.** This image is a derivative work, adapted from the following source, available under : Ottenheijm CAC, Heunks LMA, Dekhuijzen RPN. Diaphragm adaptations in patients with COPD. *Respir Res.* 2008; 9(1): 12. DOI: 10.1186/1465-9921-9-12.

459 **Wrist and hand injuries: Image A.** Metacarpal neck fracture. This image is a derivative work, adapted from the following source, available under : Bohr S, Pallua N. Early functional treatment and modern cast making for indications in hand surgery. *Adv Orthop.* 2016; 2016: 5726979. DOI: 10.1155/2016/5726979.

459 **Wrist and hand injuries: Image B.** Thenar eminence atrophy in carpal tunnel syndrome. Courtesy of Dr. Harry Gouvas.

460 **Common hip and knee conditions: Image A.** ACL tear. This image is a derivative work, adapted from the following source, available under : Chang MJ, Chang CB, Choi J-Y, et al. Can magnetic resonance imaging findings predict the degree of knee joint laxity in patients undergoing anterior cruciate ligament reconstruction? *BMC Musculoskelet Disord.* 2014;15:214. DOI: 10.1186/1471-2474-15-214. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

460 **Common hip and knee conditions: Images B (prepatellar bursitis) and C (Baker cyst).** This image is a derivative work, adapted from the following source, available under : Hirji Z, Hunhun JS, Choudur HN. Imaging of the bursae. *J Clin Imaging Sci.* 2011;1:22. DOI: 10.4103/2156-7514.80374. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

462 **Common pediatric fractures: Image A.** Greenstick fracture. This image is a derivative work, adapted from the following source, available under : Randsborg PH, Sivertsen EA. Classification of distal radius fractures in children: good inter- and intraobserver reliability, which improves with clinical experience. *BMC Musculoskelet Disord.* 2013;13:6. DOI: 10.1186/1471-2474-13-6.

462 **Common pediatric fractures: Image B.** Torus (buckle) fracture. This image is a derivative work, adapted from the following source, available under : Aksel Seyahi, et al. Tibial torus and toddler's fractures misdiagnosed as transient synovitis: a case series. *J Med Case Reports.* 2011; 5: 305. DOI: 10.1186/1752-1947-5-305.

462 **Osteoporosis.** Vertebral compression fractures of spine. This image is a derivative work, adapted from the following source, available under : Imani F, Gharaei H, Rahimzadeh P, et al. Management of painful vertebral compression fracture with kyphoplasty in a sever cardio-respiratory compromised patient. *Anesth Pain Med.* 2012 summer;2(1):42–45. DOI: 10.5812/aapm.5030.

463 **Osteopetrosis.** This image is a derivative work, adapted from the following source, available under : Kant P, Sharda N, Bhowate RR. Clinical and radiological findings of autosomal dominant osteopetrosis type II: a case report. *Case Rep Dent.* 2013;2013:707343. DOI: 10.1155/2013/707343.

463 **Osteomalacia/rickets: Image A, left.** Clinical photo. This image is a derivative work, adapted from the following source, available under : Linglart A, Biosse-Duplan M, Briot K, et al. Therapeutic management of hypophosphatemic rickets from infancy to adulthood. *Endocr Connect.* 2014;3:R13-R30. DOI: 10.1530/EC-13-0103.

463 **Osteomalacia/rickets: Image B.** Rachitic rosary on chest X-ray. This image is a derivative work, adapted from the following source, available under : Essabar L, Meskini T, Ettair S, et al. Malignant infantile osteopetrosis: case report with review of literature. *Pan Afr Med J.* 2014;17:63. DOI: 10.11604/pamj.2014.17.63.3759.

463 **Osteitis deformans.** Thickened calvarium. This image is a derivative work, adapted from the following source, available under : Dawes L. Paget's disease. [Radiology Picture of the Day Website]. Published June 21, 2007. Available at http://www.radpod.org/2007/06/21/pagets-disease/.

463 **Avascular necrosis of bone.** Bilateral necrosis of femoral head. This image is a derivative work, adapted from the following source, available under : Ding H, Chen S-B, Lin S, et al. The effect of postoperative corticosteroid administration on free vascularized fibular grafting for treating osteonecrosis of the femoral head. *Sci World J.* 2013;708014. DOI: 10.1155/2013/708014. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

465 **Primary bone tumors: Image A.** Osteochondroma. This image is a derivative work, adapted from the following source, available under : Lucien Monfils. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

465 **Primary bone tumors: Image B.** Osteoid osteoma. This image is a derivative work, adapted from the following source, available under : Jankharia B, Burute N. Percutaneous radiofrequency ablation for osteoid osteoma: how we do it. *Indian J Radiol Imaging.* 2009 Feb; 19(1): 36–42. DOI: 10.4103/0971-3026.44523.

465 **Primary bone tumors: Image C.** Giant cell tumor. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.

465 **Primary bone tumors: Image D.** This image is a derivative work, adapted from the following source, available under : Xu SF, Yu XC, Zu M, et al. Limb function and quality of life after various reconstruction methods according to tumor location following resection of osteosarcoma in distal femur. *BMC Musculoskelet Disord.* 2014; 15: 453. DOI: 10.1186/1471-2474-15-453.

465 **Primary bone tumors: Image E.** Starburst pattern in osteosarcoma. This image is a derivative work, adapted from the following source, available under : Ding H, Yu G, Tu Q, et al. Computer-aided resection and endoprosthesis design for the management of

(c) ketabton.com: The Digital Library

malignant bone tumors around the knee: outcomes of 12 cases. *BMC Musculoskelet Disord.* 2013; 14: 331. DOI: 10.1186/1471-2474-14-331.

466 Osteoarthritis vs rheumatoid arthritis: Image A. Histology of rheumatoid nodule. This image is a derivative work, adapted from the following source, available under CC BY: Gomez-Rivera F, El-Naggar AK, Guha-Thakurta N, et al. Rheumatoid arthritis mimicking metastatic squamous cell carcinoma. *Head Neck Oncol.* 2011;3:26. DOI: 10.1186/1758-3284-3-26.

467 Gout: Image B. Uric acid crystals under polarized light. This image is a derivative work, adapted from the following source, available under CC BY-SA: Robert J. Galindo. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

467 Gout: Image C. Podagra. This image is a derivative work, adapted from the following source, available under CC BY: Roddy E. Revisiting the pathogenesis of podagra: why does gout target the foot? *J Foot Ankle Res.* 2011;4:13. DOI: 10.1186/1757-1146-4-13.

467 Calcium pyrophosphate deposition disease. Calcium phosphate crystals. This image is a derivative work, adapted from the following source, available under CC BY: Dieppe P, Swan A. Identification of crystals in synovial fluid. *Ann Rheum Dis.* 1999 May;58(5):261–263.

468 Sjögren syndrome: Image A. Lymphocytic infiltration. Courtesy of the Department of Health and Human Services.

468 Sjögren syndrome: Image B. Dry tongue. This image is a derivative work, adapted from the following source, available under CC BY: Negrato CA, Tarzia O. Buccal alterations in diabetes mellitus. *Diabetol Metab Syndr.* 2010;2:3. DOI: 10.1186/1758-5996-2-3.

468 Septic arthritis. Joint effusion. This image is a derivative work, adapted from the following source, available under CC BY-SA: Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

469 Seronegative spondyloarthropathies: Image C, left. Bamboo spine. This image is a derivative work, adapted from the following source, available under CC BY-SA: Stevenfruitsmaak. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

469 Seronegative spondyloarthropathies: Image C, right. Bamboo spine. Courtesy of Heather Hawker.

471 Polymyositis/dermatomyositis: Image A. Groton papules of dermatomyositis. This image is a derivative work, adapted from the following source, available under CC BY: *Pan Afr Med J.*2015; 21: 89. DOI: 10.11604/pamj.2015.21.89.6971.

472 Raynaud phenomenon. This image is a derivative work, adapted from the following source, available under CC BY-SA: Jamclaassen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

474 Epithelial cell junctions: Image A. Intracellular membrane. This image is a derivative work, adapted from the following source, available under CC BY: Tang VW. Proteomic and bioinformatic analysis of epithelial tight junction reveals an unexpected cluster of synaptic molecules. *Biol Direct.* 2006; 1: 37. DOI: 10.1186/1745-6150-1-37.

474 Epithelial cell junctions: Image B. Large, electron-dense actin structures within adherens junction. This image is a derivative work, adapted from the following source, available under CC BY: Taylor RR, Jagger DJ, Saeed SR, et al. Characterizing human vestibular sensory epithelia for experimental studies: new hair bundles on old tissue and implications for therapeutic interventions in ageing. *Neurobiol Aging.* 2015 Jun;36(6):2068–2084. DOI: 10.1016/j.neurobiolaging.2015.02.013.

474 Epithelial cell junctions: Image C. Desmosome. This image is a derivative work, adapted from the following source, available under CC BY: Massa F, Devader C, Lacas-Gervais S, et al. Impairement of HT29 cancer cells cohesion by the soluble form of neurotensin receptor-3. *Genes Cancer.* 2014 Jul; 5(7-8):240–249. DOI: 10.18632/genesandcancer.22.

474 Epithelial cell junctions: Image D. Gap junction. This image is a derivative work, adapted from the following source, available under CC BY: Shu X, Lev-Ram V, Deerinck TJ. A Genetically encoded tag for correlated light and electron microscopy of intact cells, tissues, and organisms. *PLoS Biol.* 2011 Apr; 9(4): e1001041. DOI: 10.1371/journal.pbio.1001041.

474 Epithelial cell junctions: Image E. Hemidesmosome. This image is a derivative work, adapted from the following source, available under CC BY: Nguyen NM, Pulkkinen L, Schlueter JA, et al. Lung development in laminin gamma2 deficiency: abnormal tracheal hemidesmosomes with normal branching morphogenesis and epithelial differentiation. *Respir Res.* 2006 Feb 16;7:28. DOI: 10.1186/1465-9921-7-28.

476 Seborrheic dermatitis. This image is a derivative work, adapted from the following source, available under CC BY-SA: Roymishali.

477 Common skin disorders: Image O. Urticaria. This image is a derivative work, adapted from the following source, available under CC BY-SA: Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under CC BY-SA.

478 Vascular tumors of skin: Image C. Glomus tumor under fingernail. This image is a derivative work, adapted from the following source, available under CC BY: Hazani R, Houle JM, Kasdan ML, et al. Glomus tumors of the hand. *Eplasty.* 2008;8:e48. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

479 Skin infections: Image C. Erysipelas. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of Klaus D. Peter.

480 Autoimmune blistering skin disorders: Image D. Bullous pemphigoid on immunofluorescence. This image is a derivative work, adapted from the following source, available under CC BY: Courtesy of M. Emmanuel.

484 Skin cancer: Image D. Basal cell palisading nuclei. This image is a derivative work, adapted from the following source, available under CC BY: Yuri T. Jadotte, MD, et al. Superficial spreading basal cell carcinoma of the face: a surgical challenge. *Eplasty.* 2010; 10: e46. Published online 2010 Jun 21.

Neurology and Special Senses

491 Holoprosencephaly: Image A. This image is a derivative work, adapted from the following source, available under CC BY: Alorainy IA, Barlas NB, Al-Boukai AA. Pictorial essay: infants of diabetic mothers. *Indian J Radiol Imaging.* 2010 Aug;20(3):174-81. DOI: 10.4103/0971-3026.69349.

492 Posterior fossa malformations: Image A. Chiari I malformation. This image is a derivative work, adapted from the following source, available under CC BY: Toldo I, De Carlo D, Mardari R, et al. Short lasting activity-related headaches with sudden onset in children: a case-based reasoning on classification and diagnosis. *J Headache Pain.* 2013;14(1):3. DOI: 10.1186/1129-2377-14-3.

492 Posterior fossa malformations: Image B. Dandy-Walker malformation. This image is a derivative work, adapted from the following source, available under CC BY: Krupa K, Bekiesinska-Figatowska M. Congenital and acquired abnormalities of the corpus callosum: a pictorial essay. *Biomed Res Int.* 2013;2013:265619. DOI: 10.1155/2013/265619.

FAS1_2019_20_ImageAck.indd 740 11/8/19 4:50 PM

(c) ketabton.com: The Digital Library

492 **Syringomyelia.** Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.

494 **Myelin.** Myelinated neuron. Courtesy of the Electron Microscopy Facility at Trinity College.

495 **Chromatolysis.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonnert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

499 **Limbic system: Image A.** This image is a derivative work, adapted from the following source, available under : Schopf V, Fischmeister FP, Windischberger C, et al. Effects of individual glucose levels on the neuronal correlates of emotions. *Front Hum Neurosci.* 2013 May 21;7:212. DOI: 10.3389/fnhum.2013.00212.

499 **Cerebellum.** This image is a derivative work, adapted from the following source, available under : Jarius S, Wandinger KP, Horn S, et al. A new Purkinje cell antibody (anti-Ca) associated with subacute cerebellar ataxia: immunological characterization. *J Neuroinflammation.* 2010;7: 21. DOI: 10.1186/1742-2094-7-21.

500 **Basal ganglia.** This image is a derivative work, adapted from the following source, available under : Rudger P, Jaunmuktane Z, Adlard P, et al. Iatrogenic CJD due to pituitary-derived growth hormone with genetically determined incubation times of up to 40 years. *Brain.* 2015 Nov; 138(11): 3386–3399. DOI: 10.1093/brain/awv235.

502 **Cerebral arteries—cortical distribution.** Cortical watershed areas. This image is a derivative work, adapted from the following source, available under : Isabel C, Lecler A, Turc G, et al. Relationship between watershed infarcts and recent intra plaque haemorrhage in carotid atherosclerotic plaque. *PLoS One.* 2014;9(10):e108712. DOI: 10.1371/journal.pone.0108712.

503 **Dural venous sinuses.** This image is a derivative work, adapted from the following source, available under : Cikla U, Aagaard-Kienitz B, Turski PA, et al. Familial perimesencephalic subarachnoid hemorrhage: two case reports. *J Med Case Rep.* 2014;8. DOI: 10.1186/1752-1947-8-380. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

508 **Spinal cord and associated tracts: Image A.** Spinal cord cross-section. This image is a derivative work, adapted from the following source, available under : Regents of University of Michigan Medical School.

512 **Neonatal interventricular hemorrhage.** This image is a derivative work, adapted from the following source, available under : Shooman D, Portess H, Sparrow O. A review of the current treatment methods for posthaemorrhagic hydrocephalus of infants. *Cerebrospinal Fluid Res.* 2009;6:1. DOI: 10.1186/1743-8454-6-1.

513 **Intracranial hemorrhage: Images A and B.** Axial CT of brain showing epidural blood. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

513 **Intracranial hemorrhage: Image C.** Subdural hematoma. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

513 **Intracranial hemorrhage: Image E.** Subarachnoid hemorrhage. This image is a derivative work, adapted from the following source, available under : Hakan T, Turk CC, Celik H. Intra-operative real time intracranial subarachnoid haemorrhage during glial tumour resection: a case report. *Cases J.* 2008;1:306. DOI: 10.1186/1757-1626-1-306. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

515 **Effects of strokes: Image A.** Large abnormality of the left MCA territory. This image is a derivative work, adapted from the following source, available under : Hakimelahi R, Yoo AJ, He J, et al. Rapid identification of a major diffusion/perfusion mismatch in distal internal carotid artery or middle cerebral artery ischemic stroke. *BMC Neurol.* 2012 Nov 5;12:132. DOI: 10.1186/1471-2377-12-132. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

515 **Effects of strokes: Image B.** MRI diffusion weighted image shows a hypersensitive lesion on posterior limb of internal capsular. This image is a derivative work, adapted from the following source, available under : Zhou L, Ni J, Yao M, et al. High-resolution MRI findings in patients with capsular warning syndrome. *BMC Neurol.* 2014;14:16. DOI: 10.1186/1471-2377-14-16.

515 **Effects of strokes: Image C.** This image is a derivative work, adapted from the following source, available under : Nouh A, Remke J, Ruland S. Ischemic posterior circulation stroke: a review of anatomy, clinical presentations, diagnosis, and current management. *Front Neurol.* 2014 Apr 7;5:30. DOI: 10.3389/fneur.2014.00030.

515 **Effects of strokes: Image D.** This image is a derivative work, adapted from the following source, available under : Mittal P, Kalia V, Dua S. Pictorial essay: Susceptibility-weighted imaging in cerebral ischemia. *Indian J Radiol Imaging.* 2010 Nov; 20(4): 250–253. DOI: 10.4103/0971-3026.73530.

515 **Diffuse axonal injury.** Moenninghoff C, Kraff O, Maderwald S, et al. Diffuse axonal injury at ultra-high field MRI. *PLoS One.* 2015;10(3):e0122329. DOI: 10.1371/journal.pone.0122329.

516 **Aneurysms.** This image is a derivative work, adapted from the following source, available under : Kayhan A, Koc O, Keskin S. The role of bone subtraction computed tomographic angiography in determining intracranial aneurysms in non-traumatic subarachnoid hemorrhage. *Iran J Radiol.* 2014 May; 11(2): e12670. DOI: 10.5812/iranjradiol.12670.

521 **Neurodegenerative disorders: Image A.** Lewy body in substantia nigra. This image is a derivative work, adapted from the following source, available under : Werner CJ, Heyny-von Haussen R, Mall G, et al. Parkinson's disease. *Proteome Sci.* 2008;6:8. DOI: 10.1186/1477-5956-6-8. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

521 **Neurodegenerative disorders: Image B.** Gross specimen of normal brain. This image is a derivative work, adapted from the following source, available under : Niedowicz DM, Nelson PT, Murphy MP. Alzheimer's disease: pathological mechanisms and recent insights. *Curr Neuropharmacol.* 2011 Dec;9(4):674-84. DOI: 10.2174/157015911798376181.

521 **Neurodegenerative disorders: Images C (brain atrophy in Alzheimer disease) and F (atrophy in frontotemporal dementia).** This image is a derivative work, adapted from the following source, available under : Niedowicz DM, Nelson PT, Murphy MP. Alzheimer's disease: pathological mechanisms and recent insights. *Curr Neuropharmacol.* 2011 Dec;9(4):674-84. DOI: 10.2174/157015911798376181.

521 **Neurodegenerative disorders: Image G.** Frontotemporal dementia: Pick bodies in frontotemporal demendia (Pick disease). This image is a derivative work, adapted from the following source, available under : Neumann M. Molecular neuropathology of TDP-43 proteinopathies. *Int J Mol Sci.* 2009 Jan; 10(1): 232–246. DOI: 10.3390/ijms10010232.

521 **Neurodegenerative disorders: Image H.** Spongiform changes in brain in Creutzfeld-Jacob disease. This image is a derivative work, adapted from the following source, available under : DRdoubleB. The

(c) ketabton.com: The Digital Library

image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

522 Hydrocephalus: Image B. Communicating hydrocephalus. This image is a derivative work, adapted from the following source, available under : Torres-Martin M, Pena-Granero C, Carceller F, et al. Homozygous deletion of *TNFRSF4*, *TP73*, *PPAP2B* and *DPYD* at 1p and *PDCD5* at 19q identified by multiplex ligation-dependent probe amplification (MLPA) analysis in pediatric anaplastic glioma with questionable oligodendroglial component. *Mol Cytogenet.* 2014;7:1. DOI: 10.1186/1755-8166-7-1.

522 Hydrocephalus: Image C. Ex vacuo ventriculomegaly. This image is a derivative work, adapted from the following source, available under : Ghetti B, Oblak AL, Boeve BF, et al. Frontotemporal dementia caused by microtubule-associated protein tau gene (*MAPT*) mutations: a chameleon for neuropathology and neuroimaging. Neurophathol Appl Neurobiol. 2015 Feb;41(1):24-46. DOI: 10.1111/nan.12213.

523 Multiple sclerosis. Periventricular plaques. This image is a derivative work, adapted from the following source, available under : Dooley MC, Foroozan R. Optic neuritis. *J Ophthalmic Vis Res.* 2010 Jul;5(3):182–187.

524 Other demyelinating and dysmyelinating diseases: Image B. Progressive multifocal leukoencephalopathy. This image is a derivative work, adapted from the following source, available under : Garrote H, de la Fuente A, Ona R, et al. Long-term survival in a patient with progressive multifocal leukoencephalopathy after therapy with rituximab, fludarabine and cyclophosphamide for chronic lymphocytic leukemia. *Exp Hematol Oncol.* 2015;4:8. DOI: 10.1186/s40164-015-0003-4.

524 Other demyelinated and dysmyelinating disorders: Image A. Central pontine myelinolysis. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

525 Neurocutaneous disorders: Image A. Sturge-Weber syndrome and port wine stain. This image is a derivative work, adapted from the following source, available under : Babaji P, Bansal A, Krishna G, et al. Sturge-Weber syndrome with osteohypertrophy of maxilla. *Case Rep Pediatr.* 2013. DOI: 10.1155/2013/964596.

525 Neurocutaneous disorders: Image B. Leptomeningeal angioma in Sturge-Weber syndrome. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.

525 Neurocutaneous disorders: Image C. Tuberous sclerosis. This image is a derivative work, adapted from the following source, available under : Fred H, van Dijk H. Images of memorable cases: case 143. Connexions Web site. December 4, 2008. Available at: http://cnx.org/content/m14923/1.3/.

525 Neurocutaneous disorders: Image D. Ash leaf spots in tuberous sclerosis. This image is a derivative work, adapted from the following source, available under : Tonekaboni SH, Tousi P, Ebrahimi A, et al. Clinical and para clinical manifestations of tuberous sclerosis: a cross sectional study on 81 pediatric patients. *Iran J Child Neurol.* 2012;6:25-31. PMCID PMC3943027.

525 Neurocutaneous disorders: Image E. Angiomyolipoma in tuberous sclerosis. This image is a derivative work, adapted from the following source, available under : KGH. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

525 Neurocutaneous disorders: Image F. Café-au-lait spots in neurofibromatosis. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

525 Neurocutaneous disorders: Image G. Lisch nodules in neurofibromatosis. Courtesy of the Department of Health and Human Services.

525 Neurocutaneous disorders: Image H. Cutaneous neurofibromas. This image is a derivative work, adapted from the following source, available under : Kim BK, Choi YS, Gwoo S, et al. Neurofibromatosis type 1 associated with papillary thyroid carcinoma incidentally detected by thyroid ultrasonography: a case report. *J Med Case Rep.* 2012;6:179. DOI: 10.1186/1752-1947-6-179.

525 Neurocutaneous disorders: Image I. Cerebellar hemangioblastoma histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

525 Neurocutaneous disorders: Image J. Brainstem and spinal cord hemangioblastomas in von Hippel-Lindau disease. This image is a derivative work, adapted from the following source, available under : Park DM, Zhuang Z, Chen L, et al. von Hippel-Lindau disease-associated hemangioblastomas are derived from embryologic multipotent cells. *PLoS Med.* 2007 Feb;4(2):e60. DOI: 10.1371/journal.pmed.0040060.

526 Adult primary brain tumors: Image A. This image is a derivative work, adapted from the following source, available under : Rossmeisl JH, Clapp K, Pancotto TE. Canine butterfly glioblastomas: A neuroradiological review. *Front Vet Sci.* 2016; 3: 40. DOI: 10.3389/fvets.2016.00040.

526 Adult primary brain tumors: Image B. Glioblastoma multiforme histology. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

526 Adult primary brain tumors: Image C. Oligodendroglioma in frontal lobes. This image is a derivative work, adapted from the following source, available under : Celzo FG, Venstermans C, De Belder F, et al. Brain stones revisited—between a rock and a hard place. *Insights Imaging.* 2013 Oct;4(5):625-35. DOI: 10.1007/s13244-013-0279-z.

526 Adult primary brain tumors: Image D. Oligodendroglioma, "fried egg" cells. This image is a derivative work, adapted from the following source, available under : Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

526 Adult primary brain tumors: Image E. Meningioma with dural tail. This image is a derivative work, adapted from the following source, available under : Smits A, Zetterling M, Lundin M, et al. Neurological impairment linked with cortico-subcortical infiltration of diffuse low-grade gliomas at initial diagnosis supports early brain plasticity. *Front Neurol.* 2015;6:137. DOI: 10.3389/fneur.2015.00137.

526 Adult primary brain tumors: Image F. Meningioma, psammoma bodies. This image is a derivative work, adapted from the following source, available under : Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

526 Adult primary brain tumors: Image G. Cerebellar hemangioblastoma. This image is a derivative work, adapted from the following source, available under : Park DM, Zhengping Z, Chen L, et al. von Hippel-Lindau disease-associated hemangioblastomas are derived from embryologic multipotent cells. *PLoS Med.* 2007 Feb;4(2):e60. DOI: 10.1371/journal.pmed.0040060.

526 Adult primary brain tumors: Image H. Minimal parenchyma in hemangioblastoma. This image is a derivative work, adapted from the following source, available under : Marvin 101. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

(c) ketabton.com: The Digital Library

527 **Adult primary brain tumors: Image I.** Field of vision in bitemporal hemianopia. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

527 **Adult primary brain tumors: Image J.** Prolactinoma. This image is a derivative work, adapted from the following source, available under : Wang CS, Yeh TC, Wu TC, et al. Pituitary macroadenoma co-existent with supraclinoid internal carotid artery cerebral aneurysm: a case report and review of the literature. *Cases J.* 2009;2:6459. DOI: 10.4076/1757-1626-2-6459.

527 **Adult primary brain tumors: Image K.** Schwannoma at cerebellopontine angle. Courtesy of MRT-Bild.

527 **Adult primary brain tumors: Image L.** Schwann cell origin of schwannoma. This image is a derivative work, adapted from the following source, available under : Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

528 **Childhood primary brain tumors: Image A.** MRI of pilocytic astrocytoma. This image is a derivative work, adapted from the following source, available under : Hafez RFA. Stereotaxic gamma knife surgery in treatment of critically located pilocytic astrocytoma: preliminary result. *World J Surg Oncol.* 2007;5:39. doi 10.1186/1477-7819-5-39.

528 **Childhood primary brain tumors: Image C.** CT of medulloblastoma. Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

528 **Childhood primary brain tumors: Image D.** Medulloblastoma histology. This image is a derivative work, adapted from the following source, available under : KGH. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

528 **Childhood primary brain tumors: Image E.** MRI of ependymoma. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

528 **Childhood primary brain tumors: Image F.** Ependymoma histology. This image is a derivative work, adapted from the following source, available under : Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

528 **Childhood primary brain tumors: Image G.** CT of craniopharyngioma. This image is a derivative work, adapted from the following source, available under : Garnet MR, Puget S, Grill J, et al. Craniopharyngioma. *Orphanet J Rare Dis.* 2007;2:18. DOI: 10.1186/1750-1172-2-18.

528 **Childhood primary brain tumors: Image H.** Craniopharyngioma histology. This image is a derivative work, adapted from the following source, available under : Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

531 **Friedreich ataxia: Image A.** Clinical kyphoscoliosis. This image is a derivative work, adapted from the following source, available under : Axelrod FB, Gold-von Simson. Hereditary sensory and autonomic neuropathies: types II, III, and IV. *Orphanet J Rare Dis.* 2007;2:39. DOI: 10.1186/1750-1172-2-39.

531 **Friedreich ataxia: Image B.** Radiograph showing kyphoscoliosis. This image is a derivative work, adapted from the following source, available under : Bounakis N, Karampalis C, Tsirikos AI. Surgical treatment of scoliosis in Rubinstein-Taybi syndrome type 2: a case report. *J Med Case Rep.* 2015; 9: 10. doi 10.1186/1752-1947-9-10.

532 **Facial nerve lesions.** Facial nerve palsy. This image is a derivative work, adapted from the following source, available under : Socolovsky M, Paez MD, Di Masi G, et al. Bell's palsy and partial hypoglossal to facial nerve transfer: Case presentation and literature review. *Surg Neurol Int.* 2012;3:46. DOI: 10.4103/2152-7806.95391.

533 **Cholesteatoma.** This image is a derivative work, adapted from the following source, available under : Welleschik. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

534 **Normal eye anatomy.** This image is a derivative work, adapted from the following source, available under : Jan Kaláb. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

534 **Conjunctivitis.** This image is a derivative work, adapted from the following source, available under : Baiyeroju A, Bowman R, Gilbert C, et al. Managing eye health in young children. *Community Eye Health.* 2010;23:4-11.

535 **Cataract.** Juvenile cataract. This image is a derivative work, adapted from the following source, available under : Roshan M, Vijaya PH, Lavanya GR, et al. A novel human CRYGD mutation in a juvenile autosomal dominant cataract. *Mol Vis.* 2010;16:887-896. PMCID PMC2875257.

536 **Glaucoma: Image C.** Closed/narrow angle glaucoma. This image is a derivative work, adapted from the following source, available under : Low S, Davidson AE, Holder GE, et al. Autosomal dominant Best disease with an unusual electrooculographic light rise and risk of angle-closure glaucoma: a clinical and molecular genetic study. *Mol Vis.* 2011;17:2272-2282. PMCID PMC3171497. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

536 **Glaucoma: Image D.** Acute angle closure glaucoma. This image is a derivative work, adapted from the following source, available under :. Courtesy of Dr. Jonathan Trobe.

536 **Uveitis: Image A.** This image is a derivative work, adapted from the following source, available under : Weber AC, Levison AL, Srivastava, et al. A case of *Listeria monocytogenes* endophthalmitis with recurrent inflammation and novel management. *J Ophthalmic Inflamm Infect.* 2015;5(1):28. DOI: 10.1186/s12348-015-0058-8.

536 **Age-related macular degeneration.** Courtesy of the Department of Health and Human Services.

537 **Diabetic retinopathy.** This image is a derivative work, adapted from the following source, available under : Sundling V, Gulbrandsen P, Straand J. Sensitivity and specificity of Norwegian optometrists' evaluation of diabetic retinopathy in single-field retinal images – a cross-sectional experimental study. *BMC Health Services Res.* 2013;13:17. DOI: 10.1186/1472-6963-13-17.

537 **Hypertensive retinopathy.** This image is a derivative work, adapted from the following source, available under : Diallo JW, Méda N, Tougouma SJB, et al. Intérêts de l'examen du fond d'œil en pratique de ville: bilan de 438 cas. *Pan Afr Med J.* 2015;20:363. DOI: 10.11604/pamj.2015.20.363.6629.

537 **Retinal vein occlusion.** This image is a derivative work, adapted from the following source, available under : Alasil T, Rauser ME. Intravitreal bevacizumab in the treatment of neovascular glaucoma secondary to central retinal vein occlusion: a case report. *Cases J.* 2009;2:176. DOI: 10.1186/1757-1626-2-176. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

537 **Retinal detachment.** Courtesy of EyeRounds.

538 **Retinitis pigmentosa.** Courtesy of EyeRounds.

538 **Leukocoria.** This image is a derivative work, adapted from the following source, available under : Aerts I, Lumbroso-Le Rouic L,

(c) ketabton.com: The Digital Library

Gauthier-Villars M, et al. Retinoblastoma. *Orphanet J Rare Dis.* 2006 Aug 25;1:31. DOI: 10.1186/1750-1172-1-31.

540 **Ocular motility.** Testing ocular muscles. This image is a derivative work, adapted from the following source, available under : Au.yousef. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

541 **CN III, IV, VI palsies: Image A.** Cranial nerve III damage. This image is a derivative work, adapted from the following source, available under : Hakim W, Sherman R, Rezk T, et al. An acute case of herpes zoster ophthalmicus with ophthalmoplegia. *Case Rep Ophthalmol Med.* 1012; 2012:953910. DOI: 10.1155/2012/953910.

541 **CN III, IV, VI palsies: Image B.** Cranial nerve IV damage. This image is a derivative work, adapted from the following source, available under : Mendez JA, Arias CR, Sanchez D, et al. Painful ophthalmoplegia of the left eye in a 19-year-old female, with an emphasis in Tolosa-Hunt syndrome: a case report. *Cases J.* 2009; 2: 8271. DOI: 10.4076/1757-1626-2-8271.

541 **CN III, IV, VI palsies: Image C.** Cranial nerve VI damage. This image is a derivative work, adapted from the following source, available under : Jordi March i Nogué. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

Psychiatry

563 **Trichotillomania.** Courtesy of Robodoc.

Renal

578 **Potter sequence (syndrome).** Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

580 **Kidney anatomy and glomerular structure.** This image is a derivative work, adapted from the following source, available under : Ramidi GA, Kurukumbi MK, Sealy PL. Collapsing glomerulopathy in sickle cell disease: a case report. *J Med Case Reports.* 2011; 5: 71. DOI: 10.1186/1752-1947-5-71.

581 **Course of ureters.** This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

581 **Glomerular filtration barrier.** This image is a derivative work, adapted from the following source, available under : Feng J, Wei H, Sun Y, et al. Regulation of podocalyxin expression in the kidney of streptozotocin-induced diabetic rats with Chinese herbs (Yishen capsule). *BMC Complement Altern Med.* 2013;13:76. DOI: 10.1186/1472-6882-13-76.

594 **Casts in urine: Image B.** WBC casts. This image is a derivative work, adapted from the following source, available under : Perazella MA. Diagnosing drug-induced AIN in the hospitalized patient: a challenge for the clinician. *Clin Nephrol.* 2014 Jun; 81(6): 381-8. DOI: 10.5414/CN108301.

594 **Casts in urine: Image D.** Fatty casts. This image is a derivative work, adapted from the following source, available under : Li S, Wang ZJ, Chang TT. Temperature oscillation modulated self-assembly of periodic concentric layered magnesium carbonate microparticles. *PLoS One.* 2014;9(2):e88648. DOI:10.1371/journal.pone.0088648

596 **Nephritic syndrome: Image A.** Histology of acute poststreptococcal glomerulonephritis. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

596 **Nephritic syndrome: Image B.** This image is a derivative work, adapted from the following source, available under : Immunofluorescence of acute poststreptococcal glomerulonephritis. Oda T, Yoshizawa N, Yamakami K, et al. The role of nephritis-associated plasmin receptor (naplr) in glomerulonephritis associated with streptococcal infection. *Biomed Biotechnol.* 2012;2012:417675. DOI 10.1155/2012/417675.

596 **Nephritic syndrome: Image C.** Histology of rapidly progressive glomerulonephritis. Courtesy of the Department of Health and Human Services and Uniformed Services University of the Health Sciences.

596 **Nephritic syndrome: Image D.** This image is a derivative work, adapted from the following source, available under : Kiremitci S, Ensari A. Classifying lupus nephritis: an ongoing story. *Scientific World Journal.* 2014; 2014: 580620. DOI: 10.1155/2014/580620.

597 **Nephrotic syndrome: Image A.** This image is a derivative work, adapted from the following source, available under : Teoh DCY, El-Modir A. Managing a locally advanced malignant thymoma complicated by nephrotic syndrome: a case report. *J Med Case Reports.* 2008; 2: 89. DOI: 10.1186/1752-1947-2-89.

597 **Nephrotic syndrome: Image B.** Histology of focal segmental glomerulosclerosis. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

597 **Nephrotic syndrome: Image D.** Diabetic glomerulosclerosis with Kimmelstiel-Wilson lesions. This image is a derivative work, adapted from the following source, available under : Doc Mari. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

598 **Kidney stones: Image A.** Nair S, George J, Kumar S, et al. Acute oxalate nephropathy following ingestion of *Averrhoa bilimbi* juice. *Case Rep Nephrol.* 2014; 2014: DOI: 10.1155/2014/240936.

598 **Kidney stones: Image B.** This image is a derivative work, adapted from the following source, available under : Joel Mills. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

599 **Hydronephrosis.** Ultrasound. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

600 **Pyelonephritis: Image A.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

600 **Pyelonephritis: Image B.** CT scan. Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

602 **Acute tubular necrosis: Image A.** Muddy brown casts. This image is a derivative work, adapted from the following source, available under : Dr. Serban Nicolescu.

602 **Renal papillary necrosis.** Courtesy of the Department of Health and Human Services and William D. Craig, Dr. Brent J. Wagner, and Mark D. Travis.

604 **Renal cyst disorders: Image C.** Ultrasound of simple cyst. This image is a derivative work, adapted from the following source, available under : Nevit Dilmen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

605 **Renal cell carcinoma: Image A.** Histoilogy. This image is a derivative work, adapted from the following source, available under :

(c) ketabton.com: The Digital Library

Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

605 **Renal cell carcinoma: Image C.** CT scan. This image is a derivative work, adapted from the following source, available under : Behnes CL, Schlegel C, Shoukier M, et al. Hereditary papillary renal cell carcinoma primarily diagnosed in a cervical lymph node: a case report of a 30-year-old woman with multiple metastases. *BMC Urol.* 2013;13:3. DOI: 10.1186/1471-2490-13-3.

605 **Renal cell carcinoma: Image B.** Gross specimen. Courtesy of Dr. Ed Uthman.

605 **Renal oncocytoma: Image A.** Gross specimen. This image is a derivative work, adapted from the following source, available under : Courtesy of M. Emmanuel.

605 **Renal oncocytoma: Image B.** Histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

606 **Nephroblastoma.** This image is a derivative work, adapted from the following source, available under : Refaie H, Sarhan M, Hafez A. Role of CT in assessment of unresectable Wilms tumor response after preoperative chemotherapy in pediatrics. *Sci World J.* 2008;8:661-669. doi 10.1100/tsw.2008.96.

606 **Urothelial carcinoma of the bladder: Image A.** This image is a derivative work, adapted from the following source, available under : Geavlete B, Stanescu F, Moldoveanu C, et al. NBI cystoscopy and bipolar electrosurgery in NMIBC management—an overview of daily practice. *J Med Life.* 2013;6:140-145. PMCID PMC3725437.

Reproductive

615 **Fetal alcohol syndrome.** This image is a derivative work, adapted from the following source, available under : Courtesy of Teresa Kellerman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

618 **Umbilical cord: Image A.** Cross-section of normal umbilical cord. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

618 **Meckel diverticulum: Image B.** This image is a derivative work, adapted from the following source, available under : Mathur P, Gupta R, Simlot A, et al. Congenital pouch colon with double Meckel's diverticulae. *J Neonatal Surg.* 2013 Oct-Dec; 2(4): 48.

623 **Uterine (Müllerian) duct anomalies: Images A-D.** This image is a derivative work, adapted from the following source, available under : Ahmadi F, Zafarani F, Haghighi H, et al. Application of 3D ultrasonography in detection of uterine abnormalities. *Int J Fertil Steril.* 2011; 4:144-147. PMCID PMC4023499.

626 **Female reproductive epithelial histology.** Transformation zone. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

628 **Seminiferous tubules.** This image is a derivative work, adapted from the following source, available under : Dr. Anlt Rao. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

641 **Pregnancy complications.** This image is a derivative work, adapted from the following source, available under : Li W, Wang G, Lin T, et al. Misdiagnosis of bilateral tubal pregnancy: a case report. *J Med Case Rep.* 2014;8:342. DOI: 10.1186/1752-1947-8-342.

642 **Hydatidiform mole: Image A.** Cluster of cluster of grapes appearance in complete hydatidiform mole. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman

642 **Choriocarcinoma: Image B.** "Cannonball" metastases. This image is a derivative work, adapted from the following source, available under : Lekanidi K, Vlachou PA, Morgan B, et al. Spontaneous regression of metastatic renal cell carcinoma: case report. *J Med Case Rep.* 2007;1:89. DOI: 10.1186/1752-1947-1-89.

644 **Vulvar pathology: Image A.** Bartholin cyst. Courtesy of the Department of Health and Human Services and Susan Lindsley.

644 **Vulvar pathology: Image B.** Lichen sclerosis. This image is a derivative work, adapted from the following source, available under : Lambert J. Pruritus in female patients. *Biomed Res Int.* 2014;2014:541867. DOI: 10.1155/2014/541867.

644 **Vulvar pathology: Image C.** Vulvar carcinoma. This image is a derivative work, adapted from the following source, available under : Ramli I, Hassam B. Carcinome épidermoïde vulvaire: pourquoi surveiller un lichen scléro-atrophique. *Pan Afr Med J.* 2015;21:48. DOI: 10.11604/pamj.2015.21.48.6018.

644 **Vulvar pathology: Image D.** Extramallary Paget disease. This image is a derivative work, adapted from the following source, available under : Wang X, Yang W, Yang J. Extramammary Paget's disease with the appearance of a nodule: a case report. *BMC Cancer.* 2010;10:405. DOI: 10.1186/1471-2407-10-405.

645 **Polycystic ovarian syndrome.** This image is a derivative work, adapted from the following source, available under : Kopera D, Wehr E, Obermayer-Pietsch B. Endocrinology of hirsutism. Int J Trichology. 2010;2(1):30–35. doi:10.4103/0974-7753.66910

647 **Dysgerminoma: Image B.** This image is a derivative work, adapted from the following source, available under : Montesinos L, Acien P, Martinez-Beltran M, et al. Ovarian dysgerminoma and synchronic contralateral tubal pregnancy followed by normal intra-uterine gestation: a case report. *J Med Rep.* 2012;6:399. DOI: 10.1186/1752-1947-6-399.

647 **Ovarian neoplasms: Image D.** Mature cystic teratoma. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

647 **Ovarian neoplasms: Image E.** Yolk sac tumor. This image is a derivative work, adapted from the following source, available under : Jensflorian. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

647 **Ovarian neoplasms: Image F.** Call-Exner bodies. This image is a derivative work, adapted from the following source, available under : Katoh T, Yasuda M, Hasegawa K, et al. Estrogen-producing endometrioid adenocarcinoma resembling sex cord-stromal tumor of the ovary: a review of four postmenopausal cases. *Diagn Pathol.* 2012;7:164. DOI: 10.1186/1746-1596-7-164.

648 **Uterine conditions: Image A.** Endometrial tissue found outside the uterus. This image is a derivative work, adapted from the following source, available under : Hastings JM, Fazleabas AT. A baboon model for endometriosis: implications for fertility. *Reprod Biol Endocrinol.* 2006;4(suppl 1):S7. DOI: 10.1186/1477-7827-4-S1-S7.

648 **Uterine conditions: Image B.** Endometritis with inflammation of the endometrium. This image is a derivative work, adapted from the following source, available under : Montesinos L, Acien P, Martinez-Beltran M, et al. Ovarian dysgerminoma and synchronic contralateral tubal pregnancy followed by normal intra-uterine

(c) ketabton.com: The Digital Library

gestation: a case report. *J Med Rep.* 2012;6:399. DOI: 10.1186/1752-1947-6-399.

648 Uterine conditions: Image C. Endometrial carcinoma. This image is a derivative work, adapted from the following source, available under : Izadi-Mood N, Yarmohammadi M, Ahmadi SA, et al. Reproducibility determination of WHO classification of endometrial hyperplasia/well differentiated adenocarcinoma and comparison with computerized morphometric data in curettage specimens in Iran. *Diagn Pathol.* 2009;4:10. DOI:10.1186/1746-1596-4-10.

648 Uterine conditions: Image D. Leiomyoma (fibroid), gross specimen. This image is a derivative work, adapted from the following source, available under : Courtesy of Hic et nunc.

648 Uterine conditions: Image E. Leiomyoma (fibroid) histology. This image is a derivative work, adapted from the following source, available under : Londero AP, Perego P, Mangioni C, et al. Locally relapsed and metastatic uterine leiomyoma: a case report. *J Med Case Rep.* 2008;2:308. DOI: 10.1186/1752-1947-2-308. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

649 Benign breast disease: Image A. Fibroadenomas. This image is a derivative work, adapted from the following source, available under : Gokhale S. Ultrasound characterization of breast masses. *Indian J Radiol Imaging.* 2009 Aug;19(3):242-7. DOI: 10.4103/0971-3026.54878.

649 Benign breast disease: Images B (phyllodes tumor on ultrasound) and C (phyllodes cyst). This image is a derivative work, adapted from the following source, available under : Muttarak MD, Lerttumnongtum P, Somwangjaroen A, et al. Phyllodes tumour of the breast. *Biomed Imaging Interv J.* 2006 Apr-Jun;2(2):e33. DOI: 10.2349/biij.2.2.e33.

650 Breast cancer: Image A. Mammography of breast cancer. This image is a derivative work, adapted from the following source, available under : Molino C, Mocerino C, Braucci A, et al. Pancreatic solitary and synchronous metastasis from breast cancer: a case report and systematic review of controversies in diagnosis and treatment. World J Surg Oncol. 2014;12:2. DOI:10.1186/1477-7819-12-2

650 Breast cancer: Image C. Comedocarcinoma. This image is a derivative work, adapted from the following source, available under : Costarelli L, Campagna D, Mauri M, et al. Intraductal proliferative lesions of the breast—terminology and biology matter: premalignant lesions or preinvasive cancer? *Int J Surg Oncol.* 2012;501904. DOI: 10.1155/2012/501904. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

650 Breast cancer: Image D. Paget disease of breast. This image is a derivative work, adapted from the following source, available under : Muttarak M, Siriya B, Kongmebhol P, et al. Paget's disease of the breast: clinical, imaging and pathologic findings: a review of 16 patients. *Biomed Imaging Interv J.* 2011;7:e16. DOI: 10.2349/biij.7.2.e16.

650 Breast cancer: Image E. Invasive lobular carcinoma. This image is a derivative work, adapted from the following source, available under : Franceschini G, Manno A, Mule A, et al. Gastro-intestinal symptoms as clinical manifestation of peritoneal and retroperitoneal spread of an invasive lobular breast cancer: report of a case and review of the literature. *BMC Cancer.* 2006;6:193. DOI: 10.1186/1471-2407-6-193.

650 Breast cancer: Image E. Peau d'orange of inflammatory breast cancer. This image is a derivative work, adapted from the following source, available under : Levine PH, Zolfaghari L, Young H, et al. What Is inflammatory breast cancer? Revisiting the case definition. *Cancers (Basel).* 2010 Mar;2(1):143–152. DOI: 10.3390/cancers2010143.

651 Penile pathology: Image A. Peyronie disease. This image is a derivative work, adapted from the following source, available under : Tran VQ, Kim DH, Lesser TF, et al. Review of the surgical approaches for Peyronie's disease: corporeal plication and plaque incision with grafting. *Adv Urol.* 2008; 2008: 263450. DOI: 10.1155/2008/263450.

651 Penile pathology: Image B. Squamous cell carcinoma. This image is a derivative work, adapted from the following source, available under : Antônio JR, Antônio CR, Trídico LA. Erythroplasia of queyrat treated with topical 5-fluorouracil. *An Bras Dermatol.* 2016 Sep-Oct; 91(5 Suppl 1): 42–44. DOI: 10.1590/abd1806-4841.20164595.

651 Cryptorchidism. This image is a derivative work, adapted from the following source, available under : Pandey A, Gangopadhyay AN, Kumar V. High anorectal malformation in a five-month-old boy: a case report. *J Med Case Reports.* 2010; 4: 296. DOI: 10.1186/1752-1947-4-296.

651 Varicocele. This image is a derivative work, adapted from the following source, available under : Mak CW, Tzeng WS. Sonography of the scrotum. DOI: 10.5772/27586.

652 Scrotal masses: Image A. Congenital hydrocele. This image is a derivative work, adapted from the following source, available under : Leonardi S, Barone P, Gravina G, et al. Severe Kawasaki disease in a 3-month-old patient: a case report. *BMC Res Notes.* 2013;6:500. DOI: 10.1186/1756-0500-6-500.

Respiratory

661 Alveolar cell types: Image A. Electron micrograph of type II pneumocyte. This image is a derivative work, adapted from the following source, available under : Fehrenbach H, Tews S, Fehrenbach A, et al. Improved lung preservation relates to an increase in tubular myelin-associated surfactant protein A. *Respir Res.* 2005 Jun 21;6:60. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

661 Alveolar cell types: Image B. Micrograph of type II pneumocyte. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Thomas Caceci.

661 Neonatal respiratory distress syndrome: Image A. This image is a derivative work, adapted from the following source, available under : Alorainy IA, Balas NB, Al-Boukai AA. Pictorial essay: infants of diabetic mothers. *Indian J Radiol Imaging.* 2010;20:174-181. DOI: 10.4103/0971-3026.69349.

663 Lung anatomy: Image A. X-ray of normal lung. This image is a derivative work, adapted from the following source, available under : Namkoong H, Fujiwara H, Ishii M, et al. Immune reconstitution inflammatory syndrome due to *Mycobacterium avium* complex successfully followed up using 18 F-fluorodeoxyglucose positron emission tomography-computed tomography in a patient with human immunodeficiency virus infection: A case report. *BMC Med Imaging.* 2015;15:24. DOI 10.1186/s12880-015-0063-2.

663 Lung anatomy: Image B. CT scan of the chest. This image is a derivative work, adapted from the following source, available under : Wang JF, Wang B, Jansen JA, et al. Primary squamous cell carcinoma of lung in a 13-year-old boy: a case report. *Cases J.* 2008 Aug 22;1(1):123. DOI: 10.1186/1757-1626-1-123. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

667 Cyanide vs carbon monoxide poisoning. This image is a derivative work, adapted from the following source, available under : Subhaschandra S, Jatishwor W, Suraj Th. Isolated symmetrical bilateral basal ganglia T2 hyperintensity in carbon monoxide poisoning. *Ann Indian Acad Neurol.* 2008 Oct-Dec; 11(4): 251–253. DOI: 10.4103/0972-2327.44563.

671 Rhinosinusitis. This image is a derivative work, adapted from the following source, available under : Strek P, Zagolski O, Sktadzien

(c) ketabton.com: The Digital Library

J. Fatty tissue within the maxillary sinus: a rare finding. *Head Face Med.* 2006;2:28. DOI: 10.1186/1746-160X-2-28.

671 **Deep venous thrombosis.** This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

672 **Pulmonary emboli: Image B.** CT scan. This image is a derivative work, adapted from the following source, available under : Dr. Carl Chartrand-Lefebvre. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

672 **Mediastinal pathology.** This image is a derivative work, adapted from the following source, available under : Aga Z, Avelino J, Darling GE. An unusual case of spontaneous esophageal rupture after swallowing a boneless chicken nugget. *Case Rep Emerg Med.* 2016; 2016: 5971656. DOI: 10.1155/2016/5971656.

675 **Obstructive lung diseases: Image A.** Lung tissue with enlarged alveoli in emphysema. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonnert.

675 **Obstructive lung diseases: Image B.** CT of centriacinar emphysema. Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr.

675 **Obstructive lung diseases: Image C.** Emphysema histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

675 **Obstructive lung diseases: Image D.** Barrel-shaped chest in emphysema. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

675 **Obstructive lung diseases: Image E.** Mucus plugs in asthma. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

675 **Obstructive lung disease: Image F.** Curschmann spirals. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under . Dr. James Heilman.

675 **Obstructive lung diseases: Image G.** Charcot-Leyden crystals on bronchalverolar lavage. This image is a derivative work, adapted from the following source, available under : Gholamnejad M, Rezaie N. Unusual presentation of chronic eosinophilic pneumonia with "reversed halo sign": a case report. *Iran J Radiol.* 2014 May;11(2):e7891. DOI: 10.5812/iranjradiol.7891.

675 **Obstructive lung disease: Image H.** Bronchiectasis in cystic fibrosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

675 **Restrictive lung diseases: Image A.** Pulmonary fibrosis. This image is a derivative work, adapted from the following source, available under : Walsh SLF, Wells AU, Sverzellati N, et al. Relationship between fibroblastic foci profusion and high resolution CT morphology in fibrotic lung disease. *BMC Med.* 2015;13:241. DOI: 10.1186/s12916-015-0479-0.

676 **Sarcoidosis: Images A.** Kajal B, Harvey J, Alowami S. Melkerrson-Rosenthal Syndrome, a rare case report of chronic eyelid swelling. *Diagn Pathol.* 2013; 8: 188. DOI: 10.1186/1746-1596-8-188.

676 **Sarcoidosis: Images B (X-ray of the chest) and C (CT of the chest).** X-ray of the chest) and C CT of the chest. This image is a derivative work, adapted from the following source, available under : Lønborg J, Ward M, Gill A, et al. Utility of cardiac magnetic resonance in assessing right-sided heart failure in sarcoidosis. *BMC Med Imaging.* 2013;13:2. DOI: 10.1186/1471-2342-13-2.

676 **Inhalational injury and sequelae: Images A (18 hours after inhalation injury) and B (11 days after injury).** This image is a derivative work, adapted from the following source, available under : Bai C, Huang H, Yao X, et al. Application of flexible bronchoscopy in inhalation lung injury. *Diagn Pathol.* 2013;8:174. DOI: 10.1186/1746-1596-8-174.

677 **Pneumoconioses: Image A.** Pleural plaques in asbestosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

677 **Pneumoconioses: Image B.** CT scan of asbestosis. This image is a derivative work, adapted from the following source, available under : Miles SE, Sandrini A, Johnson AR, et al. Clinical consequences of asbestos-related diffuse pleural thickening: a review. *J Occup Med Toxicol.* 2008;3:20. DOI: 10.1186/1745-6673-3-20.

677 **Pneumoconioses: Image C.** Ferruginous bodies in asbestosis. This image is a derivative work, adapted from the following source, available under : Dr, Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

677 **Pneumoconioses: Image D.** Berylliosis: non-caseous granuloma. *Ann Saudi Med.*2009 Nov-Dec; 29(6): 485–486. DOI: 10.4103/0256-4947.57175.

678 **Mesothelioma.** This image is a derivative work, adapted from the following source, available under : Weiner SJ, Neragi-Miandoab S. Pathogenesis of malignant pleural mesothelioma and the role of environmental and genetic factors. *J Carcinog.* 2008;7:3. DOI: 10.1186/1477-3163-7-3.

678 **Acute respiratory distress syndrome: Image A.** This image is a derivative work, adapted from the following source, available under : Pires-Neto RC, Del Carlo Bernardi F, de Araujo PA. The expression of water and ion channels in diffuse alveolar damage is not dependent on DAD etiology. *PLoS One.* 2016; 11(11): e0166184. DOI: 10.1371/journal.pone.0166184.

678 **Acute respiratory distress syndrome: Image B.** Bilateral lung opacities. This image is a derivative work, adapted from the following source, available under : Imanaka H, Takahara B, Yamaguchi H, et al. Chest computed tomography of a patient revealing severe hypoxia due to amniotic fluid embolism: a case report. *J Med Case Reports.* 2010;4:55. DOI: 10.1186/1752-1947-4-55.

680 **Atelectasis.** This image is a derivative work, adapted from the following source, available under : Khan AN, Al-Jahdali H, Al-Ghanem S, et al. Reading chest radiographs in the critically ill (Part II): Radiography of lung pathologies common in the ICU patient. Ann Thorac Med. 2009;4(3):149–157. DOI:10.4103/1817-1737.53349

681 **Pleural effusions: Images A and B.** This image is a derivative work, adapted from the following source, available under : Toshikazu A, Takeoka H, Nishioka K, et al. Successful management of refractory pleural effusion due to systemic immunoglobulin light chain amyloidosis by vincristine adriamycin dexamethasone chemotherapy: a case report. *Med Case Rep.* 2010;4:322. DOI: 10.1186/1752-1947-4-322.

682 **Pneumothorax: Image A.** This image is a derivative work, adapted from the following source, available under : Miura K, Kondo R, Kurai M, et al. Birt-Hogg-Dubé syndrome detected incidentally by asymptomatic bilateral pneumothorax in health screening: a

(c) ketabton.com: The Digital Library

case of a young Japanese woman. *Surg Case Rep.* 2015 Dec; 1: 17. DOI: 10.1186/s40792-015-0014-8.

682 **Pneumothorax: Image B.** This image is a derivative work, adapted from the following source, available under : Rosat A, Díaz C. Reexpansion pulmonary edema after drainage of tension pneumothorax. *Pan Afr Med J.* 2015; 22: 143.DOI: 10.11604/pamj.2015.22.143.8097.

683 **Pneumonia: Image A.** This image is a derivative work, adapted from the following source, available under : Yoon BW, Song YG, Lee SH. Severe community-acquired adenovirus pneumonia treated with oral ribavirin: a case report. *BMC Res Notes.* 2017; 10: 47. DOI: 10.1186/s13104-016-2370-2.

683 **Pneumonia: Image B.** Lobar pneumonia, gross specimen. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

683 **Pneumonia: Image C.** Acute inflammatory infiltrates in bronchopneumonia. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

683 **Pneumonia: Image D.** Bronchopneumonia, gross specimen. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

683 **Pneumonia: Image E.** This image is a derivative work, adapted from the following source, available under : Allen CM, AL-Jahdali HH, Irion KL, et al. Imaging lung manifestations of HIV/AIDS. *Ann Thorac Med.* 2010 Oct-Dec; 5(4): 201–216. DOI: 10.4103/1817-1737.69106.

684 **Lung cancer: Image B.** Adenocarcinoma histology. Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

684 **Lung cancer: Image C.** Squamous cell carcinoma. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

684 **Lung cancer: Image E.** Large cell lung cancer. This image is a derivative work, adapted from the following source, available under : Jala VR, Radde BN, Haribabu B, et al. Enhanced expression of G-protein coupled estrogen receptor (GPER/GPR30) in lung cancer. *BMC Cancer.* 2012;12:624. doi 10.1186/1471-2407-12-624.

685 **Lung abscess: Image A.** Gross specimen. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .

685 **Lung abscess: Image B.** X-ray. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Yale Rosen.

685 **Pancoast tumor.** This image is a derivative work, adapted from the following source, available under : Manenti G, Raguso M, D'Onofrio S, et al. Pancoast tumor: the role of magnetic resonance imaging. *Case Rep Radiol.* 2013; 2013:479120. DOI: 10.1155/2013/479120.

685 **Superior vena cava syndrome: Images A (blanching of skin with pressure) and B (CT of chest).** This image is a derivative work, adapted from the following source, available under : Shaikh I, Berg K, Kman N. Thrombogenic catheter-associated superior vena cava syndrome. *Case Rep Emerg Med.* 2013; 2013: 793054. DOI 10.1155/2013/793054.

(c) ketabton.com: The Digital Library

Index

A

A-a gradient
  by age, 668
  with oxygen deprivation, 669
  restrictive lung disease, 675
Abacavir, 203
Abciximab
  Glycoprotein IIb/IIIa inhibitors, 438
  therapeutic antibodies, 122
  thrombogenesis and, 411
Abdominal aorta
  atherosclerosis in, 302
  bifurcation of, 663
  branches, 363
Abdominal aortic aneurysm, 302
Abdominal pain
  bacterial peritonitis, 390
  Budd-Chiari syndrome, 392
  diabetic ketoacidosis and, 347
  ectopic pregnancy, 641
  Henoch-Schönlein purpura, 315
  hypercalcemia, 591
  hyperparathyroidism, 345
  intussusception, 385
  irritable bowel syndrome, 383
  Meckel diverticulum, 618
  pancreas divisum, 360
  pancreatic cancer, 398
  polyarteritis nodosa, 314
  porphyria, 425
  postprandial, 363
  RLQ pain, 384
  RUQ pain, 396
Abdominal wall
  anatomy, 369
  inguinal hernias, **369**
  ventral defects, 358
Abducens nerve (CN VI), 506
  ocular motility, 540
  palsy, 541
Abduction
  arm, 446
  hand, 450
  hip, 451, 453
  passive, abnormal, 454
Abductor digiti minimi muscle, 450
Abductor pollicis brevis muscle, 450
Abetalipoproteinemia, **94**, 414
Abnormal uterine bleeding (AUB)
  non-structural causes (COEIN), **633**
  structural causes (PALM), 633
ABO blood classification, 405
  hemolytic disease of the newborn, 405
Abortion
  ethical situations, 268
  methotrexate for, 440
Abruptio placentae, 640
  cocaine use, 614
  preeclampsia, 643
Abscesses, 479
  acute inflammation and, 214
  brain, 156, 177, **180**
  calcification with, 211
  cold staphylococcal, 116
  frontal lobe, 153
  Klebsiella spp, 145
  liver, 155, 179
  lung, 685
  necrosis with, 209
  Staphylococcus aureus, 135
  Toxoplasma gondii, 177
  treatment of lung, 192
  in unvaccinated children, 186
Absence seizures
  characteristics of, 517
  drug therapy for, 544
Absolute risk reduction (ARR), 258
AB toxin, 132
Abuse
  child, 269, **556**
  intimate partner violence, 269
Acalculia, 511
Acalculous cholecystitis, 396
Acanthocytes, 414
Acantholysis, 475, 480
Acanthosis, 475
  psoriasis, 477
Acanthosis nigricans, 228, 482
  stomach cancer, 379
Acarbose, 353
Accessory nerve (CN XI), 506
  arm abduction, 446
  lesion of, 532
Accessory pancreatic duct, 360, 368
Accommodation, eye, 506, 535
Accuracy (validity), 259
Acebutolol, 245, 319
ACE inhibitors, **610**
  acute coronary syndromes, 307
  C1 esterase inhibitor deficiency, 107
  dilated cardiomyopathy, 308
  dry cough, 251
  heart failure, 309
  hypertension, 316
  naming convention for, 253
  preload/afterload effects, 284
  teratogenicity, 614
Acetaldehyde, 72
Acetaldehyde dehydrogenase, 72
Acetaminophen, **485**
  vs aspirin for pediatric patients, 485
  free radical injury and, 210
  hepatic necrosis from, 249
  for osteoarthritis, 466
  toxicity effects, 485
  toxicity treatment for, 248
Acetazolamide, 252, 552, **608**
  pseudotumor cerebri, 521
Acetoacetate metabolism, 90
Acetone breath, 347
Acetylation
  chromatin, 34
  drug metabolism, 232
  histones, 34
  posttranslation, 45
Acetylcholine (ACh)
  anticholinesterase effect on, 240
  change with disease, 495
  Clostridium botulinum inhibition of release, 138
  opioid analgesics, 551
  pacemaker action potential and, 292
Acetylcholine (ACh) receptor agonists, 551
Acetylcholine receptors, 228, **236**
Acetylcholinesterase (AChE)
  in amniotic fluid, 491
  malathion and, 200
  neural tube defects and, 491
Acetylcholinesterase (AChE) inhibitors
  naming convention for, 253
  for neuromuscular junction disease, 472
  toxicity treatment for, 248
Acetyl-CoA carboxylase
  fatty acid synthesis, 73
  vitamin B_7 and, 68
Achalasia, **376**
  esophageal cancer and, 378
  LES tone in, 371
Achilles reflex, 510
Achilles tendon xanthomas, 301
Achlorhydria
  stomach cancer, 379
  VIPomas, 371
Achondroplasia, **462**
  chromosome disorder, 64
  endochondral ossification in, 458
  inheritance, 60
Acid-base physiology, **592**
Acid-fast oocysts, 177
Acid-fast organisms, 125, 155
Acidic amino acids, 81
Acid maltase, 86
Acidosis, **592**
  acidemia diuretic effect on, 609
  cardiac contractility in, 284
  hyperkalemia with, 590
  metabolic, 85, 349
  renal tubular, 592
Acid phosphatase in neutrophils, 406
Acid reflux
  H_2 blockers for, 399
  proton pump inhibitors for, 399
Acid suppression therapy, **398**
Acinetobacter baumannii
  highly resistant bacteria, 198
  nosocomial infections, **142**
Acinetobacter spp
  nosocomial infections, 185
Acne, 475, 477
  danazol, 658
  tetracyclines for, 192
Acquired hydrocele (scrotal), 652
Acrodermatitis enteropathica, 71
Acromegaly, **339**
  carpal tunnel syndrome, 459
  GH, 329
  octreotide for, 400
  somatostatin analogs for, 328
Actin
  cytoskeleton, 48
  muscular dystrophies, 61
Acting out, 554
Actinic keratosis, 482
  squamous cell carcinoma, 484
Actinomyces israelii
  culture requirements of, 127
  oral infections, 186
  pigment production, 128
Actinomyces spp
  effects and treatment of, **139**
  penicillin G/V for, 187
Action/willpower stage, substance addiction, 568
Activated carrier molecules and form, **75**
Active errors, 274
Active immunity, 110
Acute chest syndrome, 422
Acute cholestatic hepatitis
  drug reactions and, 249
  macrolides, 193
Acute coronary syndrome
  ADP receptor inhibitors for, 437
  heparin for, 436
  nitrates for, 318
  treatments for, **307**
Acute cystitis, 594, **600**
Acute disseminated (postinfectious) encephalomyelitis, 524
Acute dystonia, 569
  treatment of, 241
Acute gastritis, 379
Acute hemolytic transfusion reactions, 114
Acute hemorrhagic cystitis, 164
Acute inflammation, **214**

(c) ketabton.com: The Digital Library

Acute inflammatory demyelinating polyradiculopathy, **524**
Acute intermittent porphyria, 425
Acute interstitial nephritis, **601**
Acute kidney injury, **601**
Acute laryngotracheobronchitis, **170**
Acute lymphoblastic leukemia (ALL), 432
  methotrexate for, 440
  oncogenes and, 224
Acute mesenteric ischemia, 386
Acute myelogenous leukemia (AML), 432
  chromosomal translocations, 434
  cytarabine for, 440
  myelodysplastic syndromes, 432
Acute pancreatitis, **397**
  hyperparathyroidism, 345
  necrosis and, 209
Acute pericarditis, **313**
Acute-phase proteins, 108, 213
Acute-phase reactants, **213**
  IL-6, 108
Acute poststreptococcal glomerulonephritis, 596
Acute promyelocytic leukemia
  vitamin A for, 66
Acute pulmonary edema
  opioid analgesics, 551
Acute pyelonephritis, 600
  WBC casts in, 594
Acute respiratory distress syndrome (ARDS), **678**
  eclampsia and, 643
Acute stress disorder, 564
Acute transplant rejection, 119
Acute tubular necrosis, **602**
Acyclovir, **201**
Adalimumab, 122, 487
  for Crohn disease, 382
Adaptive immunity, 99
Addiction, stages of change in overcoming, 568
Addison disease, 349
  HLA subtype association, 100
Additive effect
  of drugs, 235
Adduction
  arm (rotator cuff), 446
  hand, 450
  hip, 451
  passive, abnormal, 454
  thigh, 452
Adductor brevis, 452
Adductor longus, 451, 452
Adductor magnus, 452
Adenine
  Shiga/Shiga-like toxins and, 132
Adenocarcinomas
  carcinogens causing, 225
  esophagus, 378
  gastric, 216, 226
  lung, 224, 684
  nomenclature for, 220
  nonbacterial thrombotic endocarditis and, 228
  pancreas, 226, 368, 398
  paraneoplastic syndromes, 228
  pectinate line, 366
  prostatic, 654
  stomach, 379
Adenohypophysis, 327
  embryologic derivatives, 613
  hypothalamus and, 498
Adenomas
  bone, 464
  colorectal, 389
  nomenclature for, 220
  salivary gland, 376
  thyroid, 342
Adenomatous colonic polyps, 387
Adenomyosis (endometrial), 648
  uterine bleeding from, 633
Adenopathy
  Kawasaki disease, 314
Adenosine
  as antiarrhythmic drug, 324
  blood flow regulation, 297
  pacemaker action potential and, 292
Adenosine deaminase deficiency, 37, 117
Adenosine triphosphate (ATP)
  electron transport chain, 78
  production of, 74, 78
  in TCA cycle, 77
  in urea cycle, 82
Adenosine triphosphate (ATP) synthase inhibitors, 78
Adenoviruses
  characteristics of, 164
  conjunctivitis, 534
  pneumonia, 683
Adherens junctions, 474
Adhesions, 386
Adipose lipolysis, 320
Adipose tissue
  estrogen production, 630
  in starvation, 91
Adjustment disorder, 564
Adnexal (ovarian) torsion, **625**
Adoption studies, 256
ADPKD (Autosomal dominant polycystic kidney disease)
  saccular aneurysms and, 516
ADP receptor inhibitors, **437**
ADP ribosyltransferases, 132
Adrenal adenomas
  Cushing syndrome, 348
  hyperaldosteronism, 349
Adrenal carcinomas
  Li-Fraumeni syndrome, 224
  P-glycoprotein in, 227
Adrenal cortex, **327**
  progesterone production, 630
Adrenal hemorrhage
  Waterhouse-Friderichsen syndrome, 349
Adrenal hyperplasia
  Cushing syndrome, 348
  hyperaldosteronism and, 349
Adrenal insufficiency
  adrenoleukodystrophy, 47
  anovulation with, 645
  fludrocortisone for, 354
  mechanism and types of, **349**
  vitamin B_5 deficiency, 67
Adrenal medulla, **327**
  neuroblastomas of, 350
  pheochromocytomas in, 350
Adrenal steroids, **335**
Adrenal zona fasciculata, 336
Adrenocortical atrophy
  Addison disease, 349
  exogenous corticosteroids, 348
Adrenocortical insufficiency
  drug reaction and, 249
Adrenocorticotropic hormone (ACTH)
  adrenal cortex regulation of, 327
  in Cushing syndrome, 228, 348
  secretion of, 327
  signaling pathways of, 337
Adrenoleukodystrophy, 47, 524
Adults
  common causes of death, 272
  primary brain tumors, **526**
Adult T-cell leukemia, 226
Adult T-cell lymphoma, 430
Advance directives, **266**
Aedes mosquitoes
  yellow fever transmission, 168
Aerobes, **126**
Aerobic metabolism
  ATP production, 74
  fed state, 91
  vitamin B_1 (thiamine), 66
Aerobic organisms
  culture requirements, 126
Afferent arteriole, 580
  ANP/BNP effect on, 588
  constriction of, 583
Afferent nerves, 296
Aflatoxins, 153
  as carcinogen, 225
AFP, 117, 653
African sleeping sickness, 156
Afterload
  cardiac output, 284
  hydralazine, 318
  in shock, 310
Agammaglobulinemia
  chromosome affected, 64
Agars (bacterial culture), 126
Agenesis, 613
  Müllerian, 622
  uterovaginal, 639
Age-related amyloidosis, 212
Age-related macular degeneration (ARMD), **536**
Aging changes, 270
Agnosia, 511
Agonists
  indirect cholinomimetic, 234
  indirect general, 242
  indirect sympathomimetics, 242
  partial, 234
  potency and efficacy, 234
Agoraphobia, 563
Agranulocytosis, 573
  drug reaction and, 250
  sulfa drug allergies, 252
  thionamides, 354
Agraphia, 511
AIDS (acquired immunodeficiency syndrome)
  bacillary angiomatosis, 478
  brain abscess, 180
  Candida albicans, 153
  cryptococcal meningitis, 199
  Cryptosporidium, 155
  Cytomegalovirus (CMV), 165
  human herpesvirus 8, 165
  marijuana for, 571
  mycobacteria, 140
  Pneumocystis jirovecii, 154
  primary central nervous system lymphoma, 430
  retinitis, 165
  retroviruses, 167
  time course (untreated), 176
Air emboli, 672
Airways (conducting zone), 662
Akathisia, 499, 519
  antipsychotic drugs and, 573
Akinesia, 520
ALA dehydratase, 419, 425
Alanine
  ammonia transport, 82
  gluconeogenesis in starvation, 91
  pyruvate dehydrogenase complex deficiency, 77
Alanine aminotransferase (ALT), 77
  hepatitis viruses, 172
  in liver damage, 390
  toxic shock syndrome, 135
Alar plate, 490
Albendazole
  cestodes, 160
Albinism, 476
  locus heterogeneity, 57
  ocular, 61
Albumin, 213
  calcium and, 333
  as liver marker, 390
Albuminocytologic dissociation (CSF), 524
Albuterol, 242
  asthma, 687
Alcohol dehydrogenase, 72
Alcohol exposure
  in utero, 300
Alcoholic cirrhosis, **391**
  cholelithiasis and, 396
Alcoholic hepatitis, 391
Alcoholic liver disease, **391**
Alcoholism, 145
  anemia, 420
  cataracts and, 535
  cirrhosis and, 389
  common organisms affecting, 179
  esophageal cancer, 378
  ethanol metabolism and, 72
  folate deficiency, 420
  gastritis in, 379
  hepatitis, 367
  hypertension and, 300
  ketone bodies in, 90
  Korsakoff syndrome, 558
  liver serum markers in, 390
  magnesium levels in, 332
  Mallory-Weiss syndrome in, 377
  osteonecrosis in, 463
  pancreatitis, 249, 397
  porphyria, 425
  sideroblastic anemia, 419
  vitamin B_1 deficiency, 66
  vitamin B_9 deficiency, 68
Alcohol-related disorders
  readmissions with, 272
Alcohol use
  essential tremor, 519
  gout and, 467
  head and neck cancer, 671
  intoxication and withdrawal, 570
  sleep, 497
  teratogenic effects, 614
Alcohol use disorder, **571**
Alcohol withdrawal, 570
  drug therapy, 546, 572
  hallucinations in, 559, 570
Aldesleukin, 121
Aldolase B, 80
Aldose reductase, 81
Aldosterone, 588, 590
  adrenal cortex secretion of, 327
  SIADH, 338
  signaling pathways for, 337
Aldosterone antagonists, 316
Aldosterone resistance, 593
Aldosterone synthase, 335

(c) ketabton.com: The Digital Library

Alemtuzumab, 122
Alendronate, 486
Alexia, 515
Alirocumab, 320
Aliskiren, **610**
Alkaline phosphatase (ALP), 390, 463
  bone disorder lab values, 464
  hyperparathyroidism and, 345
  Paget disease of bone, 463
  in thyroid storm, 342
  as tumor marker, 226
Alkalosis, **592**
  contraction, 60
  diuretic effects, 609
  hypokalemia with, 590
  metabolic, 349
Alkaptonuria, **84**
ALK gene, 224
  lung cancer, 684
Alkylating agents, **441**
  as carcinogens, 225
  targets of, 438
  teratogenicity of, 614
Allantois, 618
Allelic heterogeneity, 57
Allergic bronchopulmonary aspergillosis (ABPA), 153
Allergic contact dermatitis, 477
Allergic reactions, 112
  blood transfusion, 114
All-*trans* retinoic acid
  for promyelocytic leukemia, 66
Allopurinol
  for gout, 487
  kidney stones, 598
  Lesch-Nyhan syndrome, 37
  rash with, 250
Alopecia
  doxorubicin, 439
  etoposide/teniposide, 442
  minoxidil for, 658
  syphilis, 147
  tinea capitis, 152
  vitamin A toxicity, 66
  vitamin B_5 deficiency, 67
  vitamin B_7 deficiency, 68
α-1,4-glucosidase
  glycogen metabolism, 86
α_1-antagonists
  BPH treatment, 654
α_1-antitrypsin, 52
α_1-antitrypsin deficiency, 51, 392
  emphysema, 674
α_1-blockers
  tamsulosin, example of, 237
α_1 selective blockers, 244
α_2-agonists, 243
α_2-antagonists, 576
α_2 selective blockers, 244
α-adrenergic agonists, 686
α-agonists
  glaucoma treatment, 552
  muscle spasm treatment, 551
α-amanitin
  RNA polymerase inhibition, 42
α-amylase, 373
α-antagonists
  for pheochromocytomas, 350
α-blockers, 244
  Beers criteria, 247
  for cocaine overdose, 571
  nonselective, 244
α cells, 328
  glucagonomas in, 351
  glucagon production by, 333
α-dystroglycan
  muscular dystrophy, 61
α-fetoprotein
  as tumor marker, 226
  in hepatocellular carcinoma, 392
  neural tube defects, 491
α-galactosidase A
  Fabry disease, 88
α-glucosidase inhibitors, 353
α-hemolytic cocci
  viridans group streptococci, 136
α-hemolytic bacteria
  Streptococcus pneumoniae, 136
α-intercalated cells, 593
α-ketoglutarate
  hyperammonemia and, 82
α-ketoglutarate dehydrogenase
  metabolic pathways, 74
  TCA cycle, 77
  vitamin B_1 and, 66
α-methyldopa, 243
  anemia and, 423
α-oxidation, 47
Alpha rhythm (EEG), 497
α-synuclein, 520
α-thalassemia, 418
Alpha toxin, 133
α-toxin
  Clostridium botulinum, 138
α (type I) error, 263
Alport syndrome, 596
  cataracts and, 535
  collagen deficiency in, 50
Alprazolam, 546
Alteplase (tPA), 413, 437
Alternative hypothesis, 262
Alternative splicing, 43
Altitude sickness, 670
Altruism, 555
Aluminum hydroxide, 399
Alveolar cell types, **661**
Alveolar dead space, 664
Alveolar gas equation, **668**
Alveolar macrophages, 661, 662
Alveolar PO_2, 668
Alveolar sacs, 662
Alveolar stage (development), 660
Alveolar ventilation, 664
Alveoli, 660
  pneumocytes, 661
Alzheimer disease, 520
  amyloidosis in, 212
  drug therapy for, 240, 549
  neurotransmitters for, 495
Amanita phalloides
  necrosis caused by, 42, 249
  RNA polymerase inhibition, 42
Amantadine, 548
Ambiguous genitalia
  46,XY DSD, 639
  ovotesticular disorder of sex development, 638
  placental aromatase deficiency, 639
Amebiasis
  Entamoeba histolytica amebiasis, 155
Amenorrhea
  antiandrogens, 658
  cystic fibrosis, 60
  ectopic pregnancy and, 641
  functional hypothalamic, 645
  menopause diagnosis, 636
  Müllerian agenesis, 622
  pituitary prolactinomas, 328
Amides (local anesthetics), 550
Amikacin, 191
Amiloride, 609
  for diabetes insipidus, 338
Amines
  MAO inhibitors, 575
Amine whiff test, 148
Amino acids
  blood-brain barrier and, 496
  branched, 84
  classification of, **81**
  codons for, 37
  derivatives of, **83**
  genetic code for, 37
  in histones, 34
  metabolism, 90
  purine synthesis, 35
  tRNA, 44
  urea cycle, 82
Aminoacyl-tRNA, 45
Aminoglycosides, **191**
  magnesium levels and, 332
  pregnancy use, 204
  Pseudomonas aeruginosa, 143
  teratogenicity, 614
  toxicity of, 251
Aminopenicillins
  mechanism and use, 188
Amiodarone, 323
  hypothyroidism, 249
  hypothyroidism with, 341
  photosensitivity, 250
  pulmonary fibrosis, 251
Amitriptyline, 575
  migraine headaches, 518
Amlodipine, 318
Ammonia
  Ornithine transcarbamylase deficiency, 83
Ammonia transport, 82
Ammonium chloride
  overdose treatment, 233
Ammonium magnesium phosphate (kidney stones), 598
Amnesia
  brain lesions, 511
  classification of, **558**
Amnionitis
  Listeria monocytogenes, 139
Amniotic fluid abnormalities, **641**
Amniotic fluid emboli, 672
Amniotic fluid tests
  AChE in, 491
  with neural tube defects, 491
Amoxapine, 575
Amoxicillin
  clinical use, 188
  Haemophilus influenzae, 142
  Helicobacter pylori, 146
  Lyme disease, 146
  prophylaxis, 198
Amphetamines, 242
  intoxication and withdrawal, 570
  narcolepsy treatment, 568
  norepinephrine and, 239
  as weak bases, 233
Amphotericin B
  clinical use, **199**
  Cryptococcus neoformans, 153
  Naegleria fowleri, 156
  opportunistic fungal infections, 153
  systemic mycoses, 151
Ampicillin
  Clostridium difficile, 138
  Listeria monocytogenes, 139
  mechanism and use, 188
  meningitis, 180
  prophylaxis, 198
Ampulla of Vater, 368
Amygdala
  lesions of, 511
  limbic system, 499
Amylase in pancreatitis, 397
Amylin analogs, 353
Amyloid angiopathy
  intraparenchymal hemorrhage, 513
Amyloidosis
  cardiomyopathy with, 308
  carpal tunnel syndrome, 459
  classification, **212**
  kidney deposition in, 597
  with rheumatoid arthritis, 466
Amyloid precursor protein (APP), 520
Amyotrophic lateral sclerosis (ALS)
  drug therapy for, 549, 551
  spinal cord lesions, 530
Anabolic steroids
  hepatic adenomas and, 392
Anaerobic metabolism
  glycolysis, 74
  pyruvate metabolism, 77
Anaerobic organisms
  aspiration and, 179
  clindamycin, 192
  Clostridia (with exotoxins), 138
  culture requirements, **127**
  glycyclines, 192
  metronidazole, 195
  Nocardia vs *Actinomyces*, 139
  overgrowth in vagina, 148
  pneumonia caused by, 179
Anal atresia, 614
Anal cancer
  HIV and, 177
  oncogenic microbes and, 226
Anal fissures, 366
Anal wink reflex, 510
Anaphase, 46
Anaphylaxis, 112
  blood transfusion, 114
  complement and, 106
  epinephrine for, 242
  IgA-containing products, 116
  shock with, 310
Anaplasma spp
  Gram stain, 125
  transmission, 146, 149
Anaplasmosis
  Anaplasma spp, 150
Anaplastic thyroid carcinomas, 343
Anastomoses, 503
Anastrozole, 656
Anatomic dead space, 664
Anatomic snuff box, 449
Anatomy
  endocrinal, 327–328
  gastrointestinal, 360–369
  of heart, 276, **283**
  hematologic/oncologic, 406–409
  musculoskeletal, 446–454
  nervous system, 493–510
  renal, 580
  reproductive, 624–627
  respiratory, 662–663
"anchovy paste" exudate, 155
Ancylostoma, 159
  diseases associated with, 161
  infection routes, 158
  microcytic anemia, 161
Androblastoma, 653

(c) ketabton.com: The Digital Library

Androgen-binding protein
Sertoli cell secretion, 628
Androgenetic alopecia, 658
Androgenic steroid abuse, 636
Androgen insensitivity syndrome, **639**
Androgen receptor defect, 639
Androgens, source and functions, **636**
Androstenedione, 335, 636
Anemia, **417**
amphotericin B, 199
Ancylostoma, 161
babesiosis, 157
bacterial endocarditis, 311
blood oxygen content, 666
blood transfusion therapy, 429
blood viscosity in, 286
cephalosporins, 189
chloramphenicol, 192
colorectal cancer, 388
cytarabine and, 440
dapsone, 194
Diphyllobothrium latum, 160
drug reaction and, 250
Escherichia coli, 145
ESR in, 214
extrinsic hemolytic, 423
G6PD deficiency, 79
hookworms, 159
in hypertensive emergency, 300
intrinsic hemolytic, 422
isoniazid, 197
kwashiorkor, 71
lab values, 419
macrocytic, 419
malaria, 157
megaloblastic, 420
microcytic, hypochromic, 418
nonhemolytic, normocytic, 421
normocytic, normochromic, 421
NRTIs, 203
oxygen deprivation and, 669
penicillin G, V, 189
pernicious anemia, 372, 379
pregnancy and, 633
pure red cell aplasia, 228
recombinant cytokines for, 121
renal failure, 603
sideroblastic, 67, 419
spherocytes in, 415
in sulfa drug allergies, 252
taxonomy, 417
thioamides causing, 354
trimethoprim, 194
tropical sprue, 381
vitamin B_{12} deficiency, 69
vitamin B_9 deficiency, 68
Weil disease, 147
Wilson disease, 395
Anemia of chronic disease, 421
rheumatoid arthritis, 466
Anencephaly, 491
polyhydramnios and, 641
Anergy, **110**
Anesthetics
general principles, **549**
inhaled, 550
intravenous, 550
local, 550
Aneuploidy, 638
Aneurysms, **516**
atherosclerosis, 302
coarctation of aorta, 299
Ehlers-Danlos syndrome, 51
superior vena cava syndrome, 685
ventricular, 305, 307
Angelman syndrome
chromosome association, 64
Angina
aortic stenosis, 291
atherosclerosis, 302
cocaine causing, 571
contraindicated drugs, **323**
drug therapy for, 318, 324
glycoprotein IIb/IIIa inhibitors for, 438
ischemic disease and, 304
unstable/NSTEMI treatment, 307
Angina, "intestinal," 386
Angina pectoris
β-blockers for, 245
Angiodysplasia, 386
Angioedema, 610
C1 esterase inhibitor deficiency, 107
scombroid poisoning, 247
Angiogenesis
bevacizumab and, 442
wound healing and, 216
Angiokeratomas, 88
Angiomas, 117
Angiosarcomas, 392, 478
carcinogens causing, 225
nomenclature for, 220
Angiotensin-converting enzyme (ACE) inhibitors, **610**
Angiotensin II, 588, 590
ACE inhibitor effects on, 610
signaling pathways for, 337
Angiotensin II receptor blockers, **610**
heart failure, 309
hypertension, 316
naming convention for, 253
preload/afterload effects, 284
Anhedonia, 561
Anhidrosis
Horner syndrome, 540
Anidulafungin, 200
Aniline dyes, 606
transitional cell carcinoma and, 606
Anisocytosis, 407
Anitschkow cells, 312
Ankle sprains, **455**
Ankylosing spondylitis, 469
HLA-B27 and, 100
therapeutic antibodies for, 122
TNF-α inhibitors for, 487
Annular pancreas, 360
Anopheles mosquito, 157
Anopia
visual field defects, 542
Anorectal varices
portal circulation, 365
Anorexia
hypothalamus and, 498
liver cancer/tumors, 392
pancreatic adenocarcinoma, 398
renal failure, 603
Anorexia nervosa
anovulation with, 645
characteristics of, 567
Anosmia
zinc deficiency, 71
ANOVA tests, 264
Anovulation
common causes, 645
Antacids, **399**
Antagonists
ADH, 254
endothelin receptor, 254
ethanol antidote, for, 235
H_2, 254
nonselective, 245
of drugs, 235
Anterior cerebral artery
cingulate herniation, 529
cortical distribution, 502
stroke, 514
Anterior circulation strokes, 514
Anterior communicating artery
saccular aneurysm, 516
Anterior corticospinal tract, 508
Anterior cruciate ligament (ACL) injury
anterior drawer sign in, 454
"unhappy triad," 460
Anterior drawer sign, 454
Anterior hypothalamus, 498
Anterior inferior cerebellar artery, 514
Anterior inferior tibiofibular ligament, 455
Anterior pituitary gland, 327, 331
Anterior spinal artery
complete occlusion, 530
stroke, 514
Anterior talofibular ligament, 455
Anterograde amnesia, 558
benzodiazepines, 550
brain lesions, 511
Anthracosis, 677
Anthracyclines, 439
cardiomyopathy from, 248
Anthrax, 132, 137
Anthrax toxin
Bacillus anthracis and, 137
Anti-ACh receptor antibody, 115
Antiandrogen drugs, **658**
Antianginal therapy, 307, 318, **319**, 324
Antiapoptotic molecule
oncogene product, 224
Antiarrhythmic drugs
mechanisms and clinical uses, **322–324**
torsades de pointes, 248
Antibiotics, 153
acne treatment, 477
Clostridium difficile with, 138
Jarisch-Herxheimer reaction with, 148
long QT interval, 294
selective growth media, 126
torsades de pointes, 248
Antibodies
in adaptive immunity, 99
antibody diversity generation, 104
antibody specificity generation, 104
exo- and endotoxins, 131, 133
hepatitis viruses, 174
hypersensitivity mediation, 112
structure and function, **104**
therapeutic, 122
Antibody-dependent cell-mediated cytotoxicity, 101
Anticardiolipin
antiphospholipid syndrome, 470
Anticardiolipin antibody, 115
Anti-CCP antibody, 115
Anti-centromere antibodies
scleroderma, 473
Anticentromere autoantibody, 115
Anticholinergic drugs
delirium with, 558
toxicity treatment for, 248
Anticholinesterase drugs, 240
Anticholinesterase poisoning, **240**
Anticipation (genetics), 56
Anticoagulant drugs
acute coronary syndromes, 307
antiphospholipid syndrome, 470
atrial fibrillation, 295
coagulation cascade and, 412
Anticoagulation
coagulation cascade and, 413
Anticonvulsant drugs, 471
Antidepressant drugs, **574**–576
atypical, 576
for fibromyalgia, 471
long QT interval with, 294
torsades de pointes, 248
Anti-desmoglein (anti-desmosome) autoantibody, 115
Anti-digoxin Fab fragments, 248
for cardiac glycoside toxicity, 321
Antidiuretic hormone (ADH), **329**, 588, 590
antagonist naming conventions, 254
antagonists, 338, 354
in diabetes insipidus, 338
function of, 328
hypothalamus synthesis, 498
pituitary gland and, 327
SIADH and, 338
signaling pathways of, 337
Anti-DNA topoisomerase I autoantibody, 115
Anti-dsDNA antibody, 115
Antiemetic drugs
aprepitant, 401
long QT interval with, 294
marijuana, 571
metoclopramide, 400
ondansetron, 400
torsades de pointes, 248
Antiepileptic drugs
Cytochrome P-450 interactions, 252
for fibromyalgia, 471
rash from, 250
teratogenicity, 614
Antifungal drugs
griseofulvin, 48
mechanism and use, **198**–200
seborrheic dermatitis, 476
tinea versicolor, 152
Antigenic shift/drift, 169
Antigen-presenting cells (APCs)
B cells as, 409
CD28, 110
MHC I and II and, 100
naive T-cell activation, 103
in spleen, 98
Antigens
active immunity, 110
antibody structure and function, 104
chronic mucocutaneous candidiasis, 116
for self, 102
HLA I and II, 100
lymphocyte recognition of, 98
type and memory, **105**
Anti-glomerular basement membrane autoantibody, 115
Anti-glutamic acid decarboxylase autoantibody, 115
Antigout drugs
colchicine, 48
Anti-HBc, 174
Anti-HBe, 174
Anti-HBs, 174
Anti-helicase autoantibody, 115

(c) ketabton.com: The Digital Library

Antihelminthic drugs, **200**
  mebendazole, 48
  naming convention, 253
Anti-hemidesmosome autoantibody, 115
Antihistamines, 686
  for scombroid poisoning, 247
Anti-histone antibody, 115
Antihypertensive drugs
  hypertension in pregnancy, 643
Anti-IgE monoclonal therapy, 687
Anti-IL-5 monoclonal therapy, 687
Anti-inflammatory drugs, 485
Anti-intrinsic factor autoantibody, 115
Anti-La/SSB autoantibody, 115
Antileukotrienes
  for asthma, 687
Antimetabolites, **440**
Antimicrobial drugs
  highly resistant bacterial treatment, 198
  mechanisms of action summary, **187**–204
  naming conventions for, 253
  pregnancy contraindications, **204**
  prophylaxis, **198**
Antimicrosomal autoantibody, 115
Anti-mite/louse therapy, **200**
Antimitochondrial autoantibody, 115
Antimuscarinic drugs
  Parkinson disease, 548
  reactions to, 251
  toxicity treatment for, 248
Antimycin A
  electron transport chain, 78
Antimycobacterial therapy, **196**
Anti-NMDA receptor paraneoplastic syndrome encephalitis, 228
Antinuclear (ANA) antibody, 115
  Sjögren syndrome, 468
Antioxidants
  free radical elimination by, 210
Antiparasitic drugs
  naming convention for, 253
Antiparietal cell autoantibody, 115
Anti-phospholipase A2 receptor autoantibody, 115
Antiphospholipid syndrome, **470**
  autoantibody in, 115
Antiplatelet antibodies
  abciximab as, 122
Antiplatelet drugs
  for acute coronary syndromes, 307
Anti-presynaptic voltage-gated calcium channel autoantibody, 115
Antiprogestin drugs, **657**
Antiprotozoan drugs, **200**
Antipseudomonal drugs
  cephalosporins, 189
  fluoroquinolones, 195
  penicillins, **188**
Antipsychotic drugs
  adverse effects/events, 573
  antimuscarinic reaction, 251
  atypical, 573
  dopaminergic pathways, 499
  dystonia with, 569
  long QT interval with, 294
  Parkinson-like syndrome, 251
  tardive dyskinesia, 251
  torsades de pointes, 248
  Tourette syndrome, 572
  typical, 573
Antiribonucleoprotein antibodies
  Sjögren syndrome, 468
Anti-Ro/SSA autoantibody, 115
Anti-Scl-70 autoantibody, 115
Anti-Smith autoantibody, 115
Anti-smooth muscle autoantibody, 115
Antisocial personality disorder, 565
  early-onset disorder, 557
Antispasmodics, 551
Anti-SRP autoantibody, 115
Anti-streptolysin O (ASO) titers, 312
Antisynthetase autoantibody, 115
Antithrombin
  coagulation cascade and, 413
  deficiency of, 428
Antitoxins
  as passive immunity, 110
Anti-TSH receptor autoantibody, 115
Antitumor antibiotics, **439**
Anti-U1 RNP antibodies, 115, 470
Antiviral therapy
  hepatitis C, 203, 204
  mechanism and use, **201**
Anti-β_2 glycoprotein
  antiphospholipid syndrome, 470
  autoantibody, 115
Anxiety
  drug therapy, 546, 563, 575
  neurotransmitters, 495
Aorta
  branches, 363
  coarctation of, 299, 300
  congenital heart disease, 298
  diaphragm, 663
  ECG and, 293
  retroperitoneal, 360
  in syphilitic heart disease, 312
  traumatic rupture of, 303
  "tree bark" appearance, 312
Aortic aneurysm, 302
  Ehlers-Danlos syndrome, 51
  hypertension, 300
  Marfan syndrome, 300
  syphilitic heart disease, 312
Aortic arch derivatives, **619**
Aortic arch receptors, 296
Aortic dissection, **303**
  hypertension, 300
  Marfan syndrome, 300
Aortic insufficiency
  syphilitic heart disease, 312
Aorticopulmonary septum, 281
  embryologic derivatives, 613
Aortic regurgitation, 288
  heart murmurs with, 291
  Marfan syndrome, 300
Aortic root dilation
  heart murmur with, 291
Aortic stenosis, 288
  heart murmurs, 291
  macroangiopathic anemia, 423
  Williams syndrome, 300
Aortic valve
  cardiac cycle, 287
  embryological development of, 281
Aortitis
  syphilis, 147, 184
Apalutamide, 658
APC gene, 224
  adenomatous colonic polyps and, 387
  colorectal cancer and, 389
  familial adenomatous polyposis and, 387
"Ape hand," 447, 451
Apgar score, **634**
Aphasia
  MCA stroke, 514
  types of, **516**
Aphthous stomatitis
  Crohn disease, 382
Apixaban
  factor Xa inhibitors, 437
Aplasia, 613
Aplasia cutis
  methimazole, 354
Aplastic anemia, 421
  chloramphenicol, 192
  drug reaction and, 250
  neutropenia with, 424
  thionamides, 354
Aplastic crisis
  hereditary spherocytosis, 422
  sickle cell anemia, 422
Apolipoproteins
  functions of, 93
Apoptosis, **208**
  corticosteroids, 424
Appendages (bacterial), 124
Appendicitis, **383**
  mittelschmerz vs, 631
Appetite regulation, **336**, 371
"Apple core" lesion (X-ray), 388
Apraclonidine, 552
Aprepitant, **401**
Aqueous humor pathway, **535**
Arabinofuranosyl cytidine, 440
Arabinogalactan synthesis, 196
Arabinosyltransferase, 197
Arachidonic acid pathway, **485**
Arachnodactyly, 52
Arachnoid granulations, 503, 504, 522
Arachnoid mater
  meninges, 496
  meningioma, 526
  ventricular system, 504
Arcuate fasciculus
  aphasia and, 516
  diagram, 501
Area postrema, 498
Area under the curve, 231
Arenaviruses, characteristics of, 167
Argatroban, 435
Arginine
  classification, 81
  cystinuria, 85
  kidney stones and, 598
Argininosuccinate,, 82
Argyll Robertson pupils
  in syphilis, 184
  tabes dorsalis, 530
Aripiprazole, 573
Arm abduction, **446**
Armadillos (disease vectors), 149
Aromatase, 636
  in pathway, 335
Aromatase inhibitors, **656**
Aromatic amines
  carcinogenicity of, 225
Arrhythmias
  amphotericin B, 199
  diabetic ketoacidosis, 347
  diphtheria, 139
  drug reactions and, 248
  hypokalemia and, 591
  local anesthetics and, 550
  macrolides, 193
  McArdle disease, 87
  muscular dystrophy, 61
  myocardial infarction and, 305, 307
  shock caused by, 310
  sleep apnea and, 679
  stimulants and, 570
  TCA toxicity, 569
  thyroid hormones and, 354
Arsenic
  angiosarcomas, 392, 478
  carcinogenicity of, 225
  glycolysis and, 74
  toxicity symptoms, 76
  toxicity treatment, 248
Artemether, 200
Arterial oxygen saturation, 666
Arterial PCO_2, 668
Arteries, anatomy of, 283
Arteriolosclerosis, 301, 346
Arteriosclerosis, **301**
Arteriovenous malformations (AVMs)
  hereditary hemorrhagic telangiectasia, 316
Arteriovenous shunts, 463
Arteritis
  giant cell (temporal), 314, 518
  headaches, 518
Artesunate
  malaria, 157, 200
Arthralgias
  alkaptonuria, 84
  coccidiomycosis, 151
  Henoch-Schönlein purpura, 315
  hepatitis viruses, 172
  in alkaptonuria, 84
  rubella, 169, 182
  serum sickness, 113
  vitamin A toxicity, 66
Arthritis, 457, 466
  azathioprine for, 440
  Campylobacter jejuni, 145
  carpal tunnel syndrome and, 459
  celecoxib for, 486
  chlamydiae, 148, 184
  Crohn disease, 382
  gonococcal, 468
  gonorrhea, **142**, 180, 184
  immunosuppressants, 120
  inflammatory bowel disease, 100
  lupus, 470
  Lyme disease, 146
  psoriatic, 469
  reactive arthritis, 469
  septic, 468
  Staphylococcus aureus, 135
  Takayasu arteritis, 314
  therapeutic antibodies, 122
  ulcerative colitis, 382
Arthropathy
  hemochromatosis, 395
Arthus reaction, 113
Arylsulfatase A
  metachromatic leukodystrophy, 88
Arytenoids, 620
Asbestos
  carcinogenicity, 225
Asbestosis, **677**, 678
Ascaris lumbricoides, 159
Ascaris spp, 158
Ascending cholangitis, 397
Ascending colon, 360
Aschoff bodies, 312
Ascites
  Budd-Chiari syndrome, 392
  hepatocellular carcinoma, 392
  spontaneous bacterial peritonitis, 390
Ascorbic acid, 69
Asenapine, 573
Aseptic meningitis
  mumps, 170
  picornaviruses, 167

(c) ketabton.com: The Digital Library

Asherman syndrome, 648
Ashkenazi Jews
  disease incidence, 88
ASO titer, 136
Aspart, 352
Aspartame, 84
Aspartate
  urea cycle, 82
Aspartate aminotransferase (AST), 390
  hepatitis, 172
  toxic shock syndrome, 135
Aspartic acid, 81
Aspergillosis
  Aspergillus fumigatus, 153
  bronchiectasis, 674, 675
  echinocandins, 200
Aspergillus fumigatus, 153
  HIV-positive adults, 177
Aspergillus spp
  chronic granulomatous disease, *109*
Aspiration
  ARDS and, 678
  in utero "breathing," 660
  lung abscess, 685
  reflux-related, 359, 377
  Zenker diverticulum, 384
Aspiration pneumonia
  alcoholics, 179
  clindamycin, 192
  nosocomial infections, 185
Aspirin, **486**
  acute coronary syndromes, 307
  cyclooxygenase, 411
  hemolysis in G6PD deficiency, 250
  for ischemic stroke, 512
  Kawasaki disease, 314
  thrombogenesis and, 411
  uncoupling agent, 78
  as weak acid, 233
  zero-order elimination of, 232
Asplenia
  Streptococcus pneumoniae, 136
  target cells, 415
Asterixis, 519
  hepatic encephalopathy, 391
  renal failure, 603
Asteroid bodies, 676
Asthma, 674
  albuterol for, 242
  β-blockers and, 245
  breast milk and, 636
  cromolyn sodium for, 408
  drug therapy, **687**
  epinephrine for, 242
  gastroesophageal reflux disease, 377
  hypertension treatment with, 316
  immunosuppressants, 120
  muscarinic antagonists for, 241
  omalizumab for, 122
  pulsus paradoxus in, 310
  salmeterol for, 242
  type I hypersensitivity, 112
Astigmatism, 535
Astrocytes, **493**
  foot processes, 496
  origin of, 490
Ataxia
  abetalipoproteinemia, 94
  cerebellar hemisphere lesions, 511
  Friedreich, 62, 64, 531
  hypnotics, 546
  lithium toxicity, 569
  metachromatic leukodystrophy, 88
  normal pressure hydrocephalus, 522
  opsoclonus-myoclonus syndrome, 228
  prion disease, 178
  psychoactive drug intoxication, 570
  streptomycin, 197
  syphilis, 147
  tabes dorsalis, 530
  trinucleotide repeat expansion disease, 62
  truncal, 511
  vitamin B_{12} deficiency, 530
  vitamin E deficiency, 70
  Wernicke-Korsakoff syndrome, 511, 571
Ataxia-telangiectasia, 40, **117**
Atazanavir, 203
Atelectasis, **680**
Atenolol, 245, 323
Atherosclerosis, **302**
  abdominal aortic aneurysms and, 302
  diabetes mellitus and, 346
  familial dyslipidemias, 94
  homocystinuria, 84
  renovascular disease, 604
  stable angina with, 304
  transplant rejection, 119
Athetosis, 511, 519
Atomoxetine, 557
Atonic seizures, 517
Atopic dermatitis (eczema), 477
Atopic reactions, 112
Atovaquone
  babesiosis, 157
  malaria, 157
  P. falciparum, 200
  for *Pneumocystis jirovecii*, 154
ATPase, 395
ATP production, **74**
Atresia
  anal, 614
  duodenal, 359
  esophageal, 359
  intestinal, 359
  jejunal/ileal, 359
Atria
  cardiac tumors, 316
  depolarization/repolarization of, 293
  embryologic development of, 281
Atrial fibrillation
  β-blockers for, 323
  calcium channel blockers for, 324
  cardiac glycosides for, 321
  ECG tracing of, 295
  hypertension, 300
  jugular venous pulse in, 287
  potassium channel blockers for, 323
Atrial flutter
  β-blockers for, 323
  ECG tracing of, 295
  potassium channel blockers for, 323
"Atrial kick," 287
Atrial natriuretic peptide (ANP), **296**, 588, 590
  in SIADH, 338
  signaling pathways for, 337
Atrial septal defect (ASD), 299
  congenital rubella, 300
  Down syndrome, 300
  fetal alcohol syndrome, 300
Atrioventricular (AV) block
  β-blockers, 245, 323
  calcium channel blockers, 318, 324
  ECG tracings, 295
  Lyme disease, 146
Atrioventricular canals, 281
Atrioventricular node
  AV node, 292
  conduction pathway, 293
  ECG and, 293
Atrioventricular valves
  embryologic development of, 281
Atrophic gastritis
  gastrin in, 371
Atrophy, 206
  motor neuron signs, 529, 531
  neurodegenerative disorders, 520
  optic disc/nerve, 536
  ventriculomegaly, 522
Atropine, **241**
  antimuscarinic reaction, 251
  for β-blocker overdose, 323
  effects of, 241
  toxicity treatment, 248
Attention-deficit hyperactivity disorder (ADHD), 557
  drug therapy for, 557, 572
  Tourette syndrome, 557
Attributable risk (AR), 258
Atypical antidepressants, **576**
Atypical antipsychotic drugs, **573**
  postpartum psychosis, 562
Atypical depression, 575
Atypical pneumonias
  chlamydiae, 148
  macrolides, 193
  typical organisms, 683
Auditory cortex
  diagram, 501
  thalamic relay, 498
Auditory hallucinations, 559
Auditory physiology, **533**
Auerbach plexus, 376, 384
Auer rods
  in AML, 432
Auramine-rhodamine stain, 125
Auscultation of heart, **290**
Auspitz sign, 477
Autism spectrum disorder, 557
  double Y males and, 638
  fragile X syndrome, 62
Autoantibodies, **115**
Autoclaves
  disinfection/sterilization, 204
Autodigestion, 397
Autoimmune diseases
  acute pericarditis, 313
  blistering skin, **480**
  collagen and, 50
  Dressler syndrome, 307
  myocarditis with, 313
  rheumatoid arthritis, 466
  self-antigen in, 102
  Sjögren syndrome, 468
  SLE, 470
Autoimmune gastritis, 379
Autoimmune hemolytic anemia, 423
  cephalosporins, 189
Autoimmune hepatitis type 1
  autoantibody, 115
Autoimmune hypothyroidism, 173
Autoimmune lymphoproliferative syndrome, 208
Autoimmune regulator (AIRE), 102
Autoimmune thrombocytopenia, 121
Autonomic drugs, **236–245**
  actions of, 239
  bladder dysfunction, action on, 237
  naming conventions for, 253
Autonomic ganglia, 236
Autonomic insufficiency, 242
Autonomic nervous system (ANS)
  delirium tremens, 569
  dysregulation in inflammatory demyelinating polyradiculopathy, 524
  limbic system in, 499
  male sexual response, **627**
  receptors in, **236**
  in serotonin syndrome, 569
Autonomy (ethics), 265
Autoregulation of blood flow, **297**
Autosomal dominant diseases, **60**
  ADPKD, 516
  Brugada syndrome, 294
  Charcot-Marie-Tooth disease, 524
  elastin syndrome, 52
  Huntington disease, 520
  hyper-IgE syndrome, 116
  hypertrophic cardiomyopathy, 308
  malignant hyperthermia susceptibility, 550
  neurofibromatosis, 525
  porphyrias, 425
  Romano-Ward syndrome, 294
  tuberous sclerosis, 525
  von Hippel-Lindau disease, 525
Autosomal dominant polycystic kidney disease (ADPKD), 604
  chromosome association, 64
Autosomal dominant tubulointerstitial kidney disease, 604
Autosomal recessive diseases, **60**
  abetalipoproteinemia, 94
  adenosine deaminase deficiency, 117
  alkaptonuria, 84
  Chédiak-Higashi syndrome, 117
  cystic fibrosis, 60
  5α-reductase deficiency, 639
  Friedreich ataxia, 531
  hemochromatosis, 395
  hereditary hyperbilirubinemias, 394
  IL-12 receptor deficiency, 116
  Jervell and Lange-Nielsen syndrome, 294
  Kartagener syndrome, 49
  leukocyte adhesion deficiency, 117
  maple syrup urine disease, 84
  severe combined immunodeficiency, 117
  spinal muscular atrophy, 530
  Wilson disease, 395
Autosomal recessive polycystic kidney disease (ARPKD), 604
  Potter sequence caused by, 578
Autosomal trisomies, **63**
  Down syndrome (trisomy 21), 63
  Edwards syndrome (trisomy 18), 63
  karyotyping for, 55
  Patau syndrome (trisomy 13), 63
Avascular necrosis
  femoral head, 461
Avascular necrosis, **463**
  scaphoid bone, 449
  sickle cell anemia, 422
Aversive stimulus (positive punishment), 554
Avoidant personality disorder, 566
Axilla/lateral thorax, 455
Axillary nerve
  arm abduction, 446
  injury presentation, 447
  neurovascular pairing, 455

(c) ketabton.com: The Digital Library

Axonal injury, 495
Axonal trafficking, 48
Axonemal dynein, 49
Azathioprine
  antimetabolites, 440
  for Crohn disease, 382
  immunosuppressant, 120
  pancreatitis caused by, 249
Azithromycin
  atypical pneumonia treatment, 148
  babesiosis, 157
  chlamydiae, 148
  in cystic fibrosis, 60
  gonorrhea treatment, 142
  macrolides, 193
  Mycobacterium avium-intracellulare, 140, 196
  prophylaxis in HIV, 198
Azoles, 153, **199**
  vaginal infections, 181
Azotemia
  acute interstitial nephritis, 601
Aztreonam, 190

B

Babesia spp, 146, 157
Babesiosis, 157
Babinski reflex, 635
  motor neuron signs, 529
  primitive reflexes, 510
Bachmann bundle, 293
Bacillary angiomatosis, 478
  animal transmission, 149
  HIV-positive adults, 177
Bacillus anthracis, **137**
  exotoxin production, 132
Bacillus cereus, **138**
  food poisoning, 178
Bacitracin
  gram-positive antibiotic test, 134
  sensitivity to, 134, 136
Baclofen, 551
  multiple sclerosis, 523
Bacteremia
  brain abscesses, 180
  cutaneous anthrax, 137
  daptomycin, 195
  Staphylococcus gallolyticus, 137
  Streptococcus bovis, 137
Bacteria
  biofilm-producing, 128
  culture requirements, 126
  encapsulated, 127
  genetics, **130**, 131
  hemolytic, 135
  highly resistant, 198
  normal flora, 178
  phage infection of, 130
  pigment-producing, 128
  spore-forming, 129
  stains for, 125
  structures and functions, **124**
  virulence factors, 127, **129**, 135, 143, 144, 145
  zoonotic, 149
Bacterial capsules, 124
Bacterial endocarditis, **311**
  daptomycin, 195
  Staphylococcus aureus, 135
Bacterial exotoxin mechanisms
  increase fluid secretion, **132**
  inhibit phagocytic ability, 132
  inhibit protein synthesis, 132
  lyse cell membranes, 133
  superantigens causing shock, 133
Bacterial infections
  with immunodeficiency, 118
  myocarditis with, 313
  skin, 479
Bacterial peritonitis (spontaneous), 389, 390
Bacterial toxin mechanisms
  inhibit release of neurotransmitter, 132
  lysogenic phage encoding of, 130
Bacterial vaginosis
  characteristics of, 158, 181
  Gardnerella vaginalis, 148
Bacteroides fragilis, 178
Bacteroides spp
  alcoholism, 179
  clindamycin, 192
  culture requirements of, 127
  metronidazole, 195
  nosocomial infections, 185
"Bag of worms," 651
Baker cyst, 460
  tibial nerve injury, 453
BAK protein, 208
Balancing (quality measurement), 273
Bamboo spine, 469
Band cells, 406
Barbiturates
  intoxication and withdrawal, 570
  intravenous anesthetics, 550
  mechanism and use, **546**
  naming convention for, 253
  sleep alterations, 497
Barlow maneuver, 461
Baroreceptors, **296**
Barr bodies, 34
  in x-inactivation, 61
Barrett esophagus, **378**
Bartholin cyst/abscess, 644
Bartonella henselae
  bacillary angiomatosis, 478
Bartonella quintana, 161
Bartonella spp
  animal transmission, 149
  Gram Stain, 125
Bartter syndrome, 586
  markers in, 591
Basal cell carcinomas, 484
  5-fluorouracil for, 440
Basal ganglia, **500**
  intraparenchymal hemorrhage, 513
  lesions in, **511**
  movement disorders, 519
  thalamic connections, 498
Basal lamina, 50
Basal nucleus of Meynert, 495
Basal plate, 490
Base excision repair, 40
Basement membrane, 98
  blood-brain barrier, 496
  collagen in, 50
  glomerular filtration barrier, 581
Basic amino acids, 81
Basilar artery
  herniation syndromes, 529
  stroke effects, 515
Basilar membrane (cochlea), 533
Basiliximab
  immunosuppressant, 120
Basophilia, 408
Basophilic stippling, 416
  lead poisoning, 419
  sideroblastic anemia, 419
Basophils, **408**
  IgE antibody, 105
BAX protein, 208
B-cell lymphomas
  HIV-positive adults, 177
B cells, **409**
  activation, 103, 105
  adaptive immunity, 99
  anergy, 110
  cell surface proteins, 110
  class switching, 103
  disorders of, 116, 117
  function of, 409
  functions of, 101
  glucocorticoid effects, 120
  immunodeficiency infections, 118
  in lymph node, 96
  neoplasms, 430
  non-Hodgkin lymphoma, 429
  sirolimus effect, 120
  spleen, 98
BCG vaccine
  false positives from, 140
  IL-12 receptor deficiency and, 116
BCL-2 gene, 224
Bcl-2 protein, 208
BCR-ABL gene, 224
Bead-like costochondral junctions, 463
Becker muscular dystrophy, 61
Beck triad (cardiac tamponade), 310
Beckwith-Wiedemann syndrome, 358, 606
Beers criteria, **247**
Behavioral therapy, 572
Behavior modulation
  frontal lobe lesions and, 511
  limbic system and, 499
Behçet syndrome, 314
Bell palsy, 532, 676
Bell-shaped distribution, 262
Bendazoles, 159
Bends, 463
Beneficence (ethics), 265
Benign breast disease, **649**
Benign paroxysmal positional vertigo (BPPV), 534
Benign prostatic hyperplasia (BPH), 237, **654**
  α_1-blockers for, 244
  epididymitis and orchitis with, 654
  tamsulosin for, 658
  treatment of, 237
Benign tumors, 220
  bones, 464
  breast, 649
Benralizumab, 687
Benzathine penicillin G, 198
Benzene
  aplastic anemia, 250
Benzidine as carcinogen, 225
Benznidazole, 158
Benzocaine, 550
Benzodiazepines
  addictive risk, 546
  alcohol withdrawal, 572
  Beers criteria, 247
  clinical use and adverse effects, **546**
  cocaine overdose, 571
  epilepsy treatment, 544
  intoxication and withdrawal, 570
  naming convention for, 253
  phobias, 563
  sleep effects, 497
  toxicity treatment for, 248
Benzoyl peroxide for acne, 477
Benztropine, 241, 548
Berger disease, 596
Beriberi
  vitamin B_1 deficiency, 66
Berkson bias, 260
Bernard-Soulier syndrome, 411, 427
Berry aneurysm, 516
Berylliosis, 677
β_1-blockade, 284
β_2-agonists
  naming convention for, 253
β_2-agonists
  asthma, 687
  insulin and, 334
β-adrenergic agonist
  potassium shifts, 590
β-blockers, 245, **323**
  acute coronary syndromes, 307
  angina, 319
  aortic dissections, 303
  Cardiomyopathy (hypertrophic), 245
  cocaine overdose, 571
  diabetes and, 245
  dilated cardiomyopathy, 308
  essential tremor, 519
  for cocaine overdose, 242
  for pheochromocytomas, 350
  for thyroid storm, 342
  glaucoma treatment, 552
  heart failure, 309
  hyperkalemia, 590
  hypertension, 316
  hypertrophic cardiomyopathy, 308
  juxtaglomerular apparatus effects, 589
  migraine headaches, 518
  naming convention for, 253
  overdose treatment, 323
  phobias, 563
  selectivity, 245
  toxicity treatment for, 248
β cells, 328
  diabetes mellitus type 1 and 2, 347
  insulinomas of, 351
  insulin production by tumors, 351
  insulin secretion by, 334
β-dystroglycan, 61
β-galactosidase, 144
β-glucan, 200
β-glucuronidase, 406
β-hCG, 653
  as tumor marker, 226
β-hemolysis, 133
β-hemolytic bacteria, 135
  common colonization sites, 135
  Staphylococcus aureus, 135
  Staphylococcus epidermidis, 135
  Staphylococcus saprophyticus, 136
  Streptococcus agalactiae (Group B strep), 137
  Streptococcus pyogenes (Group A strep), 136
β-hydroxybutyrate, 90
β-interferon
  multiple sclerosis, 523
β-lactam antibiotics, 187
β-lactamase inhibitors, 189
β_2-microglobulin
  MHC I and II and, 100
β-oxidation of very-long-chain fatty acids (VLCFA), 47
β-prophage
  Corynebacterium exotoxin encoding, 139
Beta rhythm (EEG), 497
β-thalassemia, 418
  allelic heterogeneity, 57
Betaxolol, 245, 552
Bethanechol, 240

(c) ketabton.com: The Digital Library

Bevacizumab, 122, **442**
Bezafibrate, 320
Bias and study errors, **260**–262
Bicalutamide, 658
Bicarbonate
  carbon dioxide transport, 670
  GI secretion, 372
  overdose treatment, 233
  pancreatic insufficiency, 381
  salicylate toxicity, 248
  TCA toxicity, 233, 248
Biceps brachii muscle
  Erb palsy, 448
Biceps femoris, 452, 453
Biceps reflex, 510
Biceps tendon, 446
Bicornuate uterus, 623
Bicuspid aortic valve
  aortic dissection and, 303
  coarctation of aorta and, 299
  heart murmur with, 291
  thoracic aortic aneurysms and, 302
  Turner syndrome, 300
Bifid ureter, 579
Bifurcation external landmarks, 663
Biguanide drugs, 353
Bilaminar disc, 612
Bilateral adenopathy, 676
Bilateral renal agenesis
  oligohydramnios and, 641
  Potter sequence, 578
  pulmonary hypoplasia and, 660
Bile, **374**
  hereditary hyperbilirubinemias, 394
  secretin effect on, 371
Bile acid resins, 320
  lipid transport and, 92
  reabsorption of, 320
  synthesis of, 47
Bile canaliculus, 367
Bile ducts, 367, 368
Bile salts, 374
  in cholelithiasis, 396
Biliary atresia, **393**
Biliary cholangitis, primary
  autoantibody, 115
Biliary cirrhosis, 389, 393
  cystic fibrosis, 60
Biliary colic, 396
Biliary structures, **368**
Biliary tract disease, **395**
  Clonorchis sinensis, 161
  gallstones, 368
  hyperbilirubinemia with, 393
Biliary tract infections
  Enterococci, 137
Bilirubin, **375**
  bile, 374
  cholelithiasis, 396
  hereditary hyperbilirubinemias, 394
  liver marker, 390
  toxic shock syndrome, 135
Bimatoprost, 552
Bimodal distribution, 262
Binge-eating disorder, 567, 575
Binge-eating/purging, anorexia nervosa, 567
Bioavailability, 231
  area under the curve from, 231
Biochemistry
  cellular, 46–52
  genetics, 56
  laboratory techniques, 52–94
  metabolism, 72–94
  molecular, 34
  nutrition, 65
Biochemistry laboratory techniques
  blotting procedures, 53
  Cre-lox system, 56
  CRISPR/Cas9, 53
  enzyme-linked immunosorbent assay, 54
  flow cytometry, 54
  fluorescence in situ hybridization, 55
  free light chain (FLC) assay, 431
  gene expression modifications, 56
  karyotyping, 55
  microarrays, 54
  molecular cloning, 55
  polymerase chain reaction, 52
  reverse transcriptase polymerase chain reaction, 52
  RNA interference, 56
  serum protein electrophoresis, 431
Biofilm-producing bacteria, 128
  Staphylococcus epidermidis, 135
Biologic agents
  naming conventions for, **254**
Biomarkers
  α-fetoprotein, 491
  astrocytes, 493
  neurons, 493
Biostatistics/epidemiology, 256–262
Biotin, 68
Bipolar disorder, **561**
  drug therapy for, 572
  lithium for, 574
Birbeck granules
  Langerhans cell histiocytosis, 434
"Bird's beak" sign (X-ray), 376
Birds (disease vectors), 148, 149
Bismuth, **399**
Bisoprolol, 245
Bisphosphonates, 462, **486**
  esophagitis with, 249
  naming convention for, 254
  osteogenesis imperfecta treatment, 51
Bitemporal hemianopia, 542
  craniopharyngioma, 528
  hypopituitarism, 339
  optic chiasm compression, 516
Bitot spots, 66
Bivalirudin, 435
BK virus, 164
Black eschar, 137
Black lung disease, 677
Bladder, 160
  bethanechol effect on, 240
  BPH and, 654
  development of, 618
  exstrophy, 624
  outlet obstruction, 579
  placenta percreta invasion, 640
  spasm treatment, 241
  urachus, 618
  urgency in cystitis, 241
Bladder cancer
  cisplatin/carboplatin for, 442
  hematuria with, 594
  hypercalcemia and, 228
  Schistosoma haematobium, 161
"Blast crisis," 433
Blastocyst implantation, 612
Blastomyces spp
  amphotericin B, 199
  itraconazole, 199
Blastomycosis, 151
Bleeding, 642
  adenomatous polyps, 387
  direct factor Xa inhibitors, 437
  direct thrombin inhibitors, 435
  diverticulosis, 383
  essential thrombocythemia, 433
  glycoprotein IIb/IIIa inhibitors, 438
  heparin, 436
  inflammatory bowel disease, 382
  peptic ulcer disease, 380
  thrombolytics, 437
  variceal, 371
  warfarin, 436
Bleeding time, 427
Bleomycin
  antitumor antibiotics, 439
  pulmonary fibrosis, 251
  targets of, 438
  toxicity, 444
Blepharospasm, 519
Blindness
  Chlamydia trachomatis, 149
  giant cell arteritis, 314
  neonatal, 142
  Onchocerca volvulus, 159
  Toxocara canis, 159
Blistering skin disorders, 480
Blood
  chocolate-colored, 666
  coagulation and kinin pathways, 412
  hCG detection in, 633
  oxygen content, 666
  in placenta, 617
  umbilical cord, 618
  viscosity of, 668
Blood-brain barrier
  anesthetics, 549
  astrocytes, 493
  function and mechanism, **496**
  at hypothalamus, 498
  L-DOPA, 549
Blood flow
  autoregulation, 297
  exercise response, 670
Blood groups, **405**
Blood-nerve permeability barrier, 495
Blood pH
  diuretic effects on, 609
Blood pressure
  angiotensin II effects, 588
  antianginal therapy, 319
  cortisol effect on, 336
  fenoldopam and, 318
  renal disorders and, 591
  sympathomimetic effect on, 243
Blood-testis barrier, 628
Blood transfusions, **429**
  reactions, **114**
  therapy, 429
Blood vessels
  collagen in, 50
  Ehlers-Danlos syndrome, 50
  hereditary hemorrhagic telangiectasia, 316
Blood volume
  atrial natriuretic peptide release, 296
  regulation, 588
Bloody diarrhea, 179
  Campylobacter jejuni, 145, 149
  Shigella, 144
  ulcerative colitis, 382
Bloody stool, 366
Blotting procedures, **53**
Blown pupil, 541
  CN III damage, 541
  saccular aneurysms, 516
"Blue babies," 298
Blueberry muffin rash
  cytomegalovirus, 182
  rubella, 169, 182
  Toxoplasma gondii, 182
"Blue bloater," 674
"Blue kids," 299
Blue sclerae, 51
Blumer shelf, 379
BMPR2 gene, 679
Body compartments, 231
Body dysmorphic disorder, 563
Boerhaave syndrome, 377
Bombesin, 350
Bone cancer, 464
  primary bone tumors, 464
Bone cell biology, 458, 459
Bone crises, 88
Bone disorders
  adult T-cell lymphoma and, 430
  lab values in, 464
  Langerhans cell histiocytosis, 434
  lytic ("punched out"), 431
  osteogenesis imperfecta, 51
Bone formation, **458**
Bone-in-bone (x-ray), 463
Bone marrow
  cytokine stimulation of, 121
  lymphoid functions of, 96
  suppression, 199
Bone mineral density scan, 462
Bone morphogenetic protein (BMP), 490
Bones
  collagen in, 50
  cortisol effect on, 336
  lytic/blastic metastases, 223
  primary bone tumors, 464
  PTH effect on, 332
  renal osteodystrophy, 603
Borderline personality disorder, 565
Bordetella pertussis, **143**
  culture requirements, 126
  exotoxin production, 132
  macrolides, 193
  vaccines, 143
Bordet-Gengou agar, 126
Borrelia burgdorferi
  animal transmission, 149
  coinfection with, 157
  facial nerve palsy, 186
  Lyme disease, 146
  tetracyclines, 192
Borrelia recurrentis
  animal transmission, 149
  vectors, 161
Borrelia spp, 146
Bortezomib, **443**
Bosentan, 686
Botulinum toxin
  lysogenic transduction, 130
  passive antibodies for, 110
  symptoms of, 138
  toxin effects, 132
Bovine spongiform encephalopathy (BSE), 178
Bowel stenosis, 383
Bowen disease, 651
Bowenoid papulosis, 651
Bow legs (genu varum), 463

(c) ketabton.com: The Digital Library

Bowman capsule, 583
Boxer's fracture, 459
BPH (benign prostatic hyperplasia)
  hydronephrosis in, 599
Brachial artery, 455
Brachial plexus
  Pancoast tumor, 685
Brachial plexus lesions, **448**
Brachiocephalic syndrome, 685
Brachiocephalic vein, 685
Brachioradialis reflex, 510
Bradycardia
  amiodarone and, 323
  atropine for, 241
  β-blockers and, 245, 323
  dopamine for, 242
  on ECG, 293
  hypermagnesemia, 591
  reflex, 588
  sympatholytic drugs and, 243
Bradykinesia
  with antipsychotic drugs, 573
Bradykinin
  ACE inhibitors and, 610
  C1 esterase inhibitor deficiency, 107
Bradykinin, 610
BRAF gene, **224**, 387
  melanomas and, 484
  papillary thyroid carcinoma and, 343
  vemurafenib and, 444
Brain
  blood flow autoregulation, 297
  embryologic derivatives, 613
  embryology of, 490
  infarcts, 209
  ischemia in, 210
  metastasis to, 223
Brain abscesses
  bacteremia, 180
  HIV-positive adults, 180
  otitis media, 180
  Staphylococcus aureus, 180
  Toxoplasma gondii, 177
  Viridans streptococci, 180
Brain cysts, 161
Brain death, 269, 501, 502
Brain injury
  gastritis with, 379
  hypopituitarism from, 339
Brain lesions (common), **511**
Brain natriuretic peptide (BNP), 296, 588
  in SIADH, 338
  signaling pathways for, 337
Brain stem
  dorsal view, **504**
  ventral view, **504**
Brain stem/cerebellar syndromes
  multiple sclerosis, 523
Brain tumors
  adult primary, **526–527**
  childhood primary, 528
  hallucinations with, 559
  incidence and mortality, 222
  metastatic source, 223
  nitrosoureas for, 441
Branched-chain ketoacid dehydrogenase, 66
Branching enzyme (glycogen metabolism), 86
BRCA1/BRCA2 genes, 224
  DNA repair in, 40
  tumor suppressor genes, 224
Breastfeeding, 636
Breast milk
  IgA antibodies in, 105
  oxytocin's role in, 328
  prolactin and, 330
Breast/ovarian cancer
  BRCA1 mutation, 64
  BRCA2 mutation, 64
  incomplete penetrance, 56
Breast pathology, **649**
  benign disorders, 649
  invasive carcinomas, **650**
  noninvasive carcinomas, **650**
Breast cancer
  aromatase inhibitors for, 656
  breastfeeding and, 636
  hormonal contraception contraindication, 657
  hypercalcemia and, 228
  incidence/mortality of, 222
  oncogenes and, 224
  paclitaxel for, 441
  paraneoplastic cerebellar degeneration and, 228
  tamoxifen for, 443
  trastuzumab for, 443
  tumor suppressor genes and, 224
Breathing
  mechanics of, 675
  with pneumothorax, 682
Breath sounds
  bronchial, 680, 682
  diminished, 682
  physical findings, 680
Brenner tumor, 646
Brief psychotic disorder, 560
Brimonidine, 552
Brittle bone disease, gene defects in, 51
Broad-base budding (blastomycosis), 151
Broad ligament, 625
Broca area, 501
  aphasia, 516
  MCA stroke, 514
Bromocriptine, 548
Bronchi, 662
Bronchial carcinoid tumor, 684
Bronchiectasis
  Aspergillus fumigatus, 153
  cystic fibrosis, 60
  Kartagener syndrome, 49
Bronchioles, 662
  histamine receptors and, 238
Bronchiolitis obliterans, 119, 683
  organizing pneumonia (BOOP), 683
Bronchitis
  cystic fibrosis, 60
  Haemophilus influenzae, 142
Bronchoconstriction, 687
Bronchodilation, 687
  sympathetic receptors and, 238
Bronchogenic carcinomas
  asbestosis and, 677
  carcinogens causing, 225
Bronchogenic cysts, 660
Bronchopneumonia, 683
Bronchopulmonary dysplasia, 210
  free radical injury, 210
  neonatal respiratory distress syndrome, 661
Brown-Séquard syndrome, **531**
  Horner syndrome, 531
"Brown tumors," 464
Brucella spp, 127
  transmission and treatment of, **143**
  zoonotic infections, 149
Brucellosis, 149
Brugada syndrome, **294**, 304
Bruising
  scurvy, 69
Brunner glands
  bicarbonate product, 372
  duodenum, 362
Bruton agammaglobulinemia, 61, **116**
Bruxism, 497
BTK gene, 116
B-type natriuretic peptide, **296**
Buckle (torus fracture), 462
Budd-Chiari syndrome, **392**
Budesonide, 687
Buerger disease, 314
Bugs
  affecting unvaccinated children, **186**
  causing diarrhea, **179**
  causing food-borne illness, **178**
  hints, **186**
Bulbar (spongy) urethra injury, 627
Bulbus cordis, 281
Bulimia nervosa, 567
  anovulation and, 645
  drug therapy for, 572
  laxative abuse by, 401
  Mallory-Weiss syndrome and, 377
  SSRIs for, 575
Bulk-forming laxatives, 401
Bullae, 475
  impetigo, 479
  necrotizing fasciitis, 479
  skin lesions, 475
Bull neck lymphadenopathy, 132
Bullous impetigo, 479
Bullous pemphigoid, 475, 480
  autoantibody, 115
Bulls-eye erythema, 146
Bumetanide, 608
BUN (blood urea nitrogen)
  ornithine transcarbamylase deficiency, 83
Bundled payment, 271
Bundle of His, 293
Bundle of Kent, 294
Bunyaviruses, 167–168
Bupivacaine, 550
Buprenorphine, 551
  heroin detoxification, 576
  morphine and, 234
Bupropion, 576
  seizures with, 251
Burkholderia cepacia
  cystic fibrosis, 179
Burkitt lymphoma, 430
  chromosomal translocations and, 434
  EBV, 165
  oncogenes and, 224
  oncogenic microbes and, 226
Burnout (medical errors), **274**
Burns
  classification, **483**
  shock with, 310
  sunburn, 482
  testosterone/methyltestosterone for, 658
"Burr cells," 414
Bursitis
  prepatellar, 460
Burton line
  lead poisoning, 419
Buspirone
  mechanism and clinical use, **574**
Busulfan, 441
  pulmonary fibrosis and, 251
  toxicity, 444
Butorphanol, 551, 552

C

C1 esterase inhibitor deficiency, 107
C3 deficiency, 107
C5a receptor, 406
C5-C9 deficiencies, 107
CA 15-3/CA27-29 (tumor markers), 226
CA 19-9 (tumor marker), 226, 398
CA 125 (tumor marker), 226
CAAT box, 41
Cachexia, **227**
  TNF-α and, 108
Café-au-lait spots
  McCune-Albright syndrome, 57
Caffeine intoxication and withdrawal, 570
Cahill cycle, 82
Calcarine sulcus
  thalamic relay to, 498
Calciferol (vitamin D), 589
Calcification, 211
  dystrophic, 227
Calcineurin, 120
Calcinosis cutis, 473
Calcitonin, **333**
  medullary thyroid carcinoma production, 343
  tumor marker, 226
Calcitriol, 589
Calcium
  in bone disorders, 464
  calcitonin and, 333
  in cardiac muscle, 292
  in osteomalacia/rickets, 463
  in Paget disease of bone, 463
  Vitamin D and, 337
Calcium carbonate, 399
Calcium channels
  ethosuximide effect on, 544
  glucose and, 334
  Lambert-Eaton myasthenic syndrome, 228
  myocardial action potential, 292
  opioid effect on, 551
  pacemaker action potential, 292
Calcium channel blockers
  angina, 318
  antiarrhythmic drugs, **324**
  contractility in, 284
  cutaneous flushing, 248
  gingival hyperplasia, 250
  hypertension, 316
  hypertrophic cardiomyopathy, 308
  mechanism and clinical use, 318
  migraine headaches, 518
  Raynaud phenomenon, 472
Calcium homeostasis, 333
Calcium (kidney stones), 598
  calcium oxalate nephrolithiasis, 69
Calcium pyrophosphate deposition disease, **467**
Calcium-sensing receptor (CaSR), 355
Calculation
  bioavailability of, 231
  reabsorption and secretion rate, **584**
Calculous cholecystitis, 396
Caliciviruses, 163
  characteristics of, 167

(c) ketabton.com: The Digital Library

California encephalitis, 167
Calluses (dermatology), 475
cAMP (cyclic adenosine monophosphate)
  endocrine hormone messenger, 337
  fructose bisphosphatase-2 and, 76
  heat-labile/heat-stable toxin effects, 132
  hyperparathyroidism, 345
  PTH effect on, 332
  Vibrio cholerae, 146
CAMP factor, 137
Campylobacter spp
  animal transmission, 149
  bloody diarrhea, 179
  jejuni, **145**
  reactive arthritis and, 469
Canagliflozin, 353
Canalicular stage (development), 660
Cancer
  common metastases, 223
  deaths from, 272
  ESR in, 214
  hallmarks of, **221**
  immune evasion in, 221
  mortality of, 222
  pneumoconioses, 676, 677
Cancer drugs
  cell cycle, **438**
  targets, **438**
Cancer epidemiology, **222**
Candesartan, 610
Candida albicans, 153
  HIV-positive adults, 177
  T cell dysfunction, *116*
Candida spp
  amphotericin B, 199
  azoles, 199
  echinocandins, 200
  immunodeficiency infections, 118
  osteomyelitis, 180
  tricuspid valve endocarditis and, 311
  vulvovaginitis, 181
Candidiasis
  Candida albicans, 153
  chronic mucocutaneous, 116
  cortisol and, 336
  nystatin, 199
Cannibalism, 178
"Cannonball" metastases, 642
Capecitabine
  5-F-dUMP, 36
"Cape-like" sensory loss, 492
Capillary fluid exchange, **297**
Capitate bone, 449
Capitation, 271
Caplan syndrome, 466, 677
Capsules (bacterial), 124
Captain's wheel
  Paracoccidioidomycosis, 151
Captopril, 610
Caput medusae, 365
Carbachol, 240, 552
Carbamazepine
  agranulocytosis, 250
  aplastic anemia, 250
  cytochrome P-450 and, 252
  epilepsy, 544
  SIADH and, 249
Carbamoyl phosphate, 82
Carbamoyl phosphate synthetase, 73
Carbapenems
  mechanism and use, **190**
  Pseudomonas aeruginosa, 143
Carbidopa, **549**
Carbohydrate absorption, **373**
Carbohydrate metabolism
  inborn errors of, 80
Carbol fuchsin, 125
Carbon dioxide (CO_2)
  production in tissues, 127
  retention, 679
  transport, **670**
Carbonic anhydrase, 670
Carbon monoxide (CO)
  vs cyanide poisoning, 667
  electron transport inhibition, 78
  poisoning, 666
  teratogenicity, 614
  toxicity treatment, 248
Carbon tetrachloride
  free radical injury and, 210
Carboplatin
  mechanism and clinical use, **442**
  toxicities of, 444
Carboplatin toxicity, 444
Carboxylases, 73
Carboxypeptidase, 373
Carcinoembryonic antigen (CEA) (tumor marker), 226
Carcinogens, **225**
  griseofulvin, 200
Carcinoid syndrome, **352**
  bronchial carcinoid tumors, 684
  somatostatin in treatment, 371
Carcinoid tumors
  biomarkers for, 226
  immunohistochemical stains for, 227
  octreotide for, 400
  stomach, 379
Carcinoma in situ, 219
  cervical dysplasia, 645
  ductal, 650
  neoplastic progression, 219
  penis, 651
  vulvar, 644
Carcinomas
  bone, 464
  colorectal, 388
  invasive, 219
  metastases of, 219, 223
  nomenclature of, 220
  thyroid, 343
Cardiac arrest
  antacid adverse effects, 399
  hypermagnesemia, 591
Cardiac cycle, 287
Cardiac depression, 318
Cardiac function curves, **286**
Cardiac glycosides
  mechanism and clinical use, **321**
  sodium-potassium pump inhibition, 49
Cardiac looping, 280
Cardiac output
  exercise and, 670
  in pregnancy, 633
  variables in, **284**
  V/Q mismatch and, 669
Cardiac output equations, **285**
Cardiac pressures (normal), 297
Cardiac tamponade, **310**
  aortic dissection and, 303
  jugular venous pulse in, 287
  MI, 305, 307
  shock, 310
Cardiac therapy, **317**
Cardiac tumors, **316**
Cardinal veins, 281
Cardiogenic shock, 310
  etiology, 310
Cardiomegaly
  Pompe disease, 87
Cardiomyopathy, **308**
  β-blockers, 245
  Chagas disease, 158
  drug reaction and, 248
  familial amyloid, 212
  glycogen storage diseases, 87
  heart failure with, 309
  hemochromatosis and, 395
  hypertrophic, 245
  Kussmaul sign in, 316
  Starling curves, 285
  sudden cardiac death, 304
Cardiotoxicity
  doxorubicin, 439
  drugs causing, 444
  methylxanthines, 687
  TCA adverse effects, 575
  trastuzumab, 443
Cardiovascular drugs
  naming conventions for, 253
  reactions to, 248
Cardiovascular system, 281–323
  anatomy, 283
  embryology, 281–283
  pathology, 298–313
  pharmacology, 316–322
  physiology, 284–299
  sclerosis of, 473
Carditis
  Lyme disease, 146
  rheumatic fever, 312
Carfilzomib, **443**
Carina (trachea), 663
Carmustine, 441
  pulmonary fibrosis, 251
Carnitine acyltransferase I, 73, 89
Carotid artery
  atherosclerosis in, 302
  cavernous sinus, 542
  emboli from, 538
  embryonic development, 619
  giant cell arteritis and, 314
Carotid massage, 296
Carotid sinus, 296
Carpal bones, 449
Carpal tunnel syndrome, 459
  lunate dislocation, 449
  nerve injury, 447
  rheumatoid arthritis, 466
Carteolol, 552
Cartilage
  collagen in, 50
  fluoroquinolone damage to, 250
Cartilage damage, 204
Carvedilol, 245, 323
Casal necklace, 67
Caseating granulomas
  in tuberculosis, 140
Case-control studies, 256
Caseous necrosis, 209
Caspases, 208
Caspofungin
  echinocandins, 200
Casts in urine, **594**
Catabolism of amino acids, 82
Catalase, 210
Catalase-positive organisms, **128**
Cataplexy, 568
Cataracts, **535**
  corticosteroid toxicity, 120
  diabetes mellitus and, 346
  galactosemia, 80
  muscular dystrophy, 61
  rubella, 182
  sorbitol, 81
Catecholamines
  adrenal medulla secretion, 327
  amphetamines and, 242
  contractility effects of, 284
  ephedrine and, 242
  heart contractility, 284
  pacemaker action potential, 292
  pheochromocytoma and, 350
Catecholamine synthesis
  tyrosine catabolism, **83**
Cats (disease vectors)
  Campylobacter jejuni, 145
  Cat scratch disease, 149
  Pasteurella multocida, 149, 186
  Tinea corporis, 152
  Toxoplasma gondii, 156, 182
Cauda equina, 507
Cauda equina syndrome, 530
Caudal fold closure defects, 358
Caudal regression syndrome, 614
Caudate
  basal ganglia, 500
  Huntington disease, 520
Cavernous hemangiomas
  liver, 392
Cavernous hemangiomas (liver), 392
Cavernous sinus, **542**
  dural venous sinuses, 503
  syndrome, 542
  thrombosis with mucormycosis, 153
CCR5 protein
  HIV and, 175
  maraviroc, 203
  viral receptor, 166
CD4+ cell count
  disease associations by levels, 177
CD4 protein, 101
  viral receptor, 166
CD4+ T cells (HIV), 176
CD5 protein, 432
CD8 protein, 101
CD16 protein, 101
CD20 protein, 110
  in CLL, 432
CD21 protein, 110
  viral receptor, 166
CD25 protein
  cell surface protein, 110
  regulatory T cells and, 102
CD28 protein, 110
CD34 protein, 110
  leukocyte extravasation and, 215
CD40 protein, 110
CDKN2A gene, 224
CEA tumor marker, 388
Cefaclor, 189
Cefazolin
  mechanism and use, 189
  prophylaxis, 198
Cefepime
  mechanism and use, 189
  Pseudomonas aeruginosa, 143
Cefotaxime, 189
Cefotetan
  mechanism and use, 189
Cefoxitin
  mechanism and use, 189
Cefpodoxime
  mechanism and use, 189
Ceftaroline
  mechanism and use, 189
  MRSA, 198

(c) ketabton.com: The Digital Library

Ceftazidime
  mechanism and use, 189
  Pseudomonas aeruginosa, 143
Ceftriaxone
  Chlamydia spp, 148
  for gonococci, 142
  for *Haemophilus influenzae*, 142
  mechanism and use, 189
  meningitis, 180
  meningococci, 142
  typhoid fever, 144
Cefuroxime
  mechanism and use, 189
Celecoxib, 252, **486**
Celiac artery
  mesenteric ischemia, 386
  structures supplied, 364
Celiac disease, 381
  autoantibody, 115
  HLA genes and, 100
  IgA deficiency, 116
Celiac sprue, 381
Celiac trunk, **364**
Cell biology of bone, **459**
Cell cycle phases, **46**
Cell envelope (bacterial), **124**
Cell injury, **207**
Cell lysis, 590
Cell membrane in apoptosis, 208
Cell surface proteins
  association and functions, **110**
  leukocyte adhesion deficiency, 117
  T cells and, 101
Cell trafficking, **47**
Cell types
  labile, 46
  permanent, 46
  stable (quiescent), 46
Cellular biochemistry, 46–52
Cellular injury
  cellular adaptations, **206**
  irreversible, 207
  reversible, 207
Cellulitis, 136, 479
  Pasteurella multocida, 149
Cell walls (bacterial), 124
Central clearing
  nuclei, 343
  rash, 152
Central diabetes insipidus, 338
Central/downward transtentorial herniation, 529
Central nervous system (CNS)
  anesthetic principles for, 549
  antiarrhythmic adverse effects, 322, 323
  cancer epidemiology, 222
  cell types in, 493–494
  depression, 546
  drug name conventions, 253
  nitrosoureas effect on, 441
  origins of, **490**
  posterior fossa malformations, 491, 492
  regional specification of, 506
  shock from injury, 310
Central nervous system stimulants, **572**
Central pontine myelinolysis, 524
Central post-stroke pain syndrome, **515**
Central precocious puberty, 637
Central retinal artery occlusion, **538**
Central sleep apnea, 679
Central sulcus, 501
Central tendency measures, 262
Central tendon (diaphragm), 663
Central vertigo, 534
Centriacinar emphysema, 674
Cephalexin
  mechanism and use, 189
Cephalosporins
  disulfiram-like reaction, 251
  mechanism and use, **189**
  pseudomembranous colitis, 249
  Pseudomonas aeruginosa, 143
Ceramide trihexoside
  in sphingolipidoses, 88
Cerebellar degeneration
  paraneoplastic, 228
Cerebellar lesions
  hemisphere, 511
  lateral, 499
  medial, 499
  tonsillar herniation, 492, 529
  vermis lesions, 511
Cerebellum
  development of, 490
  input/output of, **499**
  tonsils, 492
Cerebral aqueduct of Sylvius, 504
Cerebral artery distributions, **502**
Cerebral cortex
  aphasia, 514
  arterial distribution, **502**
  dominant parietal lesions, 511
  functional areas of, **501**
  hemineglect, 514
  nondominant parietal lesions, 511
  visual field defects, 514
Cerebral edema
  diabetic ketoacidosis and, 347
  therapeutic hyperventilation, 501
Cerebral hemispheres, 490
Cerebral palsy, 551
Cerebral perfusion pressure (CPP), **501**
"Cerebriform" nuclei, 430
Cerebrospinal fluid (CSF)
  albuminocytologic dissociation, 524
  blood-brain barrier and, 496
  circulation of, 496, 503
  findings in meningitis, **180**
  Guillain-Barré syndrome, 524
  hydrocephalus, 522
  multiple sclerosis, 523
  neurodegenerative disorders, 521
  origins, 490
  poliomyelitis, 531
  production, 493
  ventricular system, 504
Cerebrovascular disease
  diabetes mellitus, 346
Certolizumab, 487
Ceruloplasmin
  free radical elimination by, 210
Cervical cancer, **645**
  carcinogens causing, 225
  epidemiology of, 643
  HIV-positive adults, 177
  hydronephrosis with, 599
  oncogenic microbes and, 226
  papillomaviruses, 164
Cervical rib, 448
Cervicitis
  sexually transmitted infections, 184
Cervix
  anatomy of, 625
  epithelial histology, 626
  lymphatic drainage of, 624
  pathology of, 645
Cestodes, **160**
Cetirizine, 686
Cetuximab, **442**
CFTR gene
  chronic pancreatitis and, 397
cGMP (cyclic guanosine monophosphate)
  atrial natriuretic peptide and, 296
  endocrine hormone messenger, 337
  male sexual response, 627
Chagas disease, 158
  achalasia in, 376
Chalk-stick fractures, 463
Chancroids, 184
Changes in the elderly, **270**
Chaperone protein, **45**
Charcoal yeast extract culture
  Legionella pneumophila, 126, 143
Charcot-Bouchard microaneurysm, 516
Charcot joints
  syphilis, 147
  tabes dorsalis and, 530
Charcot-Leyden crystals, 674
Charcot-Marie-Tooth disease, 524
Charcot triad, 397
Charging, tRNA, 44
Chédiak-Higashi syndrome, 117
Cheilosis, 67
Chelation
  hemochromatosis, 395
  lead poisoning, 419
Chemokines, 108
  delayed hypersensitivity, 112
Chemoreceptors, **296**
Chemoreceptor trigger zone (CTZ), 496
Chemotherapy
  AML and, 432
  MDR1 and responsiveness to, 227
  neutropenia with, 424
  ondansetron, 400
  paclitaxel, 48
  pancreatic cancer, 398
  readmissions with, 272
  treatments for vomiting, 401
  vincristine/vinblastine, 48
Chemotoxicities, 444
Cherry hemangiomas, 478
"Cherry red" epiglottis, 142
Cherry-red spot (macula), 538
  lysosomal storage disease, 88
Chest pain
  panic disorder, 563
  pneumothorax, 682
Chest wall
  elastic properties, 665
Chest X-rays
  aortic dissections on, 303
  balloon heart on, 308
  eggshell calcification, 677
  notched ribs on, 299
  Wegener granulomatosis on, 315
  widened mediastinum on, 137
Cheyne-Stokes respirations, 679
Chiari malformations, 492
Chickenpox
  rash, 183
  VZV, 165
Chief cells (parathyroid), 332
Chief cells (stomach), 372
Child abuse, **556**
  osteogenesis imperfecta and, 51
  reporting requirements, 269
Childbirth
  brachial plexus injury in, 448
  Budd-Chiari syndrome and, 392
  Graves disease and, 342
  low birth weight, 635
  misoprostol induction, 399
  neonatal flora, 178
  oxytocin and uterine contractions, 328, 636
  postpartum mood disturbances, 562
  preterm, as common cause of death, 272
  progesterone levels after, 630
  Sheehan syndrome after, 339
  stress incontinence and, 599
Childhood disorders, **557**
Childhood musculoskeletal conditions, **461**
Childhood primary brain tumors, **528**
Child neglect, **556**
Children
  causes of death, 272
Chi-square tests, 264
Chlamydia spp, **148**, 184
  atypical infections, 179
  diagnostic tests for, 148
  Giemsa stain, 125
  Gram stain, 125
  intracellular organism, 127
  macrolides, 193
  pneumonia, 683
  reactive arthritis, 469
  sulfonamides for, 194
  tetracyclines, 192
Chlamydia trachomatis
  eosinophilia, 149
  pelvic inflammatory disease, 149
  pneumonia, 179
  serotypes, **149**
  UTIs, 600
Chlamydophila pneumoniae, 148
  pneumonia, 179
Chlamydophila psittaci
  atypical pneumonia, 148
  transmission, 149
Chloasma (melasma), 476
Chloramphenicol
  aplastic anemia and, 250
  gray baby syndrome, 250
  mechanism and use, **192**
  pregnancy contraindications, 204
  protein synthesis inhibition, 191
Chlordiazepoxide, 546
  alcohol withdrawal, 572
Chlorhexidine, 204
Chloride channels
  cystic fibrosis, 60
Chlorine, 204
Chloroprocaine, 550
Chloroquine, **200**
  malaria, 157
Chlorpheniramine, 686
Chlorpromazine, 573
Chlorthalidone, 609
Chocolate agar
  Haemophilus influenzae, 126, 142
Cholangiocarcinomas
  Clonorchis sinensis, 160, 161
  hyperbilirubinemia, 393
  oncogenic microbes and, 226
  sclerosing cholangitis, 395
Cholangitis, 368, 382, 393, 397
Cholecalciferol, 70
Cholecystectomy, 396, 397
Cholecystitis, 396

(c) ketabton.com: The Digital Library

Cholecystokinin (CCK)
  secretory cell location, 373
Choledocholithiasis, 396
Cholelithiasis, 395, **396**, 397
  acute pancreatitis, 397
  bile ducts and, 368
  biliary cirrhosis and, 396
  Crohn disease, 382
  hyperbilirubinemia and, 393
  octreotide and, 400
Cholera toxin
  lysogenic phage infection, 130
  mechanism, 132
Cholestasis serum markers, 390
Cholesteatomas, **533**
Cholesterol
  atherosclerosis, 302
  in bile, 374
  cholelithiasis and, 396
  lipid-lowering agents, 320
  synthesis of, 47, 73, 79
  vitamin B_3 effects, 67
Cholesteryl ester transfer protein, 93
Cholestyramine, 320
Cholinergic agonists, 253
Cholinergic effects, 321
  cardiac glycosides, 321
Cholinesterase inhibitors
  diarrhea with, 249
  neuromuscular blockade reversal, 551
Cholinomimetic agents, **240**
  glaucoma treatment, 552
Chondrocalcinosis, 467
Chondrocytes, 458, 462
  osteoarthritis, 466
Chondroma, 464
Chondrosarcoma, 465
Chordae rupture, 291
Chorea
  brain lesions, 511
  Huntington disease, 520
  movement disorders, 519
Choriocarcinoma, **642**
  hCG in, 633
  methotrexate for, 440
  testicular, 653
  theca-lutein cysts and, 646
Chorionic plate, 617
Chorionic somatomammotropin, 634
Chorionic villi
  hydatidiform moles, 642
  placenta, 617
Chorioretinitis
  congenital toxoplasmosis, 182
Choristomas, 220
Choroid layer (ophthalmology)
  inflammation, 536
  neovascularization, 536
Choroid plexus (CNS), 504
Christmas tree distribution, 482
Chromaffin cells
  diagram, 327
  pheochromocytomas, 350
Chromatin structure, **34**
Chromatolysis, **495**
Chromogranin, 226, 684
Chromosomal translocations, **434**
Chromosome abnormalities
  congenital microdeletion, 64
  gene associations with, 64
  hemochromatosis, 395
  karyotyping for, 55
  nephroblastoma, 606
  nondisjunction (meiosis), 63
  omphaloceles, 358
  polyposis syndrome, 387
  renal cell carcinoma, 605
  Robertsonian translocation, 64
  von Hippel-Lindau disease, 525
  Wilson disease, 395
Chronic bronchitis, 674
Chronic disease, anemia of, 421
Chronic gastritis, 379
Chronic granulomatous disease (CGD)
  catalase-positive microbes, 186
  immunodeficiencies and, 117
  recombinant cytokines for, 121
  respiratory burst in, 109
Chronic inflammation, **216**
Chronic ischemic heart disease, 304
Chronic kidney disease
  erythropoietin in, 589
  hypertension and, 300
Chronic lymphocytic leukemia (CLL), 432
  immunosuppressants, 120
  rituximab for, 443
Chronic mesenteric ischemia, 386
Chronic mucocutaneous candidiasis, 116
Chronic myelogenous leukemia (CML), 433
  basophilia caused by, 408
  busulfan for, 441
  chromosomal translocations and, 434
  imatinib for, 443
  oncogenes and, 224
Chronic myeloproliferative disorders, **433**
Chronic obstructive pulmonary disease (COPD)
  albuterol for, 242
  β-blockers and, 245
  muscarinic antagonists for, 241
  salmeterol for, 242
Chronic pancreatitis, **397**
  pancreatic insufficiency from, 381
Chronic placental insufficiency, 578
Chronic pyelonephritis, 600
Chronic renal failure, 603
  hyperphosphatemia with, 345
Chronic respiratory diseases
  bronchitis, 674
  with chronic inflammatory diseases, 683
  death in children, 272
  obstructive diseases, 674
  pneumoconioses, 675
  sinusitis, 674
  thromboembolism, 679
Chronic thromboembolic pulmonary hypertension, 679
Chronic transplant rejection, 119
Churg-Strauss syndrome, 315
  autoantibody, 115
Chvostek sign
  hypocalcemia, 591
  hypoparathyroidism, 344
Chylomicrons, 92, 94
Chymotrypsin, 373
Cidofovir, **202**
Cigarette smoke (carcinogen), 225
Ciguatoxin, 247
Cilastatin
  imipenem and, 190
  seizures with, 251
Ciliary ganglia, 539
Cilia structure, **49**
Ciliated cells, 662
Cimetidine, 399
  cytochrome P-450 and, 252
  gynecomastia and, 649
Cinacalcet, **355**
Cinchonism
  antiarrhythmic causing, 322
  neurologic drug reaction, 251
Cingulate gyrus
  limbic system, 499
Cingulate (subfalcine) herniation, 529
Ciprofloxacin
  for Crohn disease, 382
  cytochrome P-450 and, 252
  fluoroquinolones, 195
  meningococci, 142
  Mycobacterium avium-intracellulare, 196
  prophylaxis, 198
  Pseudomonas aeruginosa, 143
Circadian rhythm
  hypothalamic control, 498
  sleep physiology, 497
Circle of Willis, **503**
  saccular aneurysms, 516
Circulatory system
  fetal, 282
  kidneys and, 580
Circumflex femoral artery, 463
Circumoral pallor
  group A streptococcal pharyngitis, 136
Cirrhosis
  α_1-antitrypsin deficiency, 392
  alcoholic, 71, 391
  bacterial peritonitis (spontaneous), 390
  cholelithiasis and, 396
  cystic fibrosis, 60
  encephalopathy with, 391
  esophageal varices and, 377
  fructose intolerance, 80
  gynecomastia, 649
  hemochromatosis, 395
  with hepatitis, 389
  hepatocellular carcinomas, 392
  hyperbilirubinemia in, 393
  non-alcoholic fatty liver disease, 391
  portal hypertension, **389**
  serum markers for, 390
Cisplatin
  mechanism and clinical use, **442**
  targets of, 438
  toxicities of, 251, 444
Citalopram, 575
Citrate synthase, 74
Citrulline, 82
c-KIT gene, 224
CK-MB, 304, 306
Cladribine, **440**
  for hairy cell leukemia, 432
Clara cells, 661, 662
Clarithromycin
  Helicobacter pylori, 146
  HIV prophylaxis, 198
  macrolides, 193
  Mycobacterium avium-intracellulare, 196
  pregnancy use, 204
Clasp knife spasticity, 529
Class IA antiarrhythmics, 322
Class IB antiarrhythmics, 322
Class IC antiarrhythmics, 322
Class II antiarrhythmics, 323
Class III antiarrhythmics, 323
Class IV antiarrhythmics, 324
Classical conditioning, **554**
Class switching
  CD40, 103
  thymus-dependent antigens, 105
Clathrin, 47
Claudication
  atherosclerosis, 302
  Buerger disease, 314
  giant cell arteritis, 314
Clavicle fractures, **460**
Clavulanate
  Haemophilus influenzae, 142
Clavulanic acid, 189
Clawing (hand), 451
  Klumpke palsy, **448**
Clearance (CL) of drugs, 231
Clear cell adenocarcinoma, 644
  DES and, 656
Cleavage in collagen synthesis, 50
Cleft lip and palate
  development, **621**
  Patau syndrome, 63
  Pierre Robin sequence, 620
Clevidipine, 318
  for hypertensive emergency, 318
Clindamycin
  bacterial vaginosis, 148
  Clostridium difficile and, 138
  mechanism and use, **192**
  metronidazole vs, 192
  protein synthesis inhibition, 191
  pseudomembranous colitis with, 249
Clinical reflexes, **510**
Clinical trials, **256**
Clinical vignette strategies, 23
"Clock-face" chromatin, 409, 431
Clofazimine
  Hansen disease, 141
  Mycobacterium leprae, 196
Clomiphene
  estrogen receptor modulators, 656
  hot flashes with, 249
Clomipramine, 575
Clonidine, 240, 243
Cloning methods (laboratory technique), 55
Clonorchis sinensis
  cholangiocarcinoma, 226
  diseases association, **161**
  trematodes, 160
Clopidogrel, 437
  acute coronary syndromes, 307
  for ischemic stroke, 512
  thrombogenesis and, 411
Closed-angle glaucoma, 536
  pilocarpine for, 240
Clostridium botulinum
  exotoxin production, 132
  food poisoning, 178
  therapeutic uses, 138
Clostridium difficile
  antibiotic use, 185
  metronidazole, 195
  nosocomial infection, 185
  PPI association, 138
  proton pump inhibitor use, 399
  toxins and effects of, 138
  vancomycin, 190
  watery diarrhea, 179
Clostridium perfringens
  clindamycin, 192
  exotoxin production, 133

(c) ketabton.com: The Digital Library

food poisoning, 178
toxins produced, 138
traumatic open wound, 186
watery diarrhea, 179
Clostridium spp, **138**
anaerobic organism, 127
exotoxins, 138
Clostridium tetani, 138
exotoxin production, 132
Clotrimazole, 199
Clotting factors, 71
Clozapine, 573
agranulocytosis with, 250
Clubbing (digital), 675
cystic fibrosis, 60
Eisenmenger syndrome, 299
Club cells, **661**
Clue cells
bacterial vaginosis, 148, 181
Cluster A personality disorders
characteristics of, **565**
schizoid, 565
schizotypal, 565
Cluster B personality disorders
antisocial, **565**
borderline, 565
histrionic, 565
narcissistic, 565
Cluster C personality disorders
avoidant, **566**
dependent, 566
obsessive-compulsive, 566
Cluster headaches, 518, 547
c-MYC gene, 224
CN III, IV, VI palsies, **541**
CNS (central nervous system)
cancer epidemiology, 222
CNS lymphomas
HIV-positive adults, 177
oncogenic microbes and, 226
CO_2
production in tissues, 127
Coagulation, 71
Coagulation disorders, **426**
hemophilia, 426
hereditary thrombosis syndromes, 428
mixed platelet/coagulation, 428
vitamin K and, 426
Coagulation pathways, **412**
Coagulative necrosis, 209
MI, 305
Coagulopathy
postpartum hemorrhage, 641
preeclampsia, 643
uterine bleeding with, 633
Coal workers' pneumoconiosis, 677
CoA production, 67, 72
Coarctation of aorta, 299, 300
Cobalamin, 69
Cobblestone mucosa, 382
Cocaine
β-blockers and, 245
coronary vasospasm, 248
intoxication and withdrawal, 571
liver processing of, 367
sympathomimetic action, 242
teratogenicity, 614
Coccidioides spp
silver stain, 125
treatment, 199
Coccidioidomycosis, 151
erythema nodosum and, 482
Coccobacilli, 141
Cocci bacteria
antibiotic tests, 134
Cochlea
CN VIII, 506
inner ear, 533
presbycusis, 533
Codeine, 551
Codman triangle (x-ray), 465
Codominance, 56
Codons
amino acid specification, 37
genetic code features, 37
start and stop, 44
Cofactors
apolipoproteins, 93
biotin, 68, 73
cobalamin, 69
copper, 51
Menkes disease, 51
pantothenic acid, 67
phenylketonuria, 84
pyridoxine, 67
pyruvate dehydrogenase complex, 76
riboflavin, 67
TCA cycle, 77
thiamine, 74
vitamin K, 71
"Coffee bean sign" (X-ray), 386
Cognitive behavioral therapy (CBT), 557, 572
ADHD, 557
anxiety disorders, 562
atypical depression, 561
bipolar disorder, 561
body dysmorphic disorder, 563
obsessive-compulsive disorder, 563
panic disorder, 563
phobias, 563
postpartum depression, 562
PTSD, 563
Cohort studies, 256
Coin lesion (X-ray), 684
x-ray signs, 684
Cola-colored urine, 596
Colchicine, 55
agranulocytosis, 250
calcium pyrophosphate deposition disease, 467
diarrhea with, 249
gout, 467, 487
microtubules and, 48
myopathy with, 250
"Cold enrichment," 139
Colectomy
adenomatous polyposis, 387
inflammatory bowel disease, 382
Colesevelam, 320
Colestipol, 320
Colistin
polymyxin E, 193
Pseudomonas aeruginosa, 143
Colitis
Clostridium difficile, 138
oral vancomycin, 190
pseudomembranous, **179**, 188, 192
Collagen, **50**
decreased/faulty production, 51
epithelial cell junctions and, 474
osteoblast secretion of, 459
polyostotic fibrous dysplasia and, 57
scar formation, 218
synthesis and structure, **50**
in systemic sclerosis, 473
types of, 50
vitamin C in synthesis, 69
wound healing, 216
Collagenase in neutrophils, 406
Collapsing pressure (alveoli), 661
Collecting tubules
potassium-sparing diuretics and, 609
Colles fracture, 462
Colliculi, 504
Colon
histology of, 362
ischemia in, 210
Colon cancer, **388**
adenomatous polyposis and, 387
bevacizumab for, 442
cetuximab for, 442
5-fluorouracil for, 440
incidence/mortality in, 222
irinotecan/topotecan for, 442
Lynch syndrome, 40
molecular pathogenesis of, 389
oncogenes and, 224
serrated polyps and, 387
Staphylococcus gallolyticus and, 137
tumor suppressor genes and, 224
Colonic ischemia, 363, 386
Colonic polyps, **387**
Colony stimulating factor, 121
Colorado tick fever, 167
Color blindness, 197
Colovesical fistulas, 383
Coltivirus, 167
Coma
hepatic encephalopathy, 391
herniation syndromes, 529
hyponatremia, 591
rabies, 171
thyroid storm, 342
Toxocara canis, 159
Trypanosoma brucei, 156
Comedones, 477
Commaless genetic code, 37
Comma-shaped rods, 141
Common bile duct, 361, 368
Common cold, 168
Common peroneal nerve, 452, 453
Common variable immunodeficiency (CVID), 116
Communicating hydrocephalus, 522
Communication with patient, 268
Compartment syndrome, 461
Competence (bacterial genetics), 130
Competitive agonists, 234
Competitive antagonist, 234
Competitive inhibitors, 230
Complement
activation pathways and functions, **106**
binding of, 104
disorders of, **107**–122
eculizumab, 122
endotoxin activation, 133
immunodeficiency infections, 118
immunoglobulin isotypes, 105
innate immunity, 99
splenic dysfunction, 98
transplant rejection, 119
Complement activation inhibition
β-hemolytic bacteria, 135
Complementation (viral), 162
Complete (third-degree) AV block, 295
Complex partial seizures, 517
Complex renal cysts, 604
Compliance (lung), 665
Comprehensive Basic Science Examination (CBSE), 11
Comprehensive Basic Science Self-Assessment (CBSSA), 11
Computer-Based Test (CBT)
environment of, 3–4
exam schedule for, 7–8
structure of, 3
COMT inhibitors, 548
Conditioning (psychological), 554
Conduct disorder, 557
Conducting zone (respiratory tree), 662
Conduction aphasia, 516
Conductive hearing loss, 533
Condylomata acuminata, 477
sexual transmission, 184
Condylomata lata
syphilis, 147, 184
Confidence intervals, **263**
Confidentiality, **267**
in abuse, 269
behavioral science ethics, 265
exceptions to, 267
Confluence of the sinuses, 503
Confounding bias, 261
Congenital adrenal hyperplasias, **335**
Congenital heart disease, **298**–300
defect associations, **300**
maternal phenylketonuria, 84
pulmonary arterial hypertension, 679
rubella, 182
Congenital hydrocele (scrotal), 652
Congenital hypothyroidism (cretinism), 341
Congenital long QT syndrome, 294
Congenital lung malformations, **660**
Congenital malformation mortality, 272
Congenital nevus, 475
Congenital rubella
cardiac defect associations, 300
heart murmur, 291
Congenital solitary functioning kidney, **579**
Congenital syphilis, 147
Congestion (respiratory)
with lobar pneumonia, 683
Congo red stain, 212
Conivaptan, 354
SIADH, 354
Conjoined tendon, 369
Conjugated (direct) hyperbilirubinemia, 393
Conjugate vaccines, 127
Conjugation (bacterial genetics), 130
Conjunctival suffusion/injection
eye disorders, 147
Kawasaki disease, 314
Conjunctivitis, 148, **534**
adenovirus, 164
chlamydia, 184
gonococcal prophylaxis, 198
Haemophilus influenzae, 142
reactive arthritis, 469
rubeola, 170, 183, 186
Zika virus, 171
Connective tissue diseases
aortic dissection and, 303
pulmonary arterial hypertension, 679
thoracic aortic aneurysms and, 302
Connective tissue drug reactions, 250
Conn syndrome, 349, 591
Consensual light reflex, 539
Consent
healthcare proxy, 269
minors, **265**, 268

(c) ketabton.com: The Digital Library

Consolidation (lung finding), 680
  lobar pneumonia, 683
Constipation
  anal fissures, 366
  calcium channel blockers, 318
  Hirschsprung disease, 384
  irritable bowel syndrome, 383
  laxative treatments, 401
  loperamide, 400
  ondansetron, 400
  ranolazine, 319
  vincristine, 441
Constrictive pericarditis
  jugular venous pulse in, 287
  Kussmaul sign, 316
Contact dermatitis, 113
Contemplation stage, substance addiction, 568
Continuous heart murmurs, 291
Contraception
  isotretinoin teratogenicity, 614
  methods for, **657**
  parental consent for minors and, 265
  progestins for, 657
Contractility in cardiac output, 284
Contraction alkalosis, 60, 587, 588, 609
Coombs-positive hemolysis
  α-methyldopa, 243
  anemia with, 423
Coombs test, 423
Cooperative kinetics, 230
COPI/COPII proteins, 47
Copper deficiency, 419
Copper intrauterine device, **657**
Copper metabolism
  Wilson disease, 395
Copper toxicity, 248
Coprolalia, 557
Copy number variations (CNV), 54
Cord factor, 140
Cori cycle, 82
Cori disease, 87
Corkscrew fibers, 528
"Corkscrew" hair, 69
Cornea
  astigmatism, 535
  collagen in, 50
Corneal arcus
  familial hypercholesterolemia, 94
  hyperlipidemia, 301
Corneal reflex, 507
Corneal vascularization, 67
Corniculate cartilage, 620
Coronary arteries
  anatomy of, **283**
  atherosclerosis in, 302
Coronary artery disease
  atrial fibrillation and, 295
  diabetes mellitus and, 346
  hormonal contraception with, 657
  hypertension and, 300
  sudden cardiac death, 304
Coronary sinus
  anomalous pulmonary return, 298
  development, 281
Coronary steal syndrome, 304
Coronary vasospasm, 248
  triptans and, 547
Coronaviruses
  characteristics of, 167
  genomes of, 163
Cor pulmonale, 309, 668
  from obstructive lung disease, 674
  pneumoconioses, **677**
  pulmonary hypertension, 679
  right ventricular failure, 668
Corpus cavernosum
  lymphatic drainage of, 624
Corpus luteum
  hCG and, 633
  progesterone production, 630
Correlation coefficient, 264
Cortical signs, 514
Corticopapillary osmotic gradient, 588
Corticosteroid-binding globulin, 336
Corticosteroids
  asthma, 687
  cataracts, 535
  Crohn disease, 382
  Cushing syndrome, 348
  giant cell arteritis, 314
  hyperglycemia with, 249
  hypopituitarism, 339
  lymphopenia with, 424
  microscopic polyangiitis, 315
  neutrophilia from, 424
  osteonecrosis, 463
  osteoporosis with, 250
  pancreatitis with, 249
  for polymyalgia rheumatica, 470
  Takayasu arteritis, 314
  thyroid storm, 342
  Wegener granulomatosis, 315
Corticotropin-releasing hormone (CRH)
  adrenal cortex regulation of, 327
  cortisol regulation, 336
  function of, 328
  signaling pathways of, 337
Cortisol, **336**
  congenital adrenal hyperplasias, 335
  in Cushing syndrome, 348
  signaling pathways for, 337
Corynebacterium diphtheriae
  culture requirements for, 126
  exotoxin effects, **139**
  exotoxin production, 132
Costovertebral angle tenderness, 601
Co-transporter 2 (SGLT2) inhibitors, 353
Cough
  ACE inhibitors, 251, 610
  chronic bronchitis, 674
  gastroesophageal reflux disease, 377
  guaifenesin, 686
  hypersensitivity pneumonitis, 675
  lung cancer, 684
  nonproductive, 140
  staccato, 149
  Wegener granulomatosis, 315
  whooping, 132, 143
Councilman bodies
  yellow fever, 168
Countertransference, 554
Courvoisier sign
  pancreatic cancer, 398
Cowpox, 164
Coxiella burnetii
  animal transmission, 149
  Q fever, 150
Coxiella spp
  intracellular organisms, 127
Coxsackievirus
  acute pericarditis, 313
  picornavirus, 168
  presentation, 167
  rashes of childhood, **183**
C-peptide
  insulin and, 334
  with insulinomas, 351
CpG island methylator phenotype (CIMP), 387
Crackles (physical findings), 680
Cranial nerve palsies
  osteopetrosis, 463
Cranial nerves, **506**
  common lesions, **532**
  functions of, 506
  locations of, 504
  nerve and vessel pathways, **505**
  nuclei of, **505**
  pharyngeal arch derivation, 620
  reflexes of, **507**
Craniopharyngioma, 528, 613
  hypopituitarism with, 339
Craniotabes, 463
C-reactive protein (CRP), 213
  innate immunity and, 99
Creatine kinase, 203
Creatinine
  ACE inhibitor effects, 610
  glomerular filtration rate and, 582
  proximal convoluted tubules, 587
Creatinine clearance, 582
Cre-lox system, 56
Cremasteric muscle and fascia
  inguinal canal and, 369
Cremasteric reflex, 452, 510
Crepitus
  esophageal perforation, 377
  in necrotizing fasciitis, 479
  soft tissue, 138
Crepitus in necrotizing fasciitis, 479
Crescentic glomerulonephritis, 596
CREST syndrome, 473
  autoantibody, 115
  sclerodermal esophageal dysmotility, 377
Creutzfeldt-Jakob disease, 178, 521
"Crew cut" (skull X-ray), 422
Cricoid cartilage, 620
Cricothyroid muscle, 620
Cri-du-chat syndrome, **64**
Crigler-Najjar syndrome, 393, 394
Crimean-Congo hemorrhagic fever, 167
CRISPR/Cas9, **53**
Crohn disease, 382
  azathioprine, 120
  B_{12} deficiency, 420
  cholelithiasis and, 396
  natalizumab, 122
  spondyloarthritis and, 469
  sulfasalazine for, 400
  vitamin B_{12} deficiency, 69
Cromolyn, 687
Cross-linking in collagen synthesis, 50
Cross-over study, **256**, 261
Cross-sectional studies, **256**
Croup, **170**
  paramyxoviruses, 167, 169
  pulsus paradoxus in, 310
CRP and ESR, 214
Crust (skin), 475
  basal cell carcinoma, 484
  impetigo, 479
  varicella zoster virus, 479
Cryoprecipitate, 429
Crypt hyperplasia, 381
Cryptococcal meningitis, 199
Cryptococcosis, 153
Cryptococcus neoformans, 153
  HIV-positive adults, 177
  stains for, 125
Cryptococcus spp
  meningitis, 180
  treatment, 199
  urease-positive, 127
Cryptogenic organizing pneumonia, 683
Cryptorchidism, **651**
  Sertoli cells and, 628
Cryptosporidium spp, 155
  fluorescent antibody stain, 125
  watery diarrhea, 179
Crypts of Lieberkühn, 362
C-section deliveries
  neonatal flora, 178
  neonatal flora with, 178
  risk factors after, 640
Culture requirements
  bacteria, 126
Cuneiform cartilage, 620
Curling ulcers
  gastritis, 379
"Currant jelly" stools, 385, 386
"Currant jelly" sputum
  Klebsiella spp, 186
Curschmann spirals, 674
Cushing disease, 348
Cushing-like symptoms, 203
Cushing reflex, 296
Cushing syndrome, **348**
  anovulation with, 645
  corticosteroids, 120
  eosinopenia, 424
  paraneoplastic syndrome, 228
  small cell lung cancer, 684
Cushing ulcers
  gastritis, 379
Cutaneous anthrax, 137
  edema toxin, 132
Cutaneous flushing
  drugs causing, 248
Cutaneous larva migrans, 159
Cutaneous leishmaniasis, 158
Cutaneous mycoses, **152**
Cutaneous paraneoplastic syndromes, 228
Cutaneous small-vessel vasculitis, 314
Cutis aplasia
  Patau syndrome, 63
CXCR4
  viral receptor, 166
CXCR4/CCR5 protein
  presence on cells, 110
Cyanide
  electron transport chain, 78
Cyanide poisoning
  vs carbon monoxide poisoning, **667**
  induced methemoglobinemia, 666
  nitroprusside, 318
  treatment for, 248
Cyanosis
  "blue babies," 298
  "blue kids," 299
  Eisenmenger syndrome, 299
  esophageal atresia, 359
  methemoglobinemia, 666
  patent ductus arteriosus, 299
  pulmonary hypertension, 679
  tetralogy of Fallot, 298
Cyclins, 46
Cyclobenzaprine, 551
Cyclooxygenase
  aspirin effect on, 411

(c) ketabton.com: The Digital Library

Cyclooxygenase inhibition
  irreversible, 486
  reversible, 485, 486
  selective, 486
Cyclophosphamide, 441
  hemorrhagic cystitis with, 249
  microscopic polyangiitis, 315
  SIADH with, 249
  toxicities of, 444
  transitional cell carcinoma and, 606
  Wegener granulomatosis, 315
Cycloplegia
  atropine, 241
  muscarinic antagonists for, 241
Cyclosporine
  gingival hyperplasia, 250
  gout, 250
  immunosuppression, 120
Cyclothymic disorder, 561
Cystathionine, 67
Cystathionine synthase deficiency, 84
Cyst disorders (renal), 604
Cysteine, 85
Cystic duct, 368
Cystic fibrosis, **60**
  Aspergillus fumigatus, 153
  bronchiectasis, 675
  chromosome association, 64
  common organisms, 179
  meconium ileus and, 386
  N-acetylcysteine, 686
  pancreatic insufficiency, 381
  vitamin deficiencies and, 65
Cystine, 598
Cystinuria, **85**
Cystitis
  acute bacterial, 594, 600
  squamous cell carcinoma risk, 606
Cytarabine, 440
Cytochrome C, 208
Cytochrome P-450
  azoles, 199
  barbiturates and, 546
  cimetidine and, 399
  griseofulvin, 200
  interactions with, **252**
  macrolides, 193
  phenobarbital effect on, 544
  rifamycins, 196
  ritonavir, 203
Cytokeratin, 227
Cytokines, 101, 108
  corticosteroids and, 120
  Graves disease and, 342
  rejection reactions, 119
  type IV hypersensitivity, 113
Cytokinesis, 46
Cytomegalovirus (CMV)
  AIDS retinitis, 165
  cholecystitis and, 396
  clinical significance, 165
  esophagitis and, 377
  HIV-positive adults, 177
  hyper-IgM syndrome and, 117
  immunodeficient patients, 118
  pneumonia, 683
  TORCH infection, 182
  treatment, 202
  viral receptor, 166
Cytoplasm
  cell cycle phase, 46
  cytoskeletal elements, 48
  glycolysis, 76
  HMP shunt, 79
  metabolism in, 72
Cytoplasmic membrane (bacterial), 124
Cytoplasmic processing bodies (P-bodies), 41
Cytoskeletal elements, **48**
Cytotoxic T cells, **102**
  cell surface proteins, 110
  MHC I and II, 100
Cytotrophoblast, 617

D

Dabigatran, 435
Dabrafenib, **444**
Daclizumab
  targets of, 121
Dacrocytes, 414
Dactinomycin, 439
  RNA polymerase inhibition, 42
  targets of, 438
Dactylitis
  seronegative spondyloarthritis, 469
  sickle cell anemia, 422
Dalfopristin
  VRE, 198
Danazol, **658**
  pseudotumor cerebri, 521
"Dancing eyes, dancing feet," 228
Dandy-Walker syndrome, 492
Dantrolene, 550, 551
Dapagliflozin, 353
Dapsone
  Hansen disease, 141
  hemolysis in G6PD deficiency, 250
  mechanism and use, **194**
  Mycobacterium leprae, 196
  Pneumocystis jirovecii, 154
Daptomycin
  mechanism and use, **195**
  MRSA, 198
Darkfield microscopy
  for Treponema, 146
Darunavir
  HIV therapy, 203
Dasabuvir, 204
Dasatinib, 443
Datura, 241
Daunorubicin, 439
  dilated cardiomyopathy, 248
DCC gene, 224
Deacetylation
  histones, 34
Dead space (lung), 664
Deafness
  Alport syndrome, 596
  congenital long QT syndrome, 294
  congenital syphilis, 147
  rubella, 182
  syphilis, 182
Deamination
  base excision repair, 40
Death
  aortic dissection in, 303
  children, explaining to, 269
  common causes, **272**
  sudden cardiac death, 304
  thyroid storm, 342
Death receptor pathway, 207, 208
Debranching enzyme
  Cori disease, 87
  glycogen metabolism, 86
Decay-accelerating factor (DAF), 106
Deceleration injury, 303
Decerebrate (extensor) posturing, 511
Decidua basalis, 617
Decision-making capacity, **266**
Decorticate (flexor) posturing, 511
Decussation
  in spinal tracts, 509
Deep brachial artery, 455
Deep inguinal lymph nodes, 624
Deep venous thrombosis (DVT), **671**
  direct factor Xa inhibitors for, 437
  glucagonomas and, 351
  heparin for, 436
  tamoxifen/raloxifene and, 443
Deer flies (disease vectors), 159
Defense mechanisms
  immature, 554–555
  mature, 555
Defensins, 99
Deferasirox, 248
Deferiprone, 248
Deferoxamine, 248
Deformation, 613
Degarelix, **656**
Degenerate/redundant genetic code, 37
Degmacytes, 414
  G6PD deficiency, 79
Dehydration
  diabetic ketoacidosis, 347
  filtration changes and, 583
  gout exacerbation, 467
  loop diuretics and, 608
  mannitol and, 607
  osmotic laxatives, 401
  salivary stones with, 376
  shock, 310
Dehydrogenases, 73
Delavirdine
  HIV therapy, 203
Delirium, **558**
  barbiturate withdrawal, 570
  diabetic ketoacidosis, 347
  PCP, 571
  thyroid storm, 342
Delirium tremens (DTs), **569**, 570
δ cells
  endocrine pancreas
  somatostatin production, 371
  somatostatinomas of, 351
Delta rhythm (EEG), 497
Delta virus, 167
Deltoid muscle
  axillary nerve injury, 447
  Erb palsy, 448
Delusional disorder, 560
Delusions, 559
  mesolimbic pathway, 499
Demeclocycline
  diabetes insipidus and, 249, 338
Dementia
  HIV-positive adults, 177
  metachromatic leukodystrophy, 88
  neurodegenerative disorders, 520–521
  prion disease, 178
  vitamin B_3 deficiency, 67
Demyelinating/dysmyelinating disorders, **523**
  lead poisoning (adult), 425
  metachromatic leukodystrophy, 88
  progressive multifocal leukoencephalopathy, 524
  vitamin B_{12} deficiency, 530
Dendritic cells, **408**
  IL-10, 108
  innate immunity, 99
  T- and B-cell activation, 102, 103
Dengue, 167
Denial, 554
Denosumab, 122
  for osteoporosis, 462
De novo synthesis
  pyrimidine and purine, **36**, 73
Dense deposit disease, 596
Dental plaque
  normal flora, 178
  viridans streptococci, 128
Dentate line, 366
Dentate nucleus, 499
Dentin
  collagen in, 50
  osteogenesis imperfecta, 51
Dentinogenesis imperfecta, 51
Denys-Drash syndrome, 606
Dependent personality disorder, 566
Depersonalization/derealization disorder, 558
  panic disorder, 563
Depolarizing neuromuscular blocking drugs, 551
Depressants, intoxication and withdrawal, 570
Depression
  atypical features in, **561**
  benzodiazepine withdrawal, 570
  dissociative identity disorder, 558
  drug therapy, 572
  electroconvulsive therapy, 562
  hyperparathyroidism, 345
  metoclopramide, 400
  mirtazapine for, 244
  neurotransmitters for, 495
  postpartum, 562
  seasonal pattern with, 561
  serotonin-norepinephrine reuptake inhibitors (SNRIs) for, 575
  SSRIs for, 575
Deprivation effects (infants), 556
De Quervain tenosynovitis, 461
De Quervain thyroiditis, 341
Dermacentor tick (disease vector), 149
Dermatitis
  B-complex deficiency, 65
  drug reactions and, 250
  glucagonomas, 351
  IPEX syndrome, 102
  type IV hypersensitivity reaction, 113
  vitamin B_5 deficiency, 67
  vitamin B_7 deficiency, 68
Dermatitis herpetiformis, 481
  celiac disease and, 381
Dermatologic lesion terms, **475**
Dermatomes
  landmarks, 510
Dermatomyositis, 228
  autoantibody, 115
Dermatomyositis/polymyositis, 471
Dermatophytes, **152**
Dermis, 473
Descending colon, 360
Desert bumps, 151
Desert rheumatism
  Coccidioidomycosis, 151
Desflurane, 550
Desipramine, 575
Desloratadine, 686
Desmin, 48, 227
Desmopressin
  central DI, 329
  DI treatment, 338
  for hemophilia, 426
Desmosome, 474
Desquamation
  staphylococcal toxic shock syndrome, 135
Desvenlafaxine, 575
Detached retina, 537

(c) ketabton.com: The Digital Library

Developmental delay
fetal alcohol syndrome, 615
renal failure and, 603
Dexamethasone
Cushing syndrome diagnosis, 348
Dexlansoprazole, 399
Dexrazoxane, 439
Dextroamphetamine, 572
Dextrocardia, 280
x-ray, 49
Dextromethorphan, 551, **686**
DHT (dihydrotestosterone), 622, 636, 639
Diabetes insipidus, **338**
antidiuretic hormone in, 329
drug reaction and, 249
lithium, 574
lithium toxicity, 569
potassium-sparing diuretics for, 609
thiazides for, 609
Diabetes mellitus, **346**–347
atherosclerosis and, 302
β-blockers and, 245
carpal tunnel syndrome, 459
cataracts and, 535
chronic renal failure and, 603
CN III damage, 541
diabetic ketoacidosis, 347
diabetic retinopathy, 537
Friedreich ataxia, 531
fungal infections, 186
glucagonomas, 351
glucosuria in, 584
hemochromatosis, 395
hepatitis C, 173
hypertension and, 300, 316
management of, 352–353
naming conventions for, 253
nephropathy with, 597
opportunistic infections, 153
pancreatic cancer, 398
polyhydramnios and, 641
in pregnancy, 300
pyelonephritis and, 600
readmissions with, 272
tacrolimus and, 120
teratogenic potential of maternal, 614
therapy management, **352**
type 1 vs type 2, 347
urinary tract infections, 181
UTIs and, 600
Diabetes mellitus type 1, 347
autoantibody, 115
HLA subtypes with, 100
Diabetes mellitus type 2, 347
amyloidosis, 212
Diabetic ketoacidosis (DKA), **347**
DM type 1 and, 346
ketone bodies, 90
Diabetic nephropathy
ACE inhibitors for, 610
Diabetic neuropathy, 575
Diabetic retinopathy, **537**
Diagnosis errors, 274
Diagnostic criteria, psychiatric
grief, 562
panic disorder, 563
symptom duration and, **564**
Diagnostic maneuvers/signs
Gower sign, 61
Diagnostic test evaluation, 257
Diagnostic tests/maneuvers
laboratory tests in bone disorders, 464
lower extremity, 491
wrist and hand injury, 459
Dialectical behavioral therapy, 565, 572
Dialysis-related amyloidosis, 212
Diamond-Blackfan anemia, 420
Diapedesis, 215
Diaper rash
Candida albicans, 153
nystatin, 199
Diaphoresis, 305
acromegaly, 339
Diaphragmatic hernia, 370
Diaphragm structures, **663**
Diaphysis, 465
Diarrhea, 117
B-complex deficiency, 65
bismuth/sucralfate for, 399
Campylobacter jejuni, 145
cholera toxin, 132
clindamycin, 192
Clostridium difficile, 138
Cryptosporidium, 155
drug reaction and, 249
ezetimibe, 320
giardiasis, 155
graft-versus-host disease, 119
HIV-positive adults, 177
inflammatory bowel diseases, 382
irritable bowel syndrome, 383
lactase deficiency, 81
lactose intolerance, 381
as laxative adverse effect, 401
leflunomide, 486
loperamide for, 400
magnesium deficiency from, 332
magnesium hydroxide, 399
malabsorption syndromes, 381
metoclopramide, 400
misoprostol, 399
opioids for, 551
organisms causing, 179
pellagra, 67
rice-water, 132
rotavirus, 168
Salmonella, 144
Shigella, 144
thyroid storm and, 342
Vibrio cholerae, 146
VIPomas, 371
vitamin C toxicity, 69
watery, 132
Diastole
cardiac cycle, 287
heart failure and, 309
heart murmurs of, 291
heart sounds of, 287
Diazepam, 546
alcohol withdrawal, 572
Diclofenac, 486
Dicloxacillin
mechanism and use, 188
Dicrotic notch, 287
Dicyclomine, 241
Didanosine, 203
HIV therapy, 203
pancreatitis, 249
Diencephalon, 490
Diethylcarbamazine
antihelminthic, 200
nematode infections, 159
Diethylstilbestrol (DES), 656
teratogenicity, 614
vaginal tumors, 644
Differential media, 126
Diffuse axonal injury, **515**
Diffuse cortical necrosis, **602**
Diffuse esophageal spasm, 377
Diffuse gastric cancer, 379
Diffuse glomerular disorders, 594
Diffuse large B-cell lymphoma (DLBCL), 430, 432
Diffuse partial seizures, 517
Diffuse proliferative glomerulonephritis (DPGN), 596
Diffuse scleroderma, 473
Diffuse stomach cancer, 379
Diffusion-limited gas exchange, 668
DiGeorge syndrome, 344
lymph node paracortex in, 96
thymic shadow in, 98
Digestion
malabsorption syndromes, 381
secretory products for, 372–374
ulcerative colitis and, 382
Digestive tract
anatomy and histology, **362**
Digitalis
arrhythmias induced by, 322
contractility effects, 284
hyperkalemia and, 590
toxicity treatment for, 248
Digoxin
contractility effects of, 285, 286
for dilated cardiomyopathy, 308
mechanism and clinical use, 321
sodium-potassium pump inhibition, 49
therapeutic index of, 234
toxicity treatment, 324
Dihydroergotamine, 518
Dihydrofolic acid, 194
Dihydroorotate dehydrogenase
leflunomide effects, 36, 486
Dihydropyridine calcium channel blockers, 253
Dihydropyridine receptor, 456
Dihydrorhodamine test, 117
Dihydrotestosterone (DHT)
5α-reductase deficiency, 639
function, 636
sexual determination, 622
Dihydroxyacetone-P, 80
Dilated cardiomyopathy, 308, 309
doxorubicin, 439
drug reaction and, 248
hemochromatosis, 395
muscular dystrophy, 61
wet beriberi, 66
with myocarditis, 313
Diltiazem, 318
Dimenhydrinate, 686
Dimercaprol
for arsenic toxicity, 248
for lead poisoning, 248, 419
for mercury poisoning, 248
Dinitrophenol, 78
Diphenhydramine, 686
Diphenoxylate, 551
Diphtheria
Corynebacterium diphtheriae, 139
exotoxins, 130, 131, 132
unvaccinated children, 186
vaccine for, 139
Diphyllobothrium latum
disease association, 161
presentation, 160
vitamin B_{12} deficiency, 69, 420
Diplococci, 142
Diplopia
brain stem/cerebellar syndromes, 523
central vertigo, 534
drug toxicity, 544
in botulism, 138
intracranial hypertension, 521
myasthenia gravis, 472
osmotic demyelination syndrome, 524
Dipyridamole
for coronary steal syndrome, 304
Direct bilirubin, 375
Direct cholinomimetic agonists, 240
Direct (conjugated) hyperbilirubinemia, 393
Direct Coombs test
Type II hypersensitivity, 112
Direct factor Xa inhibitors, **437**
Direct fluorescent antibody (DFA) microscopy
for Treponema, 146
Direct inguinal hernia, 370
Direct light reflex, 539
Direct sympathomimetics, 242
Direct thrombin inhibitors, **435**
Discharge planning, 272
Discolored teeth, 204
Discounted fee-for-service, 271
Disease associations, **161**
Disease prevention, **270**
Disease vectors
Aedes mosquitoes, 168
Anopheles mosquito, 157
armadillos, 149
birds, 148, 149
cats, 149
dogs, 145, 149
fleas, 149, 150
flies, 144, 149
horse flies, 159
Ixodes ticks, 146
rodents, 167
ticks, 146, 150
zoonotic bacteria, 149
Disinfection/sterilization methods, **204**
Disinhibited behavior
Klüver-Bucy syndrome, 511
Disinhibited social engagement, 556
Disopyramide, 322
Disorganized thought, 559
Dispersion measures, 262
Displacement, 554
Disruption (morphogenesis), 613
Disruptive mood dysregulation disorder, 557
Disseminated candidiasis, 153
Disseminated intravascular coagulation (DIC), 428
acute myelogenous leukemia, 432
Ebola, 171
endotoxins, 131, 133
meningococci, 142
microangiopathic anemia, 423
Waterhouse-Friderichsen syndrome, 349
Dissociation, 554
Dissociative disorders, **558**
amnesia, 558
identity disorder, 558
Distal humerus, 455
Distal interphalangeal (DIP) joints, 451
Distal renal tubular acidosis (type 1), 593
Distributive shock, 310
Disulfiram
alcoholism treatment, 571
ethanol metabolism and, 72
disulfiram-like reaction, 251

(c) ketabton.com: The Digital Library

Diuresis
  atrial natriuretic peptide, 296
  for shock, 310
Diuretics
  dilated cardiomyopathy, 308
  electrolyte changes, **609**
  glaucoma treatment, 552
  hypertension treatment, 316
  magnesium levels and, 332
  mechanism and clinical use, **607**
  pancreatitis, 249
  for SIADH, 338
Diverticula, **383**
Diverticulitis, 383
Diverticulosis, 383
Diverticulum, 383
Dizygotic ("fraternal") twins, 616
Dizziness
  AChE inhibitors, 549
  calcium channel blockers, 318
  dihydropyridine, 318
  nitrates, 318
  ramelteon, 547
  ranolazine, 319
  sacubitril, 319
  vertigo and, 534
DMPK gene, 61
DNA
  cloning methods, 55
  free radical effect on, 210
  introns vs exons, 43
  laddering in apoptosis, 208
  methylation in, 34
  mutations in, 39
  plasmid transfer, 130
  repair of, **40**
DNA ligase
  action of, 38
DNA mutations
  types of, 39
DNA polymerase inhibitors, 253
DNA polymerases
  action of, 38
DNA repair, **40**
DNA replication, **38**
DNA topoisomerases, 38
DNA transcription
  deacetylation, 34
DNA viruses, **164**
  characteristics, **163**
  genomes, 163
Dobutamine, 242
Dofetilide, 323
Dogs (disease vectors), 145, 149
Dolutegravir, 203
Dominant inheritance, 59
Dominant negative mutations, 57
Donepezil, 240, 549
Do not resuscitate (DNR) order, 266
Dopamine, 242
  basal ganglia, 500
  bupropion effect, 576
  changes with disease, 495
  function of, 328
  Huntington disease, 520
  kidney functions and, 589
  L-DOPA, 548, 549
  MAO inhibition, 549, 575
  Parkinson disease, 548
  PCT secretion of, 589
  pheochromocytoma secretion, 350
  vitamin B_6 and, 67
Dopamine agonists, 330, 339, 548
Dopamine receptors, 238
Dopaminergic pathways, **499**
Dornase alfa (DNAse), 60
Dorsal columns (spinal cord), 508, 509
  thalamic relay for, 498
Dorsal interossei muscle, 450
Dorsal motor nucleus, 506
Dorsal optic radiation, 542
Dorsal pancreatic bud, 360
Dorsiflexion
  common peroneal nerve injury, 453
Dosage calculations, **231**
Dosage interval, 231
Double-blinded studies, 256
Double stranded viruses, 163
Double Y males, 638
Down syndrome, 63
  ALL and AML in, 432
  cardiac defect association, 300
  cataracts and, 535
  chromosome association, 64
  hCG in, 633
  Hirschsprung disease and, 384
  intestinal atresia and, 359
Doxazosin, 244
Doxepin, 575
Doxorubicin
  dilated cardiomyopathy, 248
  targets, 438
  toxicities, 444
Doxycycline
  chlamydiae, 148
  in gonorrhea treatment, 142
  lymphogranuloma venereum, 149
  MRSA, 198
  Mycoplasma pneumoniae, 150
  rickettsial/vector-borne disease, 150
  tetracyclines, 192
Doxylamine, 686
DPP-4 inhibitors, 353
Dressler syndrome, 305, 307, 313
DRESS syndrome, 250
Drooling treatment, 241
Drop metastases, 528
Drop seizures, 517
Drug dosages, 234
  calculations, 231
  lethal median, 234
  liver disease, 231
  loading, 231
  maintenance dose, 231
  median effective, 234
  renal disease, 231
  toxic dose, 234
Drug elimination, 231
Drug-induced lupus, 250
Drug interactions
  additive, type, 235
  antagonistic, type, 235
  permissive, type 235
  potentiation, type, 235
  synergistic, type, 235
  tachyphylactic, type, 235
Drug metabolism, **232**
  cytochrome P-450 dependent, 232
  geriatric patients in, 232
Drug name conventions, **253**–255
Drug overdoses
  of weak acids, 233
  of weak bases, 233
Drug reactions
  cardiovascular, **248**
  endocrine/reproductive, **249**
  gastrointestinal, **249**
  hematologic, **250**
  multiorgan, **251**
  musculoskeletal, **250**
  neurologic, **251**
  renal/genitourinary, **251**
  respiratory, **251**
Drug reaction with eosinophilia and systemic symptoms (DRESS), 250
Drug-related disorders
  myocarditis, 313
Drug resistance
  plasmids in, 131
Drugs
  cholinomimetic agents, 240
  efficacy vs potency, 233
  elimination of, 232, 233
  errors in, 274
  patient difficulty with, 268
  phase I metabolism, 232
  reactions to, 248–251
  therapeutic index, 234
  toxicities, 248
Drug safety
  therapeutic index, measurement, 234
Drug trials, 256
Drunken sailor gait, 511
Drusen, 536
Dry beriberi, 66
Dry cough with ACE inhibitors, 251
Dry mouth
  Lambert-Eaton myasthenic syndrome, 472
Dry skin, 66
Dubin-Johnson syndrome, 393, 394
Duchenne muscular dystrophy, 61
  inheritance, 61
Ductal adenocarcinomas, 368
Ductal carcinoma in situ (DCIS), 650
Ductal carcinomas (invasive), 650
Ductus arteriosus, 282, 619
Ductus deferens
  embryology, 622
Ductus venosus, 282
Duloxetine, 575
Duodenal atresia, 359
Duodenal ulcer, 380
Duodenum
  basal electric rhythm, 362
  biliary structures and, 368
  histology of, 362
  location, 360
  secretory cells, 373
Duplex collecting system, **579**
Dural venous sinuses, **503**
Dura mater, 496
Dwarfism
  achondroplasia, 462
D-xylose test, 381
Dynein
  movement of, 48
Dynein motors, 171
Dysarthria, 516
  Friedreich ataxia as, 531
  in botulism, 138
  osmotic demyelination syndrome, 524
Dysbetalipoproteinemia, 94
Dysentery
  Entamoeba histolytica, 179
  Escherichia coli, 145
  Shigella spp, 132, 144, 179
Dysfunctional uterine bleeding, 633
Dysgerminoma, 647
Dysgeusia, 71
Dyskinesia
  tardive, 251
Dyslipidemia
  β-blocker adverse effects, 323
  β-blockers, **245**
  familial, 94
  renal failure and, 603
  vitamin B_3 for, 67
Dysmenorrhea
  copper IUD, 657
  primary, 645
Dysmetria
  central vertigo, 534
  with strokes, 514
Dyspareunia, 567
Dysphagia
  achalasia, 376
  esophageal pathologies and, 377–378
  in botulism, 138
  osmotic demyelination syndrome, 524
  Plummer-Vinson syndrome, 418
  stroke effects, 514
  Zenker diverticulum, 384
Dysplasia
  bronchopulmonary, 210
  cellular adaptive response, 206
  cervical, 645
  neoplastic progression, 219
Dysplastic kidney
  multicystic, 578, 579
Dyspnea
  α_1-antitrypsin deficiency, 392
  aortic stenosis, 291
  heart failure, 309
  hypersensitivity pneumonitis, 675
  hypertrophic cardiomyopathy, 308
  in botulism, 138
  Wegener granulomatosis, 315
Dystonia
  acute, 241
  antipsychotics/antiepileptics, 569
  benztropine for, 241
  Lesch-Nyhan syndrome, 37
  movement disorders, 519
  nigrostriatal pathway and, 499
  treatment of, 241
Dystrophic calcification, 211, 227
Dystrophin gene, 61
Dysuria, 654
  cystitis, 181
  urinary catheterization, 185
  UTIs causing, 600

E

Ear
  pharyngeal pouch derivation, 621
Eardrum, 533
Early complement deficiencies (C1-C4), 107
Eating disorders
  anovulation and, 645
  binge-eating disorder, 567
  bulimia nervosa, 567
  characteristics of, **567**
  pica, 567
Eaton agar
  culture requirements, 126
Ebola virus, 167, **171**
Ebstein anomaly, 281, 298, 300
  lithium side effect, 574
E-cadherin, 219, 474
ECF
  body compartments of, 231
Echinocandins, **200**

(c) ketabton.com: The Digital Library

Echinococcus granulosus
cestodes, 160
disease association, 161
Echinocytes, 414
Echothiophate, 552
Echovirus
picornavirus, 167, 168
Eclampsia, 300, 643
Ecthyma gangrenosum, 143
Pseudomonas spp, 143
Ectocervix, 626
Ectoderm
branchial clefts, 619
derivatives of, 613
Ectoparasites, **161**
Ectopic pregnancy, 641
appendicitis differential diagnosis, 383
Chlamydia trachomatis, 149
hCG in, 633
Kartagener syndrome, 49
methotrexate for, 440
salpingitis and, 185
Eculizumab, 122
Eczema
atopic dermatitis, 477
eczematous dermatitis, 475
phenylketonuria, 84
skin scales in, 475
type I hypersensitivity, 112
Wiskott-Aldrich syndrome, 117
Edema
acute poststreptococcal glomerulonephritis, 596
Arthus reaction, 113
calcium channel blockers, 318
capillary fluid exchange and, 297
danazol, 658
diabetic ketoacidosis, 347
endotoxins, 133
fludrocortisone, 354
heart failure and, 309
with hyperaldosteronism, 349
immunosuppressants, 120
Kawasaki disease and, 314
kwashiorkor, 71
loop diuretics for, 608
periorbital, 159, 161
peripheral, 309, 610
pitting, 309
pseudoephedrine/phenylephrine, 686
pulmonary hypertension, 668
Trichinella spiralis, 159, 161
trichinosis, 159
vasogenic, 496
wet beriberi, 66
Edema factor
Bacillus anthracis, 132
Edinger-Westphal nuclei, 539
Edrophonium, 240
Edwards syndrome, 63
cataracts and, 535
chromosome association, 64
Efavirenz
HIV-positive adults, 203
Effective refractory period
Class IA antiarrhythmic effect, 322
Class IC antiarrhythmic effect, 322
myocardial action potential, 292
Effective renal plasma flow, **582**
Efferent/afferent nerves, 296
Efferent arteriole, 580
ANP/BNP, 588
constriction of, 583
Efficacy vs potency of drugs, **233**
EGFR gene, 684
"Eggshell" calcification, 677
Ego defenses, **554**, 555
Ego-dystonic behavior, 563
Egophony, 680
Ego-syntonic behavior, 565, 566
Ehlers-Danlos syndrome
aneurysm association with, 516
collagen in, 50
collagen synthesis in, **51**
heart murmur with, 291
Ehrlichia spp
animal transmission, 149
Ehrlichia chaffeensis, 149
Gram stain, 125
rickettsial/vector-borne, **150**
Eisenmenger syndrome, 299
Ejaculation
innervation of, 627
sperm pathway in, 626
Ejaculatory ducts, 626
embryology of, 622
Ejection fraction
equation for, 285
Elastase, 373
activity in emphysema, 674
Elastic recoil (lung and chest wall), 665
Elastin
characteristics of, **52**
Elbow injuries, 459, 461
Elderly
changes in, **270**
Electrocardiograms (ECGs), **293**
acute pericarditis on, 313
cardiac tamponade on, 310
low-voltage, 308, 310
MI diagnosis with, 306
with pulmonary embolism, 672
tracings of, **295**
Electroconvulsive therapy (ECT)
adverse effects, **562**
postpartum psychosis, 562
Electroencephalogram (EEG)
Creutzfeldt-Jakob disease, 521
in delirium, 558
sleep stages, 497
Electrolytes
disturbances in, **591**
diuretic effects on, 609
Electron acceptors (universal), 75
Electron transport chain and oxidative phosphorylation, **78**
inhibitors of, 78
Electrophoresis
hemoglobin, 410
Elek test, 139
Elementary bodies (chlamydiae), 148
Elephantiasis, 159
11β-hydroxylase, 335
11-deoxycortisol
and metyrapone, 335
11-deoxycorticosterone, 335
Elfin facies, 64
Elimination constant, 231
Elimination of drugs, **232**
urine pH and, 233
ELISA (enzyme-linked immunosorbent assay), **54**
Elongation Factor 2
Corynebacterium diphtheriae, 139
eltrombopag (TPO receptor agonist), 121
Elvitegravir, 203
Emancipated minors, 265
EMB agar
Escherichia coli, 181
lactose-fermenting enterics, 144
Emboli
atherosclerosis, 302
atrial fibrillation, 295
atrial septal defect, 299
pulmonary, 310
stroke, 512
Emboliform nucleus, 499
Embryogenesis
genes involved in, 612
intrinsic pathway and, 208
Embryology
cardiovascular, 281–283
derivatives, **613**
erythropoiesis, 404
gastrointestinal, 358–359
gland derivations in, 621
hematologic/oncologic, 404
neurological, 490–492
pancreas and spleen, 359, 360
renal, 578–579
reproductive, 612–623
respiratory, 660
thyroid development, 326
USMLE Step 1 preparation, 276
Embryonal carcinoma, 653
Embryonic age calculation, 633
Embryonic development, 612
Embryonic morphogenic errors, 613
Embryonic stage (development), 660
Embryotoxic, 204
Emergent care proxy, 269
Emission
innervation of, 627
Emollients
laxative, 401
Emotion
neural structures and, 499
Emotionally distraught patients, 268
Emotional/social development
neglect and deprivation effects, 556
Empagliflozin, 353
Emphysema, 674
diffusion in, 668
diffusion-limited gas exchange, 668
elastin in, 52
panacinar, 392
Empty/full can test, 446
Empty sella syndrome, 339
Emtricitabine, 203
HIV-positive adults, 203
Enalapril, 610
Encapsulated bacteria, **127**
infections with immunodeficiency, 118
Encephalitis
anti-NMDA receptor, 228
Cryptococcus neoformans, 153
guanosine analogs, 201
herpesviruses, 165, 180
HSV identification, 166
Lassa fever, 167
neonatal, 182
West Nile virus, 180
Encephalomyelitis
paraneoplastic, 228
paraneoplastic syndrome, 228
Encephalopathy
hepatic, 365, 391
hypertensive emergency, 300
lead poisoning, 419
Lyme disease, 146
prion disease, 178
renal failure, 603
Wernicke, 66
Encephalotrigeminal angiomatosis, 525
Endemic typhus, 149
Endocannabinoids, 336
Endocardial cushion, 281
Endocardial fibroelastosis, 308
Endocarditis
bacterial, 311
Candida albicans, 153
coarctation of aorta, 299
Coxiella burnetii, 150
daptomycin, 195
enterococci, 137
heart murmurs, 291
heroin addiction and, 576
nonbacterial thrombotic, 228
prophylaxis, 198
Staphylococcus aureus, 135
Streptococcus bovis, 137
viridans streptococci, 128
Endocervix, 626
Endochondral ossification, 458
Endocrine functions
kidney, 589
Endocrine system
anatomy, 327–328
embryology, 326
hormones acting on kidney, 589, 590
hormone signaling pathways, 337
pathology, 338–354
pharmacology, 352–354
physiology, 328–336
Endocrine/reproductive drug reactions, 249
Endoderm
branchial pouch derivation, 619
derivatives of, 613
Endodermal sinus tumor, 647, 653
Endodermal tubules, 660
Endometrial abnormal uterine bleeding, 633
Endometrial artery, 617
Endometrial carcinoma, 648
epidemiology of, 643
estrogens and, 656
Lynch syndrome and, 388
PCOS and, 645
tamoxifen and, 443
Endometrial hyperplasia
follicular cysts, 646
Endometrial polyps
uterine bleeding with, 633
Endometrial vein, 617
Endometriosis
characteristics and treatment, 648
danazol for, 658
Endometritis, 648
Endometrium
hyperplasia, 648–649
maintenance of, 632
Endoneurium, 495
Endoplasmic reticulum, 46, 47
rough, 46
smooth, 46
Endosomes, 47
Endothelial cells
leukocyte extravasation and, 215
in wound healing, 216
Endothelin receptor antagonist, 686
naming conventions for, 254
Endothelium-derived relaxing factor (EDRF), 337

(c) ketabton.com: The Digital Library

Endotoxins
effects of, **133**
features of, 131
Enflurane, 550
seizures with, 251
Enfuvirtide, 203
HIV-positive adults, 203
Enhancers (gene expression), 41
Enoxacin, 195
Entacapone, 548
Entamoeba histolytica
amebiasis, 155
bloody diarrhea, 179
metronidazole, 195
Enteric gram ⊖ bacteria
facultative anaerobic metabolism, 127
Enteric nerves, 362, 401
Enteritis
vitamin B_{12} deficiency, 69
vitamin B_5 deficiency, 67
vitamin B_7 deficiency, 68
Enterobacter aerogenes, 189
Enterobacter spp
nosocomial infection, 185
Enterobius spp
diseases association, 161
infection routes, 158
Enterobius vermicularis, 159
Enterochromaffin-like (ECL) cells, 350, 373
Enterococci, **137**
penicillins for, 188
vancomycin, 190
vancomycin-resistant (VRE), 137
Enterococcus spp
UTIs, 181
Enterococcus faecalis, 137
Enterococcus faecium, 137
Enterocolitis
vitamin E excess, 70
Enterocolitis (necrotizing), 386
Enterohemorrhagic *Escherichia coli* (EHEC), 132, 145, 178, 179
Enteroinvasive *Escherichia coli* (EIEC), 145, 179
Enterokinase/enteropeptidase, 373
Enteropathogenic *Escherichia coli* (EPEC), 145
Enterotoxigenic *Escherichia coli* (ETEC), 132, 179
Enterovirus meningitis, 179, 180
Entorhinal cortex, 499
Enuresis
characteristics/treatment, **568**
sleep stages and, 497
TCA use for, 575
Envelopes (viral), 163
env gene, 175
Enzalutamide, 658
Enzyme kinetics, **230**
antagonists, 234
partial agonists, 234
Enzymes
glycolysis regulation, 76
lipid transport and, 91, 92
rate-determining, 73
terminology for, **73**
Eosin-methylene blue (EMB) agar
special culture, 126
Eosinopenia, 424
Eosinophilia
Aspergillus fumigatus, 153
Chlamydia trachomatis, 149
drug reaction and, 250
macrolides, 193
Eosinophilic casts (urine), 600
Eosinophilic esophagitis, 377
Eosinophilic granuloma, 675
Eosinophilic granulomatosis
autoantibody, 115
Eosinophilic granulomatosis with polyangiitis, 315
Eosinophils, **408**
corticosteroid effects, 424
in esophagus, 377
Ependymal cells, **493**
Ependymoma, 528
Ephedrine, 242
Epicanthal folds
cri-du-chat syndrome, 64
Down syndrome, 63
Epidemic typhus, 149
Epidemiology/biostatistics, 256–262
Epidermal growth factor (EGF)
signaling pathways for, 337
in wound healing, 216
Epidermis, 473
embryologic derivatives, 613
Epidermophyton, 152
Epididymis, 626
embryology of, 622
Epididymitis, 184, **654**
Epidural hematomas, 513
Epidural space, 496
Epigastric pain
chronic mesenteric ischemia, 386
Ménétrier disease, 379
pancreatitis, 397
Epigastric veins, 365
Epiglottitis
Haemophilus influenzae, 142
unvaccinated children, 186
Epilepsy
confidentiality exceptions for patients with, 267
gustatory hallucinations in, 559
hallucinations in, 559
seizures, 517
Epinephrine, 242
adrenal medulla secretion, 327
glaucoma treatment, 552
glycogen regulation by, 85
pheochromocytoma secretion, 350
unopposed secretion of, 346
vitamin B_6 and, 67
Epineurium, 495
Epiphysis
estrogen effects on, 459
slipped capital femoral, 461, 463
tumors in, 464
widening of, 463
Episcleritis
inflammatory bowel disease, 382
Epispadias, 624
Epistaxis, **671**
hereditary hemorrhagic telangiectasia, 316
Epithelial cell junctions, **474**
Epithelial cells
tumor nomenclature of, 220
Epithelial histology (female), **626**
Epithelial hyperplasia, 649
Eplerenone, 609
Epley maneuver, 534
Epoetin alfa (EPO analog), 121
Epstein-Barr virus (EBV)
aplastic anemia, 421
Burkitt lymphoma, 430
false-positive VDRL, 148
hairy leukoplakia and, 479
head and neck cancer, 671
HIV-positive adults, 177
Hodgkin lymphoma, 429
in immunodeficient patients, 118
nasopharyngeal carcinomas, 165
oncogenesis of, 226
paracortical hyperplasia in, 96
receptors for, 166
Eptifibatide, 438
thrombogenesis and, 411
Equations
half-life of, 231
Erb palsy, 448
Erectile dysfunction, 567
β-blockers and, 245, 323
cimetidine, 399
Lambert-Eaton myasthenic syndrome, 472
Peyronie disease, 651
sildenafil, 686
Erection
autonomic innervation, 627
ischemic priapism, 651
Ergocalciferol, 70
Ergosterol synthesis inhibitors, 253
Ergot alkaloids
coronary vasospasm, 248
Erlotinib, **442**
Erosions (gastrointestinal), 362, 379
Errors (medical), 274
Erysipelas, 479
Streptococcus pyogenes, 136
Erythema
complicated hernias, 370
Kawasaki disease, 314
Erythema marginatum, 312
Erythema migrans
in Lyme disease, 146
Erythema multiforme, 481
Coccidioidomycosis, 151
Erythema nodosum, 151, 482
inflammatory bowel disease, 382
Erythroblastosis fetalis, 405
Erythrocytes, **407**
blood types, 405
casts in urine, 594
Coombs test, 423
DAF deficiency and, 107
erythropoietin and, 589
glucose usage by, 334
hereditary spherocytosis, 422
myeloproliferative disorders, 433
transfusion of, 429
Erythrocyte sedimentation rate (ESR), **214**
subacute granulomatous thyroiditis, 341
Erythrocytosis, 407
oxygen-hemoglobin dissociation curve, 666
Erythrogenic toxin, 136
Erythromelalgia, 433
Erythromycin
macrolides, 193
prophylaxis, 198
protein synthesis inhibition, 191
reactions to, 249
erythroplasia of Queyrat, 651
Erythropoiesis, 679
fetal, 404
Erythropoietin (EPO), 121
anemia of chronic disease, 421
aplastic anemia, 421
high altitude, 670
with pheochromocytoma, 350
polycythemia and, 228
release of, 589
in renal failure, 603
signaling pathways for, 337
Eschar, 132
in cutaneous anthrax, 137
with mucormycosis, 153
Escherichia coli, **145**
cephalosporins, 189
culture requirements, 126
encapsulation, 127
galactosemia, 80
lac operon, 40
lactose fermentation, 144
meningitis, 180
neonatal illness, 182
nosocomial infection, 185
O157-H7, 132, 145, 178, 179
penicillins for, 188
pneumonia, 179
prostatitis, 654
urinary tract infections, 181, 600
Escitalopram, 575
E-selectin, 215
Esmolol, 245, 323
Esomeprazole, 399
Esophageal adenocarcinoma, 378
Esophageal atresia, 359
Esophageal cancer, **378**
achalasia and, 376
Esophageal dysmotility
CREST syndrome, 473
Esophageal perforation, 377
Esophageal strictures, 377
Esophageal varices, 365, 377
Esophageal webs, 418
Esophagitis, 377
bisphosphonates, 486
drug reaction and, 249
HIV-positive adults, 177
Esophagus
blood supply and innervation, 364
diaphragm, 663
histology of, 362
pathologies of, **377**
portosystemic anastomosis, 365
Essential amino acids, 81
Essential fructosuria, 80
Essential hypertension, 316
Essential mixed cryoglobulinemia, 173
Essential thrombocythemia, 433
Essential tremor, 519
Esters (local anesthetics), 550
Estrogen, **656**
androgen insensitivity syndrome, 639
androgen conversion to, 636
benign breast tumors, 649
bone formation, 459
epiphyseal plate closure, 636
gynecomastia, 649
menopause, 636
menstrual cycle, 632
osteoporosis, 462
ovulation, 631
premature ovarian failure, 636, 645
prolactin suppression of, 330
signaling pathways for, 337
source and function of, **630**
Turner syndrome, 638
Estrogen receptor modulators (selective), 656
Eszopiclone, 546
Etanercept, 487
Ethacrynic acid, 608
Ethambutol, 196, **197**

FAS1_2019_21_Index_749-806.indd 767 11/21/19 12:27 PM

(c) ketabton.com: The Digital Library

Ethanol
as carcinogen, 225
gluconeogenesis and, 72
lactic acidosis and, 72
metabolism, **72**, 232
NADPH (nicotinamide adenine dinucleotide phosphate), 72
zero-order elimination, 232
Ethics, **265–268**
confidentiality, 267
consent, 265
core principles of, **265**–267
directives, 268
religious beliefs and, 269
situations in, 268–269, 269–270
Ethinyl estradiol, 656, 657
Ethosuximide
absence seizures, 544
Ethylenediaminetetraacetic (EDTA), 419
Ethylene glycol
toxicity treatment, 248
Ethylene oxide, 204
Etonogestrel, 657
Etoposide/teniposide, **442**
targets of, 438
teniposide, 38
Euchromatin, 34
Eukaryotes
DNA replication, 38
functional gene organization, 41
mRNA start codons, 44
ribosomes in, 45
RNA polymerase in, 42
RNA processing, 41
Eukaryotic initiation factors, 45
Eukaryotic release factors, 45
Eustachian tubes
embryonic derivation, 621
Eversion, 453
Evolucumab, 320
Ewing sarcoma, 465
dactinomycin for, 439
Exanthem subitum, 165
"Excision" event, 130
Excitatory pathway, 500
Exclusive provider organization plan, 271
Executioner caspases, 208
Exemestane, 656
Exenatide, 353
Exercise
blood flow autoregulation, 297
peripheral resistance, 286
respiratory response, 670
syncope with, 308
Tetralogy of Fallot, 298
Exercise-induced amenorrhea, 645
Exocrine glands, 236
Exocytosis, 50
Exogenous corticosteroids, 336
Exons
deletions in muscular dystrophies, 61
vs introns, 43
Exotoxins
features of, 131
organisms with, 132–133
Pseudomonas aeruginosa, 132
Streptococcus pyogenes, 133
Expectorants, 686
Expiratory reserve volume (ERV), 664
Extension
hip, 451, 453
External hemorrhoids, 366
External iliac lymph nodes, 624
External oblique muscle
inguinal canal and, 369
External rotation
arm (rotator cuff), 446
hip, 451
External spermatic fascia, 369
Extinction (conditioning), 554
Extracellular fluid (ECF)
volume measurement, 581
volume regulation, 588
Extragonadal germ cell tumors, **652**
Extramammary Paget disease, 644
Extraperitoneal tissue, 369
Extravascular hemolysis, **421**
Extrinsic hemolytic anemia, **423**
Extrinsic pathway, 208
warfarin and, 436, 437
Exudate
"anchovy paste," 155
Ex vacuo ventriculomegaly, 522
Eye movements, 540
bilateral, 543
cranial nerve palsies, 541
with stroke, 515
Eyes
anatomy of, 534
aqueous humor pathway, 535
Ezetimibe, 320
diarrhea, 249

F

Fab region of antibodies, 104
Fabry disease, 61
Facial nerve (Bell) palsy, 146, 186
Facial nerve (CN VII), 146, 186
functions of, 506
inflammatory demyelinating polyradiculopathy, 524
lesions of, **532**
pharyngeal arch derivation, 620
tongue, 493
Facies
congenital syphilis, 147
elfin, 64
epicanthal folds, 63, 64
"facial plethora," 685
in fetal alcohol syndrome, 615
flat, 63
leonine (lion-like), 141
long face with a large jaw, 62
risus sardonicus, 138
twisted face, 578
Factitious disorder
characteristics of, **566**
on another, 566
on self, 566
Factor IX concentrate, 426
Factor VIII concentrate, 426
Factor V Leiden, 413, 428
venous sinus thrombosis and, 503
Factor Xa
direct inhibitors of, 437
heparin effect on, 436
inhibitors of, 413
Factor XI concentrate, 426
Facultative anaerobes
culture requirements, 127
Facultative intracellular organisms, 127
FADH (flavin adenine dinucleotide), 77
Failure mode and effects analysis, 274
Failure to thrive, 556
galactosemia, 80
orotic aciduria, 420
SCID, 117
Falciform ligament, 361
Fallopian tubes
anatomy, 625
epithelial histology, 626
fertilization, 633
False-negative rate, 257
False-positive rate, 257
Famciclovir, **201**
Familial adenomatous polyposis, 387
APC gene and, 389
chromosome association, 64
Familial amyloid cardiomyopathy, 212
Familial amyloid polyneuropathies, 212
Familial dyslipidemias, **94**
Familial hypercholesterolemia, 60, 94
Familial hypocalciuric hypercalcemia, **345**
Family discussions, 268
Family therapy
separation anxiety, 557
Famotidine, 399
Fanconi anemia, 421
DNA repair in, 40
nonhomologous end joining and, 40
Fanconi syndrome, 586
drug reaction and, 251
renal tubular acidosis, 593
Fascia
collagen in, 50
Fascia of Buck, 627
Fasciculations, 529
Fastigial nucleus, 499
Fasting plasma glucose test, 346
Fasting state, 76, 91
Fat emboli, 672
Fatigue
adrenal insufficiency, 349
heart failure and, 309
MI signs, 305
Fatigue, medical errors and, **274**
Fat necrosis, 209, 649
Fat redistribution, 250
Fat-soluble vitamins, 65
Fatty acids
gluconeogenesis, 78
metabolism of, 72, **89**, 90
oxidation of, 72, 73
synthesis, 73
Fatty acid synthase, 67
Fatty casts, 594
Fatty liver disease
hepatocellular carcinoma and, 392
nonalcoholic, 391
Fc region of antibodies, 104
Fear
anxiety disorder and, 562
panic disorder and, 563
Febrile nonhemolytic transfusion reaction, 114
Febrile pharyngitis, 164
Febrile seizures, 517
Febuxostat
gout, 487
Lesch-Nyhan syndrome, 37
Fecal elastase, 381
Fecal immunochemical testing (FIT), 388
Fecalith obstruction, 383
Fecal microbiota transplant, 138
Fecal occult blood testing (FOBT), 388
Fecal retention, 558
Feces
explosive expulsion of, 384
Federation of State Medical Boards (FSMB), 2
Fed state, 76, 91
Fee for service, 271
Felty syndrome, 466
Female genital embryology, 622
Female/male genital homologs, 623
Female reproductive anatomy, **625**
Female reproductive epithelial histology, **626**
Femoral artery, 368
Femoral head
osteonecrosis, 463
Femoral hernia, 370
Femoral nerve, 452
Femoral region, **368**
Femoral sheath, 368
Femoral triangle, 368
Femoral vein, 368
Fenestrated capillaries, 496, 581
Fenofibrate, 320
Fenoldopam, 242, 318
Fentanyl, 551
Ferritin, 213
anemia of chronic disease, 421
iron deficiency anemia, 418
lab values in anemia, 419
sideroblastic anemia, 419
Ferrochelatase, 425
Fertility
double Y males, 638
GnRH and, 328
menstrual cycle, 632
Fertilization, 631, 633
Fetal alcohol syndrome, 300, 614, **615**
holoprosencephaly in, 491
Fetal circulation, **282**
Fetal development
early, 612
placental component, 617
Fetal erythropoiesis, **404**
Fetal hypothyroidism, 341
Fetal lung maturity, 661
Fetal movement, 612
Fetal-postnatal derivatives, **282**
Fetal respiration, 660
Fetal tissue
collagen in, 50
Fever
amphotericin B, 199
childhood rashes, 183
clindamycin, 192
complicated hernias, 370
endotoxins, 131
epiglottitis, 186
exotoxins, 133
genital herpes, 184
vs heat stroke, **517**
high fever, 165, 168, 171, 183
with inflammation, 213
Jarisch-Herxheimer reaction, 148
Legionnaires' disease, 143
low-grade, 143, 171
malaria, 157
with meningococci, 142
mononucleosis, 165
neuroleptic malignant syndrome, 569
pulmonary anthrax, 137
recurring, 156
Rickettsia rickettsii, 150
Salmonella spp, 149
Salmonella typhi, 144
seizures with, 165
spiking, 158

(c) ketabton.com: The Digital Library

Tetralogy of Fallot, 298
thyroid storm causing, 342
toxic shock syndrome, 135
Trichinella spiralis, 159
tuberculosis, 140
undulant, 143
vasculitides, 314
Waterhouse-Friderichsen syndrome, 142
Weil disease, 147
Fexofenadine, 686
Fibrates, 320
hepatitis and, 249
myopathy and, 250
Fibrinogen, 213
in cryoprecipitate, 429
ESR and, 214
receptor for, 407
thrombocytes, 407
Fibrinoid necrosis, 209
Fibrinous pericarditis, 305
Fibroadenoma, 649
Fibroblast growth factor (FGF)
signaling pathways for, 337
in wound healing, 216
Fibroblast growth factor receptor (FGFR3), 462
Fibroblasts
cortisol and, 336
in wound healing, 216
Fibrocystic breast disease, 649
Fibro fog, 471
Fibroid (leiomyoma)
leuprolide for, 656
Fibromas, 647
nomenclature for, 220
Fibromuscular dysplasia, 300, 604
Fibromyalgia, 470, **471**, 575
Fibronectin
in cryoprecipitate, 429
thrombocytes, 407
Fibrosarcomas, 220
Fibrosis
diffusion-limited gas exchange, 668
silicosis, 677
Fibrous plaque in atherosclerosis, 302
Fick principle, 285
Fidaxomicin
Clostridium difficile, 138
Fifth disease
rash, 183
50S inhibitors, 191
Filgrastim (G-CSF), 121
Filoviruses
characteristics of, 167
negative-stranded, 168
Filtration fraction
glomerular dynamics, **583**
Fimbria, 124
Financial considerations in treatment, 269
Finasteride, 658
gynecomastia with, 649
reproductive hormones and, 636
Finger agnosia, 511
Finger drop, 447
Finger movements, 450
upper extremity nerve injury, 447
Fingernails, 478
Finkelstein test, 461
First-degree AV block, 295
First-order elimination, 231, **232**
Fish oil, 320
Fishy smell, 148
Fitz-Hugh-Curtis syndrome, 142, 185
5-aminosalicylic drugs, 382, 400
5α-reductase
deficiency, 622, **639**
inhibitors for BPH, 654
testosterone conversion, 636
5-fluorouracil (5-FU)
antimetabolites, 440
photosensitivity, 250
pyrimidine synthesis and, 36
targets of, 438
thymidylate synthase, 36
5-$HT_{1B/1D}$ agonists, 253
5-HT agonists, 547
5-HT
opioid effects, 551
MAO inhibitor effect on, 575
trazodone effects, 576
vilazodone effects, 576
vortioxetine effects, 576
Fixation, 555
Fixed splitting, 289
Flaccid paralysis
botulinum toxin, 138
LMN lesion, 531
motor neuron signs, 529
Flagellin, 99
Flagellum, 124
Flask-shaped ulcers, 155
Flat affect, 499
Flavin nucleotides, 75
Flaviviruses, 163, 167
Fleas (disease vectors), 149, 150
Flecainide, 322
Flexion
foot, 453
hand, 450
hip, 451
Flexor digiti minimi muscle, 450
Flexor pollicis brevis muscle, 450
Flies (disease vectors), 144, 159
Floppy baby syndrome
Clostridium botulinum, 138
spinal cord lesions, 530
Flow volume loops, **673**
Fluconazole, 151
Cryptococcus neoformans, 153
mechanism and use, 199
opportunistic fungal infections, 153
Flucytosine, **199**
Cryptococcus neoformans, 153
Fludrocortisone, **354**
Fluid compartments, **581**
Flumazenil
benzodiazepine overdose, 248, 546, 570
nonbenzodiazepine hypnotics, 546
Fluorescence in situ hybridization, **55**
Fluorescent antibody stain
bacteria, 125
Fluoroquinolones, 38
mechanism and use, **195**
Mycoplasma pneumoniae, 150
pregnancy contraindication, 204
Pseudomonas aeruginosa, 143
tendon/cartilage damage with, 250
TOP II (DNA gyrase) and TOP IV inhibition in prokaryotes, 38
typhoid fever, 144
Fluoxetine, 575
Fluphenazine, 573
Tourette syndrome, 572
Flutamide, 658
Fluticasone, 687
Fluvoxamine, 575
FMR1 gene, 62
Foam cells
in atherosclerosis, 302
Niemann-Pick disease, 88
Focal glomerular disorders, 594
Focal hepatic necrosis, 249
Focal necrotizing vasculitis, 315
Focal neurological deficits
pituitary apoplexy, 339
Focal segmental glomerulosclerosis, 597
Focal seizures, 517
Folate antagonist
teratogenicity, 614
Folate synthesis
inhibition/block, 194
Folic acid
antimicrobials and, 187
folate, 68
in pregnancy, 68
neural tube defects and, 491
Follicles (lymph), 96
Follicle-stimulating hormone (FSH)
clomiphene effect, 656
hCG and, 633
PCOS, 645
premature ovarian failure, 636
secretion of, 327
signaling pathways of, 337
Follicular conjunctivitis, 148
Follicular cysts, 646
Follicular lymphomas, **430**, 434
Follicular phase (menstrual cycle), 632
Follicular thyroid carcinomas, 343
Fomepizole
ethanol metabolism and, 72
toxicity treatment with, 248
Food-borne illness, 178
Food poisoning
Bacillus cereus, 138, 178
causes of, 178
Staphylococcus aureus, 135, 178
toxic shock syndrome toxin, 133
Food toxins, 247
Foot drop, 453
lead poisoning, 419
Foramen cecum, 326
Foramen of Magendie, 504
Foramen of Monro, 504
Foramen ovale
atrial septal defect, 299
embryology, 280
fetal circulation, 282
retained patency of, 298
Foramen primum, 280
Foramen secundum, 280
Foramina of Luschka, 504
Forced expiratory volume (FEV)
obstructive lung disease, 674
restrictive lung disease, 675
Forebrain, 490
Foregut
blood supply/innervation of, 364
development of, 358
Foreign body inhalation, 663
Formoterol, 687
Fornix (uterus), 625
45,XO, 638
47,XXY, 638
46,XX/46,XY DSD, 639
Fosamprenavir
HIV-positive adults, 203
Foscarnet, **202**
Fosphenytoin, 544
Fossa ovalis, 282
Fovea
cherry-red spot, 538
FOXP3 protein, 102
Fractures
bone diseases and, 51
chalk-stick, 463
common pediatric, 462
compartment syndrome with, 461
humerus, 447
in child abuse, 556
pathologic, 465
scaphoid, 449
vertebral compression, 462
Fragile X syndrome, **62**
chromosome association, 64
Frameshift mutations
deletions, 61
muscular dystrophy and, 61
Francisella spp
intracellular organism, **127**
Francisella tularensis
animal transmission, 149
Frataxin, 531
Free fatty acids
fast/starvation states, 91
lipid transport and, 92
Free light chain (FLC) assay
plasma cell dyscrasias, 431
Free nerve endings, 494
Free radical injury, **210**
Fremitus (tactile), 680, 682
Fresh frozen plasma, 429
"Fried egg" cells, 494, 526
Friedreich ataxia, **531**
chromosome association, 64
hypertrophic cardiomyopathy, 308
inheritance of, 60
mechanism of, 62
Frontal bossing, 339
Frontal eye fields
cortical functions, 501
lesions in, 511
Frontal lobe
lesions in, 511
stroke effects, 514
Frontotemporal dementia, 520
Fructokinase, 80
Fructose-1,6-bisphosphatase, 73
gluconeogenesis, 78
in metabolic pathways, 74
Fructose-2,6-bisphosphate, 76
Fructose intolerance, 80
Fructose metabolism
disorders, 80
pathways, 74
Fructosuria, 80
FTA-ABS, 125, 147
Fumarate, 82
Functional hypothalamic amenorrhea, **645**
Functional neurologic symptom disorder, 566
Functional residual capacity (FRC), 664
Fungal infections
IL-12 receptor deficiency, 116
infections with immunodeficiencies, 118
Fungi
culture requirements, 126
immunocompromised patients, 179
necrosis and, 209
opportunistic infections, 153
silver stain, 125
topical infections, 199
Funny current, 292
"funny" sodium channels, 324

(c) ketabton.com: The Digital Library

Furosemide, 252, 608
  gout with, 250
  pancreatitis, 249
Fusion inhibitors, 203
Fusion protein EWS-FLI1, 465
Fusobacterium spp
  alcoholism, 179
  anaerobic metabolism of, 127

G

G6PD
  deficiency, 61, **79**
  HMP shunt and, 73
  in respiratory burst, 109
G6PD deficiency, **422**
GABA
  anesthesia effects, 550
  barbiturate effects, 546
  basal ganglia and, 500
  benzodiazepine effects, 546
  changes with disease, 495
  epilepsy drugs, 544
  Huntington disease, 520
  vitamin B_6 and, 67
Gabapentin, 544
$GABA_B$ receptor agonists, 523, 551
gag gene, 175
Gag reflex, 507
Gait disorders
  "steppage," 453
  Trendelenburg sign/gait, 453
Gait disturbance
  cerebellar lesions and, 499
  Friedreich ataxia, 531
  Parkinson disease, 520
  vitamin B_{12} deficiency, 530
  waddling, 61
Galactocerebrosidase, 88
Galactocerebroside, 88
Galactokinase deficiency, 80
  cataracts and, 535
Galactorrhea
  antipsychotic drugs and, 328
  pituitary prolactinomas, 328
  tuberoinfundibular pathway, 499
Galactose-1-phosphate uridyltransferase, 80
Galactose metabolism
  disorders of, 80
Galactosemia, 80
  cataracts and, 535
Galantamine, 240, 549
Galant reflex, 510
Gallbladder
  biliary structures, 368
  blood supply and innervation of, 364
  regulatory substances, 371
Gallbladder cancer
  porcelain gallbladder and, 397
  sclerosing cholangitis and, 395
Gallstone ileus, 396
γ-glutamyltransferase (GGT)
  alcohol use, 570
  γ-glutamyl transpeptidase (GGT), 390
γ-interferon, 407
Ganciclovir, **202**
  agranulocytosis, 250
Ganglion cyst, 461
Ganglioneuromatosis
  oral/intestinal, 351
Gangrene
  Buerger disease, 314
  diabetes mellitus, 346
Gangrenous necrosis, 209
Gap junctions, 474
Gardener's pupil, 241
Gardnerella vaginalis, **148**
Gardner syndrome, 387
Gargoylism, 88
Gas gangrene
  alpha toxin, 133
  Clostridium perfringens, 138, 179
Gastrectomy, 420
Gastric acid, 372
  histamine receptors and, 238
  regulatory substances and, 371
Gastric arteries
  celiac trunk, 364
  intralimental, 361
Gastric bypass surgery
  ghrelin and, 371
  vitamin B_{12} deficiency, 69
Gastric cancer, **379**
  carcinogens causing, 225
  Helicobacter pylori, 146
  oncogenes and, 224
  oncogenic microbes and, 226
  sign of Leser-Trélat and, 228
  trastuzumab for, 443
  types of, 379
Gastric inhibitory peptide (GIP), 351
Gastric outlet obstruction, 359, 380
Gastric sclerosis, 473
Gastric ulcers, 380
  NSAID toxicity, 486
Gastric vessels, 361
Gastrin, 371, 373
  signaling pathways for, 337
  somatostatinomas and, 351
Gastrinomas, 354, 371
Gastrin-releasing peptide (GRP), 371
Gastritis, 146, **379**
  gastrin in, 371
  H_2 blockers for, 399
  proton pump inhibitors for, 399
  stomach cancer and, 379
Gastrocolic ligament, 361
Gastroenteritis
  caliciviruses, 167
  Listeria monocytogenes, 139
  rotavirus, 168
  Salmonella spp, **144**
Gastroepiploic arteries, 361, 364
Gastroesophageal reflux disease (GERD)
  Barrett esophagus, 378
  esophageal cancer and, 378
  presentation, 377
Gastrohepatic ligament, 361
Gastrointestinal bleeding
  hereditary hemorrhagic telangiectasia, 316
Gastrointestinal drug reactions, 249
Gastrointestinal stromal tumors (GISTs), 224
Gastrointestinal system
  anatomy, 360–369
  blood supply to, 363
  embryology, 358–359
  innervation of, **364**
  ligaments, 361
  pathology, 376–397
  pharmacology, 398–400
  physiology, 371–375
  regulatory substances, **371**
  secretory cells, 373
  secretory products, **372**
Gastroschisis, 358
Gastrosplenic ligament, 361
Gastrulation, 612
Gaucher disease, 88
  osteonecrosis, 463
  osteonecrosis in, 463
Gaussian distribution, 262
G cells, 371
Gemfibrozil, 320
Gemifloxacin, 195
Gender dysphoria, **567**
Gender identity, 635
Gene expression
  modifications, **56**
  regulation, 41
Generalized anxiety disorder (GAD), **563**
  buspirone, 574
  drug therapy for, 572
  Selective serotonin reuptake inhibitors (SSRIs) for, 575
  serotonin-norepinephrine reuptake inhibitors (SNRIs) for, 575
Generalized seizures, 517
Genes
  introns vs exons, 42, 43
Genetics, 56–65
  anticipation, 62
  autosomal dominant diseases, 60
  autosomal recessive diseases, 60
  autosomal trisomies, 63
  bacterial, 130–204, 131
  chromosome disorders, **64**
  code features, **37**
  inheritance modes, 59
  muscular dystrophies, 61
  terms, **56**–57
  trinucleotide repeat expansion diseases, 62
  viral, 162–163
  X-linked recessive disorders, 61
Geniculate nuclei (thalamus), 498
Genital herpes, 184
Genitalia
  ambiguous, 622, 638, 639
  embryology of, 612, **622**
  estrogen and, 630
  male/female homologs, 623
Genital tubercles, 624
Genital ulcers, 184
Genital warts, 184
Genitofemoral nerve, 452
Genitourinary/renal drug reactions, 251
Genome editing
  reverse transcriptase polymerase chain reaction, 52
Genotyping microarrays, 54
Gentamicin, 191
Genu varum (bow legs), 463
Geriatric patients
  atropine in, 241
  Beers criteria in, 247
  changes in, 270
  colonic ischemia and, 386
  colorectal cancer, 388
  common cause of death, 272
  drug-related delirium in, 558, 575
  lipofuscin in, 211
  Medicare for, 272
  nosocomial infections, 185
  osteoporosis, 462
  PPI adverse effects, 399
  respiratory system changes in, 665
  vascular skin tumors, 478
  Zenker diverticulum, 384
Germ cell tumors
  cryptorchidism risk for, 651
  hormone levels in, 653
  testicular, 652
Germinal centers of lymph nodes, 96
Germinal center (spleen), 98
Gerstmann syndrome, 511
Gestational age calculation, 633
Gestational diabetes, 634
Gestational hypertension, 643
GFAP (glial fibrillary acid proteins), 48, 227
  astrocyte marker, 493
  cytoskeletal elements, 48
  stain, 227
Ghrelin, 336, 371
Giant cells
  with Aschoff bodies, 312
  astrocytomas, 525
  Warthin-Finkeldey, 170
Giant cell (temporal) arteritis, 314, 518
  polymyalgia rheumatica, 470
Giant cell tumor, 464
Giardia spp
  fluorescent antibody stain, 125
  watery diarrhea, 179
Giardia lamblia, **155**
Giardiasis, 155
Giemsa stain, 125
  Borrelia, 146
  chlamydiae, 148
Gigantism, 329, 339
Gilbert syndrome, 393, 394
Gingival hyperplasia
  calcium channel blockers, 318
  cyclosporine, 120
  drug reaction and, 250
Gingivostomatitis, 164
Gitelman syndrome, 586
  markers in, 591
Glans penis
  lymphatic drainage of, 624
Glanzmann thrombasthenia, 427
Glaucoma, 242
  atropine, 241
  β-blockers for, 245
  carbachol for, 240
  closed-angle, 240
  diabetes mellitus and, 346
  diagnosis of, 240
  epinephrine for, 242
  open-angle, 240, 242
  pilocarpine for, 240
  types of, **536**
Glioblastoma multiforme, 526
  nitrosoureas for, 441
Glipizide, 353
Glitazones/thiazolidinediones, 353
Global aphasia, 516
Global payment, 271
Globoid cells
  Krabbe disease, 88
Globose nucleus, 499
Globus pallidus externus, 500
Glomerular disorders/disease
  etiology and presentation, 594
  nomenclature, 594
  types of, **595**
Glomerular filtration barrier, **581**
Glomerular filtration parameters, 583
Glomerular filtration rate (GFR), **582**
  ACE inhibitor effects, 610
  glomerular dynamics in, 583
  juxtaglomerular apparatus, 589

(c) ketabton.com: The Digital Library

Glomerulonephritis
azathioprine for, 120
bacterial endocarditis, 311
RBC casts in, 594
Streptococcus pyogenes, 133, 136
Wegener granulomatosis, 315
Glomerulus
dynamics of, **583**
Glomus tumors, 478
Glossitis
B-complex deficiency, 65
megaloblastic anemia, 420
vitamin B_3 deficiency, 67
vitamin B_9 deficiency, 68
Glossopharyngeal nerve (CN IX), 506
blood flow regulation, 296
pharyngeal arch derivative, 620
tongue, 493
Glossoptosis, 620
Glove and stocking neuropathy, 346
GLP-1 analogs, 353
Glucagon, **333**
for β-blocker toxicity, 323
fructose bisphosphatase-2, 76
glucagonomas and, 351
glycogen regulation, 85
insulin and, 333, 334
production of, 331
somatostatin and, 371
somatostatinomas and, 351
Glucagonomas
MEN 1 syndrome, **351**
pancreatic cell tumor, 351
somatostatin for, 354
Glucocerebrosidase
Gaucher disease, 88
Glucocerebroside
in sphingolipidoses, 88
Glucocorticoids
adrenal insufficiency, 349
calcium pyrophosphate deposition disease, 467
diabetes mellitus, 346
fat redistribution with, 250
gout, 467, 487
immunosuppression, 120
myopathy, 250
rheumatoid arthritis, 466
Glucokinase
hexokinase vs, 75
metabolic pathways, 74
Gluconeogenesis
cortisol and, 336
ethanol metabolism and, 72
irreversible enzymes, **78**
metabolic site, 72
pyruvate metabolism and, 77
rate-determining enzyme for, 73
thyroid hormone and, 331
Glucose
ATP production, 74
blood-brain barrier and, 496
clearance, **584**
glycogen metabolism, 86
insulin and, 334
metabolism of, 40
for porphyria, 425
transporters, 334
Glucose-6-phosphatase
dehydrogenase deficiency, **79**
gluconeogenesis, 78
HMP shunt, 79
Von Gierke disease, 87
Glucose-dependent insulinotropic peptide (GIP), 351, **371**, 372
polypeptide (GIP), 334
Glucosuria
glucose clearance, 584
threshold for, 584
Glucuronidation
drug metabolism, 232
Glulisine, 352
Glutamic acid
ammonia transport, 82
classification of, 81
opioid analgesics and, 551
Glutathione
acetaminophen and, 485
in G6PD deficiency, 422
Glutathione peroxidase, 109
free radical elimination by, 210
Glutathione reductase, 109
NADPH and, 75
Gluten-sensitive enteropathy, 381
Gluteus maximus muscle, 453
Gluteus minimus muscle, 451
GLUT transporters, 334
Glyburide, 353
Glycerol
starvation days and, 91
Glycogen
metabolism and storage, 73, **86**
Glycogenesis, 73
Glycogenolysis
rate-determining enzyme for, 73
thyroid hormone and, 331
Glycogen storage diseases, **87**
Glycogen synthase, 73, 86
Glycolysis
arsenic and, 74
hexokinase/glucokinase in, 75
metabolic site, 72
pyruvate metabolism and, 77
rate-determining enzyme for, 73
regulation of, **76**
Glycoprotein IIb/IIIa inhibitors, **438**
Glycoproteins
bacterial pilus/fimbria, 124
HIV, 175
Glycopyrrolate, 241
Glycosylation
collagen synthesis, 50
protein synthesis, 45
GNAQ gene, 525
Goblet cells, 362, 662
Goiter
maternal hypothyroidism from, 341
maternal iodine deficiency, 614
in Riedel thyroiditis, 341
types and causes of, 342
Golfer's elbow, 459
Golgi apparatus
in plasma cells, 409
Golgi tendon organ, 458
Golimumab, 487
Gonadal drainage, **624**
Gonadal mosaicism, 57
Gonadotropin, 646
Gonadotropin-releasing hormone (GnRH)
function of, 328
neurons producing, 498
ovulation, 631
prolactin and, 330
signaling pathways for, 337
spermatogenesis, 628
Gonads
dysgenesis of, 606
Gonococcal arthritis, 468
Gonococci, vs meningococci, 142
Gonorrhea
ceftriaxone, 189
gonococci, 142
STI, 184
Goodpasture syndrome, 50, 596
autoantibody, 115
HLA-DR2, 100
restrictive lung disease, 675
type II hypersensitivity reactions, 112
Good syndrome
paraneoplastic syndrome, **228**
thymoma and, 98
Goserelin, **656**
Gottron papules, 228, 471
Gout, **467**
drug reaction and, 250
drug therapy for, **487**
kidney stones and, 598
Lesch-Nyhan syndrome, 37
loop diuretics and, 608
Von Gierke disease, 87
Gower maneuver/sign, 61
gp41, 203
G-protein-coupled receptors, 236
G-protein-linked 2nd messengers, **238**
Gracilis, 452
Graft-versus-host disease, 119
type IV hypersensitivity, 113
Gram-negative organisms
cell wall structure, **124**
cephalosporins, 189
lab algorithm, **141**
Gram-positive organisms
antibiotic tests, **134**
cell wall structure, **124**
cephalosporins, 189
lab algorithm, **134**
vancomycin, **190**
Gram stain
peptidoglycan layer and, 125
Granular casts
acute tubular necrosis, 602
"muddy brown" in urine, 594
Granulocyte-colony stimulating factor (G-CSF), 337
Granulocytes
morulae, 150
Granulocytopenia
trimethoprim, 194
Granuloma inguinale, 184
Granulomas, 147
macrophages and, 407
in systemic mycoses, 151
TNF-α and, 110
in tuberculosis, 140
Granulomatosis infantiseptica
Listeria monocytogenes, 139
Granulomatosis with polyangiitis (Wegener)
restrictive lung disease and, 675
Granulomatous disease
catalase + organism infections with, 128
excess vitamin D in, 70
hypervitaminosis D with, 464
Granulomatous inflammation, **217**
Granulosa cells
tumors of, 647
Granzyme B
cytotoxic T cells, 101, **102**
extrinsic pathway and, 208
Grapefruit juice and cytochrome P-450, 252
Graves disease
autoantibody, 115
HLA subtype associations, 100
hyperthyroidism, 342
ophthalmopathy, 340
thyroid cellular action in, 331
type II hypersensitivity, 112
Gray baby syndrome, **192**, 204, 250
Gray hepatization, 683
Grazoprevir, 204
Greater omental sac, 361
Great vein of Galen, 503
Green twig (greenstick) fracture, 462
Grief, **562**
Griseofulvin, **200**
cytochrome P-450 interaction, 252
microtubules and, 48
pregnancy contraindication, 204
"Ground-glass" appearance (X-ray), 177, 661
Pneumocystis jirovecii, **154**
Growth factors
tumor suppressor gene mutations and, 46
Growth hormone (GH), 354
diabetes mellitus, 346
ghrelin and, 336
for hypopituitarism, 339
insulin resistance and, **329**
secretion of, 327
signaling pathways for, 337
somatostatin, 339
Growth-hormone-releasing hormone (GHRH)
function of, 328
Growth media properties, 126
Growth retardation
with renal failure, 603
GTPase, 224
GTP (guanosine triphosphate), 77
Guaifenesin, **686**
Guanfacine, 240, 243
Guanosine analogs
mechanism and use, 201
Gubernaculum, 624, 625
Guessing during USMLE Step 1 exam, 23
Guillain-Barré syndrome
acute inflammatory demyelinating polyradiculopathy, 524
Campylobacter jejuni, 145
restrictive lung disease, 675
Schwann cell injury, 494
Schwann cells, 524
Gummas
syphilis, 147, 184
Gustatory hallucinations, 559
Gustatory pathway
cranial nerves in, 532
thalamic relay for, 498
Guyon canal syndrome, 459
Gynecologic procedures
ureteric damage in, 581
Gynecologic tumor epidemiology, **643**
Gynecomastia, 649
antiandrogens for, 658
azoles, 199
cimetidine, 399
SHBG and, 337
spironolactone, 658
tuberoinfundibular pathway, 499

H

H_1 blockers, 251, **686**
H_2
production in tissues, 127

(c) ketabton.com: The Digital Library

H_2 blockers, **399**
H_2O_2 degradation
  catalase and, 128
H_2-antagonist
  naming conventions for, 254
Haemophilus ducreyi
  sexual transmission, 184
Haemophilus influenzae, **142**
  biofilm production, 128
  cephalosporins, 189
  chloramphenicol, 192
  culture requirements, 126
  encapsulation, 127
  IgA protease, 129
  influenza, 169
  meningitis, 179, 180
  penicillins for, 188
  pneumonia, 179
  postviral infection, 179
  rifamycins, 196
  transformation, 130
  type b conjugate vaccine, 127
  unvaccinated children, 186
  vaccine, 142, 180
Hair
  "kinky," 51
  Menkes disease, 51
  vitamin C deficiency, 69
Hairy cell leukemia, 227, **432**
  cladribine for, 440
Hairy leukoplakia, 479
  HIV-positive adults, 177
Half-life equation, 231
Halitosis
  fetor hepaticus, 389
  Zenker diverticulum, 384
Hallucinations, 559
  cocaine, 571
  delirium, 558
  mesolimbic pathway, 499
  pellagra, 67
  postpartum psychosis, 562
  tricyclic antidepressants, 575
Hallucinogen intoxication and withdrawal, 571
Haloperidol, 573
  torsades de pointes, **294**
Halothane, 550
  hepatic necrosis, **249**
Hamartin protein, 224, 525
Hamartomas, **220**, 525
Hamartomatous colonic polyps, **387**
Hamate bone, 449
  fracture of hook, 447
Hamman sign crepitus, 672
Hammer toes, 531
Hand
  distortions of, 451
  injuries of, 459
  muscles of, **450**
  squamous cell carcinoma, 484
Hand-foot-mouth disease, 183
Hansen disease, 141
  animal transmission, 149
  dapsone, 194
  erythema nodosum, 482
Hantavirus, 167
Haptens
  acute interstitial nephritis, 601
  amiodarone as, 323
Haptoglobin, 421
Hardy-Weinberg population genetics, **57**
Hartnup disease, 67
  vitamin B_3 deficiency, 67
Hashimoto thyroiditis, 341
  autoantibody, 115
  HLA subtype association, 100
Hassall corpuscles, 98
Hay fever
  association, 100
HbA_{1c} test, 346
HBcAg (hepatitis B core antigen), 174
HbC disease, 422
  target cells in, 415
HBsAg (hepatitis B surface antigen), 174
HDL (high-density lipoprotein), 94
Headaches, **518**
  adverse effects with drugs, 195, 199, 200
  α-blockers, 244
  bupropion toxicity, 576
  caffeine withdrawal, 570
  Chiari I malformation, 492
  cimetidine, 399
  drug adverse effects, 546
  electroconvulsive therapy, 562
  genital herpes, 184
  giant cell (temporal) arteritis, 518
  glaucoma, 536
  hydralazine, 318
  increased intracranial pressure, 521
  Jarisch-Herxheimer reaction, 148
  lead poisoning, 425
  malaria, 157
  Mucor spp and *Rhizopus* spp, 153
  nitrates, 318
  ondansetron, 400
  pituitary apoplexy, 339
  ranolazine, 319
  Rocky Mountain spotted fever, 150
  sodium-channel blockers, 322
  subarachnoid hemorrhage, **513**, 516
  triptans for, 547
  vasculitides, 314
  venous sinus thrombosis and, 503
Head and neck cancer, **671**
  cetuximab for, 442
Head size
  Paget disease of bone, 463
Healing, wound, 216
Healthcare delivery, 270–273
Healthcare payment models, **271**
Healthcare proxy, 269
Health maintenance organization plan, 271
Hearing loss, 533
  conductive, 49
  cytomegalovirus, 182
  osteogenesis imperfecta, 51
  Paget disease of bone, 463
  sensorineural deafness, 596
Heart
  autoregulation of, 297
  electrocardiograms, 293
  embryology, **281**
  fetal development, 612
  ischemia in, 210
  morphogenesis of, **280**–281
  myocardial action potential, 292
  normal pressures in, 297
  sclerosis of, 473
Heartburn, 377
Heart disease
  congenital, 63, **298–299**
  death, common causes by age, 272
  Fabry disease, 88
  ischemic, **304**
Heart failure, **309**
  ACE inhibitors for, 610
  acromegaly, 339
  amiodarone, 323
  angiotensin II receptor blockers, 610
  aortic regurgitation as precursor, 291
  atrial septal defect, 299
  β-blockers for, 245
  B-type natriuretic peptide in, 296
  calcium channel blockers, 324
  cardiac glycosides for, 321
  chronic ischemic heart disease, 304
  contractility in, 284
  diabetic ketoacidosis, 347
  disopyramide, 322
  dobutamine for, 242
  dopamine for, 242
  Ebstein anomaly, 298
  ESR in, 214
  fludrocortisone and, 354
  hypertension, **300**, **316**
  in sleep apnea, 679
  jugular venous pulse in, 287
  left heart, 309
  Paget disease of bone, 463
  potassium-sparing diuretics, 609
  readmissions with, 272
  right heart, 309
  shock caused by, 310
  thiazides for, 609
  ventricular septal defect, 299
Heart murmurs, **291**
  cardiomyopathies, 308
  patent ductus arteriosus, 299
Heart rate, 243
Heart sounds, 287
  cardiac cycle, 287
  cardiac tamponade, 310
  splitting in, **289**
Heart transplant
  dilated cardiomyopathy, 308
Heart valve development, 281
Heat-labile toxin (LT)
  Clostridium botulinum, 138
  Clostridium perfringens, 138
  Cl^- secretion in gut, 132
Heat shock proteins, 45
Heat-stable toxin (ST)
  resorption of NaCl and H_20 in gut, 132
Heat stroke, 517
Heavy menstrual bleeding (AUB/HMB), 633
Heel pain, 461
Heinz bodies, 79, 414, **416**, 422
Helicase, 38
Helicobacter pylori, **146**
  as oncogenic microbe, 226
  disease association, 379
  metronidazole, 195
  penicillins for, 188
  silver stain, 125
  urease-positive, 127
Heliotrope rash, 228
HELLP syndrome, 643
"Helmet cells," 423
Helminthic infections
  eosinophils and, 408
Helper T cells
  cell surface proteins, 110
  cytokine secretion, 108
Hemagglutinin
  influenza viruses, 169
  parainfluenza viruses, 170
Hemangioblastomas, 526
Hemangiomas, 220
  cavernous (liver), 392
  pyogenic granuloma, 478
  strawberry, 478
Hemarthroses
  hemophilias, 426
  Vitamin C deficiency, 69
Hematemesis, 377
Hematin, 126, 142
Hematochezia
  diverticulosis, 383
  intestinal disorders, 386
  Meckel diverticulum, 384, 618
Hematocrit
  polycythemia vera, 433
Hematologic abnormalities
  laboratory techniques for, 54
Hematologic disorders
  paraneoplastic syndromes, 228
Hematologic drug reactions, 250
Hematology/oncology
  anatomy, 406–409
  pathology, 414–434
  pharmacology, 435–443
  physiology, 410–413
Hematopoiesis, **406**, 432
Hematopoietic stem cells, 110
Hematuria, 595
  bladder cancer, 606
  hereditary hemorrhagic telangiectasia, 316
  interstitial nephritis, 601
  kidney stones, 598
  protease inhibitors, 203
  renal cyst disorders, 604
  renal oncocytoma and, 605
  renal papillary necrosis, 602–610
  Schistosoma haematobium, 161
  transitional cell carcinoma, 606
  UTIs, 181
  Wegener granulomatosis, 315
  Wilms tumor, 606
Heme
  bilirubin and, 375
  chloroquine, 200
  porphyria and, **425**
  synthesis of, **425**
  vitamin B_6 and, 67
Hemianopia, 515, 542
Hemiballismus, 519
  brain lesions and, 511
Hemidesmosome, 474, 480
Hemineglect, 514
Hemiparesis
  saccular aneurysms, 516
Hemochromatosis, **395**
  calcium pyrophosphate deposition disease, 467
  cardiomyopathy with, 308
  chromosome association, 64
  chronic, 426
  free radical injury, 210
  hepatocellular carcinoma and, 392
  HLA-A3 and, 100
Hemoglobin, **665**
  carbon dioxide transport, 670
  development of, 404
  kinetics of, 230
Hemoglobin electrophoresis, **410**
Hemoglobinuria
  acute tubular necrosis and, 602
  G6PD deficiency, 422
  intravascular hemolysis, 421
  paroxysmal nocturnal, 122

(c) ketabton.com: The Digital Library

Hemolysis
alpha toxin, 133
Clostridium perfringens, 138
G6PD deficiency, 250
HELLP syndrome, 643
sulfonamides, 194
Hemolytic anemia, 421
autoimmune, 189
babesiosis, 157
cephalosporins, 189
direct Coombs-positive, 250
extrinsic, 421, **423**
folate deficiency and, 420
G6PD deficiency, 79
intrinsic, 421, **422**
penicillin G, V, 187
pyruvate kinase deficiency and, 422
spherocytes in, 415
sulfa drug allergies, 252
Wilson disease, 395
Hemolytic disease of the newborn, 112, **405**
Hemolytic reactions and blood types
newborns, 405
Hemolytic-uremic syndrome (HUS)
Escherichia coli, **145**, 179
exotoxins, 132
platelet disorders, 427
Hemophagocytic lymphohistiocytosis, **435**, 544
Hemophilia, 426
X-linked recessive disorder, 61
Hemoptysis
Aspergillus fumigatus, 177
bronchiectasis, 675
choriocarcinomas, 642
lung cancer, 684
tuberculosis, 140
Wegener granulomatosis, 315
Hemorrhage
acute pancreatitis, 397
AIDS retinitis, 165
baroreceptors and, 296
bevacizumab, 442
delirium caused by, 558
Ebola virus, 171
intracranial, 513
intraventricular, 512
pulmonary, 137
shock from, 310
subarachnoid hemorrhage, **513**, 516
ulcers, 380
Weil disease, 147
Hemorrhagic cystitis
adenovirus, 164
drug reaction, 251
Hemorrhagic fever
bunyavirus, 167
filovirus, 167
Hemorrhagic stroke, 513
Hemorrhoids, 366
Hemosiderinuria, 421
Hemostasis, 407
platelet plug formation, **411**
Henderson-Hasselbalch equation, 592
Henoch-Schönlein purpura, 315
Hepadnaviruses
characteristics of, 164
genome, 163
Heparin, **436**
acute coronary syndromes, 307
in basophils, 408
in coagulation cascade, 413
mast cells and, 408
osteoporosis, 250
thrombocytopenia, 250
toxicity treatment, 248
warfarin vs, **436**, **437**
Heparin-induced thrombocytopenia (HIT), 436
Hepatic adenomas, 392
Hepatic arteries, 364, 367
Hepatic ascites, 609
Hepatic ducts, 368
Hepatic encephalopathy, **391**
Hepatic fibrosis, 367
Hepatic lipase, 93
Hepatic necrosis, **249**, 485
Hepatic steatosis, 391
Hepatitis
alcoholic, 391, 571
drug reaction and, 249
heroin addiction and, 576
hyperbilirubinemia, 393
Hepatitis A (HAV)
characteristics of, 172
picornavirus, 167, **168**
serologic markers, 174
Hepatitis antigens, 174
Hepatitis B (HBV)
characteristics of, 172
extrahepatic manifestations, 173
hepatocellular carcinomas and, 392
medical importance, 164
nosocomial infection, 185
as oncogenic microbe, 226
passive antibodies for, 110
polyarteritis nodosa and, 314
serologic markers, 174
sexually transmitted infection, 184
Hepatitis C (HCV)
characteristics of, 172
extrahepatic manifestations, 173
flaviviruses, 167
hepatocellular carcinoma and, 392
lichen planus, 482
as oncogenic microbe, 226
therapy for, **204**
Hepatitis D (HDV), 172
Hepatitis E (HEV), 172
hepevirus, 167
Hepatitis viruses, **172**
aplastic anemia, 421
serologic markers for, **174**
Hepatocellular carcinomas, **392**
Aspergillus fumigatus, 153
Budd-Chiari syndrome and, 392
carcinogens causing, 225
cirrhosis and, 389
hemochromatosis, 395
non-alcoholic fatty liver disease, 391
oncogenic microbes, 226
Hepatocytes, 86
Hepatoduodenal ligament, 361
Hepatomas, **392**
Hepatomegaly
Budd-Chiari syndrome, 392
galactosemia, 80
hepatocellular carcinoma, 392
pulmonary hypertension, 668
right heart failure, 309
Von Gierke disease, 87
Zellweger syndrome, 47
Hepatosplenomegaly
β-thalassemia and, 418
biliary tract disease, 395
graft-versus-host disease, 119
hyperchylomicronemia, 94
leishmaniasis, 158
lysosomal storage diseases, 88
mononucleosis, 165
TORCH infections, 182
Hepatosteatosis, 72
Hepatotoxicity
amiodarone, 323
bosentan, 686
danazol, 658
inhaled anesthetics, 550
isoniazid, 197
leflunomide, 486
methotrexate, 440
pyrazinamide, 197
rifamycins, 196
terbinafine, 199
thionamides, 354
valproic acid, 544
zileuton, 687
Hepcidin, 213
in anemia of chronic disease, 421
Hepeviruses
characteristics, 167
genomes, 163
naked viruses, 163
HER2/neu (c-erbB2), 224
"Herald patch" (pityriasis rosea), 482
Herceptin (trastuzumab), 443
Hereditary amyloidosis, 212
Hereditary angioedema, 658
complement disorder and, 107
Hereditary elliptocytosis, 414
Hereditary hemorrhagic telangiectasia, **316**
autosomal dominance of, 60
Hereditary hyperbilirubinemias, **394**
Hereditary motor and sensory neuropathy, 524
Hereditary spherocytosis, 422
spherocytes in, 415
Hereditary thrombosis syndromes, **428**
Hermaphrodites, 639
Hernias, **370**
Herniation syndromes, **529**
Heroin, 551
detoxification medications, 576
intoxication and withdrawal, 570
opioids for withdrawal, 551
Herpes genitalis, 164
Herpes labialis, 164
Herpes simplex virus (HSV), **164–166**
HSV-1/HSV-2, 164, 184
cidofovir, 202
clinical significance, 164
envelope, 163
foscarnet for, 202
guanosine analogs, 201
identification, 166
meningitis caused by, 180
skin infections, 479
STI, 184
TORCH infection, **182**
Herpes zoster
dorsal root latency, 165
famciclovir, 201
reactivation, 443
Herpetic whitlow, 164
Heterochromatin, 34
Heterodimer, 48
Heterodisomy, 57
Heterogeneous nuclear RNA (hnRNA), 41
Heteroplasmy, 57
Heterozygosity loss, 56
Hexokinase
glucokinase vs, **75**
metabolic pathways, 74
"HF" cells (lungs), 309
HFE gene
hemochromatosis and, 395
HGPRT (hypoxanthine guanine phosphoribosyltransferase), 37, 440
Hiatal hernias, 370
Hiccups, 519
High altitude respiratory response, 670
High-frequency recombination (Hfr) cells, 130
Highly active antiretroviral therapy (HAART), 203
High-riding prostate, 627
Hilar lymph nodes
calcification of, 677
Hilar mass, 684
Hilum (lung), 663
lymphadenopathy, 675
Hindbrain, 490
Hindgut
blood supply/innervation of, 364
development of, 358
Hip dislocation
nerve injury with, 453
Hip injuries/conditions
common, **460**
developmental dysplasia, 461
Hip muscles, **451**
Hippocampus
lesions in, 511
limbic system, 499
pyramidal cells, 210
Hippurate test, for *Streptococcus agalactiae*, 137
Hirschsprung disease, **384**
Hirsutism
cyclosporine, 120
danazol, 658
menopause, 636
SHBG and, 337
Histaminase, 408
Histamine blockers, 398, **399**
Histamine receptors, 238
Histamines
in basophils, 408
cortisol effect on, 336
location of, 373
mast cells and, 408
seafood toxins, 247
signaling pathways for, 337
vitamin B_6 and, 67
Histidine, 81
Histiocytosis (Langerhans cell), 434
Histocompatibility complex I and II, 100
Histones
acetylation, 34
amino acids in, 81
deacetylation, 34
methylation, 34
Histoplasma spp
treatment, 199
Histoplasma capsulatum
HIV-positive adults, 177
necrosis and, 209
Histoplasmosis, 151
erythema nodosum, 482
Histrionic personality disorder, 565
HIV (human immunodeficiency virus), **175**
aplastic anemia in, 421

(c) ketabton.com: The Digital Library

HIV (*continued*)
common disease associations, **177**
diagnosis, **175**
flow cytometry diagnosis, 54
hairy leukoplakia, 479
heroin addiction and, 576
Kaposi sarcoma, 165, 478
lymphopenia, 424
meningitis, 180
non-Hodgkin lymphoma and, 429
Pneumocystis jirovecii, 154
primary central nervous system lymphoma and, 430
prophylaxis for HIV patients, 198
pulmonary arterial hypertension, 679
retrovirus, 167
rifamycins in, 196
STI, 184
T cells and, 409
therapy for, **203**
TORCH infection, **182**
untreated time course, 176
viral receptor, 166
HLA-B8
graves disease and, 342
HLA-DR3
graves disease and, 342
HLA-DR4, 466
HLA genes
celiac disease and, 381
disease associations, **100**, 341
DM type 1 association, 347
seronegative spondyloarthritis, 469
HMG-CoA reductase
cholesterol synthesis, 73
metabolic pathways, 74
HMG-CoA synthase, 73
HMP shunt
metabolic site, 72
NADPH production, 75, **79**
rate-determining enzyme, 73
vitamin B_1 deficiency, 66
Hoarseness
gastroesophageal reflux disease, 377
lung cancer, 684
Ortner syndrome, 283
Pancoast tumor, 685
thyroid cancer, 343
Hodgkin lymphoma, **429**
bleomycin for, 439
non-Hodgkin vs, **429**
oncogenic microbes and, 226
paraneoplastic cerebellar degeneration and, 228
vinca alkaloids for, 441
Holistic medical therapy, 269
Holoprosencephaly, **491**
Patau syndrome, 63
Homatropine, 241
Homer-Wright rosettes, 528
Homicide, 272
Homocysteine
folate deficiency, 420
vitamin B_9 deficiency, 68
vitamin B_{12} deficiency, 69, 420
Homocysteine methyltransferase
deficiency in, 84
vitamin B_{12} and, 69
Homocystinuria
causes of, **84**
Homologous recombination repair, 40
Homovanillic acid (HVA)
in neuroblastomas, 350
Homunculus, **502**
Hookworms, 159
Hormone effects on kidney, **590**
Hormone replacement therapy, **657**
estrogens for, 656
for hypopituitarism, 339
thrombotic complications, 250
Hormone-sensitive lipase, 93
Hormones (reproductive), 655
Horn cysts, 477
Horner syndrome, 514, 518, **540**
cavernous sinus, 542
lung cancer, 684
Pancoast tumor, 685
Horse flies (disease vector), 159
Horseshoe kidney, **579**
Hospice care, **272**
Hospital readmission causes, **272**
Hot flashes
drug reaction and, 249
"Hourglass stomach," 370
Howell-Jolly bodies
postsplenectomy, 98
Hu antigens, 228
Human chorionic gonadotropin (hCG)
as tumor marker, 226
choriocarcinomas, 642
ectopic pregnancy, 641
hydatidiform moles, 642
secretion of, 612, **633**
signaling pathways, 337
source and functions of, **633**
Human factors design, **273**
Human herpesvirus 4 (HHV-4), 165
Human herpesvirus 6 (HHV-6), 165, 183
Human herpesvirus 7 (HHV-7), 165
Human herpesvirus 8 (HHV-8), 165, 177
Kaposi sarcoma, 478
as oncogenic microbe, 226
Humanized monoclonal antibodies, 110
Human papillomavirus 6 (HPV-6), 184
Human papillomavirus 11 (HPV-11), 184
Human papillomavirus 16 (HPV-16), 671
Human papillomavirus (HPV)
cervical pathology, 645
HIV-positive adults, 177
as oncogenic microbe, 226
penile cancer, 651
tumor epidemiology, 643
verrucae, 477
warts, 164
Human placental lactogen, **634**
Humerus fracture
axillary nerve and, 447
radial nerve with, 447
Humoral immune response, 409
Hunger/satiety regulation, 498
Hunter syndrome, 60, 61, **88**
Huntington disease
drug therapy for, 549
movement disorders, 519
neurodegenerative disorder, 520
neurotransmitters for, 495
trinucleotide repeat expansion diseases, 62
Hurler syndrome, 88
Hürthle cells, 341
Hutchinson teeth, 147
Hyaline arteriolosclerosis, 301
Hyaline casts (urine), 594
Hydatid cysts, 161
Hydatidiform mole, **642**
hCG in, 633
theca-lutein cysts and, 646
Hydralazine
gestational hypertension, 316
heart failure, 309
mechanism and clinical use, **318**
Hydrocele (scrotal)
acquired, 652
congenital, 652
Hydrocephalus, **522**
childhood tumors, 528
headaches with, 518
posterior fossa malformations, 492
risk for developing, 513
Toxoplasma gondii, 182
Hydrochlorothiazide (HCTZ), 609
for diabetes insipidus, 338
hyperglycemia, 249
pancreatitis, 249
Hydrogen peroxide, 204
Hydronephrosis, **599**
BPH, 654
horseshoe kidney, 579
kidney stones, 598
posterior urethral valves, 579
Hydrophobia, 171
Hydrops fetalis
parvovirus B19, 182, 183
syphilis, 182
Hydroxychloroquine
myopathy, 250
Hydroxylases, 73
Hydroxylation
collagen synthesis, 50
in protein synthesis, 45
Vitamin C and, 50
Hydroxyurea, **442**
polycythemia vera, 433
purine synthesis, 36
sickle cell anemia, 422
targets of, 438
Hyoid artery, 619
Hyoscyamine, 241
Hyperacute transplant rejection, 119
Hyperaldosteronism, **349**
hypertension with, 300
potassium-sparing diuretics for, 609
Hyperammonemia, **82**, 85
fatty acid metabolism and, 89
Hyperbilirubinemia
conjugated (direct), 393
hereditary, 394
jaundice with, 393
unconjugated (indirect), 393
Hypercalcemia
acute pancreatitis and, 397
adult T-cell lymphoma, 430
bisphosphonates for, 486
calcium carbonate antacid effects, 399
diabetes insipidus, 338
hyperparathyroidism, 345
loop diuretics for, 608
lung cancer, 684
paraneoplastic syndrome, 228
succinylcholine, 551
teriparatide, 487
thiazides, 609
Williams syndrome, 64
Hypercalciuria
hyperparathyroidism, 345
thiazides for, 609
Hypercapnia
contractility in, 284
Hypercholesterolemia, 94
familial, 60
Hyperchylomicronemia, 94
Hypercoagulability, 671
hereditary syndromes, 428
in pregnancy, 633
marantic endocarditis in, 311
venous sinus thrombosis with, 503
warfarin adverse effect, 436
Hyperemesis gravidarum, 642
Hyperemia
pseudoephedrine/phenylephrine, 686
Hypereosinophilic syndrome, 308
Hyperestrogenism, 646
Hyperglycemia
Cushing syndrome, 348
diabetic ketoacidosis, 347
diabetic retinopathy, 537
drug reaction and, 249
glucagon and, 333
hyperkalemia, 590
immunosuppressants, 120
pancreatic cell tumors, 351
protease inhibitors, 203
thiazides, 609
vitamin B_3 toxicity, 67
Hypergonadotropic hypogonadism, 639
Hypergranulosis, **475**, 482
Hyper-IgM syndrome, 117
Hyperinsulinemia, 645
Hyperkalemia
aldosterone in, 588
aliskiren, 610
angiotensin II receptor blockers, 610
cardiac glycosides, 321
causes of, 590
depolarizing neuromuscular blocking drugs, 551
diabetic ketoacidosis, 347
potassium-sparing diuretics, 609
primary adrenal insufficiency, 349
Hyperkalemic tubular acidosis (type 4, 593
Hyperkeratosis, **475**, 477
Hyperlipidemia, **301**
atherosclerosis and, 302
immunosuppressants, 120
thiazides, 609
Hyperopia, 535
Hyperosmolar hyperglycemic state, **347**
DM type 2, 346
Hyperosmolarity, 590
Hyperparathyroidism, **345**
calcium pyrophosphate deposition disease, 467
cinacalcet for, 355
lab values in, 464
renal osteodystrophy and, 603
Hyperphagia
depression with, 561
hypothalamus and, 498
Hyperphosphatemia
hyperparathyroidism (secondary), 345
hypoparathyroidism, 344
renal osteodystrophy and, 603
Hyperpigmentation
bleomycin, 439
busulfan, 441
fludrocortisone, 354
hemochromatosis, 395
Peutz-Jeghers syndrome, 387
primary adrenal insufficiency, 349

(c) ketabton.com: The Digital Library

Hyperplasia, 206
  adrenal, 348, 349
  parathyroid, **345**, 351
  uterine bleeding with, 633
Hyperplastic arteriolosclerosis, 301
Hyperplastic polyps, 387
Hyperprolactinemia, 249, 328, **527**
  anovulation, 645
  calcium channel blockers and, 318
  risperidone and, 573
Hyperpyrexia
  with TCAs, 575
Hyperresonance (chest percussion), 682
  pneumothorax, 680, **682**
Hypersensitivity reactions, **112–113**
  acute interstitial nephritis, 601
  C3 deficiency, 107
  cephalosporins, 189
  Graves disease, 342
  IgE antibodies, 105
  mast cells and, 408
  organ transplants, 119
  penicillins, 187–188
  pneumonitis, 675
  rheumatic fever, 312
  sulfonamides, 194
Hypersensitivity reaction (type II)
  rapidly progressive glomerulonephritis, 596
Hypersensitivity reaction (type III)
  acute poststreptococcal glomerulonephritis, 596
Hypersensitivity reaction type IV
  contact dermatitis, 477
Hypersomnia, 561
Hypertension, **300**, 679
  ACE inhibitors for, 610
  acromegaly and, 339
  aliskiren for, 610
  α-blockers for, 244
  angiotensin II receptor blockers for, 610
  aortic dissection and, 303
  atherosclerosis and, 302
  atrial fibrillation and, 295
  autosomal recessive polycystic kidney disease, 604
  β-blockers for, 245
  Charcot-Bouchard microaneurysms, 516
  episodic, 350
  heart failure, 316
  hyperaldosteronism, 349
  immunosuppressants, 120
  intraparenchymal hemorrhage, 513
  leflunomide, 486
  lipohyalinosis and, 514
  local anesthetics, 550
  loop diuretics for, 608
  MDMA, 571
  microangiopathic anemia, 423
  minoxidil, 658
  PCP, 571
  pheochromocytomas, 350
  polyarteritis nodosa, 314
  in pregnancy, 243, **643**
  pseudoephedrine/phenylephrine, 686
  renal cyst disorders, 604
  renal failure, 603
  renovascular disease and, 604
  sleep apnea, 679
  thiazides for, 609
  thoracic aortic aneurysm and, 302
  treatment for, **316**
  tyramine, 244, 575
Hypertensive crisis, 569
  MAO inhibitors, 575
  phenoxybenzamine for, 244
  pheochromocytoma, 350
Hypertensive emergency, 300, **318**
  RBC casts in, 594
Hypertensive nephropathy, 300
Hypertensive retinopathy, **537**
Hyperthermia
  atropine causing, 241
  ecstasy intoxication, 571
  MDMA, 571
Hyperthyroidism, 340, **342**
  amiodarone and, 323
  β-blockers in, 245
  drug reactions, 249
  hCG elevation and, 633
  hydatidiform moles, 642
Hypertriglyceridemia, 94
  acute pancreatitis and, 397
Hypertrophic cardiomyopathy, 308, 531
  Pompe disease, 87
Hypertrophic osteoarthropathy, 684
  cancer association, 228
Hypertrophic pyloric stenosis, **359**
Hypertrophic scars, 218
Hypertrophy, 206
Hyperuricemia
  drug reaction and, 250
  gout and, 467
  kidney stones and, 598
  Lesch-Nyhan syndrome, 37
  pyrazinamide, 197
  thiazides, 609
  vitamin B_3 toxicity, 67
Hyperventilation
  metabolic acidosis and alkalosis, 592
  in pregnancy, 633
  therapeutic, 501
Hypervitaminosis D, 464
Hypnagogic hallucinations, 559
  narcolepsy, 568
Hypnopompic hallucinations, 559
  narcolepsy, 568
Hypoaldosteronism, 593
Hypocalcemia, 333
  acute pancreatitis and, 397
  cinacalcet causing, 355
  hypermagnesemia and, 591
  hyperparathyroidism, 345
  hypoparathyroidism, 344
  renal osteodystrophy, 603
  thyroidectomy, 343
  tumor lysis syndrome, 435
Hypochlorhydria hypergastrinemia, 379
Hypocretin, 568
Hypodermis, 473
Hypofibrinogenemia, 214
Hypogammaglobulinemia, 228
Hypoglossal nerve (CN XII), 506
  lesion in, 532
  with stroke, 514
  tongue, 493
Hypoglycemia
  fructose intolerance, 80
  glucagon production with, 333
  gluconeogenesis and, 78
  insulinomas, 351
  neonatal, 614
  Von Gierke disease, 87
Hypogonadism
  diagnosis of, 639
  estrogens for, 656
  gynecomastia, 649
  hemochromatosis, 395
  Kallmann syndrome, 639
  pituitary prolactinomas, 328
  testosterone/methyltestosterone, 658
  zinc deficiency, 71
Hypokalemia
  antacid use, 399
  causes of, 590
  cystic fibrosis, 60
  on ECG, 293
  loop diuretics, 608
  nephrogenic DI, 338
  VIPomas and, 371
Hypomanic episodes, **561**
Hyponatremia
  MDMA, 571
  as paraneoplastic syndrome, 228
  thiazides, 609
Hypo-osmolarity, 590
Hypoparathyroidism, **344**
Hypophosphatemia
  hyperparathyroidism, 345
Hypopituitarism, **339**
Hypoplasia, 613
  pulmonary, 660
Hypopyon, 536
Hyporeflexia
  LMN lesions, 531
  magnesium hydroxide and, 399
Hypospadias, 624
Hypotension
  adrenal insufficiency, 349
  aliskiren, 610
  amphotericin B, 199
  angiotensin II receptor blockers, 610
  baroreceptors in, 296
  cardiac tamponade, 310
  endotoxins, 131
  ephedrine for, 242
  hypermagnesemia, 591
  local anesthetics, 550
  magnesium hydroxide and, 399
  metronidazole, 195
  midodrine for, 242
  norepinephrine for, 242
  orthostatic, 349
  phenylephrine for, 242
  scombroid poisoning, 247
  sympatholytic drugs and, 243
Hypothalamic/pituitary drugs
  clinical use and adverse effects of, **354**
Hypothalamic-pituitary hormones, **328**
Hypothalamus
  ADH secretion, 329
  homeostasis and, **498**
  nuclei of, 498
  reproductive hormone control, 656
  sleep cycle role of, 497
  TRH sensitivity, 331
Hypothenar muscles, 450
  Klumpke palsy, 448
Hypotheses (statistical), 262
Hypothyroidism, 340, **341**
  amiodarone and, 323
  anemia, 420
  carpal tunnel syndrome with, 459
  drug reaction and, 249
  hormone replacement, 354
  lithium, 574
Hypotonia
  poliomyelitis, 531
  Zellweger syndrome, 47
Hypoventilation, 592
Hypovolemic shock, 310
Hypoxanthine guanine phosphoribosyltransferase (HGPRT), 37
Hypoxemia
  alveolar gas equation, 668
  oxygen deprivation, 669
  vasoconstriction, 679
Hypoxia
  apoptosis caused by, 208
  contractility in, 284
  erythropoietin and, 589
  lung diseases, 679
  nocturnal, 679
  oxygen deprivation, 669
  regions susceptible to, 210
  renal, 666
  vasoconstriction/vasodilation and, 297
Hypoxia inducible factor 1a, 224
Hypoxic stroke, 512
Hypoxic vasoconstriction (pulmonary), 668
  high altitude, 670
Hysterectomy
  cardinal ligament in, 625
Hysteresis (lung and chest wall), 665

I

Iatrogenic abnormal uterine bleeding, 633
Ibandronate, 486
Ibuprofen, 486
  hemolysis in G6PD deficiency, 250
Ibutilide, 323
ICAM-1 protein
  in leukocyte extravasation, 215
  viral receptor, 166
I cells, 371
  disease, 47
Icosahedral viruses, 163
Icterohemorrhagic leptospirosis, 147
Idarucizumab, 435
Idealization, 555
Identification, 555
Idiopathic intracranial hypertension, **521**
Idiopathic pulmonary fibrosis, 675
Idiopathic thrombocytopenic purpura (ITP), 427
  rituximab for, 443
IDL (intermediate-density lipoprotein), 94
IFN-γ (Interferon-γ)
  cachexia and, 227
Ifosfamide, 441
  hemorrhagic cystitis, 251
IgA and IgG deamidated gliadin peptide autoantibody, 115
IgA antibodies, 105
  anti-endomysial autoantibody, 115
  anti-tissue transglutaminase autoantibody, 115
  ataxia-telangiectasia, 117
  breast milk, 636
  in celiac disease, 381
  hyper-IgM syndrome, 117
  passive immunity, 110
  Peyer patches and, 374
  selective deficiency in, 116
IgA nephropathy, 596
IgA protease, 129

(c) ketabton.com: The Digital Library

IgD antibodies, 105
IgE antibodies, 105
  ataxia-telangiectasia, 117
  eczema, 477
  hyper-IgM syndrome, 117
  mast cells and, 408
  type I hypersensitivity, 112
IgG antibodies, 105
  ataxia-telangiectasia, 117
  complement activation and, 106
  hepatitis A (HAV), 174
  multiple sclerosis, 523
  as passive immunity, 110
  type II hypersensitivity reaction, 480
  type III hypersensitivity reactions, 113
IgM antibodies, 105
  in biliary cirrhosis, 395
  complement activation and, 106
  hepatitis A (HAV), 174
  hyper-IgM syndrome, 117
  overproduction, 431
  in sclerosing cholangitis, 395
  splenic dysfunction, 98
Ileum, 362
  basal electric rhythm, 362
Ileus, 386
  bacterial peritonitis (spontaneous), 390
  gallstone, 396
Iliacus, 452
Iliohypogastric nerve, 452
Iliotibial band syndrome, 461
Illness anxiety disorder, 566
Iloperidone, 573
Imatinib, 433, **443**
IMG registration timeframe, 6
Imipenem
  seizures with, 251
Imipramine, 575
Immature ego defenses, 555
Immature teratoma, 647
Immune checkpoint interactions, 222
Immune complex, 113
Immune evasion
  in cancer, 221
Immune responses
  acute-phase reactants, 102
  antibody structure and function, 104–117
  antigen type and memory, 105
  Bordetella pertussis, 143
  cell surface proteins, 110
  complement, 106
  cytokines, 108
  hypersensitivity types, 114–115
  immunoglobulin, 105
  passive vs active, 110
  respiratory burst, 109
  Salmonella/*Shigella* spp, 144
  transfusion reactions, 114
Immune system organs
  cellular components, 99
  lymph nodes, **96**
  thymus, 98
Immunocompromised patients
  acyclovir/famciclovir/valacyclovir, 201
  Candida albicans in, 153
  common organisms affecting, 179
  Cryptococcus neoformans, 153
  Cryptosporidium, 155
  esophagitis in, 377
  fungal infections, 186
  Listeria monocytogenes, 139
  Pneumocystis jirovecii, 154
Immunodeficiency syndromes
  flow cytometry diagnosis, 54
  infections in, **118**
  syndromes, **116**–117
Immunoglobulin A vasculitis, 315
Immunoglobulins
  adaptive immunity and, 99
  breast milk and, 636
  isotypes of, **105**
  for Kawasaki disease, 314
Immunohistochemical stains, 227
Immunology, 96–122
  cellular components, 98
  immune responses, 104–117
  immunosuppressants, **120**–122
  lymphoid structures, 96–98
  pathogen recognition in, 99
Immunomodulator signaling pathways, 337
Immunophenotype assessment, 54
Immunosuppressants
  for aplastic anemia, 421
  targets, **121**
  transplant rejection, **120**
Immunosuppression
  vitamin A deficiency, 66
  vitamin C deficiency, 69
Immunotherapy, 121
Impaired colleague, 269
Imperforate hymen, **644**
Impetigo, 136, 475
  crusts with, 479
Imprinting disorders, 58
Inactivated (killed) vaccine, 111
Incidence vs prevalence, **259**
  in medical error corrections, 259
Inclusions
  Cowdry A, 166
  mulberry-like (morulae), 150
  Negri bodies, 171
  "owl eye," 165
  reticulate bodies, 148
Incomplete penetrance, 56
Incontinence (fecal/urinary), 453
Incus, 533, 620
Incus (ossicles)
  pharyngeal arch derivative, 620
India ink stain, 125
Indicator media, 126
Indinavir
  HIV therapy, 203
Indirect bilirubin, 375
Indirect cholinomimetic agonists, 240
Indirect Coombs test
  unbound antibody detection, 112
Indirect inguinal hernia, 370
Indirect sympathomimetics, 242
Indirect (unconjugated) hyperbilirubinemia, 393
Indomethacin, 486
  for diabetes insipidus, 338
IFN-α (Interferon-α)
  clinical uses, 121
  myopathy, 250
  natural killer cells, 101
Infant development, **635**
Infarction
  blood-brain barrier effects, 496
  of bone, 463
Infarcts
  atherosclerosis, 302
  calcification in, 211
  pituitary, 339
  regions susceptible to, 210
  types of, 210
IFN-β (Interferon-β)
  clinical uses, 121
  natural killer cells, 101
Infections
  ESR in, 214
  in immunodeficiency, 118
Inferior colliculi, 504
Inferior gluteal nerve, 453
Inferior mesenteric artery, **363**, 364
  horseshoe kidney, 579
Inferior oblique muscle, 540
Inferior rectal artery, 366
Inferior rectal vein, 365
Inferior rectus muscle, 540
Inferior sagittal sinus, 503
Inferior vena cava, 360
  diaphragm, 663
  gonadal drainage and, 624
Infertility
  clomiphene, 656
  cystic fibrosis, 60
  ectopic pregnancy, 641
  Kallmann syndrome, 639
  Kartagener syndrome, 49
  leuprolide for, 656
  mumps, 170
  salpingitis, 185
  septate uterus, 623
  varicoceles, 651
IFN-γ (Interferon-γ), **108**, 116
  clinical uses, 121
Infiltrative cardiomyopathy, 308
Inflammasome, 214
Inflammation
  acute, 214
  in atherosclerosis, 302
  cardinal signs, **213**
  chronic, **216**
  ESR in, 214
  Extrinsic (death receptor) pathway, 208
  IL-1 in, 108
  Intrinsic (mitochondrial) pathway, 208
  wound healing, 216
Inflammatory bowel disease (IBD), **382**
  azathioprine for, 440
  colorectal cancer and, 388
  erythema nodosum, 482
  infliximab/adalimumab for, 487
  methotrexate for, 440
  sclerosing cholangitis and, 395
  spondyloarthritis, 469
  therapeutic antibodies, 122
Inflammatory breast disease, **649**, 650
Inflammatory demyelinating polyradiculopathy, 524
Infliximab, 122, **487**
  for Crohn disease, 382
  for ulcerative colitis, 382
Influenza, **169**
  orthomyxovirus, 167
  pneumonia, 683
  treatment/prevention, 201
Informed consent, **265**
Infraspinatus muscle
  Erb palsy, 448
  rotator cuff, 446
Infundibulopelvic ligament, 625
Ingested seafood toxins, **247**
Inguinal canal, **369**
Inguinal hernia, 370
Inguinal ligament, 368, 369
Inguinal triangle, 370
Inhalational injury, **676**
Inhaled anesthetics, **550**
Inheritance modes, 59
Inhibin
  Sertoli cell secretion of, 628
Inhibitors of complement activation, 106
Inhibitory pathway, 500
Injury (unintentional), 272
Innate immune system
  in acute inflammation, 214
Innate immunity, **99**
Inositol trisphosphate (IP_3), 337
Inotropes, 310
Inotropy, 286
INR (international normalized ratio), 426
Insomnia
  barbiturates for, 546
  nonbenzodiazepine hypnotics, 546
  ramelteon for, 547
  stimulants causing, 570
  suvorexant, 547
Inspiratory capacity (IC), 664
Inspiratory reserve volume (IRV), 664
Insulin, **334**
  anabolic effects of, 334
  diabetic ketoacidosis, 347
  fructose bisphosphatase-2 and, 76
  GIP effect on, 371
  glucagon and, 333
  glycogen regulation, 73, **85**
  hypokalemia from, 590
  potassium shifts wit, 590
  in pregnancy, 334
  production of, 329
  secretion of, 334
  somatostatin and, 371
  somatostatinomas and, 351
Insulin deficiency, 590
  diabetes mellitus diagnosis, 346
Insulin-like growth factor 1 (IGF-1)
  acromegaly, 339
  signaling pathways for, 337
Insulinoma, 351
  insulin and C-peptide in, 334
  MEN 1 syndrome, 351
  pancreatic cell tumor, 351
Insulin preparations, 352
Insulin resistance, 633
  acanthosis nigricans and, 482
  acromegaly, 339
  cortisol, 336
  DM type 2, 347
  GH, 329
  non-alcoholic fatty liver disease, 391
  polycystic ovarian syndrome, 645
Insurance
  disregarding in treatment, 269
  Medicare/Medicaid as, 272
  types of plans, 271
Integrase inhibitors, 203
Integrins
  epithelial cells, 474
  viral receptor, 166
Intellectual disability, 557
  autism and, 557
  cri-du-chat syndrome, 64
  Down syndrome, 63
  Lesch-Nyhan syndrome, 37
  Patau syndrome, 63

(c) ketabton.com: The Digital Library

phenylketonuria, 84
Williams syndrome, 64
Intellectualization, 555
Intention tremor, 519
cerebellar lesions, 511
Interdigital tinea pedis, 152
Interferons, **109**
Interferon-γ release assay (IGRA), 140
for tuberculosis, 140
Interleukin 1 (IL-1), 108
cachexia and, 227
endotoxins, 133
Interleukin 2 (IL-2), 108
cyclosporine and, 120
natural killer cells and, 101
sirolimus and, 120
tacrolimus and, 120
Interleukin 2 receptor (IL-2R), 120
Interleukin 3 (IL-3), 108
Interleukin 4 (IL-4), 108
Interleukin 5 (IL-5), 108
Interleukin 6 (IL-6), 108
cachexia and, 227
endotoxins, 133
Interleukin 8 (IL-8), 108
neutrophils and, 406
Interleukin 10 (IL-10), 108
Interleukin 12 (IL-12), 108
natural killer cells and, 101
receptor deficiency, 116
Interleukin receptor modulators
naming conventions for, 254
Intermediate filaments
cytoskeletal element, 48
Intermenstrual bleeding (IMB), 633
Internal capsule
intraparenchymal hemorrhage, 513
stroke effects, 514
Internal carotid artery
cavernous sinus, 542
Internal hemorrhoids, 366
Internal iliac artery, 282
Internal iliac lymph nodes, 624
Internal inguinal ring, 370
Internal jugular vein, 503
Internal oblique muscle, 369
Internal rotation
arm (rotator cuff), 446
hip, 451
Internal spermatic fascia, 369
International Foundations of Medicine (IFOM), 12
Internuclear ophthalmoplegia, 511, **543**
Interossei muscles, 450
Klumpke palsy, 448
ulnar nerve, 447
Interpersonal therapy, 572
Interpreting study results, 261
Intersex, 639
Interstitial fluid, 297
Interstitial lung disease, 466, 675
Interstitial nephritis
acute, 601
as drug reaction, 251
NSAID toxicity, 486
penicillins, 188
Interstitial pneumonia, 683
Interstitium
leukocyte extravasation and, 215
Interventricular foramen, 281
Interventricular septal rupture, 305, **307**
"Intestinal angina," 386
Intestinal atresia, **359**
Intestinal gastric cancer, 379
Intestinal obstruction
hernias, 370
superior mesenteric artery syndrome, 363
Intimate partner violence, 269
Intoxication (psychoactive drugs), 570
Intracellular bacteria, **127**
Intracellular fluid (ICF), 581
Intracellular receptors
endocrine hormone and, 337
Intracranial hemorrhage, **513**
eclampsia, 643
Intracranial hypertension
vitamin A toxicity, 66
idiopathic, **521**
Intracranial pressure (ICP)
cerebral ischemia, 296
hydrocephalus, 522
papilledema, 538
superior vena cava syndrome, 685
venous sinus thrombosis, 503
Intraductal papilloma, 649
Intraepithelial adenocarcinoma, 644
Intraocular pressure (IOP), 536
Intraparenchymal hemorrhage, 513
Intrauterine device (IUD)
copper, 657
Intrauterine growth restriction (IUGR)
low birth weight, 635
substance abuse, 614
Intravascular hemolysis, 421
paroxysmal nocturnal hemoglobinuria, 107
Intravenous anesthetics, **550**
Intraventricular hemorrhage, 512
neonatal respiratory distress syndrome, 661
Intrinsic factor, **372**, 373
Intrinsic hemolytic anemias, **422**
Intrinsic pathway, 208
coagulation defects of, 426
heparin and, 437
Intrinsic renal failure, 601
Introns
splicing out, 41
vs exons, **43**
Intussusception, **385**
Meckel diverticulum, 384
Inulin
glomerular filtration rate and, 582
in proximal convoluted tubules, 587
Inulin clearance, 582
Invariant chain, 100
Invasive carcinoma, 219
cervix, 645
Invasive lobular carcinoma (breast), 650
Inversion, 453
In vivo biofilm-producing bacteria, **128**
Involuntary treatment, 267
Iodine
deficiency in, 341, 342
infection control, 204
teratogenicity, 614
Iodophors, 204
IPEX syndrome, 102
Ipratropium, 241, **687**
Irinotecan/topotecan, **442**
targets of, 438
topoisomerase (TOP) I inhibition, 38
Iritis, 536
Iron
absorption of, 69, 374
anemia, 388, **418**
anemia of chronic disease, 421
colorectal cancer, 388
excess, 67
in hemochromatosis, 395
lab values in anemia, 419
sideroblastic anemia, 419
toxicity of, 69
toxicity treatment, 248
Iron poisoning, **426**
Iron studies
interpretation of, 419–444
Irritable bowel syndrome (IBS)
antispasmodic drugs, 241
criteria and symptoms for, **383**
Isavuconazole
mucormycosis treatment, 153
Ischemia, **210**, 669
acute tubular necrosis from, 602
atherosclerosis, 302
colonic, 386
digital, 472
in gastrointestinal tract, 386
mesenteric, 386
necrosis and, 209
watershed areas, 210
Ischemic brain disease, **512**
Ischemic heart disease
contraindicated antiarrhythmics, 322
heart murmurs in, 291
manifestations of, **304**
Ischemic priapism, 651
Islet cell cytoplasmic antibodies, 115
Islets of Langerhans, 325
Isocarboxazid, 575
Isocitrate dehydrogenase
metabolic pathways, 74
rate determining enzyme, 73
Isodisomy, 57
Isoflurane, 550
Isolation of affect, 555
Isoleucine
classification of, 81
maple syrup urine disease and, 84
Isoniazid, **197**
cytochrome P-450, 252
hemolysis in G6PD deficiency, 250
hepatitis, 249
Mycobacterium tuberculosis, 196
seizures, 251
Isoproterenol
norepinephrine vs, 243
sympathomimetic action, 242
Isosorbide dinitrate, 318
Isosorbide mononitrate, 318
Isotretinoin
cystic acne, 66
teratogenicity, 614
Isovolumetric contraction, 287
Isovolumetric relaxation, 287
Itraconazole
azoles, 199
Sporothrix schenckii, 154
systemic mycoses, 151
Ivabradine, **324**
IV drug use
common organisms, 179
Ivermectin, 200
"Ivory white" plaques, 677
IV phlebitis, 199
Ixodes ticks, 146, 149

J

JAK2 gene, 224
myeloproliferative disorders, 433
Janeway lesions, 311
Jarisch-Herxheimer reaction, **148**
Jaundice, **393**
biliary tract disease, 395
cholangitis, 368, 397
drug reaction and, 249
fructose intolerance, 80
galactosemia, 80
graft-versus-host disease, 119
hepatitis B, 184
hepatocellular carcinoma, 392
hereditary hyperbilirubinemias, 394
with leptospirosis, 147
newborn hemolytic disease, 405
pancreatic cancer, 398
TORCH infections, 182
yellow fever, 168
Jaw jerk reflex, 507
JC virus (John Cunningham virus)
HIV-positive adults, 177
immunocompromised patients, 118
polyomaviruses, 164
Jejunal and ileal atresia, 359
Jejunum, 362
Jervell and Lange-Nielsen syndrome, 294
Jimson weed, 241
Job syndrome, 116
Jod-Basedow phenomenon, 342
Joint hypermobility, 51, 62
J point in ECG, 293
Jugular foramen, 503
Jugular venous distention (JVD), **309**, 685
Jugular venous pulse, 287
Justice (ethics), 265
Juvenile polyposis, 387
Juxtaglomerular apparatus (JGA), **589**
renin secretion, 588
Juxtaglomerular cells
tumors in, 349

K

Kala-azar, 158
Kallikrein
C1 esterase inhibitor deficiency, 107
neutrophil chemotaxis and, 406
Kallmann syndrome, 498, **639**
Kaposi sarcoma, 478
AIDS and, 184
bacillary angiomatosis vs, 478
HHV-8, 165
HIV-positive adults, 177
oncogenic microbes and, 226
Kartagener syndrome, **49**, 280
obstructive lung disease, 675
Karyotyping, **55**
KatG, 197
Kawasaki disease, 314
Kayser-Fleischer rings, 395
K cells, 371
K complexes/sleep spindles, 497
Kegel exercises, 599
Keloid scars, 218
Keratinocytes, 216
Keratin pearls, 684
Keratoacanthoma, 484
Keratoconjunctivitis, 164

(c) ketabton.com: The Digital Library

Keratoconjunctivitis sicca, 468
Keratomalacia, 66
Keratosis
  actinic, 482
  hyperkeratosis, 475
  parakeratosis, 475
  seborrheic, 477
Kernicterus, 194, 204, **393**–394
Ketamine, 550
Ketoacidosis, 72, **90**
Ketoconazole, 658
  cytochrome P-450, 252
  gynecomastia from, 649
  mechanism and clinical use, 199
Ketogenesis
  diabetic ketoacidosis, 347
  metabolic site, 72
  rate-determining enzyme for, 73
Ketone bodies, **90**
  brain metabolism, 334
  in diabetic ketoacidosis, 347
  production of, 90
Ketorolac, 486
Kidney disease
  acute injury, 601
  hypertension, 300
  prenatal diagnosis of, 579
Kidneys
  anatomy, **580**
  blood flow regulation, 297
  calcification in, 211
  chronic graft nephropathy, 119
  embryology of, **578**
  endocrine functions, **589**
  glomerular structure, **580**
  hormones acting on, 590
  ischemia in, 210
  retroperitoneal location of, 360
  sclerosis, 473
  solitary functioning, 579
  transplant prophylaxis, 120
Kidney stones
  Crohn disease association, 382
  electrolyte disturbances, 591
  hematuria with, 594
  hydronephrosis, 599
  hyperparathyroidism, 345
  presentation and findings with, **598**
  risk factors for, 593
  UTIs, 181
Kiesselbach plexus, 671
Killian triangle, 384
Kinases, 73
Kinesin
  movement of, 48
Kinin cascade/pathways, 412
Klebsiella pneumoniae
  cephalosporins, 189
  encapsulation, 127
  UTIs caused by, 181
Klebsiella spp, 145
  alcoholism, 179
  currant jelly sputum, **145**, 186
  nosocomial infections, 185
  pneumonia, 683
  urease-positive, 127
  urinary tract infections, 600
Klinefelter syndrome, 638
  chromosome association, 64
  gynecomastia, 649
Klumpke palsy, 448
Klüver-Bucy syndrome, 511
Knee examination, **454**
Knee injuries/conditions
  Baker cyst, 460
  iliotibial band syndrome, 461
  ligament and meniscus, 460
  Osgood-Schlatter disease, 461
  prepatellar bursitis, 460
Knees
  common conditions of, **460**
Knock-out/Knock-in genes, 56
KOH preparation, 152
Koilocytes
  condylomata acuminata, 184
Koilocytosis, 477
Koplik spots, 183
Korsakoff syndrome, 66, 558, **571**
Krabbe disease, **88**, 524
KRAS gene, 224
  adenomatous colonic polyps and, 387
  lung cancer and, 684
Krukenberg tumors, 379
Kübler-Ross grief model, 562
Kulchitsky cells, 684
Kuru, 178
Kussmaul respirations
  in diabetic ketoacidosis, 347
Kussmaul sign, **316**
Kwashiorkor, 71
Kyphoscoliosis, 531
Kyphosis
  in homocystinuria, 84
K_m, 230

L

Labetalol, 245
  hypertension in pregnancy, 316
  hypertensive emergency, 318
Labia, 625
Labile cells, 46
Lachman test, 454
Lac operons, **40**
Lacrimation reflex, 507
Lactase deficiency, **81**
Lactation, **636**
  oxytocin's role in, 328
  progesterone and, 630
  prolactin and, 330
  Sheehan syndrome and, 339
Lactational mastitis, 649
Lactic acid dehydrogenase, 77
Lactic acidosis
  ethanol metabolism and, 72
  exercise and, 670
  MELAS syndrome, 59
  pyruvate dehydrogenase complex deficiency, 77
Lactoferrin
  in neutrophils, 406
  in respiratory burst, 109
Lactose-fermenting enteric bacteria, 126, **144**
Lactose intolerance, 381
Lactose metabolism
  genetic response to environmental change, 40
Lactulose
  for hepatic encephalopathy, 391
Lacunar infarcts, 514
Ladd bands, 385
Lambert-Eaton myasthenic syndrome, 472
  autoantibody, 115
  as paraneoplastic syndrome, 228
  small cell lung cancer, 684
Lamina propria
  Peyer patches in, 374
  in Whipple disease, 381
Lamins, 48
Lamivudine, 203
  HIV therapy, 203
Lamotrigine
  epilepsy, 544
  rash caused by, 250
Lancet-shaped diplococci, 136
Landmarks (anatomical)
  for dermatomes, **510**
  midclavicular line, 663
  pudendal nerve block, 453
Langerhans cell histiocytosis, **434**
  pulmonary, 675
Lansoprazole, 399
Laplace law, **284**, 661
Large cell carcinoma, 684
Large-vessel vasculitis
  presentation and pathology, 314
Larva migrans, 159
Laryngopharyngeal reflux, 377
Laryngospasm, 359
  drug-induced, 569
Larynx, 662
Larynx muscles, 620
Lassa fever encephalitis, 167
Latanoprost, 552
Latent errors, 274
Lateral cerebellar lesions, 499
Lateral collateral ligament (LCL) injury, 454
Lateral corticospinal tract, 508, **509**, 514
Lateral epicondylitis, 459
Lateral femoral cutaneous nerve, 452
Lateral geniculate nucleus (LGN), 498
Lateral medullary syndrome, 514
Lateral nucleus (hypothalamus), 498
Lateral pterygoid muscle, 507
Lateral rectus muscle, 540
Lateral spinothalamic tract, 508
Lateral thoracic artery, 455
Lateral ventricles
  optic radiation, 542
  ventricular system, 504
Laxatives, **401**
LD50 (lethal median dose), 234
LDH
  tumor burden indicator, 226
LDL (low-density lipoprotein), 94
  PCSK9 enzyme, 93
  receptor binding, 93
Leaden paralysis, 561
Lead poisoning
  anemia with, 419
  foot drop, 419
  mechanism, 425
  memory loss with, 425
  sideroblastic anemia, 419
  treatment of, 248, **419**
Lead-time bias, 261
Leber hereditary optic neuropathy, 59
Lecithinase, 133, 138
Lecithin-cholesterol acetyltransferase (LCAT)
  activation of, 93
Lecithin-cholesterol acyltransferase, 93
Lectin pathway (complement activation), 106
Ledipasvir, 204
Leflunomide, **486**
  dihydroorotate dehydrogenase inhibition, 36
Left anterior descending artery
  coronary circulation, 283
  myocardial infarction and, 305
Left bundle branch, 293
Left circumflex coronary artery, 283
Left heart disease, 679
Left heart failure, 309
Left horn of sinus venosus, 281
Left main coronary artery, 283
Left marginal artery, 283
Left shift, 424
Left-to-right shunts, 299
Legg-Calvé-Perthes disease, **461**, 463
Legionella pneumophila, **143**
Legionella spp
  atypical organism, 179
  culture requirements, 126
  facultative intracellular organisms, 127
  Gram stain of, 125
  macrolides, 193
  nosocomial infection, 185
  intracellular organism, 127
  pneumonia, 683
  silver stain, 125
Legionnaires' disease, 143
Leiomyoma (fibroid), 648
  nomenclature for, 220
  uterine bleeding with, 633
Leiomyosarcoma, 220, **648**
Leishmaniasis, **158**, 200
Length-time bias, 261
Lens
  collagen in, 50
  infantile cataracts, 80
Lens subluxation
  Elastin syndrome, 52
  in homocystinuria, 52, **84**
Lenticulostriate artery, 514
Lentiform nucleus, 500
Leonine facies, 141
Leprosy (Hansen disease), **141**
Leptin, 336
  hypothalamus, 498
Leptospira spp
  Gram stain of, 125
  spirochete, 146
  zoonotic infections, 149
Leptospira interrogans, **147**
Leptospirosis, 149
  symptoms of, 147
Lesch-Nyhan syndrome
  inheritance, 61
  purine salvage deficiency, 37
Leser-Trélat sign, **228**, 477
  stomach cancer, 379
Lesser omental sac, 361
Letrozole, 656
Leucine
  classification of, 81
  maple syrup urine disease and, 84
Leucovorin, 440
Leukemias, **432**
  carcinogens, 225
  cell type, 220
  cyclophosphamide for, 441
  cytarabine for, 440
  doxorubicin for, 439
  epidemiology, 222
  etoposide/teniposide for, 442
  immunohistochemical stain for, 227
  lymphoma comparison, **429**
  mucormycosis, 153
  nomenclature for, 220
  oncogenic microbes, 225
  suppressor genes, 224
  TRAP tumor marker, 227
  vinca alkaloids for, 441

(c) ketabton.com: The Digital Library

Leukocoria, **538**
Leukocyte adhesion deficiency, **117**, 215
Leukocyte alkaline phosphatase (LAP), 406
Leukocyte esterase, 181, 600
Leukocytes
  extravasation, 214, **215**
  leukemias, **432**
  in urine, **181**, 594, 600
Leukocytoclastic vasculitis, 173
Leukocytosis, 213
  diabetic ketoacidosis, 347
  nosocomial infections, 185
Leukodystrophies, 494, 524
Leukoerythroblastic reaction, 424
Leukopenias, **424**
  cytarabine, 440
  ganciclovir, 202
  trimethoprim, 194
Leukoplakia, 479
Leukotrienes
  basophils and, 408
  cortisol effects, 336
Levator veli palatini muscle, 620
Levetiracetam, 544
Levodopa, 548, **549**
Levofloxacin
  fluoroquinolones, 195
  Pseudomonas aeruginosa, 143
Levomilnacipran, 575
Levonorgestrel, 657
Levothyroxine, **354**
Lewy bodies, 520, 521
  dementia, **521**
Leydig cells
  cryptorchidism, 651
  endocrine function, **628**, 638
  genital embryology, 622
  tumors of, 653
LFA-1 antigens, 215
Libido
  in geriatric patients, 270
  testosterone and, 636
Libman-Sacks endocarditis, 470
Lice
  disease vectors, 149
  head/scalp, 161
  treatment, 200
Lichen planus, 173, 475, **482**
Lichen sclerosus, 644
Lichen simplex chronicus, 644
Liddle syndrome, 586
  markers in, 591
Lidocaine, 322, 550
Life support
  withdrawal, 269
Li-Fraumeni syndrome
  osteosarcomas, 465
  tumor suppressor genes in, 46, 224
Ligaments
  female reproductive anatomy, 625
  gastrointestinal anatomy, 361
Ligamentum arteriosum, 282
Ligamentum teres hepatis, 282, **361**
Ligamentum venosum, 282
Ligand receptors, 208
Lightheadedness, 534
Likelihood ratio (LR), **257**
Limbic system, **499**
Limited scleroderma (CREST syndrome), 115, 473
Linagliptin, 353
Lindane, 200
Linea alba, 369
Linear ulcers, 377
Linear viruses, 163
Lines of Zahn, 672
Lineweaver-Burk plot, 230
Linezolid, **193**
  highly resistant organisms, 198
  protein synthesis inhibition, 191
Lingula (lung), 663
Linkage disequilibrium, 57
Liothyronine (T3), **354**
Lipase
  pancreatic secretions, 373
  in pancreatitis, 397
Lipid-lowering agents
  mechanism and adverse effects, **320**–321
Lipid metabolism
  fatty acids, 73
Lipids
  key enzymes in, 93
  transport of, **92**–93
Lipodystrophy
  protease inhibitors, 203
  tesamorelin for, 328
Lipofuscin, **211**
Lipoic acid, 76
Lipolysis
  cortisol and, 336
  insulin and, 334
  sympathetic receptors and, 238
  thyroid hormone and, 331
Lipomas, 220
Lipophilic drug
  drug metabolism of, 232
Lipoprotein lipase, 93
Lipoproteins, 93
  functions of, **94**
Liposarcomas, 220
Lipoteichoic acid
  cytoplasmic membrane, 124
Liquefactive necrosis, 209
Liraglutide, 353
Lisch nodules, 525
Lisdexamfetamine, 572
Lisinopril, 610
Lispro, 352
Lissencephaly, **491**
Listeria monocytogenes
  β-hemolysis, 135
  neonates, 182
  penicillins for, 188
  transmission of, **139**
Listeria spp
  facultative intracellular organisms, 127
  intracellular organism, 127
Lithium
  diabetes insipidus and, 249, 338
  hypothyroidism, 341
  mechanism and use, **574**
  prenatal exposure, **298**, 300
  teratogenicity, 614
  therapeutic index of, 234
  thyroid functions with, 249
  toxicity of, 569
Live attenuated vaccines, 111
Liver
  blood supply and innervation of, 364
  in gastrointestinal anatomy, 361
  lipid transport and, 92
  tissue architecture, **367**
Liver/biliary disease
  acanthocytes in, 414
  alcoholic, 391
  anemia, 420
  autoimmune, 389, 392, 395
  cirrhosis, 71, 80, **389**
  cystic fibrosis, 60
  hepatosteatosis, 72
  hereditary, 394
  ischemia in, 210
  metastases to, 223
  serum markers, 390
  target cells in, 415
  Wilson disease and, 395
Liver failure
  Budd-Chiari syndrome and, 392
  movement disorder in, 519
Liver fluke
  hyperbilirubinemia with, 393
  as oncogenic microbe, 226
Liver function tests
  cholestatic pattern of, 395
  serum markers for, 390
Liver markers
  in alcohol use, 570
Liver tumors, 392
Living wills, 266
Loa loa, **159**
Loading dose, 231
Lobar pneumonia
  natural history of, 683
  organisms and characteristics, 683
  physical findings with, 680
Lobular carcinoma (breast), 650
Local anesthetics, **550**
Localized amyloidosis, 212
Locked-in syndrome
  osmotic demyelination syndrome, 524
  stroke, 515
Locus ceruleus, 495
Locus heterogeneity, 57
Löffler endocarditis, 308
Löffler medium, 126
  Corynebacterium diphtheriae, 139
Lomustine, 441
Lone Star tick (disease vector), 149
Long-chain fatty acid (LCFA)
  metabolism of, 89
Long QT syndrome
  congenital, 294
  ranolazine, 319
  sudden cardiac death, 304
Long thoracic nerve
  arm abduction, 446
  neurovascular pairing, 455
Loop diuretics, **608**
  for heart failure, 309
  toxicity of, 251
Loop of Henle, 608
  Bartter syndrome and, 586
  ethacrynic acid effect on, 608
"loose associations," 559
Looser zones (osteomalacia), 463
Loperamide, **400**, 551
Lopinavir
  HIV therapy, 203
Loratadine, 686
Lorazepam
  alcohol withdrawal, 572
Losartan, 610
Low birth weight, **635**
Löwenstein-Jensen agar/medium, 126
Lower esophageal sphincter (LES)
  achalasia and, 376
  in Barrett esophagus, 377
  nitric oxide and, 371
Lower extremity nerves, **452**–453
Lower left quadrant (LLQ) pain, 383
Lower motor neuron (LMN) lesions, 530, 531
  facial nerve, 532
LPS endotoxin, 124, 131, **133**, 145
LTB_4 (Leukotriene B4), 406, 485
Lumbar puncture, 507, 521
Lumbosacral radiculopathy, 455
Lumbrical muscles, 450
  Klumpke palsy and, 448
  median and ulnar nerves, 447
Lumefantrine, 200
Lunate bone, 449
Lung abscesses, **685**
Lung and chest wall expansion, 665
Lung cancer
  asbestosis and, 677
  carcinogens causing, 225
  cisplatin/carboplatin for, 442
  erlotinib for, 442
  hypercalcemia and, 228
  incidence/mortality in, 222
  metastases to, 223
  non–small cell, 684
  oncogenes and, 224
  paraneoplastic syndromes and, 228
  presentation and complications, **684**
  small cell, 684
  topotecan for, 442
Lung diseases
  obstructive, 674–675
  restrictive, 675
Lungs
  anatomical relationships, **663**
  blood flow regulation, 297
  development of, **660**
  physical findings, 680
  sclerosis of, 473
  transfusion-related injury, 114
Lung volumes, **664**
Lung zones, 669
Lupus
  autoimmune hemolytic anemia and, 423
  azathioprine for, 440
  drug-induced, 115
  isoniazid, 197
  lymphopenia, 424
  marantic endocarditis in, 311
  microangiopathic hemolytic anemia,, 423
  nephritis, 470
  neutropenia, 424
Lupus anticoagulant, 115
Lupus-like syndrome
  α-methyldopa, 243
  hydralazine, 318
  procainamide, 322
Lurasidone, 573
Luteal phase of menstrual cycle, 632
Luteinizing hormone (LH)
  clomiphene effect, 656
  contraception, 657
  estrogen/progesterone, 630
  hCG and, 617
  ovulation, 631
  PCOS, 645
  premature ovarian failure, 636
  secretion of, 327
  sex development disorders, 639
  signaling pathways of, 337
  spermatogenesis, 628
  testosterone, 658
Lyme disease
  animal transmission, 149
  AV block in, 295
  Borrelia burgdorferi, **146**
  ceftriaxone, 189

(c) ketabton.com: The Digital Library

Lymphadenopathy
Corynebacterium diphtheriae, 132
hilar, 675–676
Lymphogranuloma venereum, 184
mediastinal, 676
mononucleosis, 165
rubella, 169, 182–183
serum sickness, 113
syphilis, 147, 184
tinea capitis, 152
Toxoplasma gondii, 182
Trypanosoma brucei, 156
in viral infections, 96
Lymphatic filariasis (elephantiasis)
Wuchereria bancrofti, 159
Lymph drainage
gonadal, 624
superficial inguinal nodes, 624
Lymph nodes
absent or scanty, 116
drainage sites, 96–**97**
structure and function, **96**
T cell differentiation, 102
tumor metastases, 223
Lymphocyte-depleted lymphoma, 429
Lymphocyte-rich lymphoma, 429
Lymphocytes, **409**
breast milk and, 636
CLL/small cell lymphocytic lymphoma, 432
corticosteroid effect on, 424
lymph nodes, 96
non-Hodgkin lymphoma, 430
spleen, 98
thymus, 98
Lymphocytic choriomeningitis virus (LCMV), 167
Lymphocytic infiltrates
Bordetella pertussis, 143
Lymphocytosis, 98
Lymphogranuloma venereum, 184
Chlamydia trachomatis, 149
Lymphoid hyperplasia, 383
Lymphoid neoplasms, 432
Lymphoid structures, 96–97
Peyer patches, 362, 374
Lymphomas
carcinogens causing, 225
celiac disease and, 381
cyclophosphamide for, 441
cytarabine for, 440
doxorubicin for, 439
EBV and, 165
etoposide/teniposide for, 442
Hodgkin, 429
hypercalcemia and, 228
leukemia comparison, 429
methotrexate for, 440
nomenclature for, 220
non-Hodgkin, 430
oncogene for, 208, 224
oncogenic microbes, 226
paraneoplastic syndromes with, 228
of stomach, 379
testicular, 653
Lymphopenias, 424
ataxia-telangiectasia, 117
corticosteroid effect on, 424
Lynch syndrome, **388**
mismatch repair and, 40
Lyonization (x-inactivation)
Barr body formation, 61
Lysergic acid diethylamide (LSD), 571
Lysine
classification of, 81
in cystinuria, 85
kidney stones, 598
for pyruvate dehydrogenase complex deficiency, 77
Lysogenic phage infection, 130
Lysosomal storage diseases, 47
causes and effects of, **88**
Lysosomal trafficking regulator gene, 117
Lysosomal α-1,4-glucosidase, 87
Lysozyme
innate immunity, 99
in neutrophils, 406
LYST gene, 117
Lytic bone lesions
adult T-cell lymphoma and, 430
Langerhans cell histiocytosis, 434

M

MacConkey agar, 126, 144
"Machine-like" murmur, 291
Macroangiopathic hemolytic anemia, 423
Macrocytic anemia, **420**
Macroglobulinemia, 431
Macrolides
cytochrome P-450 and, 252
hypertrophic pyloric stenosis and, 359
Legionella pneumophila, 143
mechanism and use, **193**
Mycoplasma pneumoniae, 150
naming convention for, 253
protein synthesis inhibition, 191
torsades de pointes, 248
Macroorchidism, 62
Macro-ovalocytes, 415
Macrophages, **407**
alveolar, 662
apoptosis and, 208
binding of, 104
breast milk and, 636
cell surface proteins, 110
cytokine secretion, 108
endotoxin activation, 133
innate immunity, 99
in lymph node, 96
lymphocyte interaction, **102**
in MI, 305
necrosis and, 209
pneumoconioses, 677
splenic, 98
in wound healing, 216
Macrosomia, 614
Macula densa
juxtaglomerular apparatus, 589
Macular cherry-red spot, 88, 538
Macular degeneration, 536
Macules, 475
melanocytic nevus, 477
Maculopapular rash
graft-versus-host disease, 119
measles, 170
syphilis, 147
Magnesium
antacid use, 399
antiarrhythmic treatment, 324
cardiac glycoside toxicity, 321
in laxatives, 401
PTH regulation, 332
torsades de pointes and, 294
Magnesium hydroxide, 399
Magnesium sulfate
preeclampsia/eclampsia, 643
Maintenance dose, 231
Maintenance stage, substance addiction, 568
Major basic protein (MBP), 408
Major depressive disorder (MDD)
diagnostic symptoms for, **561**
peripartum onset, 562
vortioxetine use, 576
Malabsorption syndromes, **381**–382
fat-soluble vitamin deficiencies, 65
inflammatory bowel disease, 382
Malaria
artesunate for, 200
Plasmodium, 157
quinidine/quinine for, 200
Malassezia spp, 152, 476
Malathion, 200
Male/female genital homologs, **623**
Male genital embryology, 622
Male reproductive anatomy, **626**
Male sexual response, 627
Malformation, 613
Malignancy
marantic endocarditis in, 311
uterine bleeding with, 633
Malignancy/hyperplasia
uterine bleeding with, 633
Malignant hypertension
microangiopathic hemolytic anemia, 423
Malignant hyperthermia, 550–551
Malignant mesothelioma, 227
Malignant tumors, 220
bones, 465
Malingering, **566**
Malleus (ossicles), 533, 620
Mallory bodies
in alcoholic hepatitis, 391
Mallory-Weiss syndrome, 377
Malnutrition, 71
superior mesenteric artery syndrome and, 363
Malrotation, **385**
"Maltese cross" appearance, 157, 594
MALT lymphomas
Helicobacter pylori, 146
oncogenic microbes and, 226
Sjögren syndrome, 468
Mammary glands, 613
Mammillary bodies, 511
limbic system, 499
Wernicke-Korsakoff syndrome, 571
Mandibular process, 620
Mango flies (disease vector), 159
Manic episode, **560**
Man-in-the-barrel syndrome, 502
Mannitol, **607**
Mantle cell lymphomas, 430, 434
Mantle zone
spleen, 98
Maple syrup urine disease, **84**
Marantic endocarditis, 228, 311
Marasmus, 71
Maraviroc, 203
Marburg hemorrhagic fever, 167
Marcus Gunn pupils, **539**
multiple sclerosis, 523
Marfanoid habitus
homocystinuria, 84
MEN 2B syndrome and, 351
Marfan syndrome
aortic dissection and, 303
cardiac defect association, 300
cataracts, 535
chromosome association, 64
elastin and, 52
heart murmur with, 291
thoracic aortic aneurysm and, 302
Marginal zone lymphoma, 430, 434
Marijuana
intoxication and withdrawal, 571
Marine omega-3 fatty acids, 320
Masseter muscle, 507
Mast cells, **408**
IgE antibody and, 105
Mast cell stabilizers, 687
Mastectomy, 448
Mastication muscles, **507**
Mastoid air cells, 621
Mastoiditis
brain abscesses, 180
Wegener granulomatosis, 315
Maternal diabetes
cardiac defect association, 300
Maternal PKU, 84
Maternal (postpartum) blues, 562
Maternal pregnancy complication, 272
Mature cystic teratoma, 647
Mature ego defenses, 555
Maxillary artery, 619
Maxillary process, 620
Mayer-Rokitansky-Küster-Hauser syndrome, 622
McArdle disease, 87
McBurney point, 383
McCune-Albright syndrome, 57
McMurray test, 454
MDMA (ecstasy), 571
Mean (statistics), 262
Mean arterial pressure, 501
equation for, 285
Measles, 183
paramyxovirus, 167, 169
rubeola virus, **170**
unvaccinated children, 186
vitamin A for, 66
Measurement bias, 260
Measures of central tendency, 262
Measures of dispersion, 262
Mebendazole, 200
microtubules and, 48
mecA gene
penicillin resistance and, 135
Meckel diverticulum, **384**, 618
Meconium ileus, 386
cystic fibrosis, 60
MECP2 gene, 62
Medial calcific sclerosis, 301
Medial cerebellar lesions, 499
Medial collateral ligament (MCL) injury
in "unhappy triad," 460
Medial epicondylitis, 459
Medial femoral circumflex artery, 463
Medial geniculate nucleus (MGN), 498
Medial lemniscus, 514
Medial longitudinal fasciculus (MLF), 511, 543
Medial malleolus, 455
Medial medullary syndrome, 514
Medial meniscal tear, 460
Medial pterygoid muscle, 507
Medial rectus muscle, 540
Medial tibial stress syndrome, 461
Medial umbilical ligament, 282, 369
Median (statistics), 262
"Median claw," 451
Median nerve
carpal tunnel syndrome, 459
injury to, 447
neurovascular pairing, 455
Median umbilical ligament, 369

(c) ketabton.com: The Digital Library

Mediastinal lymphadenopathy, 676
Mediastinal pathology, **672**
Mediastinitis, 672
  in pulmonary anthrax, 137
Medical abortion
  ethical situations, 268
  methotrexate for, 440
Medical errors
  analysis of, **274**
  assessment of, 268
  types of, 274
Medical insurance plans, 271
Medical power of attorney, 266
Medicare/Medicaid, **272**
Medication errors, 274
Medication noncompliance, 268
Medium-chain acyl-CoA dehydrogenase deficiency, 89
Medium-vessel vasculitis
  presentation and pathology, 314
Medroxyprogesterone, 657
Medulla (brain)
  adrenal cortex and, **327**
  cranial nerves and nuclei, 505
  development of, 490
  spinal tracts and, 509
  strokes in, 514–515
Medulla (lymph nodes)
  lymph nodes, 96
  thymus, 102
Medullary breast carcinomas, 650
Medullary cystic kidney disease, 604
Medullary thyroid carcinomas, 343, 351
Medulloblastoma, 350, 528
"Medusa head" appearance, 137
Mefloquine, 157
Megacolon
  Chagas disease, 158
  in Hirschsprung disease, 384
Megakaryocytes in essential thrombocythemia, 433
Megaloblastic anemia, 420
  cytarabine, 440
  Diphyllobothrium latum, 160
  drug reaction and, 250
  orotic aciduria, 420
  trimethoprim, 194
  tropical sprue, 381
  vitamin B_9 deficiency, 68
  vitamin B_{12} deficiency, 69
Megestrol, 657
Meglitinides, 353
Meissner corpuscles, 494
Meissner plexus, 384
Melanocytes
  tumor nomenclature in, 220
Melanocyte-stimulating hormone (MSH)
  function of, 328
  secretion of, 327
  signaling pathways of, 337
Melanocytic nevus, 477
Melanoma, 484
  common metastases, 223
  immunohistochemical stain for, 227
  nomenclature for, 220
  oncogene, 224
  origin of, 220
  recombinant cytokines for metastatic, 121
  tumor suppressor gene, 224
Melarsoprol, 156, 200
Melasma (chloasma), 476
MELAS syndrome, 59
Melatonin
  circadian rhythms and, 497
Melatonin receptor agonist
  Ramelteon as, 547
Melena
  Meckel diverticulum, 384, 618
  polyarteritis nodosa, 314
Meloxicam, 486
Memantine, 549
Membrane attack complex (MAC), 104
  complement and, 106
Membranoproliferative glomerulonephritis (MPGN), 596
  hepatitis B and C, 173
Membranous glomerular disorders, 594
  hepatitis B and C, 173
Membranous interventricular septum, 281
Membranous nephropathy, 597
  primary autoantibody, 115
Membranous ossification, 458
Membranous urethra injury, 627
Membranous ventricular septum, 281
Memory
  neural structures and, 499
Memory loss
  anti-NMDA receptor encephalitis, 228
  lead poisoning, 425
  Wernicke-Korsakoff syndrome, 511
MEN1 gene, 224, 352
Ménétrier disease, **379**
Ménière disease, 534
Menin, 224
Meninges, **496**
Meningiomas, 526
  Psammoma bodies in, 227
Meningitis
  ceftriaxone, 189
  chloramphenicol, 192
  coccidioidomycosis, 151
  common causes, **180**
  Cryptococcus neoformans, 153
  CSF findings in, 180
  fluconazole, 199
  flucytosine, 199
  Haemophilus influenzae, 142
  headaches with, 518
  HIV-positive adults, 177
  Listeria monocytogenes, 139
  meningococci, 142
  mumps, 170
  in neonates, 182
  rifamycin prophylaxis, 196
  Streptococcus pneumoniae, 136
  Streptococcus agalactiae, 137
  unvaccinated children, 186
Meningocele, 491
Meningococcal prophylaxis, 198
Meningococcal vaccine, 127
Meningococcemia
  endotoxins, 131
Meningococci, 131, 142
Meningoencephalitis
  HSV-2, 182
  Naegleria fowleri, 156
  West Nile virus, 167
Meningomyelocele, 491
Menkes disease
  mechanism and symptoms, **51**
  protein (ATP7A), 51
Menopause, **636**
  hormone replacement therapy, 657
  primary ovarian insufficiency, 645
  Turner syndrome, 638
Menorrhagia, 633
  anemia with, 418
Menstrual cycle
  estrogens for, 656
  heavy bleeding (AUB/HMB), 633
  phases of, **632**
Meperidine, 551
Mepivacaine, 550
Mercury poisoning, 248
Merkel discs, 494
Merlin protein, 224
MERS (Middle East respiratory syndrome), 167
Mesalamine, 382
Mesangial cells
  juxtaglomerular apparatus, 589
Mesencephalon, 490
Mesenchymal tumors
  nomenclature of, 220
Mesenteric arteries, 363
Mesenteric ischemia, 386
Mesocortical pathway, 499
Mesoderm, 490
  branchial arches derivation, 619
  derivatives of, 613
Mesolimbic pathway, 499
Mesometrium, 625
Mesonephric (Wolffian) duct, 622
Mesonephros, 578
Mesosalpinx, 625
Mesothelioma, **678**
  carcinogens causing, 225
  Psammoma bodies in, 227
Mesovarium, 625
Mestranol, 656
Meta-analysis, 263, **264**
Metabolic acidosis, 592
  adrenal insufficiency, 349
  renal failure, 603
  symptoms of, 593
Metabolic alkalosis, 592
  hyperaldosteronism, 349
  in hypertrophic pyloric stenosis, 359
  loop diuretics, 608
  nephron transport, 586
  thiazides, 609
Metabolic disorders
  fructose, 80
  galactose, 80
  glycogen storage, 87
  lysosomal storage diseases, 88–89
Metabolic fuel use, **91**
Metabolic syndrome
  with antipsychotic drugs, 573
  non-alcoholic fatty liver disease and, 391
Metabolism, 72–94
  amino acid derivatives, 83
  amino acids, 81
  apolipoproteins, 93
  cellular sites of, **72**
  disorders of, 81, 84–85
  drugs, 232
  dyslipidemias, 94
  ethanol, 72, 232
  fatty acid, 89
  fuel use, 91
  gluconeogenesis, 78
  glycogen and, 86
  lipoprotein functions, 93, 94
  pyruvate, 77
  rate-determining enzymes, 73
  summary of pathways, 74
  TCA cycle, 77
  tyrosine catabolism, 83
  urea cycle, 82
Metacarpal neck fracture, 459
Metacarpophalangeal (MCP) joints, 451
Metachromatic granules
  Corynebacterium diphtheriae diagnosis, 139
Metachromatic leukodystrophy, 88, 524
Metalloproteinases, 216
Metal storage diseases, 210
Metanephros, 578
Metaphase, 46
Metaplasia, 206
  benign breast disease, 649
  esophagus, 378
  intestinal, 379
  specialized intestinal, 378
Metastases
  common, **223**
Metastasis, 219, 223
  gastric cancer, 379
  heart tumors from, 316
  liver cancer, 392
  lung, cancer, 684
Metastatic calcification, 211
Metastatic melanoma
  vemurafenib for, 444
Metatarsophalangeal (MTP) joints
  gout, 467
Metencephalon, 490
Metformin, 353
  diarrhea with, 249
Methacholine, 240
Methadone, 551
  heroin addiction, 576
  intoxication and withdrawal, 570
Methamphetamine, 572
Methanol toxicity, 248
Methemoglobin, **666**
  toxicity treatment, 248
Methemoglobinemia, 666
  local anesthetics and, 550
Methicillin, 249
Methimazole, 354
  agranulocytosis, 250
  aplastic anemia, 250
  teratogenicity, 614
Methionine
  classification of, 81
  start codons, 44
  sulfonamides and, 194
Methotrexate, 440
  folate deficiency, 420
  hydatidiform moles, 642
  megaloblastic anemia, 250
  pulmonary fibrosis, 251
  pyrimidine synthesis and, 36
  rheumatoid arthritis, 466
  targets of, 438
  teratogenicity, 614
  vitamin B_9 deficiency, 68
  as weak acid, 233
Methoxyflurane, 550
Methylation
  histones, 45
Methyldopa
  Coombs-positive hemolytic anemia, 250
  hypertension in pregnancy, 316
Methylene blue, 248

(c) ketabton.com: The Digital Library

Methylenetetrahydrofolate reductase (MTHFR) deficiency, 84
Methylmalonic acid
  vitamin B_{12} deficiency, 69
  vitamin B_9 deficiency, 68
Methylmalonyl-CoA mutase, 69
Methylmercury teratogenicity, 614
Methylnaltrexone, 551
Methylphenidate
  ADHD, 557, 572
  CNS stimulant, 572
Methyltestosterone, 658
Methylxanthines, 687
Metoclopramide, **400**
  Parkinson-like syndrome, 251
  tardive dyskinesia, 251
Metolazone, 609
Metoprolol, 245, 323
Metronidazole
  bacterial vaginosis, 148
  clindamycin vs, 192
  Clostridium difficile, 138
  for Crohn disease, 382
  disulfiram-like reaction, 251
  Giardia lamblia, 155
  Helicobacter pylori, 146
  mechanism and use, **195**
  vaginal infections, 181
  vaginitis, 158
Mevalonate synthesis, 320
Mexiletine, 322
Meyer loop, 542
MHC (major histocompatibility complex) I and II, **100**
Micafungin, 200
Michaelis-Menten kinetics, 230
Miconazole, 199
Microalbuminuria, 346
Microangiopathic hemolytic anemia (MAHA), 423
  hypertensive emergency and, 300
  intravascular hemolysis in, 421
Microarrays, **54**
Microbiology, 123–204
  antimicrobials, 187–204
  bacteriology, 124–134
  clinical bacteriology, 134–150
  mycology, 151–154
  oncogenic organisms, 226
  parasitology, 155–161
  systems, 178–186
  virology, 162–177
Microbiome
  in innate immunity, 99
Microcephaly, 63
  cri-du-chat syndrome, 64
  fetal alcohol syndrome, 615
  maternal phenylketonuria, 84
  maternal X-ray exposure, 614
  Patau syndrome, 63
  with zika virus, 171
Microcytic, hypochromic anemia, **418–419**
  Ancylostoma, 161
Microcytosis, 214
Microdeletion
  congenital, 64
  fluorescent in situ hybridization and, 55
  22q11, 116
Microfilaments, 48
Microglia, 490, **493**
Micrognathia
  Edwards syndrome, 63
  Pierre Robin sequence, 620
Micromelia, 614
Microphthalmia, 63
MicroRNA (miRNA), 42, **56**
Microscopic polyangiitis, 315
Microsporum, 152
Microtubule inhibitors, **441**
Microtubules, **48**
  drugs acting on, 48
  structure and function of, 48
Micturition center, **237**
Midazolam, 546, 550
Midbrain
  brain, 490
  cranial nerve nuclei of, 505
  lesions in, 511
Middlebrook medium, 126
Middle cerebral artery (MCA)
  cortical distribution, 502
  saccular aneurysms, 516
  stroke effects, 514
Middle meningeal artery
  epidural hematoma and, 513
Middle rectal vein, 365
Midgut
  blood supply/innervation of, 364
  development of, 358
Midgut volvulus, 386
Midodrine, 242
Mifepristone, 657
Miglitol, 353
Migraine headaches, 518
  hormonal contraception contraindication, 657
  triptans for, 547
Migrating motor complexes (MMC), 371
Migratory polyarthritis, 312
Milnacipran, 575
Mineralocorticoids
  adrenal insufficiency, 349
  adrenal steroids and, 335
Mineral oil, 65
Minimal alveolar concentration, 549
Minocycline, 192
Minors, consent for, 265
Minoxidil, **658**
Minute ventilation, 664
Miosis
  cholinomimetic agents, 552
  Horner syndrome, 531, 540
  pupillary control, 539
  sympatholytic drugs, 243
Mirabegron, 242
Mirtazapine, 244, 576
  major depressive disorder, 561
Mismatch repair, 40
Misoprostol, **399**
Missense mutations, 39
Mites/louse treatment, 200
Mitochondria
  high altitude and, 670
  metabolism in, 72
Mitochondrial encephalopathy, 59
Mitochondrial inheritance, 59
Mitochondrial myopathies, 59
Mitosis, 46
  griseofulvin, 200
Mitral regurgitation, 288
  in MI, 305
  murmurs caused by, 290, 291
Mitral stenosis, 288
  murmurs caused by, 291
Mitral valve
  in cardiac cycle, 287
  regurgitation in, 312
Mitral valve prolapse, 291
  fragile X syndrome, 62
  Marfan syndrome, 52
  renal cyst disorders and, 604
Mittelschmerz, 631
Mixed cellularity lymphoma, 429
Mixed connective tissue disease, 470
  autoantibody, 115
  Raynaud phenomenon, 472
Mixed cryoglobulinemia, 315
Mixed transcortical aphasia, 516
MMR vaccine, 170
Mobitz AV blocks, 295
Moccasin distribution (tinea pedis), 152
Modafinil, 568
Mode (statistics), 262
Molecular mimicry
  autoimmune response in rheumatic fever, 129
Molecular motor proteins, 48
Molluscum contagiosum, 164, 479
Mönckeberg sclerosis, 301
"Monday disease," 318
Monoamine oxidase (MAO) inhibitors
  atypical depression, 561
  mechanism and clinical use, **575**
  Parkinson disease, 549
  selegiline/rasagiline, 549
  tyramine and, 244
Monobactams, **190**
  Pseudomonas aeruginosa, 143
Monoclonal antibodies
  drug names for, 254
Monoclonal immunoglobulin (Ig)
  overproduction, 431
Monocytes, **407**
  innate immunity, 99
  morulae in, 150
Monospot test, 165
Monozygotic ("identical") twins, 616
Montelukast, 687
Mood
  oxytocin's role in regulating, 328
Mood disorder, **560**
  readmissions with, 272
Mood stabilizing drugs, 561
Morbidly adherent placenta, 640
Moro reflex, 510, 635
Morphine, 551
  for acute coronary syndromes, 307
  intoxication and withdrawal, 570
Morphogenesis
  of heart, 280–281
  errors in, 613
Morulae, 150
Mosaic bone architecture, 464
Mosaicism, 57
Mosquitoes (disease vectors)
  lymphatic filariasis, 159
  malaria, 157
  Zika virus, 171
Motilin, 371
Motion sickness, 241
Motor cortex, 514
  descending spinal tracts, 509
  topographic representation, 502
  ventral lateral thalamus and, 498
Motor innervation
  derivation of, 620
  lower extremity, 452
  tongue, 493
Motor neuron signs, **529**
Movement disorders, **519**
  dopaminergic pathways and, 499
Moxifloxacin, 195
M phase, 46
MPO-ANCA/p-ANCA autoantibody, 115
M protein
  rheumatic fever and, 136
  as virulence factor, 129
mRNA
  aminoglycosides, 191
  hepatitis viruses, 172
  pre-mRNA splicing, 42
  processing, 41
  protease inhibitors, 203
  stop codons, 44
MRSA (methicillin-resistant *Staphylococcus aureus*)
  cephalosporins, 189
  highly resistant, 198
  nosocomial infections, 135
  oxazolidinones, 193
  vancomycin, 190
mTOR
  sirolimus (rapamycin) inhibition of, 120
Mucicarmine stain
  polysaccharide capsule staining, 125
Mucinous cystadenocarcinoma, 646
Mucinous cystadenoma, 646
Mucociliary escalator, 662
Mucocutaneous lymph node syndrome, 314
Mucoepidermoid carcinomas, 376
Mucopolysaccharides
  Periodic acid-Schiff stain, 125
Mucopolysaccharidoses, 88
Mucormycosis
  diabetic ketoacidosis, 347
Mucor spp
  amphotericin B for, 199
  opportunistic infection, 153
Mucosa, 362
Mucosal cells, 372
Mucosal neuromas, 351
Mucosal polyps, 387
Mucositis
  methotrexate, 440
Mucus, 238
"Muddy brown" casts (urine), 594
Mulberry molars, 147
Müllerian duct
  agenesis, 622
  anomalies of, 623
  derivatives of, 622
Müllerian inhibitory factor (MIF), 622
  Sertoli cell production, 628
Multicystic dysplastic kidney, 578, 579
Multidrug resistance protein 1 (MDR1), 227
Multifactorial pulmonary hypertension, 679
Multiorgan drug reactions, 251
Multiple endocrine neoplasias (MEN syndromes), **351**
  Zollinger-Ellison syndrome, 351
Multiple gestations, 633
Multiple myeloma
  bortezomib/carfilzomib in, 443
  common metastases, 223
  ESR in, 214
  metastatic calcification, 211
  monoclonal gammopathy transition to, 431
  plasma cell dyscrasia, 409
Multiple personality disorder, 558
Multiple sclerosis, **523**
  drug therapy for, 551
  HLA-DR2 and, 100
  internuclear ophthalmoplegia, 543

(c) ketabton.com: The Digital Library

natalizumab for, 122
recombinant cytokines for, 121
Mumps, **170**
acute pancreatitis with, 397
paramyxovirus, 167, 169
Munchausen syndrome, 566
Munchausen syndrome by proxy, 566
Murphy sign, 396
Muscarinic acetylcholine (ACh) receptors, 236
Muscarinic agonists, 237
Muscarinic antagonists, 237, **241**, 687
multiple sclerosis, 523
neuromuscular blocking drugs, 550, 551
Parkinson disease, 548
Muscarinic receptor
detrusor muscle, in, 237
Muscle conduction/contraction
skeletal, 456
smooth muscle, 457
Muscle proprioceptors, **458**
Muscles
atrophy of, 42
in starvation, 91
metabolism in, 86
ragged red fibers in, 59
Muscle spindled, 458
Muscular dystrophies, **61**
frameshift mutation, 61
X-linked recessive disorder, 61
Muscularis externa, 362
Muscular ventricular septum, 281
Musculocutaneous nerve
injury presentation, 447
Musculoskeletal drug reactions, 250
Musculoskeletal paraneoplastic syndromes, 228
Musculoskeletal system
anatomy, 446–454
common conditions, **461**
pathology, 459–467
pharmacology, 485–487
Mutase, 73
Mutations
BRAF, 432
BRCA1 gene, 56
COL3A1, 51
de novo, 62
drug resistance mechanisms, 196
in HbS and HbC, 410
JAK2, 433
locus heterogeneity in, 57
muscular dystrophies, 61
myelodysplastic syndromes, 432
non-Hodgkin lymphoma, 430
in PBPs, 187
same locus, 57
STAT3, 116
thalassemia and, 418
tumor suppressor genes, 46, 525
WT1 deletion, 606
Myalgias
Ebola virus, 171
fluoroquinolones, 195
genital herpes, 184
Jarisch-Herxheimer reaction, 148
Leptospira interrogans, 147
Lyme disease, 146
meningitis, 186
Trichinella spiralis, 159, 161
trichinosis, 159
vasculitides, 314
Myasthenia gravis, 472
autoantibody, 115
diagnosis of, 240
HLA subtype association, 100
neostigmine for, 240
as paraneoplastic syndrome, 228
pyridostigmine for, 240
restrictive lung diseases, 675
MYCL1 gene, 224
MYCN gene, 224
Mycobacterial cells, 196
Mycobacterium spp, **140**
Gram stain, 125
intracellular organism, 127
Ziehl-Neelsen stain, 125
Mycobacterium avium-intracellulare, 140
HIV-positive adults, 177
prophylaxis with HIV, 198
vertebral osteomyelitis, 180
Mycobacterium leprae
animal transmission, 149
diagnosis, 141
rifamycins/dapsone, 196
Mycobacterium marinum, 140
Mycobacterium pneumoniae, 126
Mycobacterium scrofulaceum, 140
Mycobacterium spp
facultative intracellular organisms, 127
Mycobacterium tuberculosis
culture requirements for, 126
osteomyelitis, 180
reactivation site, 140
symptoms of, 140
therapeutic agents, 196, 197
Mycolic acid
isoniazid, 197
synthesis of, 196
Mycology, 151–154
Mycophenolate mofetil, 120
inosine monophosphate dehydrogenase inhibition, 36
Mycoplasma spp
atypical organisms, 179
Gram stain, 125
macrolides, 193
pneumonia caused by, **150**, 179, 683
tetracyclines, 192
Mycoses
cutaneous, 152
systemic, 151
Mycosis fungoides, 430
Mydriasis
glaucoma treatment and, 552
G-protein-linked second receptor, 238
muscarinic antagonists for, 241
pupillary control, 539
saccular aneurysm, 516
Myelencephalon, 490
Myelin, **494**
Myeloblasts (peripheral smear), 432
Myelodysplastic syndromes, **432**
sideroblastic anemia, 419
Myelofibrosis, 433
Myeloid neoplasms, 432
Myelomeningocele, 63, 492
Myeloperoxidase, 109
H_2O_2 degradation, 128
in neutrophils, 406
Myeloproliferative disorders
in AML, 432
chronic, 433
hydroxyurea for, 442
Myeloschisis, 491
Myelosuppression
alkylating agents, 441
antimetabolites, 440
drugs causing, 444
hydroxyurea, 442
irinotecan/topotecan, 442
Myenteric plexus, 362
Mylohyoid muscle, 620
Myocardial action potential, **292**
Myocardial depression, 550
Myocardial infarction (MI), 304
antiarrhythmics after, 322
β-blockers for, 245
complications of, **307**
diabetes mellitus, 346
diagnosis of, 306
on ECG, 293
ECG localization of STEMI, 306
evolution of, 305
heart failure caused by, 309
heparin for, 436
homocystinuria, 84
hypertensive emergency and, 300
shock caused by, 310
thrombolytics for, 437
Myocardial O_2 consumption/demand, 284
angina treatment, 319
Myocarditis, **313**
adenovirus, 164
coxsackievirus, 167
diphtheria, 139
picornaviruses, 167
Toxocara canis, 159
Myoclonic seizures, 517
Myoclonus, 519, 521
Myofibroblasts, 216
Myoglobin, 665
ooxygen-hemoglobin dissociation curve, 666
Myoglobinuria
acute tubular necrosis, 602
McArdle disease, 87
Myonecrosis, 138
Myopathy
daptomycin, 195
drug reaction and, 250
Myophosphorylase, 87
Myopia, 535
retinal detachment, 537
Myotonic dystrophy, **61**
cataracts and, 535
Myxedema
thyroid hormones for, 354
Myxomas, 316
Myxomatous degeneration, 291

N

N-acetylglucosaminyl-1-phosphotransferase, 47
N-acetylcysteine, **686**
for acetaminophen toxicity, 248
for cystic fibrosis, 60
NADH (reduced nicotinamide adenine dinucleotide)
electron transport chain, 78
fructose metabolism, 80
TCA cycle, 77
Nadolol, 245
NADPH (reduced nicotinamide adenine dinucleotide phosphate)
ethanol metabolism, 72
HMP shunt and, 79
respiratory burst and, 109
universal electron acceptors, 75
Naegleria fowleri, 156
Nafcillin
characteristics of, 188
Nails
clubbing, 60
glomus tumors under, 478
hemorrhages in bed of, 311
with psoriatic arthritis, 469
Naïve T cell activation, 103
Naked viral genome infectivity, **163**
Nalbuphine, 551
Naloxone
dextromethorphan overdose, 686
heroin detoxification, 576
for opioid toxicity, 248, 551
opioid toxicity, 570
Naltrexone
alcoholism, 571
heroin detoxification, 576
opioid toxicity, 551
2-naphthylamine, 225
Naproxen, 486
acute gout drugs, 487
Narcissistic personality disorder, 565
Narcolepsy
amphetamines for, 242
characteristics/treatment, **568**
hypnagogic/hypnopompic hallucinations, 568
Narrow-angle glaucoma, 536
Nasal congestion, 686
Nasal decongestion
ephedrine for, 242
Nasal polyps
cystic fibrosis, 60
Nasal septum perforation, 315
Nasopharyngeal carcinoma
EBV and, 165
oncogenic microbes and, 226
Natalizumab, 122
multiple sclerosis, 523
Nateglinide, 353
National Board of Medical Examiners (NBME), 2, 11
Natriuresis, 588
Natriuretic peptide, 296
Natural killer (NK) cells, 101, **409**
cell surface proteins, 110
function of, 409
innate immunity, 99
Nausea
adverse drug effects, 400
antiemetics for, 401
with appendicitis, 383
biliary colic, 396
with MI, 305
migraine headaches, 518
ranolazine, 319
renal failure, 603
vitamin A toxicity, 66
vitamin C toxicity, 69
Nebivolol, 245
Necator spp
disease associations, 161
infection routes, 158
Necator americanus, 159
Neck and head cancer, 671
cetuximab for, 442
Necrosis, **209**
acute pancreatitis, 397
Arthus reaction, 113
benign tumors, 220
Budd-Chiari syndrome, 392
calcification, 211
enterocolitis, 386
femoral head, 120, 461, 463
fibrinoid, 466
glioblastoma multiforme, 526
hepatic, 485

(c) ketabton.com: The Digital Library

Necrosis (*continued*)
hernias and, 370
jaw, 486
mesenteric ischemia, 386
nonalcoholic fatty liver disease, 391
saponification, 209
scaphoid avascular, 449
transplant reaction, 119
warfarin, 436
Necrotizing enterocolitis, 386
Necrotizing fasciitis, 479
Streptococcus pyogenes (Group A strep), 136
Necrotizing glomerulonephritis, 315
Negative predictive value (NPV), 257
Negative punishment, 554
Negative reinforcement, 554
Negative skew distribution, 262
Negative-stranded viruses, **168**
Neglect (child), 556
Negri bodies, 171
Neisseria gonorrhoeae
culture requirements, 126
epididymitis and orchitis, 654
osteomyelitis, 180
septic arthritis, 468
STI, 184
UTIs with, 600
Neisseria meningitidis
chloramphenicol, 192
culture requirements, 126
encapsulation, 127
meningitis, 180
penicillin G/V for, 187
Waterhouse-Friderichsen syndrome, 349
Neisseria spp
C5-C9 deficiencies, *107*
cephalosporins, 189
fluoroquinolones, 195
gram-negative algorithm, **142**
IgA protease, 129
intracellular organism, 127
transformation in, 130
Nelson syndrome, **348**
Nematodes, **159**
infection routes, **158**
Neomycin
aminoglycosides, 191
Neonatal abstinence syndrome, **615**
Neonatal conjunctivitis
Chlamydia trachomatis, 149
Neonatal pneumonia
Group B streptococci, 179
Neonatal respiratory distress syndrome (NRDS), **661**
Neonates
abstinence syndrome, 615
Apgar score, 634
Candida albicans in, 153
coagulation cascade in, 413
conjunctivitis, 142, 149
deprivation effects, 556
esophageal atresia in, 359
flora with C-section, 178
galactosemia in, 80
gastroenteritis, 168
gray baby syndrome in, 192
hemolytic anemia in, 422
herpes in, 164
hyperthermia in, 241
hypertrophic pyloric stenosis in, 359
indirect inguinal hernia in, 370
intraventricular hemorrhage, **512**
jaundice in, 393
kernicterus, 194, 204
Listeria monocytogenes in, 139
low birth weight, 635
meningitis in, 182
necrotizing enterocolitis and, 386
obesity risk factors, 636
pneumonia in, 149
primitive reflexes in, 510
sickle cell anemia in, 422
Streptococcus agalactiae in, 137
TORCH manifestations in, 182
Neoplasia
pathology of, **219**–226, 518
Neoplastic progression, **219**
Neoplastic transformation, 216
adenomatous polyps, 387
Neostigmine, 240
Nephritic-nephrotic syndrome
etiology and presentation, 595
Nephritic syndrome, **596**–597
etiology and presentation, 595
Nephritis, 608
Nephroblastoma, **606**
Nephrocalcinosis, 211
Nephrogenic diabetes insipidus, 211, 338
lithium toxicity, 569
treatment, 609
Nephrolithiasis, 606
calcium oxalate, 69
Nephron physiology, **585**
Nephropathy
diabetes mellitus, 346
hypertension and, 300
protease inhibitors, 203
transplant rejection, 119
Nephrotic syndrome, **597**
early-onset, 606
ESR in, 214
etiology and presentation, 595
fatty casts in, 594
TBG and, 331
Nephrotoxicity
aminoglycosides, 191
amphotericin B, 199
cidofovir, 202
cisplatin/carboplatin, 442
cladribine, 440
drug reaction and, 251
drugs causing, 444
immunosuppressants, 120
inhaled anesthetics, 550
streptomycin, 197
sulfonamides, 194
Nerve blockade (local anesthetics), 550
Nerve fibers, 495
Nerves
lower extremity, 452, 453
upper extremity, 446, 447
Nerve trunk, 495
Nesiritide, 296
Neural crest
derivatives of, 613
Neural crest cells, 490, 494
Neural development, **490**
Neural plate, 490
Neural tube, 490
derivatives, 613
formation, 612
Neural tube defects, **491**
maternal diabetes, 614
prevention, 68
valproic acid, 544
Neuraminidase, 169–170
Neuroblastomas, **350**
incidence and mortality, 222
oncogenes and, 224
paraneoplastic syndromes with, 228
Neurocutaneous disorders, **525**
Neurodegenerative disorders, **520**–522
drug therapy for, **549**
Neuroectoderm, 490
derivatives of, 612
pituitary gland, 327
Neuroendocrine tumors, **350**
Neurofibromatosis, 535
chromosome association, 64
inheritance, 60
tumor suppressor genes and, 224
typesI and II, 525
variable expressivity, 56
Neurofilaments
cytoskeletal element, 48
immunohistochemical stain for, 227
Neurogenic ileus, 240
Neurohypophysis, 327
hypothalamus and, 498
Neuroleptic malignant syndrome (NMS), 551, 569
Neurological signs
proximal upper and lower extremity, 502
Neurologic drug reactions, 251
Neurology and special senses, 490–544
anatomy/physiology, 493–515
embryology, 490–492
ophthalmology, 534–541
otology, **533**–**534**
pathology, 511–518
pharmacology, 544–551
Neuromuscular blocking drugs, **551**
Neuromuscular disorders
paraneoplastic syndromes, 228
Neuromuscular junction
skeletal muscle, 236
diseases of, **472**
Neurons, **493**
in ascending spinal tracts, 509
local anesthetics, 550
origins of, 490
Parkinson disease, 548
primary motor cortex, 509
Neuron-specific enolase, 226, 350
Neuropathic pain, 515
Neuropathy
diabetes mellitus, 346
Neurosyphilis, 147
Neurotoxicity
cladribine, 440
immunosuppressants, 120
methylmercury exposure, 614
methylxanthines, 687
vincristine, 441
Neurotransmitters
bacterial toxin effects, 132
changes with disease, **495**
vomiting center receptors, 496
Neurovascular pairing, **455**
Neutralization (antibody), 104
Neutropenia, 424
ganciclovir, 202
rheumatoid arthritis, 466
ticlopidine, 437
Neutrophils, **406**
chemotaxis, 44, 106, 133, 406, 485, 487, 691
corticosteroid effect on, **424**
IL-8 and, 108
innate immunity, 99
left shift, **424**
in leukocyte adhesion deficiency, 117
LTB4, 485
megaloblastic anemia, 420
in MI, 305
necrosis and, 209
nonmegaloblastic anemia, 420
pseudo-Pelger-Huet anomaly, 432
stimulation of, 44
wound healing, 216
Nevi, 220
Nevirapine
cytochrome P-450 and, 252
HIV therapy, 203
NF1/*NF2* genes, 224
N-formylmethionine (fMet), 44
NF-κB, 120
Niacin
cutaneous flushing, 248
gout, 250
hyperglycemia, 249
myopathy caused by, 250
Nicardipine, 318
Nicotinamides, 75
Nicotine
intoxication and withdrawal, **571**
teratogenicity, 614
Nicotinic acetylcholine receptors, 166, **236**
Niemann-Pick disease, 88
Nifedipine, 316, **318**, 643
Nifurtimox, 158, 200
Night terrors
benzodiazepines for, 497
Nigrostriatal pathway, 499
Nikolsky sign
blistering skin disorders, 480
scalded skin syndrome, 479
Nilotinib, 443
Nimodipine, **318**, 513
Nipple
intraductal papilloma, 649
lactational mastitis, 649
Nissl bodies, 46
Nissl substance
chromatolysis, 495
neurons, 493
Nitazoxanide, 155
Nitrates, 319
mechanism and clinical use, **318**
Nitric oxide, 371
free radical injury and, 210
Nitrites
urinary tract infections, 181
Nitroblue tetrazolium dye reduction test, 117
Nitrofurantoin
hemolysis in G6PD deficiency, 250
pulmonary fibrosis, 251
Nitroglycerin, 318
acute coronary syndromes, 307
angina, 304
Nitroprusside, 318
Nitrosamines
as carcinogens, 225
stomach cancer and, 379
Nitrosoureas, 441
Nitrous oxide, 550
Nizatidine, 399
NMDA receptor antagonist
ketamine as, 550
memantine as, 549
N-*myc* oncogene, 350

(c) ketabton.com: The Digital Library

NNRTIs, 203
Nocardia spp
  aerobic culture requirements, 126
  comparison with *Actinomyces* spp, **139**
  effects and treatment, 139
  necrosis and, 209
  sulfonamides for, 194
  urease-positive, 127
  Ziehl-Neelsen stain, 125
Nocturia, 654
Nocturnal enuresis, 329
Nodes of Ranvier, 494
Nodular phlebitis, 314
Nodular sclerosis lymphoma, 429
Noise-induced hearing loss, 533
Nonadherent patients, 268
Nonalcoholic fatty liver disease, 389, 390, **391**, 392
Nonbacterial endocarditis, 311
Nonbacterial thrombotic endocarditis, 228
Nonbenzodiazepine hypnotics, **546**
Noncaseating granulomas
  sarcoidosis, 676
Noncommunicating hydrocephalus, 522
Noncompetitive agonists, 234
Noncompetitive antagonist, 234
Noncompetitive inhibitors, 230
Noncompliant patients, 268
Nondepolarizing neuromuscular blocking drugs, 551
Nondominant parietal cortex lesions, 511
Non-frameshift mutations
  deletions, 61
Nonhemolytic, normocytic anemia, **421**
Non-Hodgkin lymphoma, **430**
  corticosteroids, 120
  Hashimoto thyroiditis and, 341
  hepatitis C, 173
  HIV-positive adults, 177
  Hodgkin lymphoma vs, 429
  oncogenes and, 224
  rituximab for, 443
  vinca alkaloids for, 441
Nonhomologous end joining, 40
Nonmaleficence (ethics), 265
Nonmegaloblastic macrocytic anemia, 420
Nonnormal distributions, 262
Nonoverlapping genetic code, 37
Nonreceptor tyrosine kinase, 337
Non-REM sleep stages, 497
Nonselective antagonists, 245
Nonsense mutations, 39
Nonsteroidal anti-inflammatory drugs (NSAIDs), **486**
  acute pericarditis, 313
  aplastic anemia, 250
  Beers criteria, 247
  calcium pyrophosphate deposition disease, 467
  colorectal cancer chemopreventative, 389
  esophagitis from, 377
  gastritis with, 379
  GFR effects of, 589
  gout, **467**, 487
  headaches, 518
  interstitial nephritis, 249
  loop diuretics and, 608
  misoprostol use with, 399
  osteoarthritis, 466
  peptic ulcer disease and, 380
  rheumatoid arthritis, 466
  for sialadenitis, 376
Non-ST-segment elevation MI (NSTEMI)
  diagnosis of, 306
  STEMI vs, 304
  treatment, 307
Noradrenergic drugs, 574
Norepinephrine (NE)
  adrenal medulla secretion, 327
  amphetamines and, 239
  bupropion effect on, 576
  changes with disease, 495
  circadian rhythm, 497
  direct sympathomimetic, 242
  isoproterenol vs, **243**
  male sexual response, 627
  MAO inhibitor effects, 575
  opioid effect on, 551
  pheochromocytoma secretion, 350
  release regulation, 239
  REM sleep and, 497
  vitamin B_6 and, 67
Norethindrone, 657
Norfloxacin, 195
Normal distribution, 262
Normal flora
  colonic, 137
  female genital tract, 136
  GI tract, 127
  neonates, **178**
  oropharynx, 136
  skin, 135
Normal pressure hydrocephalus, 522
Normal splitting, 289
Normocytic anemia, **421**
Norovirus, 167
Northern blot, 53
Nortriptyline, 575
Nosocomial infections, 274
  Acinetobacter baumannii, 142
  Ebola, 171
  enterococci, 137
  Klebsiella, 145
  MRSA, 135
  pneumonias, 179
  Pseudomonas aeruginosa, 143
  risk factors, **185**
  UTIs as, 181
Notochord, **490**, 612–613
  postnatal derivative of, 282
Novobiocin
  gram-positive antibiotic test, 134
  Staphylococcus epidermidis, 135
NRTIs, 203
NS3/4A inhibitors, 204
NS5A inhibitors, 204
NS5B inhibitors, 204
Nuchal translucency, 63
Nucleic acids
  pathogen-associated molecular pattern (PAMP), 99
Nucleosome, 34
Nucleotide excision repair, 40
Nucleotides, **35**
  deamination reactions, 35
  synthesis, 72
Nucleus accumbens, 495
Nucleus ambiguus, 506
Nucleus cuneatus, 509
Nucleus pulposus
  collagen in, 50
  fetal precursor, 282
Nucleus tractus solitarius (NTS), 496
Null hypothesis, 263
Number needed to harm (NNH), 258
Number needed to treat (NNT), 258
Nursemaid's elbow, 461
Nutcracker syndrome, 363
Nutmeg liver, 309, 392
Nutrition, 65–72
Nyctalopia, 66
Nystagmus
  cerebellum, 499
  common lesions with, 511
  Friedreich ataxia, 531
  internuclear ophthalmoplegia, 543
  PCP, 571
  phenytoin, 544
  stroke and, 514
Nystatin, **199**

O

Obesity
  amphetamine for, 242
  anovulation with, 645
  cholelithiasis and, 396
  DM type 2 and, 347
  esophageal cancer and, 378
  hypertension risk factors, 300
  hypoventilation syndrome, 679
  lateral femoral cutaneous nerve, 452
  osteoarthritis/rheumatoid arthritis, 466
  renal cell carcinoma association, 605
  sleep apnea, 679
  stress incontinence and, 599
Obligate intracellular organisms, 127
Observational studies, **256**
  errors in, 260–261
Observer-expectancy bias, 260
Obsessive-compulsive disorder (OCD)
  diagnostic criteria/treatment, **563**
  drug therapy for, 572
  SSRIs for, 575
  Tourette syndrome and, 557
  venlafaxine for, 575
Obsessive-compulsive personality disorder, 566
Obstructive jaundice, 398
Obstructive lung diseases, **674**–675
  flow volume loops in, 673
Obstructive shock, 310
Obstructive sleep apnea, 679
  pulsus paradoxus in, 310
Obturator nerve, 452
Occipital cortex, 515
Occipital lobe, 501
Occipital sinus, 503
Occult bleeding, 387
  FOBT for, 388
Octreotide, 371, **400**
  acromegaly, 339
  GH excess, 329
  hypothalamic/pituitary drugs, 354
Ocular motility, **540**
Oculomotor nerve (CN III), 506
  ocular motility, **540**
  palsy of, 513, **541**
  pupillary contraction, 539
Odds ratio (OR), 256, **258**
Ofloxacin, 195
Okazaki fragments, 38
"OK gesture," 451
Olanzapine, 573
Olfaction
  hallucinations, 559
  limbic system in, 499
Olfactory nerve (CN I), 506
Oligoclonal bands, 523
Oligodendrocytes, **494**
Oligodendroglia, 490
  in multiple sclerosis, 494
Oligodendrogliomas, 526
Oligohydramnios, 578, **641**
Oligomenorrhea, 633
Oligomycin, 78
Oligospermia, 400
Olive-shaped mass, 359
Omalizumab, 122, **687**
Ombitasvir, 204
Omental foramen, 361
Omeprazole, 399
Omphalocele, 358
Omphalomesenteric cysts, 384
Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
Oncocytoma (renal), 605
Oncogenes, **224**
Oncogenic microbes, **226**
Ondansetron, **400**
  torsades de pointes, 248
1,25-$(OH)_2D_3$
  kidney endocrine function, 589
"100-day cough," 132
Onion skin periosteal reaction, 465
Onychomycosis
  terbinafine, 199
  tinea unguium, 152
Oocysts
  acid-fast stain, 155
  Toxoplasmosis, 156
  Ziehl-Neelsen stain, 125
Oogenesis, **631**
Oophorectomy, 625
Open-angle glaucoma, 536
  carbachol for, 240
  pilocarpine for, 240
Operant conditioning, **554**
Ophthalmology, 534–541
Ophthalmoplegia
  cavernous sinus syndrome, 542
  common lesions with, 511
  internuclear, 543
  Wernicke-Korsakoff syndrome, 571
Opioid analgesics, **551**
  agonists, 551
  Beers criteria, 247
  detoxification and relapse prevention, **576**
  intoxication and withdrawal, 570
  mechanism and use, 551
  mixed agonist/antagonist analgesics, 552
  receptor binding, 234
  sleep apnea, 679
  toxicity treatment, 248
Opisthotonos
  tetanospasmin, 138
Opponens digiti minimi muscle, 450
Opponens pollicis muscle, 450
Opportunistic fungal infections, 153–154
Oppositional defiant disorder, 557
Opposition (thumb), **450**, 451
Opsoclonus-myoclonus syndrome, **228**, 350

(c) ketabton.com: The Digital Library

Opsonization, **104**, 106
  antibodies in, 112
  complement in, 106
  encapsulated organisms, 98
Optic disc
  papilledema in, 538
Optic nerve (CN II), 506
  embryologic derivation, 613
Optic neuritis, 523
Optic neuropathy, 197
Optochin
  gram-positive antibiotic test, 134
Oral advance directives, 266
Oral contraceptives (OCPs)
  hepatic adenomas and, 392
  prolactin effects on, 330
  reproductive hormones, 656
  SHBG effects on, 337
  venous sinus thrombosis with, 503
Oral glucose tolerance test, 346
Oral hairy leukoplakia, 177
Oral/intestinal ganglioneuromatosis, 351
Oral rehydration therapy, 146
Oral thrush, 177
Orange body fluids, 196
Orchiectomy, 651
Orchiopexy, 651
Orchitis, 170, **654**
Orexigenic effect, 336
Orexin, 568
  hypocretin receptor antagonist, Suvorexant as, 547
Organ failure, in acute pancreatitis, 397
Organic acidemias, **85**
Organ of Corti, 533
Organogenesis
  embryologic derivatives, 612, **613**
  errors in, 613
  fetal development, 612
  teratogens, 614
Organophosphates
  toxicity treatment, 248
Organ transplants
  azathioprine for, 440
  cytomegalovirus, 186
  hairy leukoplakia and, 479
  kidneys, 580
  WBC casts, 594
Organum vasculosum of the lamina terminalis (OVLT), 498
Orientation, **557**
Origin of replication, 38
Orlistat, **400**
  diarrhea, 249
Ornithine
  cystinuria, 85
  kidney stones and, 598
  urea cycle, 82
Ornithine transcarbamylase, 74
Ornithine transcarbamylase deficiency, 61, **83**
Orotic acid, 83
Orotic aciduria, 420
"Orphan Annie" eyes (nuclei), 343
Orthomyxoviruses, 168
  characteristics of, **167**, 168
  influenza viruses, 169
Orthopedic conditions, 460
  lower extremity, 460, 461
Orthopnea, 309
Orthostatic hypotension
  adrenal insufficiency, 349
  α-blockers, 244
  phenoxybenzamine, 244
Ortolani maneuver, 461
Oseltamivir, **201**
Osgood-Schlatter disease, 461
Osler nodes, 311
Osler-Weber-Rendu syndrome, 316
Osmotic demyelination syndrome, 524
  SIADH and, 338
Osmotic diarrhea, 381
Osmotic laxatives, 401
Ossicles, 533
  conductive hearing loss and, 51
Ossification, 458
Osteitis deformans, **463**
Osteitis fibrosa cystica, **345**, 459, 464
Osteoarthritis, 466
  celecoxib for, 486
  vs rheumatoid arthritis, **466**
Osteoarthropathy, hypertrophic cancer association, 228
Osteoblastoma, 464
Osteoblasts, 459
  bone formation, 458
  cortisol effect on, 336
  Paget disease of bone, 463
  teriparatide effect on, 487
Osteochondroma, 464
Osteoclasts, 459
  bisphosphonate effects, 486
  bone formation, 458
  osteopetrosis, 463
  Paget disease of bone, 463
Osteodystrophy
  renal, 345, **603**
Osteogenesis imperfecta
  bisphosphonates, 486
  collagen and, 50
  collagen synthesis and structure in, **51**
Osteogenic sarcoma, 463, 465
Osteoma, 220, 464–465
  nomenclature for, 220
Osteomalacia
  hypophosphatemia, 591
  lab values with rickets, 464
  rickets, **463**
Osteomyelitis, **180**
  Pseudomonas aeruginosa, 143
  sickle cell anemia, 422
  Staphylococcus aureus, 135
Osteonecrosis, **463**, 486
Osteopenia, 463
Osteopetrosis, **463**, 464
Osteophytes, 466
Osteoporosis, **462**
  bisphosphonates, 486
  corticosteroids, 120
  denosumab, 122
  drug reaction and, 250
  estrogen, 459
  Gaucher disease, 88
  heparin, 436
  homocystinuria, 84
  hormone replacement therapy, 657
  lab values in, 464
  pituitary prolactinomas, 328
  raloxifene, 443, 656
  teriparatide for, 487
  thiazides for, 609
Osteosarcoma, 220, 464–465
Otitis media
  brain abscesses with, 180
  Haemophilus influenzae, 128, **142**
  Langerhans cell histiocytosis, 434
  Streptococcus pneumoniae, 136
  Wegener granulomatosis and, 315
Otology, 533
Ototoxicity
  aminoglycosides, **191**, 204, 614
  cisplatin/carboplatin, 442
  drug reaction and, 251
  ethacrynic acid, 608
  loop diuretics, 608
Ouabain
  sodium-potassium pump and, 49
Outcome (quality measurement), 273
Outer membrane, 124
Outflow tract formation, 281
Outpatient follow-up, 272
"Oval fat bodies," 594
Ovarian artery, 625
Ovarian cancer
  breastfeeding and, 636
  cisplatin/carboplatin for, 442
  epidemiology of, 643
  hypercalcemia and, 228
  irinotecan/topotecan for, 442
  Lynch syndrome and, 388
  oncogenes and, 224
  paclitaxel for, 441
  Psammoma bodies in, 227
  tumor suppressor genes and, 224
Ovarian cycle, 632
Ovarian cysts, **646**
Ovarian ligament, 625
Ovarian neoplasms
  classification and characteristics, **646**
  epithelial tumors, 646
  germ cell tumors, 647
  sex cord stromal tumors, 647
Ovarian teratomas
  paraneoplastic syndrome, 228
Ovaries
  anatomy of, 625
  descent of, 624
  epithelial histology, 626
  estrogen production, 630
  lymphatic drainage, 624
Overflow incontinence, 599
Overuse injury
  elbow, **459**
  knee, 461
  radial nerve, 447
  wrist, 459
Ovotesticular disorder, 638
Ovulation
  anovulation causes, 645
  progesterone and, 630
  prolactin effect on, 330
  regulation of, **631**
Ovulatory uterine bleeding, 633
"Owl eyes" inclusions, 165, 429
Oxacillin
  characteristics of, 188
Oxaliplatin, **442**
Oxazepam, 546
Oxazolidinones
  mechanism and use, 193
Oxidative burst, 109
Oxidative phosphorylation, 78
  metabolic site, 72
  poisons, 78
Oxybutynin, 237, **241**
Oxygen
  deprivation, **669**
  in blood, **666**
  for carbon monoxide poisoning, 248
  cluster headaches, 518
  exercise and, 670
  hemoglobin, 665
  toxicity, 210
Oxygen-hemoglobin dissociation curve, **666**
Oxytocin
  functions of, 328
  hypothalamus production, 498
  lactation and, 636
  pituitary gland and, 327
  signaling pathways for, 337

P

P-450, 197
P2Y12 receptor
  inhibitors and, 437
  thrombogenesis and, 411
P53 gene mutation
  dominant negative mutation, 56
Pacemaker action potential, **292**
Pacinian corpuscles, 494
Paclitaxel, 441
  microtubules and, 48
  targets of, 438
Paget disease
  in breast, 650
Paget disease (extramammary), 644
Paget disease of bone, 463
  bisphosphonates, 486
  lab values in, 464
  osteosarcomas and, 465
  woven bone in, 458
Pain
  with headaches, 518
  loss in syringomyelia, 492
  post-stroke syndrome, 515
  receptors for, 494
  spinal tracts for, 509
  thalamic nuclei and, 498
  treatment in multiple sclerosis, 523
  unilateral visual loss and, 523
Pale (anemic) infarct, 210
Paliperidone, 573
Palivizumab, 122
  pneumonia prophylaxis, 169
Pallor in aplastic anemia, 421
Palmar interossei, 450
Palmar reflex, 510
PALM-COEIN uterine bleeding classification, 633
PALP, 653
Panacinar emphysema, 392, **674**
p-ANCA
  sclerosing cholangitis and, 395
  ulcerative colitis, 382
Pancoast tumor, **685**
  lung cancer, 684
  superior vena cava syndrome, 685
  thoracic outlet syndrome, 448
Pancreas
  annular, 360
  biliary structures and, 368
  blood supply and innervation of, 364
  buds, 360
  divisum, 360
  embryology, **360**
Pancreatic cancer
  5-fluorouracil for, 440
  adenocarcinomas, **398**
  biliary cirrhosis and, 395
  carcinogens causing, 225
  hyperbilirubinemia with, 393
  oncogenes and, 224
  paraneoplastic syndromes with, 228
  tumor suppressor genes and, 224

(c) ketabton.com: The Digital Library

Pancreatic ducts, 360, 368
Pancreatic insufficiency, **381**, 397
Pancreatic islet cell tumors, **351**
Pancreatic lipase, 93
Pancreatic secretions, **373**
Pancreatitis
  acute, 397
  acute respiratory distress syndrome and, 678
  alcoholism, 571
  chronic, 397
  corticosteroids and, 249
  drug reactions and, 249
  hyperchylomicronemia, 94
  hyperparathyroidism, 345
  hypertriglyceridemia, 94
  mumps, 170
  necrosis and, 209
  NRTIs, 203
  valproic acid, 544
Pancytopenia, 421
  Chédiak-Higashi syndrome, 117
  cytarabine, 440
  Gaucher disease, 88
  leishmaniasis, 158
  osteopetrosis and, 463
  with immunosuppressants, 120
Panic disorder
  drug therapy for, 572
  SSRIs for, 563, **575**
  symptoms and treatment, **563**
  venlafaxine for, 575
Panitumumab, **442**
Panniculitis, 482
Pantoprazole, 399
Pantothenic acid, 67
Papillary carcinomas, 220
Papillary cystadenoma lymphomatosum, 376
Papillary muscle
  blood supply to, 307
  rupture, 305, **307**
Papillary thyroid carcinomas, 343
  carcinogens for, 225
  Psammoma bodies in, 227
Papilledema, 521, **538**
  hypertensive emergency and, 300
Papillomas, 220
Papillomaviruses
  characteristics of, 164
  DNA viruses, 163
  genome, 163
Pappenheimer bodies, 416
Papules, 475
  capillary, 478
  molluscum contagiosum, 479
Para-aminohippuric acid (PAH), 582
Para-aortic lymph nodes, 624
Paracoccidioidomycosis, 151
Paracortex (lymph node), 96
Paracrine, 589
Paradoxical splitting, 289
Paraesophageal hiatal hernia, 370
Parainfluenza
  croup, 170
  paramyxovirus, 167, 169
Parakeratosis, 475
Paralysis
  conversion disorder and, 566
  of face, 514
  Guillain-Barré syndrome, 524
  poliovirus, 186
  rabies, 171
  stroke effects, 514
  unvaccinated children, 186
Paralytic ileus, 441
Paramedian pontine reticular formation
  extraocular movements, 497
Paramedian pontine reticular formation lesions, 511
Paramesonephric (Müllerian) duct, 622
Paramyxoviruses, **169**
  characteristics of, 167, 168
  croup, 170
  mumps, 170
Paraneoplastic syndromes, **228**, 605
Paranoia
  LSD, 571
Parasites
  infections with immunodeficiency, 118
Parasitic infections
  myocarditis with, 313
Parasitology, 155–**161**
Parasympathetic nervous system
  cranial nerves, supply of, 236
  male erection, 627
  receptor targets, 236
  VIP and, 371
Parathyroid adenomas
  hyperparathyroidism caused by, 345
  MEN 1/MEN 2A syndromes, 351
Parathyroid disease diagnosis, 343
Parathyroid glands
  pharyngeal pouch derivation, 621
Parathyroid hormone (PTH), **332**, 590
  bone disorders, 464
  bone formation, 459
  calcitonin and, 333
  calcium homeostasis and, 333
  in hyperparathyroidism, 345
  osteomalacia/rickets, 463
  Paget disease of bone, 463
  signaling pathways of, 337
  thymic aplasia, 116
Paraumbilical vein, 365
Paraventricular nucleus, 498
Parental consent, 265
Paresthesias
  vitamin B_{12} deficiency, 69, 530
Parietal cells (stomach), 372
Parietal cortex lesions, 511
Parietal lobe, 501
Parietal peritoneum, 369
Parinaud syndrome, **511**, 528
Parkinson disease, 520
  benztropine for, 241
  dopaminergic pathways, 499
  Lewy bodies, 520
  metoclopramide contraindication, 400
  nigrostriatal pathway and, 499
  resting tremor in, 519
  seborrheic dermatitis association, 476
  therapy for, **548**
  trihexyphenidyl, 241
  ubiquitin-proteasome system in, 48
Parkinson-like syndrome, 251
Parotid gland
  embryologic derivation, 613
  enlargement of, 468
  stones in, 376
  tumors in, 376
Parotitis
  mumps, 170
Paroxetine, 575
Paroxysmal nocturnal dyspnea, 309
Paroxysmal nocturnal hemoglobinuria, 422
  CD55 deficiency, 107
  eculizumab for, 122
  flow cytometry diagnosis, 54
  intravascular hemolysis in, 421
Partial agonists, 234
Partial (focal) seizures, 517
  drug therapy for, 544
Partial thromboplastin time (PTT), 426
Parvovirus
  characteristics of, 164
  DNA viruses, 163
  naked viruses, 163
Parvovirus B19
  hereditary spherocytosis, 422
  hydrops fetalis, 182
  rash, 183
Passive aggression, 555
Passive immunity, **110**
Pasteurella multocida
  osteomyelitis, 180
  transmission, **149**, 186
Patau syndrome, 63
  cataracts, 535
  chromosome association, 64
Patches (skin)
  pityriasis rosea, 482
  psoriatic arthritis, 469
Patellar reflex, 510
Patellofemoral syndrome, 461
Patent ductus arteriosus (PDA)
  congenital rubella, 300
  fetal alcohol syndrome, 300
  heart murmur with, 291
  indomethacin for, 486
  mechanism and treatment, 299
  neonatal respiratory distress syndrome and, 661
Patent foramen ovale
  atrial septal defect vs, 299
  septal fusion failure, 280
Patent urachus, 618
Pathogen-associated molecular patterns (PAMPs), 99
Pathogen recognition features, 99
Pathologic grief, 562
Pathology
  cardiovascular, 298–312
  endocrine, 338–354
  gastrointestinal, 376–397
  hematologic/oncologic, 404–424, 414–434
  musculoskeletal/skin/connective tissue, 459–467
  neoplasia, 219–226
  neurological, 511–518
  psychiatric, 556–570
  renal, 594–605
  reproductive, 638–652
  respiratory, 671–681
  USMLE Step 1 preparation for, 277
Pautrier microabscess, 430
Pavlovian (classical) conditioning, 554
Payment models for healthcare, 271
P-bodies, 41
PCP (phencyclidine)
  intoxication and withdrawal, 571
PCSK9, 93
PCSK9 inhibitors, 320
PCV13 (pneumococcal conjugate vaccine), 127
PDSA cycle, **273**
Pearson correlation coefficient(r), **264**
Peau d'orange, 650
Pectinate line, **366**
Pectineus, 451, 452
Pectoriloquy (whispered), 680
Pediatric patients
  arthritis in, 468
  brachial plexus injury, 448
  common causes of death, 272
  common fractures, **462**
  common orthopedic conditions, 461
  cystic fibrosis, 60
  dactinomycin for, 439
  failure to thrive, 556
  growth retardation in, 603
  hemolytic disease of newborn, 405
  hyperbilirubinemia (newborns), 393
  infant deprivation effects, 556
  intraventricular hemorrhage, 512
  juvenile polyposis syndrome in, 387
  leukocoria in, 538
  Munchausen syndrome by proxy, 566
  neglect in, 556
  neuroblastomas in, 350
  precocious puberty, 57, 335
  primary brain tumors, 528
  rashes, 183
  renal malignancy in, 606
  rhabdomyomas in, 316
  scalded skin syndrome, 479
  sleep terror disorder in, 568
  strawberry hemangiomas in, 478
  tetracycline side effects, 192
  unvaccinated, 186
  Wilms tumors in, 606
Pegloticase, 487
Pegvisomant, 339
Pellagra, 67
Pelvic inflammatory disease (PID), **185**
  Actinomyces spp, 139
  chlamydia, 148, 184
  Chlamydia trachomatis, 149
  clinical features, 185
  copper IUD, 657
  ectopic pregnancy, 641
  gonorrhea, 184
  Neisseria spp, 142
Pelvic splanchnic nerves, 236
Pelvis
  fracture and nerve injury, 452
  nerve injury with surgery, 452
Pemphigus vulgaris, 480
  acantholysis and, 475
  autoantibody, 115
  type II hypersensitivity, 112
Pencil-in-cup deformity (X-ray), 469
Penicillamine
  for lead poisoning, 248
  myopathy, 250
  for Wilson disease, 395
Penicillin
  Actinomyces spp, 139
  antipseudomonal, 188
  Coombs-positive hemolytic anemia, 250
  penicillinase-resistant, 188
  penicillinase-sensitive, 188
  prophylaxis, 198
  rash, 250
  for rheumatic fever, 312
Penicillinase-resistant penicillins, **188**

(c) ketabton.com: The Digital Library

Penicillinase-sensitive penicillins, **188**
Penicillin-binding proteins (PBPs)
in bacteria, 124
Penicillin G, V, **187**
meningococci, 142
prophylaxis, 198
Penile cancer, 226
Penis
congenital abnormalities, **624**
female homolog, 624
lymphatic drainage, 624
pathology of, **651**
Pentamidine, 154
Pentazocine, 551, **552**
Pentobarbital, 546
Pentose phosphate pathway
functions of, **79**
Pentostatin, 432
PEP carboxykinase, 74
Pepsin, 372
Pepsinogen
location of, 373
somatostatin and, 371
Peptic ulcer disease, **380**
glycopyrrolate for, 241
H pylori and, 146
H_2 blockers for, 399
misoprostol for, 399
proton pump inhibitors for, 399
Zollinger-Ellison syndrome, 352
Peptidoglycan
in gram negative bacteria, 124
Peptostreptococcus spp
alcoholism, 179
Percussion (chest), 680
Perforation (GI), 380
necrotizing enterocolitis, 386
Perforin
cytotoxic T cells and, 102
extrinsic pathway and, 208
natural killer cells and, 101
Performance anxiety, 567
Perfusion, and ventilation, 669
Perfusion-limited gas exchange, 668
Perfusion pressure regulation, 297
Periarteriolar lymphatic sheath (PALS), 98
Pericardial effusion, 684
Pericarditis
acute, 313
fibrinous, 305
jugular venous pulse in, 287
Kussmaul sign in, 316
picornaviruses, 167
postinfarction, 305, 307
pulsus paradoxus in, 310
renal failure, 603
rheumatoid arthritis, 466
Pericardium
anatomy of, 283
calcification in, 211
Perineal straddle injury, 627
Perinephric abscesses, 600
Perineurium, 495
Periodic acid-Schiff stain, 125
glycogen storage diseases, 87
Periorbital edema
thyroid disease and, 340
Trichinella spiralis, 161
Peripartum mood disturbances, **562**
Peripheral blood smear, 416
basophilic stippling, 419
postsplenectomy, 98, 423
Rouleaux formation, 431
schistocytes, 423
Schüffner stippling, 157
smudge cells, 432
spherocytes and agglutinated RBCs, 423
Peripheral edema
calcium channel blockers, 318
heart failure, 309
Peripheral facial palsy, 532
Peripheral nerves, **495**
Peripheral nervous system (PNS)
origins of, **490**
Peripheral neuropathy
alcoholism, 571
drug reactions and, 251
Fabry disease, 88
isoniazid, 197
Krabbe disease, 88
NRTIs, 203
oxazolidinones, 193
sorbitol, 81
vincristine, 444
vitamin B_6 deficiency, 67
Peripheral precocious puberty, 637
Peripheral resistance, 243
Peripheral vascular disease, 302
Peripheral vertigo, 534
Periplasm
in bacteria, 124
Peristalsis
motilin receptor agonists and, 371
visible, 359
Peritoneum, 360
hernias and, 370
irritation with Mittelschmerz, 631
Peritonitis
appendicitis, 383
diverticulitis, 383
spontaneous bacterial, 389, 390
Periventricular plaques, multiple sclerosis, 523
Permethrin, 200
for scabies, 161
Permissive action
catecholamine responsiveness, 235
Pernicious anemia, 372
autoantibody, 115
B_{12} deficiency, 69, **420**
Peroneus brevis, 453
Peroneus longus, 453
Peroxisome
metabolic processes, **47**
Persistent cervical sinus, 619
Persistent depressive disorder (dysthymia), 561
Persistent thyroglossal duct, 326
Persistent truncus arteriosus, 281, 298
Personality, **565**
Personality disorders, 565
Cluster A, 565
Cluster B, 565
Cluster C, 565, 566
Pertussis, 126
Pertussis toxin, 132, 143
Pes cavus
Friedreich ataxia, 531
Petechiae
aplastic anemia, 421
Peutz-Jeghers syndrome, 220, **387**
PEX genes, 47
Peyer patches, 362, **374**
IgA antibody production, 105
Salmonella/*Shigella* invasion, 144
Peyronie disease, 651
PGI_2, 485
P-glycoprotein, **227**
Phage
bacterial transduction, 130
Phagocytes, 117
Phagocytosis
dendritic cells, 408
eosinophils, 408
group A streptococcal inhibition, 136
M protein prevention of, 129
β-hemolytic bacteria inhibition of, 135
Phalen maneuver, 459
Pharmaceutical company sponsorship, 269
Pharmacokinetics, **231**
Pharmacology, 230–254
autonomic drugs, 236–245
cardiovascular, 316–322
endocrine, 352–354
gastrointestinal, 398–400
hematologic/oncologic, 435–443
musculoskeletal/skin/connective tissue, 485–487
neurology, 544–551
pharmacodynamics, 232–234
pharmacokinetics, 230–231
psychiatric, 572–576
renal, 607–610
reproductive, 655–658
respiratory, 686–687
toxicities and side effects, 248–251
USMLE Step 1 preparation for, 277
Pharyngeal apparatus, **619**
Pharyngeal arch derivatives, **620**
1st pharyngeal arch, 620
2nd pharyngeal arch, 620
4th–6th pharyngeal arches, 620
Pharyngeal cleft derivatives, **619**
Pharyngeal pouch derivatives, **621**
1st pharyngeal pouch, 621
2nd pharyngeal pouch, 621
4th pharyngeal pouch, 621
Pharyngitis
adenovirus, 164
diphtheria, 139
mononucleosis, 165
prophylaxis (rheumatic fever), 198
Streptococcus pyogenes, 136
unvaccinated children, 186
Pharyngoesophageal false diverticulum, 384
Pharynx, 662
blood supply and innervation of, 364
Phenacetin, 606
Phenelzine, 575
Phenobarbital, 546
epilepsy, 544
as weak acid, 233
Phenotypic mixing, 162
Phenoxybenzamine, 244
for pheochromocytomas, 350
Phentolamine, 244
Phenylalanine
classification of, 81
tyrosine catabolism, 83
Phenylephrine, 242, **686**
Phenylketones, 84
Phenylketonuria
tyrosine in, **84**
Phenytoin
cytochrome P-450 and, 252
epilepsy, 544
folate deficiency caused by, 420
gingival hyperplasia, 250
megaloblastic anemia, 250
peripheral neuropathy, 251
vitamin B_9 deficiency, 68
zero-order elimination of, 232
Pheochromocytomas, **350**
MEN 2A/MEN 2B and, 351
phenoxybenzamine for, 244
Philadelphia chromosome
in myeloproliferative disorders, 433
translocations of, 434
Phlebitis
IV amphotericin B, 199
Phlebotomy
for hemochromatosis, 395
Phobias, **563**
agoraphobia, 563
social anxiety disorder, 563
Phocomelia, 614
Phonophobia, migraine headache, 518
Phosphatases, 73
Phosphodiesterase (PDE) inhibitors, **246**, 658, 686
Phosphoenolpyruvate carboxykinase, 78
Phosphofructokinase-1 (PFK-1)
glycolysis and, 73
metabolic pathways, 74
Phospholipid bilayer sac
in bacteria, 124
Phospholipids, 374
Phosphorus in Paget disease of bone, 463
Phosphorylases, 73
Phosphorylation, 45
Photophobia
leptospirosis, 147
migraine headache, 518
rabies, 171
Photosensitivity
drugs causing, 192, 194, 250
Phototherapy for jaundice, 393
Phrenic nerve, 663
Phyllodes tumor, 649
Physical abuse (child), 556
Physician-assisted suicide, 268
Physician-patient relationship, 268
Physiologic dead space, 664
Physiologic neonatal jaundice, **393**
Physiology
cardiovascular, 284–298
endocrine, 328–336
gastrointestinal, 371–375
hematologic/oncologic, 410
neurological, 493–515
renal, 581–592
reproductive, 629–636
respiratory, 664–669
USMLE Step 1 preparation for, 276
Physostigmine
anticholinergic toxicity treatment, 248
anticholinesterase, 240
glaucoma, 552
Pia mater, 496
Pica, 567
Pick disease, 520
bodies, 520
Pickwickian syndrome, 679
Picornaviruses, 163, **168**
characteristics, 167
genomes, 163
Pierre Robin sequence, 620
Pigmented skin disorders, **476**
Pigment-producing bacteria, **128**
Pigment stones, 396
Pill-rolling tremor, 519

(c) ketabton.com: The Digital Library

Pilocarpine, **240**, 552
Pilocytic astrocytoma, 528
Pilus, 124
Pimozide, 572
Pindolol, 245, 319
Pineal gland, 504
Pinealoma, 528
Pinworms, 159
Pioglitazone, 353
Piperacillin
  characteristics of, 188
  Pseudomonas aeruginosa, 143
Piroxicam, 486
Pisiform bone, 449
Pitting edema, 309
Pituitary adenoma, 339, 527
Pituitary apoplexy, 339
Pituitary drugs, 354
Pituitary gland, **327**
Pituitary prolactinomas, 328
Pituitary tumors
  diabetes insipidus, 338
  MEN 1 and, 351
Pityriasis rosea, 482
Pityrosporum spp, 152
pKa, 233
PKD genes
  renal cyst disorders and, 604
Placebo, 256
Placenta
  estrogen production, 630
  fetal component, **617**
  hormone secretion by, 633
  maternal component, 617
  progesterone production, 630
Placenta accreta/increta/percreta, 640
Placental aromatase deficiency, **639**
Placental insufficiency
  oligohydramnios and, 641
  Potter sequence, 578
  preeclampsia, 643
Placenta previa, 640
Plague, 149
Plantar aponeurosis, 461
Plantar fasciitis, 461
Plantar flexion, 453
Plantaris, 453
Plantar reflex, 510
Plaques (skin), 475
  actinic keratosis, 482
  basal cell carcinoma, 484
  hairy leukoplakia, 479
  lichen planus, 482
  pityriasis rosea, 482
  psoriasis, 477
  seborrheic dermatitis, 476
  squamous cell carcinoma, 484
Plasma cell dyscrasias, **431**
Plasma cells, **409**
Plasmalogens, 47
Plasma membrane
  sodium-potassium pump, 49
  structure of, 49
Plasma osmolality
  DI treatment, 338
Plasmapheresis, 524, 596
Plasma protein concentration, 583
Plasma volume measurement, 581
Plasminogen, 437
Plasmodium spp
  chloroquine, 200
Plasmodium falciparum, 157, 200
Plasmodium malariae, 157
Plasmodium ovale, 157
Plasmodium vivax, 157
Platelet-activating factor, 406
Platelet-derived growth factor (PDGF)
  in wound healing, 216
  signaling pathways for, 337
Platelet disorders, 426, **427**
  transfusion for, 429
Platysma muscle, 620
Pleiotropy, 56
Pleomorphic adenomas, 376
Pleural effusion, **681**
  asbestosis, 677
  lung cancer, 684
  mesothelioma, 678
  physical findings, 680
Pleuritis, 466
Pleuroperitoneal membrane, 370
Plicae circulares, 362
Plummer-Vinson syndrome, 377, 418
Pneumatosis intestinalis, 386
Pneumococcal vaccine, 127
Pneumoconioses, 675, **677**
Pneumocystis jirovecii, **154**
  dapsone, 194
  fluorescent antibody stain, 125
  HIV-positive adults, 177
  immunocompromised patients, 179
  prophylaxis, 198
  silver stain for, 125
  TMP-SMX, 194
Pneumocytes, 661
Pneumomediastinum, 672
Pneumonia, **683**
  acute respiratory distress syndrome, 678
  adenovirus, 164
  chlamydiae, 148
  coccidioidomycosis, 151
  common causes, **179**
  Haemophilus influenzae, 142
  measles-associated death, 170
  Pneumocystis jirovecii, 154
  PPI adverse effects, 399
  Q fever, 150
  readmissions with, 272
  Staphylococcus aureus, 135
  Streptococcus pneumoniae, 136
  Streptococcus agalactiae, 137
  VZV, 165
Pneumoperitoneum, 380
Pneumothorax, 680, **682**
Podocytes
  glomerular filtration barrier and, **581**, 595
Poikilocytosis, 407
Point of service plan, 271
pol gene, 175
Poliomyelitis, **531**
  restrictive lung disease, 675
Poliovirus, 531
  immunodeficient patients, 118
  medical importance, 167
  picornavirus, 168
  unvaccinated children, 186
Polyadenylation signal, 41
Polyangiitis, microscopic
  autoantibody, 115
Polyarteritis nodosa, 173, **314**
Polyarthralgias
  gonococcal arthritis, 468
  rubella, 182
Polycystic disease (kidney), 604
Polycystic ovarian syndrome (PCOS), **645**
  anovulation, 645
  antiandrogens, 658
  clomiphene, 656
Polycythemia, **434**
  blood oxygen in, 666
  Eisenmenger syndrome, 299
  ESR in, 214
  paraneoplastic syndromes, 228
Polycythemia vera, 433
  Budd-Chiari syndrome and, 392
  hepatocellular carcinoma, 392
Polydactyly, 63
Polydipsia, 346
Polyhydramnios, 491, **641**
  esophageal atresia and, 359
Polymerase chain reaction (PCR), **52**
Polymorphic ventricular tachycardia, 294
Polymyalgia rheumatica, **470**
  ESR in, 214
  giant cell arteritis and, 314
Polymyositis
  autoantibody, 115
Polymyositis/dermatomyositis, **471**
Polymyxin B, 143, **193**, 198
Polymyxins, 198
  mechanism and use, **193**
Polyneuritis, 66
Polyneuropathy, 425
  familial amyloid, 212
Polyomaviruses
  characteristics of, 164
  DNA viruses, 163
  genome, 163
  naked viruses, 163
Polyostotic fibrous dysplasia, 57
Polyposis syndromes, **387**
Polyps
  adenomatous, 387
  APC gene, 387
  colonic, 387
  hyperplastic, 387
  inflammatory pseudopolyps, 387
  KRAS gene, 387
  mucosal, 387
  nasal (cystic fibrosis), 60
  neoplastic transformation of, 387
  non-neoplastic, 387
  serrated, 387
  submucosal, 387
  uterine, 650
Polysaccharide capsule antigens
  carrier proteins with, 127
Polyuria
  diabetes insipidus, 338
  diabetes mellitus, 346
  hyperparathyroidism, 345
  lithium, 574
Pompe disease, 87
Pons
  cranial nerve nuclei of, 505
  development of, 490
Pontiac fever, 143
Pontine syndrome, 514
"Pope's blessing" (median nerve injury), 451
Popliteal artery, 455
  atherosclerosis in, 302
Popliteal fossa, 455
Popliteus, 453
Porcelain gallbladder, 396, 397
Porphobilinogen deaminase, 425
Porphyria, 425, 546
Porphyria cutanea tarda, 425
Portal hypertension
  ARPKD, 604
  cirrhosis and, 389
  pulmonary arterial hypertension, 679
  Schistosoma spp, 161
  serum markers for, 390
  varices and, 365
Portal triad, 361, **367**
Portal vein, 361, **367**
  in fetal circulation, 282
Portosystemic anastomoses, **365**
Positive predictive value (PPV), 257, 259
Positive punishment (aversive stimulus), 554
Positive reinforcement, 554
Positive skew distribution, 262
Posterior cerebral artery, 502, 515
Posterior circulation strokes, 514
Posterior circumflex artery, 455
Posterior communicating artery
  in saccular aneurysm, 516
Posterior cruciate ligament (PCL) injury, 454
Posterior descending artery (PDA), 283
Posterior drawer sign, 454
Posterior fossa
  malformations, **492**
Posterior hypothalamus, 498
Posterior inferior cerebellar artery
  stroke effects, 514
Posterior malleolus, 455
Posterior pituitary gland, 327
Posterior tibial artery, 455
Posterior urethral valves, **579**
Postherpetic neuralgia, 165
Postinfectious encephalomyelitis, 524
Postoperative ileus, 240
Postpartum hemorrhage, 641
Postpartum mood disturbances, 562
Postpartum psychosis, 562
Postpartum thyroiditis, 341
Postprandial pain, 363
Postrenal azotemia, 601
Poststreptococcal glomerulonephritis (acute), 596
Post-traumatic stress disorder (PTSD)
  acute stress disorder, 564
  diagnostic criteria/treatment, 564
  dissociative identity disorder, 558
  drug therapy for, 572
  prazosin for, 244
  SSRIs for, 575
  venlafaxine, 575
Postural hypotension
  midodrine for, 242
  trazodone, 576
Postviral infections, 179
Potassium
  amphotericin B, 199
  in cardiac muscle, 292
  diabetic ketoacidosis, 347
  shifts in, 590
  torsades de pointes and, 294
Potassium channel blockers, **323**
Potassium channels
  myocardial action potential, 292
  opioid effect, 551
Potassium chloride, 249
Potassium iodide
  Sporothrix schenckii, 154
  for thyroid storm, 342
Potassium-sparing diuretics, **609**
Potency of drugs
  vs efficacy, 233
Potentiation
  of drugs, 235
Pott disease, 180

(c) ketabton.com: The Digital Library

Potter sequence (syndrome)
ARPKD, 604
oligohydramnios and, **578**, 641
pulmonary hypoplasia, 660
Poxvirus, 164, 479
PPD test
for tuberculosis, 140
PPSV23 (pneumococcal polysaccharide vaccine), 127
PR3-ANCA/c-ANCA autoantibody, 115
Practice tests, 21
Prader-Willi syndrome
chromosome association, 64
ghrelin in, **336**, 371
Pramlintide, 249, 353
Prasugrel, 411, 437
Praziquantel
antihelminthic therapy, 200
tapeworms, 160
trematodes, 160
Prazosin, 244
Precision vs accuracy, **259**
Precocious puberty
adrenal steroids and, 335
leuprolide, 656
McCune-Albright syndrome, 57
pinealoma, 528
types, **637**
Precontemplation stage, substance addiction, 568
Predictive value, 257
Prednisolone
for thyroid storm, 342
Preeclampsia, 643
hydatidiform moles, 642
Preferred provider organization plan, 271
Prefrontal cortex, 501
Pregnancy
advanced maternal age, 63
aliskiren contraindication, 610
amniotic fluid abnormalities, 641
anemia caused by, 418
carpal tunnel syndrome in, 459
choriocarcinomas and, 642
contraindicated antimicrobials, 204
ESR in, 214
estrogen in, 630
ethical situations, 268–269
fetal circulation, 282
fetal respiration, 660
folate deficiency caused by, 420
folic acid supplementation, 68
heparin in, 436
hypertension and treatment in, 243, 316, 643
hypothyroidism in, 341
insulin in, 334
Listeria monocytogenes in, 139
lithium in, 298, 300
maternal complications, 272
maternal phenylketonuria, 84
opiate use during, 615
parental consent and, 265
physiologic adaptations in, **633**
pituitary infarcts with, 339
posterior urethral valve diagnosis, 579
progesterone in, 630
prolactin and, 330
pyelonephritis, 600
pyogenic granulomas and, 478
sex hormone-binding globulin, 337
stillbirth, 182
Streptococcus agalactiae in, 137
syphilis in, 147
TBG in, 331
termination of, 657
TORCH infections, 182
Turner syndrome and, 638
twinning in, 616
urinary tract infections, 181
venous sinus thrombosis in, 503
vitamin B_9 deficiency, 68
Pregnancy complications, **640–641**
Prehn sign, 654
Preload in cardiac output, 284
Premature ejaculation, 575
Premature labor and delivery
cryptorchidism and, 651
low birth weight with, 635
murmur in prematurity, 291
Premature ovarian failure, 636, 645
Premotor cortex, 501
Preoptic nucleus, 498
Prepatellar bursitis, 460
Preprocollagen, 51
Preproinsulin, 334
Prerenal azotemia, 601
Presbycusis, 270, 533
Presbyopia, **535**
Preschool age development, 635
Presenilin, 520
Pressure-volume loops, **287**
Presynaptic α2-autoreceptor, 239
Pretectal nuclei, 539
Preterm birth
common cause of death, 272
Pretest probability, 257
Prevalence
diagnostic test evaluation, 257
incidence vs, **259**
observational studies, 256
Prevotella spp, 179
Priapism, 651
sickle cell anemia, 422
trazodone and, 576
Primaquine, 157
hemolysis in G6PD deficiency, 250
Primary adrenal insufficiency, 349
Primary amyloidosis, 212
Primary bacterial peritonitis, 390
Primary biliary cholangitis, 395
Primary central nervous system lymphoma, 430
Primary ciliary dyskinesia, 49
Primary disease prevention, 270
Primary glomerular disease, 594
Primary hemostasis, **411**
Primary hyperaldosteronism, 349
hypertension with, 300
markers in, 591
Primary hyperparathyroidism, 345, 464
Primary hypertension, 316
Primary hypogonadism, 639
Primary ovarian insufficiency, **645**
Primary polycythemia, 433
Primary sclerosing cholangitis, 395
ulcerative colitis, 382
Primary spontaneous pneumothorax, 682
Primase
replication initiation by, **38**
Primidone, 519
Primitive atrium, 281
Primitive pulmonary vein, 281
Primitive reflexes, **510**
Primitive ventricle, 281
Pringle maneuver, 361
PR interval, 293, 295
antiarrhythmic effects, 323, 324
prolonged, 295
shortened, 294
Prinzmetal angina
calcium channel blockers for, 318
ischemic manifestations, 304
propranolol adverse effects, 323
Prions, **178**
Privacy and confidentiality, 267
Probenecid, 252
cidofovir with, 202
gout, 487
Procainamide, 322
Procaine, 550
Procarbazine, 251, **441**
Procedure bias, 260
Process improvement model, 273
quality measurement, 273
Processus vaginalis, 624
Procoagulation, 413
Progesterone
lactation and, 636
menstrual cycle, 632
ovulation, 632
signaling pathways for, 337
source and function of, **630**
Progestins, **657**
Progressive multifocal leukoencephalopathy (PML), 494, **524**
HIV-positive adults, 177
polyomaviruses, 164
rituximab, 443
Proguanil, 200
Projection, 555
Prokaryotes
DNA replication in, 38
mRNA start codons, 44
RNA polymerases in, 42
Prolactin, **330**
circadian rhythm, 497
function of, 328
lactation and, 636
secretion of, 327
signaling pathways for, 337
tuberoinfundibular pathway, 499
Prolactin-inhibiting factor, 328
Prolactinomas
dopamine agonists for, 330
Proliferative glomerular disorders, 594
Prometaphase, 46
Promoters (gene expression), 41
Promyelocytic leukemia, 66
Pronephros, 578
Proopiomelanocortin, 327
Propafenone, 322
Proper hepatic artery, 361
Prophase, 46
Prophylaxis
antimicrobial, **197**, 198
antimycobacterial drugs for, 196
HIV/AIDS patients, **198**
in HIV/AIDS patients, 198
Pneumocystis jirovecii, 154
rabies postexposure, 171
for RSV, 169
Trichomonas vaginalis, 158
Propionyl-CoA carboxylase
metabolic pathways, 74
vitamin B_7 and, 68
Propofol, 550
Propranolol, 245, 323, 342
essential tremor, 519
Proprioception
Friedreich ataxia, 531
Propylthiouracil
agranulocytosis, 250
aplastic anemia, 250
thionamides, 354
for thyroid storm, 342
Prostacyclin analogues, 686
Prostaglandin analogs, 254
Prostaglandins
aspirin effects, 486
cortisol effect on, 336
glaucoma treatment, 552
kidney functions, 589
Prostate cancer
adenocarcinomas, **654**
incidence/mortality of, 222
leuprolide for, 656
metastases of, 223
Prostate gland
lymphatic drainage of, 624
with urethral injury, 627
Prostate-specific antigen (PSA), 227, 654
Prostatic acid phosphatase (PAP), 654
Prostatic adenocarcinoma, 654
Prostatitis, **654**
gonorrhea, 184
Prosthetic devices
Staphylococcus epidermidis, 135
Prosthetic heart valves, 423
Protamine sulfate, 248, 436
Protease inhibitors
fat redistribution, 250
HIV therapy, 203
hyperglycemia, 249
naming convention for, 253
Proteases, 373
Proteasome, **48**
Protein A, 129
Proteinases, 406
Protein C/S deficiency, 428
Protein-energy malnutrition, **71**
Protein kinase A
fructose bisphosphatase-2 and, 76
Protein metabolism
amino acids, 81
Proteins
free radical effect on, 210
Protein synthesis
elongation, 45
inhibitors, **191**, 253
initiation of, 44, **45**
insulin and, 334
metabolic site, 72
posttranslational modification, 45
sequence of, 45
termination, 45
trimming, 45
Proteinuria
ACE inhibitors for, 610
angiotensin II receptor blockers, 610
diabetes mellitus, 346
nephritic syndrome, 595
nephrotic syndrome, 595, 597
preeclampsia, 643
renal papillary necrosis and, 602
serum sickness, 113
Proteolysis
cortisol and, 336
Proteolytic processing, in collagen synthesis, 50

(c) ketabton.com: The Digital Library

Proteus spp
struvite stones with, 127
urease-positive, 127
xanthogranulomatous pyelonephritis, 600
Proteus mirabilis
cephalosporins, 189
penicillins for, 188
urinary tract infections, 181, 600
Prothrombin
complex concentrate transfusion, 429
warfarin effect on, 436
Prothrombin gene mutation, 428
Prothrombin time, 390
Protofilament, 48
Proton pump inhibitors, 254, **399**
Beers criteria, 247
for *Helicobacter pylori*, 146
gastrin and, 371
Protozoa
CNS infections, **156**
GI infections, **155**
hematologic infections, **157**
miscellaneous, **158**
watery diarrhea, 179
Proximal convoluted tubules (PCT)
in ATN, 602
defects in, 586
diuretics and, 609
dopamine secretion by, 589
glucose clearance and, 584
ischemia susceptibility, 210
relative concentrations in, 587
renal cell carcinoma and, 605
Proximal interphalangeal (PIP) joints, 451
Proximal renal tubular acidosis (type 2), 593
PRPP (glutamine-phosphoribosyl-pyrophosphate) amidotransferase, 73
Pruritus
anal, 159
atopic dermatitis, 477
biliary tract disease, 395
chloroquine, 200
cutaneous mycoses, 152
ectoparasites, 161
histamine receptors and, 238
hyperchylomicronemia, 94
lichen planus, 482
pseudofolliculitis barbae, 477
urticaria, 477
Prussian blue stain, 677
Psammoma bodies, 211, **227**
mesotheliomas, 678
papillary thyroid carcinoma, 343
Pseudoappendicitis
Yersinia enterocolitica, 144
Pseudocysts, 397
Pseudoephedrine, **686**
Pseudofolliculitis barbae, 477
Pseudofractures, 463
Pseudoglandular stage (development), 660
Pseudogout, 467
Pseudohermaphrodites, 639
Pseudohypoparathyroidism, 344
Pseudomembranous colitis
clindamycin, 192
Clostridium difficile, 138
drug reaction and, 249
penicillins, 188
vancomycin for, 190
watery diarrhea, 179
Pseudomembranous pharyngitis diphtheria, 139
Pseudomonas spp
aeruginosa, **143**
biofilm production, 128
ceftazidime, 189
culture requirements for, 126
cystic fibrosis, 60, 179
encapsulated, 127
epididymitis and orchitis, 654
exotoxin production, 132
fluoroquinolones, 195
immunodeficient patients, 118
multidrug-resistant, 198
nosocomial infection, 179, 185
osteomyelitis, 180
penicillins for, 188
pigment production, 128
pyocyanin of, 109
tricuspid valve endocarditis, 311
UTIs, 181
Pseudo-Pelger-Huet anomaly, 432
Pseudopseudohypoparathyroidism, 344
Pseudotumor cerebri, 521
drug reactions and, 251
vitamin A toxicity, 66
Pseudovirion, 162
Psittacosis, 149
Psoriasis, 477
arthritis and, 469
cyclosporine, 120
etanercept for, 487
hyperkeratosis/parakeratosis, 475
infliximab/adalimumab for, 487
methotrexate for, 440
skin lesions, 475
therapeutic antibodies, 122
Psoriatic arthritis, 469
HLA-B27 and, 100
leflunomide for, 486
psoriasis and, 477
Psychiatric emergencies
acute dystonia, 569
delirium tremens, 569
hypertensive crisis, 569
lithium toxicity, **569**
neuroleptic malignant syndrome, 569
serotonin syndrome, 569
tricyclic antidepressant overdose, 569
Psychiatry, 554–576
pathology, 556–570
pharmacology, 572–576
psychology, 554–555
Psychoactive drug intoxication/withdrawal, **570–571**
Psychological child abuse, 556
Psychology, 554–555
Psychosis, **559**
corticosteroids, 120
diabetic ketoacidosis, 347
drug therapy for, 573
LSD and, 571
major depressive disorder with, 561
PCP and, 571
postpartum, 562
Psychotherapy, **572**
oppositional defiant disorder, 557
Psychotic disorders
readmissions with, 272
PTEN gene, 224
Pterygoid muscles, 507
PTH-related peptide (PTHrP), 332
PTHrP (parathyroid hormone-related protein), 228
Ptosis
CN III damage, 541
Horner syndrome, 540
myasthenia gravis, 472
saccular aneurysm, 516
Puberty
GnRH and, 328
Kallmann syndrome and, 639
precocious, 57, 335
Tanner stages, 637
Pubic tubercle, 370
Public health sciences, 256–273
Pudendal nerve, 366, 453
Pulmonary anthrax, 137
Pulmonary arterial hypertension (PAH), 679
Pulmonary artery, 619
fetal circulation, 282
Pulmonary artery stenosis, 300
Pulmonary capillary wedge pressure (PCWP), 297, 668
Pulmonary circulation, **668**
Pulmonary edema, 309
consolidation in, 680
LV failure, 307
mannitol, 607
nitrates for, 318
opioids for, 551
preeclampsia and, 643
Pulmonary embolism, **672**
chronic thromboembolism, 679
direct factor Xa inhibitors for, 437
heparin for, 436
tamoxifen/raloxifene and, 443
thrombolytics for, 437
ventilation/perfusion with, 669
Pulmonary fibrosis
amiodarone and, 323
bleomycin, 439
busulfan, 441
diffusion in, 668
drug reaction and, 251
methotrexate, 440
restrictive lung disease, 675
Pulmonary hypertension, **679**
cor pulmonale, 668
drug therapy, **686**
Schistosoma, 160
Pulmonary hypoplasia, 660
Pulmonary infections
in immunocompromised patients, 139
Pulmonary Langerhans cell histiocytosis, 675
Pulmonary surfactant
club cells, 660, 661
NRDS, 661
Pulmonary vascular resistance (PVR), **668**
Pulmonic stenosis
wide splitting in, 289
Pulmonic valves, 281
"Pulseless disease," 314
Pulse pressure
equation for, 285
Pulsus paradoxus, 310
croup, 170
"Punched out" bone lesions (X-ray), 431
Punched-out ulcers, 377
Punishment, 554
Pupil
Argyll Robertson, 147
CN III palsy, 541
control, 507, **539**
light reflex, 539
pupillary light reflex, 539
syphilis, 147, 184
Pure motor stroke, 514
Pure red cell aplasia, 228
thymoma and, 98
Purines, 194
de novo synthesis, 36, 73
Lesch-Nyhan syndrome, 37
salvage deficiencies, **37**
structure of, 35
Purkinje cells
cerebellum, 499
of cerebellum, 210
in paraneoplastic cerebellar degeneration, 228
Purkinje fibers, 293
Purpura
aplastic anemia, 421
Pustular psoriasis, 475
Pustules, 475
acne, 477
pseudofolliculitis barbae, 477
rosacea, 477
Putamen, 500
neurodegenerative disorders, 520
Pyelonephritis, **600**
kidney stones, 598
urinary tract infections, 181, 600
WBC casts in, 594
Pygmalion effect, 260
Pyloric sphincter, 373
Pyloric stenosis, 359
Pyloromyotomy, 359
Pyoderma gangrenosum
inflammatory bowel disease, 382
Pyogenic granulomas, 478
Pyramidal cells, 210
Pyramidalis muscle, 369
Pyramidal tract demyelination, multiple sclerosis, 523
Pyrantel pamoate, 200
Pyrazinamide, **197**
gout, 250
hepatitis, 249
Mycobacterium tuberculosis, 196
Pyridostigmine, 240
myasthenia gravis treatment, 472
Pyridoxal phosphate, 67
Pyridoxine, 67
Pyrimethamine, 200
effect on purine synthesis, 36
Pyrimidine dimers, 40
Pyrimidines
de novo synthesis, 36
structure of, 35
Pyrimidine synthesis, 486
Pyruvate carboxylase, 77, 78
metabolic pathways, 74
vitamin B_7 and, 68
Pyruvate dehydrogenase
complex, **76**
deficiency, **77**
metabolic pathways, 74
vitamin B_1 and, 66
Pyruvate kinase, 74
deficiency, 422
Pyruvate metabolism, **77**
Pyuria, 601

(c) ketabton.com: The Digital Library

Q

Q fever
  rickettsial disease, 150
  transmission, 149
QRS complex, 293
QT interval
  atypical antipsychotic effect on, 573
  Class IA antiarrhythmic effects, 322
  congenital long QT syndrome, 294
  drug-induced long, 294
  ECG, 293
  ondansetron effect on, 400
  in torsades de pointes, 294
Quadrantanopia, 542
Quadriceps, 452
Quality measurements, **273**
Quantifying risk, **258**
Quaternary amines, 204
Quaternary disease prevention, 270
Quetiapine, 573
Quiescent (stable) cells, 46
Quinidine, 157, 200, 322
  cinchonism, 251
Quinine, 200
Quinolone
  Legionella pneumophila, 143
Quinupristin, 198

R

Rabies, **171**
  active and passive immunity, 110
  rhabdovirus, 167
  viral receptors, 166
Rachischisis, 491
Rachitic rosary, 463
Radial head subluxation, 461
Radial nerve, 447
  neurovascular pairing, 455
Radiation exposure
  acute myelogenous leukemia and, 432
  aplastic anemia, 421
  apoptosis caused by, 208
  as carcinogen, 225
  free radical injury caused by, 210
  hypopituitarism, 339
Radiation therapy
  acute pericarditis and, 313
  angiosarcomas, 478
  lymphopenia, 424
  neutropenia, 424
  osteosarcoma and, 465
  pancreatic cancer, 398
  papillary thyroid carcinoma risk, 343
  readmissions with, 272
Radiculopathy
  lumbosacral, 455
Radon
  as carcinogen, 225
Ragged red muscle fibers, 59
Rales, 309
Raloxifene, 443, 656
Raltegravir, 203
Ramelteon, **547**
Ramipril, 610
Ranitidine, 399
RANKL (RANK ligand), 332, 459
Ranolazine
  mechanism and clinical use, **319**
Raphe nucleus, 495
Rapid automated broth cultures, 126
Rapid-eye movement (REM) sleep, 497
Rapid filling (cardiac cycle), 287
Rapidly progressive glomerulonephritis (RPGN), 596
Rasagiline, **549**
Rasburicase, **444**
RAS gene, 343
Rashes
  "blueberry muffin," 169
  carbapenems, 190
  childhood, 183
  cytomegalovirus, 182
  desquamating, 314
  fluoroquinolones, 195
  heliotrope, 228
  macrolides, 193
  palms and soles, 150
  penicillinase-sensitive penicillins, 188
  rickettsial infections, 150
  rubella, 169, 182
  syphilis, 147, 184
  unvaccinated children, 186
Rathke pouch, 327, 528
  tumor, 613
Rationalization, 555
Raynaud phenomenon, **472**
  Buerger disease, 314
  calcium channel blockers for, 318
  scleroderma and, 473
Razor bumps, 477
RBC casts (urine), 594
RBC inclusions, **416**–444
RBC morphology (pathologic), **414**
Rb gene, 224
Reabsorption/secretion rate calculation, 584
Reaction formation, 555
Reactive arthritis, 469
  Campylobacter jejuni, 145
  chlamydia, 148, 184
  HLA-B27 and, 100
Reactive attachment disorder, 556
Readmission recurrences, 272
Reassortment (viral), 162
Recall bias in studies, 260
Receiving operating characteristic curve, **260**
Receptor binding, **234**
Receptor fusion proteins
  naming conventions for, 254
Receptor-mediated endocytosis, 47
Receptors (viral), **166**
Receptor tyrosine kinase
  hormone messenger, 337
  as oncogene product, 224
Recessive inheritance, 59
Recklinghausen disease, 525
Recombinant cytokines, **121**
Recombination (viral), 162
Recruiting study participants, 260
Rectal sparing, 382
Rectosigmoid junction
  blood supply to, 363
Rectum
  blood supply and innervation, 364
  familial adenomatous polyposis, 387
  Hirschsprung disease, 384
  ischemia susceptibility, 210
  portosystemic anastomosis, 365
Rectus abdominis muscle, 369
Recurrent branch (median nerve), 447
Recurrent laryngeal nerve
  compression of, 684
  Pancoast tumor, 685
Red cell casts, 315
Red-green color blindness, 197
Red hemorrhagic infarct, 210
Red hepatization, 683
Red nucleus (midbrain), 511–552
Redox reactions
  free radical injury and, 210
  vitamin B_2 and, 67
Red pulp (spleen), 98
Red rashes of childhood, **183**
Reduced filling (cardiac cycle), 287
Redundant/degenerate genetic code, 37
Reed-Sternberg cells, 429
Refeeding syndrome, anorexia nervosa, 567
Referred pain
  cholecystitis, 396
  from diaphragm, 663
Reflex bradycardia, 588
Reflexes
  clinical, 510
  cranial nerve, 507
  motor neuron sign, 529
  primitive, 510
Reflex tachycardia, 244
Refractive errors (vision), **535**
Refractory hypertension, 658
Refsum disease, 47
Refusing care
  minors, 269
Regadenoson, 304
Regan-Lowe medium, 126
Regional specification (brain), **490**
Registering for exam, 5–6
Regression, 555
Regulation of cell cycle
  Cyclin-dependent kinases (CDKs), 46
  p53, 46
  Tumor suppressors, 46
Regulation of gene expression, **41**
Regulatory T cells, **102**
  cell surface proteins, 110
Regurgitation
  in GERD, 377
Reheated rice syndrome, 138
Reichert cartilage, 620
Reid index, 674
Reinforcement, 554
Relapse stage, substance addiction, 568
Relapsing fever
  animal transmission, 149
  lice, 161
Relationship with patients, 268
Relative risk reduction (RRR), 258
Relative risk (RR), 256, **258**, 263
Reliability, 259
Remodeling (tissue), 216
REM sleep, 497
Renal agenesis
  bilateral, 578
  unilateral, 579
Renal artery, 580
  stenosis, 610
Renal blood flow (RBF), 363, 580
  acute injury and, 602
  NSAID effects on, 589
  renal plasma flow and, 582
Renal cell carcinoma, **605**
  bevacizumab for, 442
  carcinogens for, 225
  chromosome association, 64
  hypercalcemia and, 228
  immunohistochemical stain for, 227
  metastases of, 223
  recombinant cytokines, 121
  therapeutic antibodies, 122
Renal clearance, **582**
Renal cortex
  atrophy of, 599
Renal disorders/failure
  consequences of, **603**
  diabetes mellitus, 346
  diffuse cortical necrosis, 602
  ESR in, 214
  Fabry disease, 88
  features of, 591
  gout and, 467
  in utero, 578
  markers for, 591
  myoclonus in, 519
  NSAIDs, 589
  preeclampsia and, 643
  renal cyst disorders, **604**
  tetracycline use in, 192
  waxy casts in, 594
  Wilson disease, 395
Renal/genitourinary drug reactions, 251
Renal hypoxia, 666
Renal ischemia, 486
Renal medulla
  hydronephrosis, 599
Renal oncocytoma, **605**
Renal osteodystrophy, 345, **603**
Renal papillary necrosis, **602**
  pyelonephritis and, 600
  sickle cell anemia, 422
Renal plasma flow, 582
  glomerular dynamics and, 583
Renal sympathetic discharge, 588
Renal toxicity
  ganciclovir, 202
Renal tubular acidosis (RTA), **593**
Renal tubular defects, 585, **586**
Renal vascular smooth muscle, 238
Renin, 588
  ACE inhibitor effect on, 610
  aliskiren effect on, 610
  in hyperaldosteronism, 349
  renal disorders and, 591
  sympathetic receptors and, 238
Renin-angiotensin-aldosterone system, 327, 588
Renovascular disease, **604**
Renovascular hypertension, 349
Reoviruses
  characteristics, 167
  genome, 163
  naked viruses, 163
  segmented, 168
Repaglinide, 353
Reperfusion injury, 210
Reperfusion therapy, 307
Replication fork, 38
Reportable diseases
  confidentiality exceptions, 267
Repression, 555
Repressor proteins
  lactose effects on, 40
Reproductive/endocrine drug reactions, 249
Reproductive hormones, **655**
Reproductive system, 612–653
  anatomy, 624–627
  embryology, 612–623
  pathology, 638–652
  pharmacology, 655–658
  physiology, 629–636
Reptile (disease vectors), 149
RER staining, 493

(c) ketabton.com: The Digital Library

Rescheduling exam, 6
Reserpine
  Parkinson-like syndrome, 251
Residual volume (RV), 664
Resistance in vessels, 286
Respiration
  exercise response, 670
  high altitude response, 670
  in diabetic acidosis, 347
  Kussmaul, 347
Respiratory acidosis, 592
Respiratory alkalosis, 592
  high altitude, 670
Respiratory burst, **109**
  free radical injury and, 210
Respiratory depression
  anesthetics, 550
  barbiturates, 546, 570
  benzodiazepines, 544, 570
  epilepsy drugs, 544
  inhaled anesthetics, 550
  opioids, 551, 570
  psychoactive drug intoxication, 570
  tricyclic antidepressants, 575
Respiratory drug reactions, 251
Respiratory syncytial virus (RSV)
  paramyxovirus, 167, 169
  pneumonia, 179, 683
  prophylaxis, 122
Respiratory system, 660–683
  anatomy, 662–663
  change in elderly, **665**
  embryology, 660
  pathology, 671–681
  pharmacology, 686–687
  physiology, 664–669
Respiratory tract infections
  C3 deficiency, 107
Respiratory tree, **662**
Respiratory zone, 662
Resting tremor, 519
Restless legs syndrome, 519
Restricting type of anorexia nervosa, 567
Restrictive cardiomyopathy, 308
  hemochromatosis, 395
Restrictive lung diseases, **675**
  ankylosing spondylitis, 469
  flow volume loops, 673
  sarcoidosis, 676
Reteplase (rPA), 437
Rete testis, 652
RET gene, 224
  carcinoma risks with, 343
  Hirschsprung disease, 384
Reticular activating system, 511
Reticular fibrous framework (spleen), 98
Reticulate bodies, 148
Reticulin, 50
Reticulocyte index (RI), **417**
Reticulocytes, 407
  in aplastic anemia, 421
  intravascular hemolysis, 421
Retina
  chronic hyperglycemia, 537
  embryologic derivation of, 613
Retinal hemorrhage
  hypertensive emergency, 300
Retinal pathology
  degeneration, 536
  detachment, **537**
  hemorrhage, 537
  retinitis, 536
  vein occlusion, **537**
  visual field defects, 542
Retinal vein occlusion, 537
Retinitis
  AIDS, 165
  cidofovir, 202
  foscarnet, 202
Retinitis pigmentosa, **538**
Retinoblastoma
  chromosome association, 64
  heterozygosity loss, 56
Retinoblastomas
  osteosarcomas, 465
Retinoids, 477
Retinopathy
  Alport syndrome, 596
  chloroquine, 200
  diabetes mellitus, 346
  hypertension, 300
  of prematurity, 210, 661
  sorbitol, 81
Retrograde amnesia, 558
Retroperitoneal fibrosis, 599
Retroperitoneal structures, **360**
Retrospective studies, 260
Retroviruses
  characteristics, 167
  genomes, 163
Rett syndrome, 61, **62**
Reverse transcriptase, 175
  Telomerase, 38
Reverse transcriptase polymerase chain reaction, **52**
Reye syndrome, **390**
Reynolds pentad, 397
Rhabdomyolysis
  daptomycin, 195
Rhabdomyomas, 316
  nomenclature for, 220
Rhabdomyosarcomas
  dactinomycin for, 439
  nomenclature for, 220
  variant, 644
Rhabdoviruses
  characteristics, 167
  negative-stranded, 168
Rhagades, 147
Rh blood classification, 405
  newborn hemolysis, 405
Rheumatic fever, **312**
  chorea with, 519
  heart murmur with, 291
  myocarditis with, 313
  Streptococcus pyogenes, 136
  streptolysin O, 133
  type II hypersensitivity, 112
Rheumatoid arthritis, 457, 466
  autoantibody, 115
  azathioprine for, 440
  carpal tunnel syndrome and, 459
  celecoxib for, 486
  etanercept for, 487
  HLA-DR4 and, 100
  immunosuppressants, 120
  infliximab/adalimumab for, 487
  leflunomide for, 486
  methotrexate for, 440
  rituximab for, 443
Rheumatoid factor, 115
Rhinitis
  phenylephrine for, 242
Rhinophyma, 477
Rhinosinusitis, **671**
Rhinovirus
  picornavirus, 167, **168**
  receptors for, 166
Rhizopus spp, 153
Ribavirin, 204
  contraindicated in pregnancy, 204
  purine synthesis, 36
Riboflavin, 67
Ribose, 79
Ribosomes, 46
Rice-water diarrhea
  organisms causing, 179
  Vibrio cholerae, 146
Richter transformation, 432
Rickets, 463
  hypophosphatemic, 591, 593
  lab values in, 464
  vitamin D and, 70
Rickettsia
  Gram stain, 125
Rickettsial diseases, **150**
Rickettsia prowazekii, 150
  transmission of, 149, 161
Rickettsia rickettsii, 150
  animal transmission, 149
  chloramphenicol, 192
Rickettsia spp
  intracellular organism, 127
  tetracyclines, 192
Rickettsia typhi, 149, 150
Riedel thyroiditis, 341
Rifabutin, 196
Rifampin, 196
  cytochrome P-450 and, 252
  Hansen disease, 141
  hepatitis, 249
  Mycobacterium leprae, 196
  Mycobacterium tuberculosis, 196
  as prophylaxis, 198
  prophylaxis with *Haemophilus influenzae* for contacts, 142
  protease inhibitors and, 203
  RNA polymerase inhibition, 42
Rifamycins, **196**
Rifaximin, 391
Rift Valley fever, 167
Right anterior cardinal vein, 281
Right bundle branch, 293
Right bundle branch block, 289
Right common cardinal vein, 281
Right coronary artery (RCA)
  coronary circulation, 283
  occlusions of, 305
Right heart failure, 309
Right horn of sinus venosus, 281
Right lower quadrant (RLQ) pain, 384
Right marginal artery, 283
Right-to-left shunts, 298
Right upper quadrant (RUQ) pain, 397
Right ventricular hypertrophy (RVH)
  high altitude, 670
  pulmonary hypertension, 679
Riluzole, 549
Ringed sideroblasts, 416
ring-enhancing lesions (MRI)
  Toxoplasma gondii, 156
Ringworm
  griseofulvin, 200
  tinea corporis, 152
Risedronate, 486
Risk assessment, 258
Risk quantification, 258
Risperidone, 573
Ristocetin, 411
risus sardonicus
  tetanospasmin, 138
Ritonavir
  HIV therapy, 203
Rituximab, 122, **443**
Rivaroxaban, 437
Rivastigmine, 240, 549
RNA
  capping, 41
  interference, 56
  processing (eukaryotes), **41**
RNA polymerases
  types and functions of, **42**
RNA viruses, **167**
  genome, **163**
Robertsonian translocation, **64**
Rocker-bottom feet, 63
"Rocket tails," 139
Rocky Mountain spotted fever, 150
  animal transmission, 149
  chloramphenicol, 192
Romaña sign, 158
Romano-Ward syndrome, 294
Romberg sign, 147, 530
Romiplostim (TPO analog), 121
Root cause analysis, 274
Rooting reflex, 510
Ropivacaine, 550
Rosacea, 477
Rose gardener's disease, 154
Rosenthal fibers, 528
Roseola
  HHV-6/HHV-7, 165
  rash, 183
Rosiglitazone, 353
Rotator cuff muscles, **446**
Rotavirus, **168**
  diarrhea, 167
Rotenone, 78
Roth spots, 311
Rotor syndrome, 393, 394
Rough endoplasmic reticulum, **46**
Rouleaux formation, 431
Round ligament of uterus, 625
Rovsing sign, 383
Rubella, **169**
  cardiac defect association, 300
  cataracts, 535
  heart murmur with, 291
  rash, 183
  TORCH infection, 182
  unvaccinated children, 186
Rubeola (measles) virus, **170**
Ruffini corpuscles, 494
Ryanodine receptor, 456
RYR1 gene, 550

S

S-100, 227
Saber shins
  congenital syphilis, 147
  syphilis, 182
Sabin poliovirus vaccine, 167
Sabouraud agar, 126
Saccular aneurysms, 516
  Ehlers-Danlos syndrome, 51
  renal cyst disorders and, 604
Saccular (development stage), 660
Sacrococcygeal teratomas, 652
Sacubitril
  mechanism and clinical use, **319**
Saddle embolus, 672
Saddle nose
  syphilis, 182
Safety culture, **273**
Salicylates
  toxicity treatment for, 248
  as weak acids, 233
Salivary gland tumors, **376**
Salivary stimulation, 240
Salmeterol, 242, 687
Salmonella, 118

(c) ketabton.com: The Digital Library

Salmonella spp
animal transmission, 149
bloody diarrhea, 179
encapsulated bacteria, 127
food poisoning, 178
intracellular organism, 127
Shigella spp vs, **144**
osteomyelitis, 180
penicillins for, 188
reactive arthritis, 469
TMP-SMX for, 194
Salmonella typhi, 144
Salpingitis
ectopic pregnancy and, 641
Sampling bias, 260
"sand"(orange) in diaper, 37
Sandflies (disease vectors), 158
Sandfly fever, 167
SA node, 292
Saponification, 209
Saprophyticus
urease-positive, 127
Saquinavir, 203
Sarcoidosis
erythema nodosum, 482
restrictive lung disease, **676**
Sarcoma botryoides, 644
Sarcomas
metastases of, 223
methotrexate for, 440
nomenclature of, 220
Sarcoplasmic reticulum, 456
Sargramostim (GM-CSF), 121
SARS (sudden acute respiratory syndrome), 167
Satiety/hunger regulation, 498
Saturday night palsy, 447
"Saw-tooth" crypt pattern, 387
Saxagliptin, 353
SBLA cancer syndrome, 224
Scabies, 161, 200
Scalded skin syndrome
Staphylococcus aureus, 135
toxic shock syndrome toxin, 133
Scales (skin), 475
basal cell carcinoma, 484
pityriasis rosea, 482
psoriasis, 477
seborrheic dermatitis, 476
Scaphoid bone, 449
Scar formation, **218**
Scarlet fever
presentation, 136
rash with, 183
Streptococcus pyogenes, 136
S cells, 371
Schatzki rings, 377
Schaumann bodies, 676
Schilling test, 420
Schistocytes, 414
in extrinsic hemolytic anemias, 423
HELLP syndrome, 643
in intravascular hemolysis, 421
in microangiopathic anemia, 423
Schistosoma haematobium
bladder cancer, 226
disease association, 161
squamous cell carcinoma of bladder, 606
Schistosoma spp, 160, 161
Schistosomiasis
portal hypertension, 389
pulmonary arterial hypertension, 679
Schizoaffective disorder, 560
Schizoid personality disorder, 565
Schizophrenia, 572
atypical antipsychotics for, 573
diagnostic criteria, 560–576
neurotransmitters for, 495
readmissions with, 272
Schizophrenia spectrum disorders, **560**
Schizophreniform disorder, 560
Schizotypal personality, 560, 565
Schüffner stippling
in blood smear, 157
Schwann cells, **494**
Guillain-Barré syndrome, 524
origin of, 490
Schwannomas, 494, 527
Sciatic nerve, 452
SCID (severe combined immunodeficiency), 37, 117
adenosine deaminase deficiency, 37
lymphopenia caused by, 424
thymic shadow in, 98
Sclerae
alkaptonuria, 84
osteogenesis imperfecta, 51
Scleritis, 466
Sclerodactyly, 473
Scleroderma, **473**
Scleroderma (diffuse)
autoantibody, 115
Sclerodermal esophageal dysmotility, 377
Sclerosing adenosis, 649
Sclerosing cholangitis, 393, 395
ulcerative colitis association, 382
Scombroid poisoning, 247
Scopolamine, 241
Scoring of USMLE Step 1 exam, 7, 8–9
Scorpion sting, 397
Scotoma, 542
Scrotal hematoma, 627
Scrotum, 627
lymphatic drainage of, 624
masses in, **652**
Scurvy
collagen synthesis and, 50
vitamin C deficiency, 69
Seafood toxins, 247
Seal-like barking cough, 170
Seborrheic dermatitis, **476**
Sebum, 477
Secobarbital, 546
Secondary adrenal insufficiency, 349
Secondary amyloidosis, 212
Secondary biliary cholangitis, 395
Secondary disease prevention, 270
Secondary glomerular disease, 594
Secondary hyperaldosteronism, 349
Secondary hyperparathyroidism, 345, 464
Secondary polycythemia, 433
Secondary spontaneous pneumothorax, 682
Second-degree AV block, 295
Second messengers
G-protein linked, 238
Second-wind phenomenon, 87
Secretin
regulatory substances, 371
secretory cell location, 373
somatostatinomas and, 351
Secretion rate calculation, 584
Secretory (exported) protein synthesis, 46
Segmented viruses, **168**
Seizures
anti-NMDA receptor encephalitis, 228
barbiturates for, 546
benzodiazepine withdrawal, 546, 570
β-blockers, 245
brain injury with recurring, **517**
bupropion, 576
clozapine use and, 573
cytomegalovirus, 182
drug reaction and, 251
with eclampsia, 643
electrolyte disturbances, 591
enflurane, 251
foscarnet, 202
during heat stroke, 517
herpesviruses and, 164, 165, 183
high fever, 165
imipenem/cilastatin, 251
isoniazid, 197
medium-chain acyl-CoA dehydrogenase deficiency, 89
meropenem, 190
nitrosourea toxicity, 441
parasite infestation and, 161
PCP, 571
phenylketonuria, 84
psychoactive drug intoxication/withdrawal, 570–571
Rett syndrome, 62
Sturge-Weber syndrome, 525
Taenia solium, 160, 161
tramadol and, 552
types of, 517
venous sinus thrombosis, 503
visceral larva migrans, 159
vitamin B_6 deficiency, 67
Zellweger syndrome, 47
Selection bias, 260
Selective estrogen receptor modulators (SERMs), 443, 462, **656**
Selective IgA deficiency, 116
Selective media, 126
Selective mutism, 557
Selectivity
β-blockers, 245
Selegiline, 548, **549**, 575
Selenium sulfide
tinea versicolor, 152
Self-fulfilling prophecies, 260
Self-image of patient, 268
Self-mutilation
Lesch-Nyhan syndrome, 37
Semimembranosus, 451, 452
Seminal vesicles, 622
Seminiferous tubules, 626, **628**, 629
Seminoma, 653
Semitendinosus, 451, 452
Sensitivity (diagnostic tests), 257
Sensorineural hearing loss, 533
Sensory cortex, 514
topographic representation, 502
Sensory innervation
derivation of, 620
lower extremity, 452, 453
tongue, 493
upper extremity nerve injury, 447
Sensory loss
conversion disorder and, 566
stroke effects, 514
Sensory modalities/pathways
receptors for, **494**
spinal tracts in, 509
thalamus in, 498
Separation anxiety disorder, 557
Separation anxiety (infants), 635
Sepsis
ARDS, 678
immunodeficient patients, 118
lymphopenia with, 424
neutropenia with, 424
shock with, 310
Streptococcus agalactiae, 137
Septate uterus, 623
Septation of heart chambers, 280
Septic arthritis, **468**
gonococci, 142
Staphylococcus aureus, 135
Septicemia
Listeria monocytogenes, 139
readmissions with, 272
Waterhouse-Friderichsen syndrome, 349
Septic shock
diffuse cortical necrosis (renal), 602
macrophages and, 407
norepinephrine for, 242
Septum primum, 280
Septum secundum, 280
Sequence (morphogenesis error), 613
Serine, 224
Serologic markers
hepatitis, 174
Seronegative spondyloarthritis, **469**
Serosa, 362
Serotonergic drugs, 574
Serotonin
changes with disease, 495
vitamin B_6 and, 67
Serotonin syndrome, 400, 547, 552, 569
atypical antidepressants, 576
dextromethorphan, 686
MAOIs, 575
MDMA, 571
oxazolidinones, 193
Serous cystadenocarcinoma, 227, 646
Serous cystadenoma, 646
Serpentine cord, 140
Serrated colon polyps, 387
Serratia marcescens, 128
treatment of, 189
in immunodeficiency, 128
UTIs, 181
Serratia spp
immunodeficient patients, 118
Serratus anterior muscle, 448
Sertoli cells
secretions of, 622, 628
sexual determination, 622
tumors of, 653
Sertoli-Leydig cell tumor, 647
Sertraline, 575
Serum amyloid A, 213
Serum markers (liver pathology), **390**
Serum osmolarity
antidiuretic hormone regulation of, 329
Serum protein electrophoresis (SPEP)
in plasma cell dyscrasias, 431
Serum tumor markers, **226**
Sevelamer, **355**
17α-hydroxylase, 335
17-hydroxyprogesterone, 335
Sevoflurane, 550
Sex chromosome disorders, **638**
Sex cord stromal tumors, 647
Sex development disorders
phenotypic and gonadal disagreement in, 639
physical characteristics, 639

FAS1_2019_21_Index_749-806.indd 794 11/21/19 12:27 PM

(c) ketabton.com: The Digital Library

Sex hormone–binding globulin (SHBG), 337
Sex hormone disorder diagnosis, 639
Sex pilus (bacterial genetics), 130
Sex steroid replacement, 339
Sexual abuse, 556, 558
Sexual behavior
  hypothalamus regulation of, 498
Sexual differentiation, **622**, 636
Sexual dysfunction
  β-blockers and, 245, 323
  cimetidine, 399
  differential diagnosis of, **567**
  Lambert-Eaton myasthenic syndrome, 472
  Peyronie disease and, 651
  tuberoinfundibular pathway, 499
Sexually transmitted infections (STIs)
  clinical features, **184**
  parental consent with, 265
Sézary syndrome, 430
Shawl and face rash, 471
Sheehan syndrome, 339
Sheep (disease vectors), 160
Shiga-like toxin (SLT), 145
  cytokine release and, 132
Shiga toxin, 130, 132, 144
Shigella boydii, 144
Shigella dysenteriae, 144
Shigella flexneri, 144
Shigella sonnei, 144
Shigella spp
  bloody diarrhea, 179
  penicillinase-sensitive penicillins for, 188
  reactive arthritis, 469
  TMP-SMX, 194
  vs *Salmonella* spp, 144
Shingles, 165
Shin splints, 461
Shock, **310**
  dopamine for, 242
  Ebola, 171
  endotoxins, 131
  in pulmonary anthrax, 137
  norepinephrine for, 242
  Waterhouse-Friderichsen syndrome and, 349
short-chain fatty acids
  in anaerobic organisms, 127
SIADH, 338
  drug raction and, 249
  markers in, 591
  paraneoplastic syndrome, 228
Sialadenitis, 376
Sialolithiasis, **376**
Sialyl Lewis^x, 215
Sibling studies, 256
Sickle cell disease, **422**, 651
  ESR in, 214
  osteonecrosis and, 463
  postsplenectomy state in, 98
  Streptococcus pneumoniae, 136
Sickle cells, 415
Sideroblastic anemia
  causes and treatment, 419
  lead poisoning, 419
  vitamin B_6 deficiency, 67
Sigmoid colon, 383
Sigmoid sinus, 503
Sigmoid volvulus, 386
Signaling pathways
  endocrine hormones, **337**
  steroid hormones, **337**
Signal recognition particle (SRP), 47
Signet ring cells, 379
Sign of Leser-Trélat, 228
Sildenafil, 651
Silencer (gene expression), 41
Silicosis, 677
Silver stain, 125, 143
Simeprevir, 204
Simple partial seizures, 517
Simple pneumothorax, 680
Simple renal cysts, 604
Single nucleotide (point) mutation, 39
Single nucleotide polymorphisms (SNPs), 54
Single nucleotide substitutions, 39
Single-stranded binding proteins, 38
Sinusitis
  brain abscesses, 180
  C3 deficiency and, 107
  Kartagener syndrome, 49
  Streptococcus pneumoniae, 136
  Wegener granulomatosis, 315
Sinus venosus, 281
Sirolimus
  immunosuppressant, 120
Sister Mary Joseph nodules, 379
Sitagliptin, 353
Situs inversus, 49
6-mercaptopurine, 440
  azathioprine, 120
  for ulcerative colitis, 382
  purine synthesis, 36
  targets of, 438
Sjögren syndrome, **468**
  autoantibody, 115
  pilocarpine for, 240
  rheumatoid arthritis, 466
Skeletal muscles
  ACh receptors in, 236
  blood flow regulation in, 297
  glycogen metabolism in, 86
  somatic nerve, supply of, 236
Skewed distributions, 262
Skin
  blood flow regulation in, 297
  collagen in, 50
  normal flora of, 135
  pigmentation, 56
  wrinkles of aging, 52
Skin anatomy, 473–483
  layers of, **473**
  microscopic terms, 475
  morphology, 475
Skin cancer, **484**
  albinism and, 476
  Lynch syndrome and, 388
Skin drug reactions, 250
Skin flora, 178
Skin lesions
  autoimmune disorders, 480
  bulla, 475
  burns, 483
  café-au-lait spots, 57
  cancer, 222
  common disorders, **477**
  crust, 475
  dermatitis herpetiformis, 381
  erythema multiforme, 151
  Gottron papules, 228
  hyperlipidemia signs, 300, **301**
  hyperpigmentation, 395
  inflammatory bowel disease, 382
  Kaposi sarcoma, 165
  kwashiorkor, 71
  macule, 475
  papule, 475
  patch, 475
  petechiae, 407
  pigmentation disorders, 475, **476**
  plaque, 475
  pustule, 475
  scale, 475
  scaling, 152
  scaly, 66
  seborrheic keratoses, 228
  splinter hemorrhages, 311
  striae, 348
  T-cell lymphoma, 430
  telangiectasia, 316, 473
  ulcers, 158
  vascular tumors, 478
  vasculitides, 314
  verrucous, 151
  vesicle, 475
  wheal, 475
Skip lesions, 382
Skull thickening, 463
Slapped cheek rash, 183
Sleep
  ghrelin/leptin production, 336
Sleep disturbance
  apnea, **679**
  benzodiazepines and, 570
  β-blockers, 245
  delirium and, 558
  generalized anxiety disorder, 563
  in geriatric patients, 270
  hypnagogic hallucinations, 559
  hypnopompic, 559
  paralysis, 568
  paroxysmal nocturnal dyspnea, 309
  pulsus paradoxus in, 310
  sleep terror disorder, 567, **568**
  varenicline, 576
Sleep physiology, **497**
  stages in, 497
Sleepwalking
  sleep stages and, 497
SLE (systemic lupus erythematosus), **470**
  antiphospholipid syndrome and, 470
  autoantibodies, 115
  DPGN, 596
  HLA subtypes, 100
  kidney disease with, 596
  Raynaud phenomenon, 472
Sliding hiatal hernia, 370
Slime (S) layer, 124
Slipped capital femoral epiphysis, 461
  osteonecrosis, 463
Slow acetylators, 232
Small bowel disease, 374
Small cell carcinoma of lung, **684**
  carcinogens for, 225
  immunohistochemical stains for, 227
  oat cell carcinoma, 684
  paraneoplastic syndromes, 228
  Lambert-Eaton myasthenic syndrome, 472
  topotecan for, 442
Small interfering RNA (siRNA) **56**
Small intestine, 371
Small lymphocytic lymphoma (SLL), 432
Small molecule inhibitors
  naming conventions for, 254
Smallpox, 164
Small-vessel vasculitis
  presentation and pathology, 314
Smoking
  abdominal aortic aneurysms and, 302
  atherosclerosis and, 302
  Buerger disease and, 314
  bupropion for cessation, 576
  carcinogenicity of, 225
  cataracts, 535
  colorectal cancer and, 388
  emphysema, 674
  esophageal cancer and, 378
  head and neck cancer, 671
  hormonal contraception, 657
  Legionnaires' disease, 143
  lung cancer, 684
  pancreatic cancer and, 398
  renal cell carcinoma, 605
  saccular aneurysms, 516
  squamous cell carcinoma of bladder, 606
  stomach cancer and, 379
  teratogenic effects, 614
  transitional cell carcinoma, 606
  varenicline for cessation, 576
Smooth brain, 491
Smooth endoplasmic reticulum, **46**
Smooth muscle
  BMPR2 gene, 679
  contraction of, 457
  glomus tumors, 478
  respiratory tree, 662
  tumor nomenclature in, 220
Smooth muscle (vascular)
  in arteriolosclerosis, 301
  atherosclerosis and, 302
  calcium channel blocker action, 318
Smudge cells, 432
SNARE proteins
  in neurotransmission, 138
SNc (substantia nigra pars compacta), 495
SNRIs (serotonin-norepinephrine reuptake inhibitors)
  clinical use, 572
  major depressive disorder, 561
  mechanism and clinical use, **575**
Snuffles, 147
"Soap bubble" appearance/lesions
  giant cell tumor, 464
  Cryptococcus neoformans, 153
Social anxiety disorder, 563
  drug therapy for, 572
  SSRIs for, 575
  venlafaxine for, 575
Social engagement
  infant deprivation effects, 556
Sodium channel blockers, **322**
Sodium channels
  cystic fibrosis, 60
  epilepsy drug effects, 544
  local anesthetic effects, 550
  pacemaker action potential and, 292
  permethrin, 200
Sodium-glucose co-transporters (SGLT), 334, 353, 373, 584
Sodium oxybate (GHB)
  narcolepsy treatment, 568
Sodium-potassium channels, 236
Sodium-potassium pump, **49**
Sodium stibogluconate, 158, **200**
Sofosbuvir, 204
Solifenacin, 241
Solitary functioning kidney, 579
Solitary nucleus of medulla, 296

(c) ketabton.com: The Digital Library

Somatic hypermutation, 101
Somatic mosaicism
  Sturge-Weber syndrome, 525
Somatic mosaicism, 57
Somatic nerves
  male sexual response, 627
Somatic symptom disorder, **566**
Somatic symptoms
  characteristics of, 566
  conversion disorder, 566
  illness anxiety disorder, 566
  somatic symptom disorder, 566
Somatomedin, 329
Somatosensory cortex (primary), 501
  thalamic relays to, 498
Somatostatin
  function of, 328
  glucagon and, 333
  hypothalamic/pituitary drugs, 354
  production of, 329
  regulatory substances, 371
  secretory cell locations, 373
Somatostatinoma, 351
  pancreatic cell tumor, 351
Somatotropin, 329
Sorbitol metabolism, **81**
Sotalol, 323
Southern blot, 53
Southwestern blot, 53
Space of Disse, 367
Spaghetti and meatballs appearance, 152
Spasmolytics, **551**
Spastic paralysis
  tetanospasmin, 138
Spastic paresis, 529
Special senses
  ophthalmology, 534–543
  otology, 533–534
Specificity equation, 257
Specific learning disorder, 557
Spermatic cord, 369
Spermatocele, 652
Spermatocytes, 628
Spermatogenesis, 628
  cryptorchidism and, 651
  process of, **629**
  prolactin effect on, 330
Spermatogonia, 628
Spermiogenesis, 629
Sphenomandibular ligament, 620
Sphenoparietal sinus, 503
Spherocytosis, 415
  autoimmune hemolytic anemia, 415
  extrinsic hemolytic anemia, 423
  hereditary, 421, 422
Sphincter of Oddi, **368**, 371
Sphingolipidoses, 88
Sphingomyelin, 88
Sphingomyelinase, 88
Spina bifida
  neural tube defect, 491
Spinal cord
  embryologic derivation, 613
  lesions of, **530**
  lower extent of, **507**
  nerve nuclei of, 505
  nerves of, **507**
  tracts of, **508**, 509
Spinal cord syndromes
  multiple sclerosis, 523
Spinal muscular atrophy, 530
  splicing of pre-mRNA in, 42
Spinothalamic tract, 509
  thalamic relay for, 498
Spirochetes, **146**
Spironolactone, **609**, 658
  for heart failure, 309
Splay (glucose clearance), 584
Spleen
  bacterial clearance by, 127
  blood supply and innervation of, 364
  embryology, 360
  in gastrointestinal anatomy, 361
  ischemia susceptibility, 210
  structure and function, **98**
  thrombocytes in, 407
Splenectomy, 422
  peripheral blood smear after, 423
Splenic artery, 364
Splenic flexure
  blood supply to, 363
Splenomegaly
  anemia, 157
  cirrhosis, 389
  hairy cell leukemia, 432
  hereditary spherocytosis, 422
  malaria, 157
  myelofibrosis, 433
  rheumatoid arthritis, 466
  visceral leishmaniasis, 158
Splenorenal ligament, 361
Splicing of pre-mRNA, **42**
  alternative splicing, 43
Splinter hemorrhages, 311
Splitting, 555
Splitting of heart sounds, 288, 289
Spondyloarthritis (seronegative), 469
Spongiosis, 475
Spontaneous abortion
  antiphospholipid syndrome, 470
  Listeria monocytogenes, 139
  syphilis, 182
  vitamin A excess, 614
Spontaneous bacterial peritonitis, 389, **390**
Spontaneous pneumothorax, 682
Sporadic porphyria cutanea tarda, 173
Spore-forming bacteria, **129**, 138
Spores, 124
Sporothrix schenckii, **154**
Sporotrichosis, 154
Sprain (ankle), 455
Sprue
  fat-soluble vitamin deficiencies and, 65
  vitamin B_{12} deficiency, 69
"Spur cells," 414
Sputum
  currant jelly, 145
  Klebsiella spp, 186
  Streptococcus pneumoniae, 136
  rusty, 136
Squalene epoxidase, 199
Squamous cell carcinoma, 484
  anus and cervix, 177
  bladder, 160, **606**
  carcinogens in, 225
  cervix, 645
  esophagus, 378
  head and neck, 671
  hypercalcemia and, 228
  lungs, 684
  pectinate line and, 366
  penis, 651
  of skin, 482
Squamous epithelium, 662
SRY gene, 622
SSRIs (selective serotonin reuptake inhibitors), **575**
  anxiety disorders, 562
  atypical depression, 561
  clinical use, 572
  major depressive disorder, 561
  mechanism and clinical use, 575
  obsessive-compulsive disorder, 563
  panic disorder, 563
  phobias, 563
  postpartum depression, 562
  SIADH caused by, 249
Stable angina, 304
Stable (quiescent) cells, 46
Stab wounds and winged scapula, 448
Staghorn calculi, 598
Stains (bacterial), **125**
Standard deviation
  dispersion, 262
  variability, 259
Standard error of the mean, 262
Stapedial artery, 619
Stapedius muscle, 620
Stapes (ossicles), 533, 620
Staphylococcal scalded skin syndrome, 479
Staphylococcal toxic shock syndrome (TSS), 135
Staphylococcus aureus, **135**
  bacterial endocarditis, 311
  β-hemolytic nature of, 135
  brain abscesses, 180
  cephalosporins, 189
  cystic fibrosis, **60**, 179
  dapsone, 195
  exotoxin production, 133
  food poisoning, 178
  immunocompromised patients, 179
  influenza, 169
  IV drug use, 179
  nosocomial infection, 179, **185**
  osteomyelitis and, 180
  penicillins for, 188
  pigment production, 128
  pneumonia, 683
  postviral infection, 179
  prophylaxis for, 198
  septic arthritis, 468
  skin infections, 479
Staphylococcus epidermidis, **135**
  gram-positive testing, 134
  in vivo biofilm production, 128
  normal flora, 178
  nosocomial infection, 185
  osteomyelitis, 180
  urease-positive, 127
  vancomycin for, 190
Staphylococcus gallolyticus, 137
Staphylococcus pyogenes
  skin infections, 479
Staphylococcus saprophyticus, **136**
  gram-positive testing, 134
  urinary tract infections, 600
  UTIs, 181
Staphylococcus spp
  antibiotic tests for, 134
  facultative anaerobic metabolism, 127
Starling curves, **285**, 297
Starling forces, 297
"Starry sky" appearance of B cells, 430
Start and stop codons, **44**
Startle myoclonus, 521
Starvation phases, 91
Statins
  for acute coronary syndromes, 307
  hepatitis, 249
  myopathy, 250
Statistical distribution, **262**
Statistical hypotheses, **262**
  confidence interval, 263
  correct result, 263
  incorrect results, 263
  test for, 264
Status epilepticus, 517
  treatment, 544
Stavudine, 203
Steady state, 231
Steatohepatitis, 389
Steatorrhea
  chronic pancreatitis, 397
  cystic fibrosis, 60
  malabsorption syndromes and, 381
  octreotide effect, 400
Steatosis (hepatic), 390, **391**
Steeple sign (x-ray), 170
Stellate cells, 367
Stellate ganglion, 685
Stem cells
  in aplastic anemia, 421
  bone marrow, 108
  CD34 protein, 110
  myelodysplastic syndromes and, 431, 432
Steppage gait, 453
Sterilization/disinfection methods, 204
Steroid hormone signaling pathways, 337
Steroids
  acute pancreatitis, 397
  adrenal insufficiency, 349
  berylliosis, 677
  CRH levels in, 328
  multiple sclerosis, 523
  synthesis of, 46, 72
  TBG and, 331
Stevens-Johnson syndrome, 194, **481**, 544
  atypical variant of, 150
  drug reaction and, 250
  sulfa drug allergies, 252
Stimulants
  for ADHD, 557
  intoxication and withdrawal, 570
  laxative, 401
St. John's wort, 252
St. Louis encephalitis, 167
Stomach
  basal electric rhythm, 362
  blood supply to, 364
  cholecystokinin effect on, 371
  in gastrointestinal anatomy, 361
  histology of, 362
  regulatory substances, 371
  sclerosis of, 473
  secretin effect on, 371
Stone bone, 463
Straight sinus, 503
Stranger anxiety (infants), 635
Strategies
  clinical vignette, 23
  test-taking, 22–23
Strawberry cervix, **181**, 184
Strawberry hemangiomas, 478
Strawberry tongue, 136
  Kawasaki, 314
  scarlet fever, 136
Streak gonads, 622

(c) ketabton.com: The Digital Library

Streptococcus agalactiae (group B strep), **137**
β-hemolytic nature of, 135
encapsulated bacteria, 127
gram-positive testing, 134
in neonates, 182
prophylaxis for, 198
Streptococcus bovis, **137**
Streptococcus mutans
biofilm production, 128
normal flora, 178
Streptococcus pneumoniae, **136**
chloramphenicol, 192
cystic fibrosis, 179
encapsulated bacteria, 127
gram-positive testing, 134
IgA protease and, 129
influenza, 169
IV drug use and, 179
meningitis, 179, 180
penicillin G/V for, 187
pneumonia, 179, **683**
postviral infection, 179
transformation in, 130
α-hemolysis, 135
Streptococcus pyogenes (group A strep), 130, 133, **136**
β-hemolysis, 135
lab testing, 134
in renal disease, 596
M protein and, 129
rash, 183
treatment of, 187, 192
Streptococcus pyogenes toxic shock–like syndrome
skin infection with, 135
Streptococcus sanguinis, 128
Streptococcus spp
antibiotic tests for, 134
facultative anaerobic metabolism, 127
septic arthritis, 468
Streptogramins, 198
Streptokinase, 437
Streptolysin O, 133
Streptomycin, 191, **197**
Stress incontinence, 599
Stress-related disorders, 564
Striated muscle, 220
Striatum, **500**, 514
String sign (x-ray), 382
Stroke, 512
ADP receptor inhibitors for, 437
atrial fibrillation and, 295
central post-stroke pain syndrome, 515
direct factor Xa inhibitors for, 437
eclampsia, 643
effects of, 514–515
homocystinuria, 84
hypertension, 300
hypertensive emergency and, 300
sickle cell anemia, 422
syphilis, 147
thrombolytics for, 437
Stroke volume, 284
equation for, 285
Strongyloides spp, 158, 159
Structural quality measurement, 273
Struvite stones
with *Proteus* spp, 127
ST segment, 293
ST-segment elevation MI (STEMI)
diagnosis of, 304
Studies
error types, 256
Studying for USMLE Step 1 exam
timeline for, 16–19
Sturge-Weber syndrome, 525
Stylohyoid ligament, 620
Stylohyoid muscle, 620
Styloid process, 620
Stylopharyngeus, 620
Subacute combined degeneration (SCD), 69, 530
Subacute endocarditis
enterococci, 137
Staphylococcus gallolyticus, 137
Subacute granulomatous thyroiditis, 341
Subarachnoid hemorrhage, **513**, 518
aneurysms, 516
nimodipine for, 318
Subarachnoid space, 507
Subclavian arteries, 619
Subcutaneous fat
erythema nodosum in, 482
skin layers, 473
Subcutis, 473
Subdural hematomas, 513
Subendocardium, 210
Sublimation, 555
Sublingual gland
stones in, 376
Submandibular gland
stones in, 376
Submucosa, 362
Submucosal polyps, 387
Subscapularis muscle, 446
Substance abuse
adult T-cell lymphoma and, 430
Candida albicans, 153
delirium with, 558
dissociative identity disorder and, 558
parental consent, 265
torsades de pointes in, 294
tricuspid valve endocarditis and, 311
Substance P, 551
Substance P antagonist, 401
Substance use disorder, **568**
addiction, stages of change in overcoming, 568
Substantia nigra pars compacta (SNc), 500
Subthalamic nucleus, 500
lesions in, 511
Subunit vaccines, 111
Succimer
heavy metal toxicity, 248
lead poisoning, 419
Succinate dehydrogenase, 67, **78**
Succinylcholine, **551**, 590
Succinyl-CoA
gluconeogenesis, 78
TCA cycle, 77
Sucking reflex, 510
Sucralfate, 399
Sudden death
cardiac death, 304, 313
cocaine use, 571
cor pulmonale, 679
with myocarditis, 313
sleep apnea, 679
Sudden infant death syndrome (SIDS), 635
Suicidal patients, 268
confidentiality exceptions and, 267
elderly, 270
Suicide
deaths from, 272
physician-assisted, 268
risk factors for, 562
Sulbactam, 189
Sulfadiazine, 194
Toxoplasma gondii, 156
Sulfa drugs, **252**
megaloblastic, 250
rash, 250
Sulfamethoxazole (SMX), 194
Sulfapyridine, 400
Sulfasalazine, 252, **400**, 466
Sulfatides, 140
Sulfisoxazole, 194
Sulfonamides
cytochrome P-450 and, 252
hemolysis in G6PD deficiency, 250
hypothyroidism, 249
mechanism and use, **194**
Nocardia spp, 139
photosensitivity, 250
pregnancy contraindication, 204
trimethoprim, 194
vitamin B_9 deficiency, 68
Sulfonylureas, 353
disulfiram-like reaction, 251
insulin and, 334
Sulfur granules, 128, **139**
Sumatriptan, 547
cluster headaches, 518
coronary vasospasm with, 248
Sunburn, 482
Sunburst pattern (X-ray), 465
Superficial inguinal nodes, 624
Superior colliculi, 504
Superior gluteal nerve, 453
Superior mesenteric artery (SMA), 364
syndrome, 363
Superior oblique muscle, 540
Superior olive (nucleus), 498
Superior ophthalmic vein, 503
Superior rectal vein, 365
Superior rectus muscle, 540
Superior sagittal sinus, 503
Superior sulcus tumor, 685
Superior vena cava
embryologic development of, 281
in fetal circulation, 282
Superior vena cava syndrome, 98, **685**
lung cancer, 684
Pancoast tumor, 685
Superoxide dismutase, 109
free radical elimination by, 210
Supination
Erb palsy, 448
forearm, 447
Supportive therapy, 572
Suppression, 555
Suprachiasmatic nucleus (SCN), 497–498
sleep physiology and, 497
Supracondylar fracture, 447
Supraoptic nucleus, 498
Suprascapular nerve, 446
Supraspinatus muscle, **446**, 448
Supraventricular tachycardia
adenosine for diagnosing, 324
β-blockers for, 245, 323
calcium channel blockers for, 324
Suramin, 200
Surface F protein, 169
Surfactant (pulmonary), 661
Surgical neck of humerus, 455
Surgical procedures
readmissions with, 272
Surrogate decision-maker, **266**
Suvorexant, **547**
Swallowing
tongue movement in, 493
Swan-Ganz catheter, 297
Swarming, 181
Sweat glands, 236
embryologic derivation, 613
pilocarpine effects, 240
Swiss cheese model, **273**
Sydenham chorea, 312, **519**
Sylvian fissure, 501
Sympathetic nervous system
denervation of face, 540
male sexual response, 627
receptor targets, 236
venous return and, 286
Sympatholytic drugs, **243**
Sympathomimetics
direct, **242**
indirect, **242**
Syncope
during exercise, 308
pulsus parvus et tardus, 291
Syncytiotrophoblasts, 617, 633
β-hCG and, 226
hCG secretion by, 633
Syndrome of apparent mineralocorticoid excess, 586
markers in, 591
Syndrome of inappropriate antidiuretic hormone secretion (SIADH), **338**
Synergism
Aspirin and, 235
of drugs, 235
Syphilis, **147**
fluorescent antibody stain for, 125
STI, 184
syphilitic heart disease, **312**
tabes dorsalis, 530
testing for, 148
thoracic aortic aneurysms and, 302
TORCH infection, 182
Syringomyelia, **492**
spinal cord lesions, 530
Syrinx, 492
Systemic amyloidosis, 212
Systemic juvenile idiopathic arthritis, **468**
Systemic lupus erythematosus (SLE), **470**
Systemic mycoses, **151**
Systemic primary carnitine deficiency, 89
Systemic sclerosis, 473
Systemic senile amyloidosis, 212
Systolic ejection, 287
Systolic murmur, 308

T

T3 (liothyronine), 354
T4 (levothyroxine), 354
Tabes dorsalis, **147**, 184
spinal cord lesions, 530
Tachyarrhythmia
isoproterenol for evaluating, 242
thyroid storm, 342
Tachycardia
β-blockers, 245
drug-induced, 318
MDMA, 571

(c) ketabton.com: The Digital Library

Tachycardia (*continued*)
metronidazole, 195
with myocarditis, 313
PCP, 571
phenoxybenzamine, 244
reflex, 244
stimulants and, 570
supraventricular, 245
thyroid hormones, 354
Wolff-Parkinson-White syndrome, 294
Tachyphylaxis, 235
Tacrolimus
hyperglycemia, 249
immunosuppression, 120
Tactile hallucinations, 559
cocaine, 571
Taenia solium, 160, 161
Takayasu arteritis, 314
Tamm–Horsfall mucoprotein, 594
Tamoxifen, **443**, 656
hot flashes with, 249
Tamsulosin, 244, **658**
Tanner stages (sexual development), **637**
Tardive dyskinesia
antipsychotic drugs and, 573
metoclopramide adverse effect, 400
nigrostriatal pathway, 499
Target cells, 415
postsplenectomy, 98
Tarsal tunnel syndrome, 453
Taste
cranial nerve lesions and, 532
loss with stroke, 514
thalamic relay for, 498
TATA box, 41
Taxanes, 441
Tay-Sachs disease
lysosomal storage disease, 88
Tazobactam, 189
TCA cycle, **77**
hyperammonemia, 82
metabolic site, 72
pyruvate metabolism, 77
rate-determining enzyme for, 73
TCA toxicity
treatment of, 233
T cells, 127, **409**
activation, **103**
adaptive immunity, 99
anergy, 110
cell surface proteins, 110
corticosteroid effects, 120
cytokine production, 101, **108**
cytotoxic, 102
delayed (type IV) hypersensitivity, 101
differentiation and maturation, 98, **102**
disorders of, 116, 117
functions, 101
leflunomide effects, 486
lymph nodes, 96
macrophage interaction, 102
major functions of, 101
neoplasms, 430
polysaccharide antigens and, 127
regulatory, 102
sirolimus effect, 120
spleen, 98
thymus, 98
untreated HIV, 176
Tea-colored urine, 425
"Teardrop" RBCs, **414**, 433
Tearing stimulation, 240
Teenagers
common causes of death, 272
Teeth
congenital syphilis, 147
dentinogenesis imperfecta, 51
discoloration, 192, 204, 250, 614
Gardner syndrome, 387
osteogenesis imperfecta, 51
Sjögren syndrome and, 468
Telangiectasias
basal cell carcinomas, 484
hereditary hemorrhagic, 316
Telencephalon, 490
Tellurite agar, 126
Telomerase
action of, 38
Telophase, 46
Temazepam, 546
Temperature receptors, 494
Temperature regulation, 498
Temperature sensation
cape-like distribution loss, **492**, 530
loss with strokes, 514
Temporal arteritis, 314
Temporalis muscle, 507
Temporal lobe, **501**, 514
Temporal lobe encephalitis, 164
Tendinopathy (rotator cuff), 446
Tendinous xanthomas, 301
familial hypercholesterolemia, 94
Tendonitis
drug reaction and, 250
fluoroquinolones, 195
Tendons
collagen in, 50
Tenecteplase (TNK-tPA), 437
Teniposide, 442
Tennis elbow, 459
"Tennis rackets" (Birbeck) granules, 434
Tenofovir, 203
Tenosynovitis, 468
Tension headaches, 518
Tension pneumothorax, 680, **682**
Tensor fascia latae muscle, 453
Tensor tympani muscle, 620
Tensor veli palatini muscle, 620
Teratogens
ACE inhibitors, 610
aminoglycosides, 191
angiotensin II receptor blockers, 610
in fetal development, 612
fetal effects of, **614**
griseofulvin, **200**, 204
leflunomide, 486
lithium as, 574
methimazole as, 354
propylthiouracil in pregnancy, 354
ribavirin, 204
vitamin A, 66
warfarin as, 436
Teratoma
immature, 647
mature cystic, 647
testicular, 653
Terazosin, 244
Terbinafine, **199**
Terbutaline, 242
Teres minor, 446
Teriparatide, 462, **487**
Terminal bronchioles, 660
Terminal complement deficiencies (C5–C9), 107
Terminal deoxynucleotidyl transferase (TdT), 104
Tertiary adrenal insufficiency, 349
Tertiary disease prevention, 270
Tertiary hyperparathyroidism, 345
Tesamorelin, 328
Testes
descent of, 624
lymphatic drainage of, 624
progesterone production, 630
Testicular atrophy
alcoholism, 571
muscular dystrophy, 61
Testicular cancer, 439, 442, **653**
Testicular torsion, **651**
Testicular tumors
germ cell, **652–653**
non-germ cell tumors, 653
Testing agencies, 24
Testis-determining factor, 622
Testosterone, 636, **658**
androgen insensitivity syndrome, 639
Leydig cell secretion, 628
Sertoli cells, 628
SHBG effect on, 337
signaling pathways for, 337
spermatogenesis, 628
Testosterone-secreting tumors, 639
Testosterone synthesis, 199
Test-taking strategy, 22–23
Tetanospasmin, 132
blocks release of GABA, 138
Tetanus
exotoxins, 131
Tetanus toxin, 110
Tetany
hypocalcemia, 591
hypoparathyroidism, 344
Tetrabenazine
Huntington disease, 549
Tourette syndrome, 572
Tetracaine, 550
Tetracyclines, **192**
esophagitis, 249
Fanconi syndrome, 251
photosensitivity, 250
protein synthesis inhibition, 191
pseudotumor cerebri and, 521
teratogenicity, 204, **614**
tooth discoloration, 250
Tetrahydrobiopterin (bh4)
in phenylketonuria, 84
Tetrahydrofolic acid (THF), **68**, 194
Tetralogy of Fallot, 298
22q11 syndromes, 300
fetal alcohol syndrome, 300
outflow tract formation, 281
Tetrodotoxin, 247
TGF-β
in wound healing, 216
neural development, 490
regulatory T cells, 102
Thalamus
development of, 490
limbic system and, **498**
neuropathic pain, 515
Thalassemia, 418
target cells in, 415
Thalidomide
teratogenicity, 614
Thayer-Martin agar, 126
Theca-lutein cysts, 642, **646**
Thecoma, 647
Thenar muscles, 447, 448, 450
Theophylline, 687
therapeutic index of, 234
Therapeutic antibodies, 121, **122**
Therapeutic index (TI), **234**
Therapeutic window, 234
Thermogenin, 78
Theta rhythm (EEG), 497
Thiamine, 66
Thiazide diuretics, **609**
in gout, 250
in heart failure, 309
in hypertension, 316
Thiazides, 609
Thionamides, **354**
Thiopental, 546, 550
Thioridazine, 573
Third-degree (complete) AV block, 295
3rd pharyngeal pouch, 621
3rd pharyngeal arch, 620
Thirst
hypothalamus and, 498
30S inhibitors, 191
Thoracic aortic aneurysm, 300, **302**
Thoracic outlet syndrome, 448, 684
Threadworms, 159
Threonine, 81
Threonine kinase, 224
Thrombi
atherosclerosis, 302
mural, 305, **307**
post-MI, 305
Thrombin, 436
Thromboangiitis obliterans, 314
Thrombocytes (platelets), **407**
disorders, 426–427
function tests of, 426
heparin adverse effects, 436
leukemias, 432
liver markers, 390
mixed coagulation disorders, 428
thrombolytics and, 437
transfusion of, 421, **429**
in wound healing, 216
Thrombocythemia (essential), 433
Thrombocytopenia, 407
Class IA antiarrhythmics, 322
cytarabine, 440
drug reaction and, 250
Escherichia coli, 145
ganciclovir, 202
glycoprotein IIb/IIIa inhibitors, 438
heparin adverse effects, 436
oxazolidinones, 193
protease inhibitors, 203
recombinant cytokines, 121
sulfa drug allergies, 252
TORCH infections, 182
transfusion for, 429
Wiskott-Aldrich syndrome, 117
Thrombocytosis
postsplenectomy, 98
Thromboembolic event
atrial fibrillation, 295
Thrombogenesis, **411**
Thrombolytic drugs, 413, **437**
Thrombophlebitis
pancreatic cancer, 398
Thrombopoietin, 121
signaling pathways, 337
Thrombosis
celecoxib, 486
contraceptive and hormone replacement, 250
essential thrombocythemia, 433
homocystinuria, 84

(c) ketabton.com: The Digital Library

Thrombotic stroke, 512
Thrush, 117
  Candida albicans, 153
  hairy leukoplakia vs, 479
  HIV-positive adults, 177
  nystatin, 199
"Thumbprint" sign (imaging) colonic ischemia, 386
"Thumb sign" (X-ray) flu, 142
Thymic aplasia, 116
  chromosome association, 64
  lymphopenia with, 424
Thymic cortex
  T cell selection in, 102
Thymic hyperplasia
  myasthenia gravis association, 472
Thymic shadow, 117
Thymidine, 194
Thymidine kinase, 201
Thymidylate, 36
Thymoma
  disease associations with, 98
  myasthenia gravis and, 228, 472
  paraneoplastic syndromes, 228
Thymus
  benign neoplasm, **98**
  fetal development, 326
  pharyngeal pouch derivation, 621
  T cell differentiation, 102
  T cell origination in, 409
Thymus-dependent antigens, 105
Thymus-independent antigens, 105
Thyroglossal duct cyst, 326
Thyroid adenomas, 342
Thyroid cancer, 343
  amyloidosis in, 212
  carcinogens in, 225
  metastases to, 223
  Psammoma bodies in, 227
Thyroid cartilage, 620
Thyroid development, 326
  pharyngeal pouch derivation, 621
Thyroidectomy, 343
Thyroid hormones, **331**
  signaling pathways for, 337
  in toxic multinodular goiter, 342
Thyroiditis, 341
Thyroidization of kidney, 600
Thyroid peroxidase, 331
Thyroid-regulating hormone (TRH)
  signaling pathways for, 337
Thyroid-stimulating hormone (TSH)
  Graves disease and, 342
  secretion of, 327
  signaling pathways of, 337
Thyroid-stimulating immunoglobulin (TSI), 331
Thyroid storm, 342
Thyrotoxicosis, 331
Thyrotropin-releasing hormone (TRH), **328**, 330
Thyroxine, 339
Thyroxine-binding globulin (TBG), 331
TIBC (total iron-binding capacity)
  anemia of chronic disease, 421
  lab values in anemia, 419
  microcytic anemia, 418
Tibialis anterior, 453
Tibial nerve, 452–453
  neurovascular pairing, 455
Ticagrelor, 437
Ticarcillin
  characteristics of, 188
  Pseudomonas aeruginosa, 143
Ticks (disease vectors), 149–150
Ticlopidine, 411, **437**
Tics (Tourette syndrome), 557
Tidal volume (TV), 664
Tigecycline, **192**
Tight junctions, **474**, 496
Timolol, 245, 323, **552**
Tinea, **152**, 200
Tinea capitis, 152
Tinea corporis, 152
Tinea cruris, 152
Tinea pedis, 152
Tinea unguium, 152
Tinea versicolor, 152
Tinel sign, 459
Tinnitus
  streptomycin, 197
Tiotropium, 241, **687**
Tirofiban, 411, **438**
Tissue factor activation, 133
Tissue plasminogen activator (tPA)
  for ischemic stroke, 512
Tizanidine, 243, **551**
TMP-SMX, 194
  for *Pneumocystis jirovecii*, 154
  prophylaxis, 198
TNF-α, 108, 133, 208
TNF-α inhibitors, 487
TNF (tumor necrosis factor), 227
Tobramycin, 191
Tocolytics, 657
Toddler development, 635
Togaviruses
  characteristics of, 167
  genomes of, 163
  rubella as, 169
Toll-like receptors (TLRs), 99
Tolterodine, 241
Tolvaptan, 354
Tongue
  development, 493
  as ectopic thyroid tissue, 326
  glossoptosis, 620
  pharyngeal arch derivation, 620
Tonic-clonic seizures, 517
  drug therapy for, 544
Tonic seizures, 517
Tonsils
  agammaglobulinemia, 116
  immune system organ, 96
  pharyngeal pouch derivation, 621
Tooth abnormalities
  opalescent teeth, 51
Tophus formation, 467
Topiramate
  epilepsy, 544
  migraine headaches, 518
  pseudotumor cerebri, 521
Topoisomerases, 195
Topotecan, 442
TORCH infections, 169, **182**
  cataracts, 535
  neonatal manifestations, 182
Torsades de pointes, 294
  adenosine for, 324
  Class IA antiarrhythmics, 322
  drug reaction and, 248
  hypomagnesemia, 591
  ibutilide, 323
  sotalol, 323
Torsemide, 608
Torticollis, 519
Torus (buckle) fracture, 462
Total anomalous pulmonary venous return (TAPVR), 298
Total lung capacity (TLC), 664
Total parenteral nutrition (TPN), 396
Total peripheral resistance (TPR), 286
Tourette syndrome, 557
  drug therapy for, 572
  obsessive-compulsive disorder and, 563
  sympatholytic drugs for, 243
Toxic dose, 234
Toxicities and side effects of drugs, 120, 248–253
Toxic megacolon
  Clostridium difficile, 138
Toxic multinodular goiter, 342
Toxic shock-like syndrome, 136
Toxic shock syndrome, 133
  exotoxin A, 133
  presentation, 135
  Staphylococcus aureus, 135
  toxin, 133
Toxins
  myocarditis with, 313
  seafood (ingested), 247
Toxins (bacterial)
  anthrax, 137
  endotoxins, 132
  enterotoxins, 135
  erythrogenic, 136
  exfoliative, **133**, 135
  exotoxins, 132–133
  features of, 131
  lysogenic phage encoding, 130
  toxin-mediated disease, 135
Toxocara canis, 159
Toxocara spp, 158
Toxoid, 111
Toxoplasma gondii, 156
  HIV-positive adults, 177
  TORCH infection, 182
Toxoplasma spp, 180
Toxoplasmosis
  primary central nervous system lymphoma vs, 430
  prophylaxis, 194, **198**
  pyrimethamine, 200
TP53 gene, 224
Trabecula
  spleen, 98
Trachea
  bifurcation of, 663
  fetal development, 326
  respiratory tree, 662
Tracheal deviation, 680, **682**
Tracheoesophageal fistula (TEF)/anomalies, 359
Traction apophysitis, 461
Tractus solitarius, 506
Tramadol, 552
  seizures, 251
"Tram-track" appearance, 596
Transcortical aphasia, 516
Transcription factor, 224
Transduction (bacterial genetics), 130
Transference, 554
Transferrin, 213
  free radical elimination by, 210
  indirect measure of, 419
  lab values in anemia, 419
Transformation (bacterial genetics), 130
Transformation zone (cervix)
  dysplasia, 645
  histology of, 626
Transfusion reaction, 114
Transient arthritis
  in Lyme disease, 146
Transient ischemic attack (TIA), 512
Transitional cell carcinomas, 225
Transition metals and free radical injuries, 210
Transjugular intrahepatic portosystemic shunt (TIPS), 365
Transketolase
  metabolic pathways, 74
  vitamin B_1 and, 66
Translocation
  Down syndrome, 63
  fluorescence in situ hybridization, 55
  in protein synthesis, 45
  Robertsonian, 64
Transmural inflammation fistulas, 382
Transpeptidases, 187
Transplants
  immunosuppressants in, 120
  rejection, 101, **119**
t(8;14), 430, **434**
t(9;22) (Philadelphia), 434
t(14;18), 430, **434**
t(15;17), 434
Transposition (bacterial genetics), 131
Transposition of great vessels, 298
  embryologic development, 281
  maternal diabetes and, 300
Transposon
  in bacterial genetics, 131
Transsexualism, 567
Transversalis fascia, 369
Transverse sinus, 503
Transversion mutation, 39
Transversus abdominis, 369, 452
Transvestism, 567
Tr antigens, 228
Tranylcypromine, 575
Trapezium bone, 449
Trapezoid bone, 449
TRAP (tartrate-resistant acid phosphatase), 227, 342
Trastuzumab, 122, 444–443
Trauma
  pneumothorax, 682
  psychiatric disorders due to, 564
Traumatic aortic rupture, 303
Traumatic pneumothorax, 682
Travelers' diarrhea, 145
Trazodone, 576, 651
Treacher Collins syndrome, 620
"tree bark" appearance (aorta), 312
Trematodes, 160
Tremor, 519
  immunosuppressants, 120
Trench fever, 161
Trendelenburg sign, 453
Treponema, 146
  Gram stain, 125
Treponema pallidum
  penicillin G/V for, 187
  STI, 184
Triamterene, 609
Triazolam, 546
Triceps reflex, 510
Triceps surae, 453
Trichinella spiralis, **159**, 161
Trichinosis, 159
Trichomonas spp
  vaginitis, 181
Trichomonas vaginalis, **158**, 181, 184
Trichomoniasis, 184
Trichophyton spp, 152

(c) ketabton.com: The Digital Library

Trichotillomania, 563
Trichuris, 158
Tricuspid atresia, 281, **298**
Tricuspid regurgitation, 287
  Ebstein anomaly and, 298
  heart murmurs with, 291
Tricuspid valve endocarditis, 311
Tricyclic antidepressants (TCAs)
  antimuscarinic reaction, 251
  mechanism and clinical use, 575
  naming convention for, 253
  torsades de pointes, 248
  toxicity of, 569
  toxicity treatment, 248
  as weak bases, 233
Trientine, 395
Trifluoperazine, 573
Trigeminal nerve (CN V), 506
  lesion of, 532
  neuralgia, 518
  pharyngeal arch derivation, 620
  thalamic relay for, 498
  tongue, 493
Triglycerides
  hypertriglyceridemia, 94
  insulin and, 334
  Von Gierke disease, 87
Trihexyphenidyl, 241
  acute dystonia treatment, 241
Trimethoprim, 187, **194**
  folate deficiency with, 420
  mechanism and use, 194
  pyrimidine synthesis and, 36
  teratogenicity, 614
Trimming (protein synthesis), 45
Trinucleotide repeat expansion diseases, **62**
  Friedreich ataxia, 62
  Huntington disease, 62
  myotonic dystrophy, 62
Triose kinase, 80
Triple-blinded studies, 256
Triptans, 547
  angina and, 304
  for migraine headaches, 518
Triquetrum bone, 449
Trismus (lockjaw)
  tetanospasmin, 138
Trisomies, autosomal, 63
Trisomy 13 (Patau syndrome), **63**, 64
  disease associations with, 491
  hCG in, 633
Trisomy 18 (Edwards syndrome), **63**, 64
  disease associations with, 535
  hCG in, 633
Trisomy 21 (Down syndrome), 62, **63**, 64
  disease associations with, 226, 299–300, 384, 395, 432, 535
tRNA
  structure, 44
Trochlea, 540
Trochlear nerve (CN IV), 506
  ocular motility, 540
  palsy of, 541
Tropheryma whipplei, 125, **381**
Tropical sprue, 381
Tropicamide, 241
Troponins, 304, 306, 456
Trousseau sign, **344**, 591
Trousseau syndrome
  pancreatic cancer, 398
  paraneoplastic syndrome, 228
True-negative rate, 257
True-positive rate, 257
Truncal ataxia, 499
Truncus arteriosus
  22q11 syndromes, 300
Trypanosoma brucei, **156**, 200
Trypanosoma cruzi, 158
  achalasia and, 376
  nifurtimox for, 200
Trypsin, 373
Trypsinogen, 373
Tryptase, 408
Tryptophan, 81
TSC1/TSC2 genes, 224
Tsetse flies (disease vectors), 156
TSST-1 superantigen, 135
t-tests, 264
T-tubule membrane, 456
Tuberculosis, 140
  Addison disease, 349
  corticosteroids and, 336
  erythema nodosum, 482
  isoniazid, 197
  necrosis and, 209
  silicosis, 677
Tuberin protein, 224, **525**
Tuberoinfundibular pathway, 499
Tuberous sclerosis
  chromosome abnormalities, 525
  tumor suppressor genes and, 224
Tubular necrosis, 594, **602**
Tubulointerstitial inflammation
  WBC casts in, 594
Tubulointerstitial nephritis, 601
Tularemia, 149
Tumor grade vs stage, 220
Tumor lysis syndrome, 434–435
Tumor markers (serum), **226**
  acute lymphoblastic leukemia, 432
  colorectal cancer, 388
  pancreatic adenocarcinomas, 398
Tumors
  benign vs malignant, 220
  grade vs stage, 220
  immunohistochemical stains for, 227
  nomenclature of, 220
Tumor suppressor genes, 46, **224**
Tunica albuginea, 651
Tunica vaginalis, 624
Turcot syndrome, 387
Turner syndrome, 638
  cardiac defect association, 300
  coarctation of aorta and, 299
T wave (ECG), 293
21-hydroxylase, 335
22q11 deletion syndromes, 116, 300
Twin concordance studies, 256
Twinning, 616
Thromboxane A_2 (TXA), 411, 486
  aspirin effects, 486
  thrombogenesis, 411
Tympanic membrane, 533
Type I errors (hypothesis testing), 263
Type I hypersensitivity, 105, **112**, 408
Type II errors in hypothesis testing, 263
Type II hypersensitivity, **112**
  organ transplants, 119
Type II hypersensitivity reactions
  rheumatic fever, 312
Type III hypersensitivity reactions, 113
  organ transplants, 119
Type IV hypersensitivity
  DRESS syndrome, 250
  graft-versus-host disease, 119
Type IV hypersensitivity reactions
  contact dermatitis, 477
Typhoid fever, 144
Typhus, 150
  transmission of, **149**, 161
Tyramine, 244
Tyramine-induced hypertensive crisis
  procarbazine, 441
Tyrosinase, 476
Tyrosine
  in phenylketonuria, 84
Tyrosine kinase
  endocrine hormone messenger, 337
  insulin and, 334
  as oncogene product, 224
Tzanck test, 166

U

Ubiquitination, 45
Ubiquitin-proteasome system, 48
UDP-glucuronosyltransferase
  physiologic neonatal jaundice, 393
Ulcerative colitis, 382
  autoantibody, 115
  spondyloarthritis, 469
  sulfasalazine for, 400
Ulcers (gastrointestinal)
  bismuth/sucralfate for, 399
  complications of, 380
  Crohn disease, 382
  Curling, 379
  Cushing, 379
  esophageal, 377
  Helicobacter pylori, 146
  palatal/tongue, 151
  peptic, 379
  Zollinger-Ellison syndrome, 352
Ulcers (skin)
  Raynaud syndrome, 472
Ulipristal, 657
Ulnar claw, 447, 451
Ulnar nerve, **447**, 459
Ulnar nerve injury, 449
Ultrasonography
  fetal cardiac activity on, 612
  kidney disease/disorder diagnoses, 579
  renal cysts on, 604
Umbilical artery, 282, **618**
Umbilical cord, 618
Umbilical hernia
  congenital, 358
Umbilical vein, 618
  blood in, 282
  postnatal derivative of, 282
Umbilicus
  portosystemic anastomosis, 365
UMP synthase, 420
Unambiguous genetic code, 37
Uncal herniation, 529
Uncinate process, 360
Unconjugated bilirubin, 375
Unconjugated (indirect) hyperbilirubinemia, 393
Uncoupling agents, 78
Undifferentiated thyroid carcinomas, 343
Undulant fever, 149
"Unhappy triad" (knee injuries), 460
Unilateral renal agenesis, 579
Uniparental disomy, 57
Universal electron acceptors, 75
Universal genetic code, 37
Unnecessary procedure requests, 268–269
Unstable angina, 304
Unvaccinated children, 186
Upper extremity nerves, 447
Upper motor neuron lesion
  facial nerve, 532
Urachal cysts, 618
Urachus, 282, **618**
Urea cycle, 82
  metabolic site, 72
  ornithine transcarbamylase deficiency and, 83
  rate-determining enzyme for, 73
Ureaplasma spp, 125, 127
Urease, 181
Urease-positive organisms, 127
Uremia
  acute pericarditis, 313
  renal failure, 603
Ureter, 625
  bifid, 579
  constrictions in, 583
  course of, **581**
  embryology, 578
  gynecological exam damage to, 581
  obstruction of, 579, **599**
Ureteric bud, 579
Ureteropelvic junction
  development of, 578
Urethra
  BPH, 654
  injury to, 627
  posterior valves in, 579
Urethritis
  chlamydia, 148, **184**
  Chlamydia trachomatis, 149
  gonorrhea, 184
  reactive arthritis, 469
Urge incontinence, 599
  drug therapy for, 241
Uric acid
  Lesch-Nyhan syndrome, 37
  Von Gierke disease, 87
Uric acid (kidney stones), 598
Urinalysis
  in renal disease, 601
  reducing sugar, 80
Urinary incontinence
  drug therapy for, 241
  ephedrine for, 242
  hydrocephalus, 522
  mechanisms and associations of, 599
  urgency incontinence, 237
Urinary retention, 237
  atropine, 241
  bethanechol for, 240
  delirium, 558
  neostigmine for, 240
  post-void residual, 599
  treatment of, 237
Urinary tract infections (UTIs), 181
  antimicrobial prophylaxis for, 198
  BPH, 654
  cystitis, 600
  duplex collecting system and, 579
  enterococci, 137
  epididymitis and orchitis with, 654
  Klebsiella, 145
  pyelonephritis, 600
  Staphylococcus saprophyticus, 136
  sulfa drugs for, 252
  sulfonamides for, 194
  TMP-SMX for, 194
Urinary tract obstruction, 599
Urine
  Bence Jones proteinuria, 431
  casts in, 594

(c) ketabton.com: The Digital Library

diuretic effects on, 609
leaks with urethral injury, 627
osmolality in acute injury, 601
pregnancy test, 633
renal tubular acidosis, 593
Urine acidification, 233
Urine alkalization, 233
Urine pH and drug elimination, 233
Urobilinogen
extravascular hemolysis, 421
intravascular hemolysis, 421
Urogenital fold, 624
Urosepsis, 600
Urothelial carcinoma (bladder), 606
Urticaria, 475, **477**
ethosuximide, 544
scombroid poisoning, 247
serum sickness, 113
sulfa drug allergies, 252
as type I hypersensitivity, 112
USMLE Step 1 exam
check-in process, 7
clinical vignette strategies, 23
content areas covered in, 3
goal-setting for, 12
leaving exam early, 8
overview of, 2
passing rates for, 10
practice exams for, 11, **21–22**
registering for, 5–6
rescheduling, 6
score notifications for, 7
scoring of, 8–9
testing agencies, 24
testing locations, 6
test-taking strategies, 22–23
time budgeting during, 7–8
types of questions on, 8
Ustekinumab, 122
Uterine conditions
non-neoplastic, 648
Uterine (Müllerian duct) anomalies, 622–623
Uterine neoplasms, 648
Uterovaginal agenesis, 639
Uterus
anomalies of, 623
collagen in, 50
epithelial histology, 626
genital embryology, 622
zygote implantation, 633
Uveitis, 536
inflammatory bowel disease, 382
in sarcoidosis, 676
seronegative spondyloarthritis, 469
U wave in ECG, 293

V

Vaccines
B-cell disorders, 116
Bordetella pertussis, 143
capsular polysaccharide and protein conjugates in, 127
diphtheria, 139
encapsulated bacteria, 127
Haemophilus influenzae, **142**, 180
Poliovirus, 167
PPSV23, 105
rabies, 171
rotavirus, 168
Salmonella typhi, 144
splenectomy and, 98
thymus-independent antigens, 105
toxoids as, 131
types of, 111
Vagal nuclei, 506
Vagina
anatomy of, 626
drainage of, 624
epithelial histology of, 626
genital embryology, 622
Vaginal bleeding
cervical cancer, 645
Vaginal candidiasis
nystatin, 199
Vaginal infections, **181**
Vaginal tumors, **644**
Vaginismus, 567
Vaginitis
Trichomonas spp, **158**, 181
trichomoniasis, 184
Vagus nerve (CN X), 506
baroreceptors/chemoreceptors and, 296
cardiac glycoside effects, 321
Curling ulcers and, 379
diaphragm innervation, 663
gastrointestinal regulation substances and, 371
lesions of, 532
pharyngeal arch derivation, 620
structures innervated, 373
tongue, 493
Valacyclovir, **201**
Validity, 259
Valine
classification of, 81
maple syrup urine disease, 84
Valproic acid
cytochrome P-450, 252
epilepsy, 544
hepatic necrosis, 249
migraine headaches, 518
pancreatitis, 249
Valsartan, 610
Valvular disease
pressure-volume loops, 288
Valvular dysfunction, 310
Vancomycin, 190
Clostridium difficile, 138
cutaneous flushing, 248
meningitis, 180
MRSA, 198
toxicity of, 251
Vanillylmandelic acid (VMA)
in neuroblastomas, 350
Vanishing bile duct syndrome, 119
Varenicline, 576
Variable expressivity, 56
Variance, 262
Variant angina, 304
Variceal bleeding, 245
Varicella zoster virus (VZV), **165**, 475, 479
guanosine analogs, 201
immunodeficient patients, 118
meningitis, 180
rash, 183
vaccine, 110
Varices
Budd-Chiari syndrome, 392
Varicocelectomy, 651
Varicocele (scrotal), 628, **651**
Vasa previa, 641
Vasa vasorum
syphilis, 147
Vascular dementia, 521
Vascular function curves, 286
Vascular tumors of skin, 478
Vasculitides, **314–315**
Vasculitis
focal necrotizing, 315
immunoglobulin A, 315
intraparenchymal hemorrhage, 513
large-vessel, 314
leukoclastic, 173
medium-vessel, 314
methotrexate for, 440
small-vessel, 314
Vasculopathy
noninflammatory, 473
Vas deferens, 626
Vasoactive intestinal polypeptide (VIP), 371
Vasoconstriction, 589
Vasoconstrictors, 550
Vasodilation
sympathetic receptors, 238
Vasodilators
afterload effects, 284
aortic dissections, 303
atrial natriuretic peptide as, 296
coronary steal syndrome, 304
nitrates as, 318
Vasogenic edema, 496
Vasopressin, 329
receptors, 238
Vasopressors, 286
V(D)J recombination, 99
VDRL false positives, 148
Vector-borne illnesses, 150
Veganism and B_{12} deficiency, 420
Vegetative state
axonal injury and, 515
VEGF (vascular endothelial growth factor), 216
Velocardiofacial syndrome, 116
Vemurafenib, **444**, 484
Venlafaxine, 575
clinical use, 572
panic disorder, 563
phobias, 563
PTSD, 564
Venodilators, 284
Venous gonadal drainage, 624
Venous return, 286
Venous sinus thrombosis, 503
Venous thrombosis
heparin for, 436
paroxysmal nocturnal hemoglobinuria, 422
Ventilation, 664
high altitude, 670
perfusion and, 669
Ventilation/perfusion (V/Q) defects, 664
Ventilation/perfusion (V/Q) ratio
exercise response, 670
mismatch, 669
Ventral lateral (VL) nucleus, 498
Ventral pancreatic bud, 360
Ventral posterolateral (VPL) nucleus, 498
Ventral posteromedial (VPM) nucleus, 498
Ventral tegmentum, 495
Ventricles
contractility of, 285
embryology, 281
morphogenesis of, 281
Ventricular action potential, 292
Ventricular aneurysm
pseudoaneurysm, 307
true, 305, **307**
Ventricular fibrillation
ECG tracing, 295
torsades de pointes, 294
Ventricular filling
early diastole, 287
ECG and, 293
Ventricular free wall rupture, 307
Ventricular myocytes, 296
Ventricular noncompliance, 287
Ventricular septal defect (VSD), 299
congenital rubella, 300
cri-du-chat syndrome, 64
Down syndrome, 300
fetal alcohol syndrome, 300
heart murmurs, 291
outflow tract formation, 281
Ventricular system, 504
Ventriculomegaly, 520, **522**
Ventromedial nucleus (hypothalamus), 498
Verapamil, 308, **318**, 319, 321, 518
Verrucae, 477
Vertebral compression fractures, 462
Vertebral landmarks
diaphragm, 663
Vertigo, 534
posterior circulation stroke, 514
streptomycin, 197
Vesicles (skin), 475
varicella zoster virus, 479
Vesicourachal diverticulum, 618
Vesicoureteral reflux, 579
hydronephrosis, 599
pyelonephritis, 600
Vesicular monoamine transporter (VMAT), 549
Vesicular tinea pedis, 152
Vesicular trafficking proteins, 47
Vestibular schwannomas, 527
Vestibulocochlear nerve (CN VIII), 506
VHL gene, 224
Vibrio cholerae, 146
exotoxin production, 132
watery diarrhea, 179
Vibrio parahaemolyticus, 178
Vibrio vulnificus, 178
Vigabatrin, 544
Vilazodone, 576
Vimentin, 48, **227**
Vinblastine, 441
microtubules and, 48
Vinca alkaloids, 438
Vincristine, 441
microtubules and, 48
toxicities of, 444
Vinyl chloride
angiosarcomas, 392, **478**
as carcinogen, 225
VIPomas
MEN 1 syndrome, 351
octreotide for, 400
regulatory substances, 371
Viral envelopes, 163
Virchow nodes, 379
Viridans streptococci, 136
α-hemolysis, 135
bacterial endocarditis, 311
biofilm production, 128
brain abscesses, 180
gram-positive algorithm, 134
normal flora, 178
Virilization, 335
Virology, 162–177

FAS1_2019_21_Index_749-806.indd 801 11/21/19 12:27 PM

(c) ketabton.com: The Digital Library

Virulence factors
bacterial, 129
Bordetella pertussis, 143
Escherichia coli, 145
Salmonella/Shigella, 144
Staphylococcus aureus, 135
Streptococcus pneumoniae, 136
β-hemolytic bacteria, 135
Viruses
diarrhea with, 179
fluorescent antibody stain, 125
genetics, 162
immunocompromised patients, 179
in immunodeficiency, 118
as cause of myocarditis, 313
negative-stranded, 168
pneumonia, 179
receptors for, 166
segmented, 168
skin, 479
structure of, **162**
Visceral leishmaniasis, 158
Viscosity (blood), 286
Visual cortex, **501**, 515
Visual disturbance
drug-related, 251
Visual field defects, 542
saccular aneurysms and, 516
with stroke, 514, 515
Visual hallucinations, 559
Vital capacity (VC), 664
Vitamin A (retinol), **66**
free radical elimination by, 210
idiopathic intracranial hypertension, 251, **521**
measles morbidity and mortality, 170
teratogenicity, 614
Vitamin B_1 (thiamine), **66**
brain lesions and, 511
deficiency of, 66
functions of, 74
pyruvate dehydrogenase complex, 76
solubility of, 65
Vitamin B_2 (riboflavin), **67**
pyruvate dehydrogenase complex, 76
solubility, 65
Vitamin B_3 (niacin), **67**
pyruvate dehydrogenase complex, 76
solubility, 65
vitamin B_6 and, 67
Vitamin B_5 (pantothenic acid), **67**
pyruvate dehydrogenase complex and, 76
solubility of, 65
Vitamin B_6 (pyridoxine), **67**
deficiency, 67
isoniazid, 197
sideroblastic anemia, 419
Vitamin B_7 (biotin), **68**
activated carriers, 75
functions of, 73
pyruvate metabolism, 77, 78
solubility of, 65
Vitamin B_9 (folate), **68**
deficiency, 406, 420
functions, 68
solubility, 65
Vitamin B_{12} (cobalamin), **69**
absorption of, 374
deficiency, 160, 161
solubility, 65
spinal cord lesions, **530**
Vitamin C (ascorbic acid)
free radical elimination by, 210
functions, 69
methemoglobin treatment, 248
solubility of, 65
Vitamin D
excess, 70
functions, 70
hyperparathyroidism, 464
hypervitaminosis lab values, 464
osteomalacia/rickets, 463, 464
osteoporosis prophylaxis, 462
signaling pathways for, 337
solubility of, 65
vitamin D (calciferol)
calcitriol production, 589
Vitamin E
free radical elimination by, 210
function, 70
solubility of, 65
Vitamin K
cephalosporins, 189
coagulation cascade, 413
deficiency, **413**, 426
solubility of, 65
vitamin E interaction, 71
for warfarin toxicity, 436
Vitamin/mineral absorption, 374
Vitamins
fat-soluble, **65**
water-soluble, **65**
Vitelline duct/fistula, 618
Vitiligo, 476
Vitreous body
collagen in, 50
VLDL (very low-density lipoprotein), 94
Volume contraction
alkalemia from diuretics, 609
Volume of distribution, 231
Volumetric flow rate ($\dot{Q}$), 286
Volvulus, 385, **386**
Meckel diverticulum, 384
midgut, 386
Onchocerca, 158
sigmoid, 386
Vomiting
annular pancreas, 360
biliary colic, 396
bilious, 359, 384
diabetic ketoacidosis, 347
Ebola virus, 171
fructose intolerance, 80
glycylcyclines, 192
Histoplasma capsulatum, 177
intestinal atresia, 359
Legionella spp, 185
lithium toxicity, 569
Mallory-Weiss syndrome, 377
maple syrup urine disease, 84
metoclopramide for, 400
MI and, 305
ondansetron for, 400
posttussive, 143
pyloric stenosis, 359
receptors for, 496
Salmonella spp, 149
in stroke, 514
toxic shock syndrome, 135
treatment of, 400, 401
trichinosis, 159
vitamin C toxicity, 69
Von Gierke disease, 87
von Hippel-Lindau disease, 525
chromosome association, 64
renal cell carcinoma and, 605
tumor suppressor genes and, 224
von Willebrand disease, 411, **428**
Voriconazole, 199
Vortioxetine, 576
VRE (vancomycin-resistant enterococci)
daptomycin, 195
enterococci, 137
highly resistant, 198
oxazolidinones, 193
Vulnerable child syndrome, **556**
Vulvar carcinoma, **644**
Vulvar lymphatic drainage, 624
Vulvar pathology, 644
neoplastic, 644
non-neoplastic, 644
Vulvovaginitis, 153, **181**
vWF (von Willebrand factor)
receptor for, 407
in thrombocytes, 407
in thrombogenesis, 411
V_{max}, 230

W

WAGR complex/syndrome, 606
"Waiter's tip" (Erb palsy), 448
Waiving right to confidentiality, 267
Waldenstrom macroglobulinemia, 431
Walking milestone, 635
Wallenberg syndrome, 514
Wallerian degeneration, 495
Wall tension, 284, 285
Warburg effect, 221
Warfarin, 436
adverse effects of, 428
coagulation cascade, 413
griseofulvin and, 200
heparin vs, 436, 437
PT measurement, 426
teratogenicity, 614
therapeutic index of, 234
toxicity treatment, **248**, 429
vitamin K antagonist, 71
Warthin-Finkeldey giant cells, 170
Warthin tumors, 376
Waterhouse-Friderichsen syndrome, 349
meningococci, 142
Watershed zones, **210**, 502
Water-soluble vitamins, 65
Waxy casts (urine), 594
WBC casts (urine), **594**, 600
Weak acid overdose
treatment, 233
Weak bases overdose treatment, 233
"Wear and tear" pigment, 211
Wegener granulomatosis, 315
autoantibody, 115
restrictive lung disease, 675
Weight gain
danazol, 658
duodenal ulcer, 380
mirtazapine, 576
Weight loss
adrenal insufficiency, 349
cholelithiasis and, 396
chronic mesenteric ischemia, 386
diabetes mellitus, 346
esophageal cancer, 378
gastric ulcers, 380
glucagonoma, 351
Histoplasma capsulatum, 177
malabsorption syndromes, 381
Mycobacterium avium-intracellulare, 177
orlistat for, 400
pancreatic cancer, 398
polyarteritis nodosa, 314
polymyalgia rheumatica, 470
pseudotumor cerebri treatment, 521
renal cell carcinoma, 605
sleep apnea, 679
stomach cancer, 379
for stress incontinence, 599
tuberculosis, 140
Weil disease, 147
Well-patient care, 270–271
Wenckebach AV block, 295
Werdnig-Hoffmann disease, 530
Wernicke aphasia, 514, **516**
Wernicke area, 501
stroke effects, 514
Wernicke encephalopathy, 66, **571**
Wernicke-Korsakoff syndrome, 511, **571**
vitamin B_1 deficiency, 66
Western blot, 53
West Nile virus, **167**, 180
Wet beriberi, 66
Wharton duct, 376
Wharton jelly, 618
Wheal
urticaria, 477
Wheals, 475
Wheezing
lung cancer, 684
Whipple disease, 381
periodic acid-Schiff stain for, 125
Whipple procedure
for pancreatic cancer, 398
Whipple triad
insulinomas and, 351
Whispered pectoriloquy, 680
White matter
axonal injury, 515
demyelinating disorders, 524
glial cells in, 494
multiple sclerosis, 523
White pulp (spleen), 98
Whooping cough
Bordetella pertussis, 143
pertussis toxin, 132
Wickham striae, 482
Wide splitting, 289
Williams syndrome, 64
cardiac defect association, 300
Wilms tumor, 606
dactinomycin for, 439
neuroblastomas vs, 350
tumor suppressor genes and, 224
Wilson disease, 395
ATP7B protein in, 51
autosomal recessive inheritance, 60
chromosome association, 64
Fanconi syndrome, 586
free radical injury and, 210
Winged scapula, 448
Winters formula, 592

(c) ketabton.com: The Digital Library

"Wire looping" of capillaries, 596
"Wire lupus," 596
Wiskott-Aldrich syndrome, 117
  X-linked recessive disorder, 61
Withdrawal (psychoactive drugs), 570
Wobble, 37
Wolff-Chaikoff effect, 341
Wolffian duct, 622
Wolff-Parkinson-White syndrome, 294
Woolsorter's disease, 137
"word salad," 559
Wound healing
  keratinocytes, 216
  phases of, 216
  scar/keloid formation, 218
Woven bone, 458
Wright-Giemsa stain, 407
Wright stain
  Borrelia spp, 146
Wrinkles of aging, 52
Wrist
  injuries of, 459
  bones, 449
Wrist drop, 447
  lead poisoning, 419
Written advance directives, 266
WT1/*WT2* genes
  in renal disease, 224, **606**
  oncogenicity of, 224
Wuchereria bancrofti, 158, 159

X

Xanthine oxidase inhibitors, 467
Xanthogranulomatous pyelonephritis, 600
Xanthomas
  familial dyslipidemias, 94
  hyperlipidemia and, 301
Xeroderma pigmentosum
  DNA repair defects in, 40
Xerosis cutis, 66
Xerostomia, 240, 243, **468**
X-inactivation (lyonization)
  Barr body formation, 61
X-linked agammaglobulinemia, 116
X-linked dominant inheritance, 59
X-linked recessive disorders, **61**
  agammaglobulinemia, 116
  hyper-IgM syndrome, 117
  Menkes disease, 51
  NADPH oxidase defect, 117
  Wiskott-Aldrich syndrome, 117
X-linked recessive inheritance, 59
X-ray/imaging findings
  Apple core lesion, 388
  bamboo spine, 469
  Bird's beak sign, 376
  Bone-in-bone, 463
  Codman triangle, 465
  Coffee bean sign, 386
  Coin lesion, 684
  Crew cut (skull x-ray), 422
  kidney stones, 598
  pencil-in-cup, 469
  punched out bone lesions, 431
  Steeple sign (x-ray), 170
  String sign, 382
  Sunburst pattern, 465
  Thumbprint sign (imaging), 386
  Thumb sign, 142
X-ray teratogenicity, 614

Y

Yellow fever, 167, **168**
  liver anatomy and, 367
Yersinia enterocolitica, 179
  transmission and treatment, 144
Yersinia pestis
  animal transmission, 149
  facultative intracellular organisms, 127
Yo antigens, 228
Yolk sac tumor
  ovarian, 647
  testicular, 653
Yersinia spp
  reactive arthritis, 469

Z

Zafirlukast, 687
Zaleplon, 546
Zanamivir, 201
Zellweger syndrome, 47
Zenker diverticulum, 384
Zero-order elimination, 232
Zidovudine, 203
Ziehl-Neelsen stain, 125
Zika virus, 171
Zileuton, 687
Zinc, 71
  Wilson disease, 395
Ziprasidone, 573
Zoledronic acid, 486
Zollinger-Ellison syndrome, 352
  duodenal ulcer, 380
  gastrin in, 371
  MEN 1 syndrome, 351
  proton pump inhibitors for, 399
Zolpidem, 546
Zona fasciculata, **327**, 336
Zona glomerulosa, 327
Zona reticularis, 327
Zoonotic bacteria, 149
Zymogens, 373

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

NOTES

(c) ketabton.com: The Digital Library

About the Editors

Tao Le, MD, MHS

Tao developed a passion for medical education as a medical student. He currently edits more than 15 titles in the *First Aid* series. In addition, he is Founder and Chief Education Officer of USMLE-Rx for exam preparation and ScholarRx for undergraduate medical education. As a medical student, he was editor-in-chief of the University of California, San Francisco (UCSF) *Synapse,* a university newspaper with a weekly circulation of 9000. Tao earned his medical degree from UCSF in 1996 and completed his residency training in internal medicine at Yale University and fellowship training at Johns Hopkins University. Tao subsequently went on to cofound Medsn, a medical education technology venture, and served as its chief medical officer. He is currently chief of adult allergy and immunology at the University of Louisville.

Matthew Sochat, MD

Matthew is a third-year hematology/oncology fellow at St. Louis University in St. Louis, Missouri. He completed his internal medicine residency training at Temple University Hospital in Philadelphia. He completed medical school in 2013 at Brown University and is a 2008 graduate of the University of Massachusetts, Amherst, where he studied biochemistry and the classics. Pastimes include skiing, cooking/baking, traveling, the company of friends/loved ones (especially his wonderful wife), the Spanish language, and computer/video gaming. Be warned: Matt also loves to come up with corny jokes at (in)opportune moments.

Sarah Schimansky, MB BCh BAO

Sarah is a third-year ophthalmology resident in the UK. She grew up in Germany before moving to Dublin, Ireland, to study medicine at the Royal College of Surgeons in Ireland. She has a keen interest in medical education and is currently enrolled in a Masters in Surgical Education program at Imperial College London. An avid traveler, Sarah is always on the lookout for new destinations to explore and new countries to call home. When she is not on the road, she enjoys yoga, long walks, and red wine in the company of friends and family.

Kimberly Kallianos, MD

Originally from Atlanta, Kimberly graduated from the University of North Carolina at Chapel Hill in 2006 and from Harvard Medical School in 2011. She completed her radiology residency and fellowship at UCSF and is currently an Assistant Professor of Clinical Radiology at UCSF in the Cardiac and Pulmonary Imaging section.

Vikas Bhushan, MD

Vikas is a writer, editor, entrepreneur, and teleradiologist on extended sabbatical. In 1990 he conceived and authored the original *First Aid for the USMLE Step 1*. His entrepreneurial endeavors included a student-focused medical publisher (S2S), an e-learning company (medschool.com), and an ER teleradiology practice (24/7 Radiology). Trained on the Left Coast, Vikas completed a bachelor's degree at the University of California Berkeley; an MD with thesis at UCSF; and a diagnostic radiology residency at UCLA. His eclectic interests include technology, cryptoeconomics, information design, South Asian diasporic culture, and avoiding a day job. Always finding the long shortcut, Vikas is an adventurer, knowledge seeker, and occasional innovator. He enjoys intermediate status as a kiteboarder and father, and strives to raise his three children as global citizens.

Vaishnavi Vaidyanathan, MD

Vaishnavi is a second-year child neurology resident at Phoenix Children's Hospital in Phoenix, Arizona. She is a graduate of the University of Missouri-Kansas City School of Medicine, where she earned her bachelor's and medical degrees. Her interests include medical education and health advocacy. Outside of medicine, she loves to dance, learn new languages, and watch Bollywood movies.

Jordan Abrams

Jordan is a fourth-year medical student at St. George's University School of Medicine who hopes to pursue residency training in anesthesiology. He graduated magna cum laude from the University of Delaware, earning a bachelor's degree in neuroscience with minors in medical humanities and biological sciences. Combining his creative mindset and passion for drawing, Jordan founded theHYMedicine.com, an educational website that offers free medical study guides, tutoring, and study schedules for students worldwide. Aside from medicine, Jordan enjoys traveling, reading, and playing soccer.

(c) ketabton.com: The Digital Library

SECTION IV

Top-Rated Review Resources

"Some books are to be tasted, others to be swallowed, and some few to be chewed and digested."

—Sir Francis Bacon

"Always read something that will make you look good if you die in the middle of it."

—P.J. O'Rourke

"So many books, so little time."

—Frank Zappa

"If one cannot enjoy reading a book over and over again, there is no use in reading it at all."

—Oscar Wilde

- How to Use the Database 2
- Question Banks and Books 4
- Web and Mobile Apps 6
- Comprehensive 10
- Anatomy, Embryology, and Neuroscience 12
- Behavioral Science 14
- Biochemistry 15
- Cell Biology and Histology 16
- Microbiology and Immunology 17
- Pathology 19
- Pharmacology 21
- Physiology 22

FAS1_2019_23_BookRev_ONLINE.indd 1 11/8/19 6:49 AM

(c) ketabton.com: The Digital Library

HOW TO USE THE DATABASE

This section is a database of top-rated basic science review books, sample examination books, websites, apps, and commercial review courses that have been marketed to medical students studying for the USMLE Step 1. At the end of the section is a list of publishers and independent bookstores with addresses and phone numbers. For each recommended resource, we list (where applicable) the **Title**, the **First Author** (or editor), the **Series Name** (where applicable), the **Current Publisher**, the **Copyright Year**, the **Number of Pages**, the **ISBN**, the **Approximate List Price**, the **Format** of the resource, and the **Number of Test Questions**. We also include **Summary Comments** that describe their style and overall utility for studying. Finally, each recommended resource receives a **Rating**. Within each section, resources are arranged first by Rating and then alphabetically by the first author within each Rating group.

A letter rating scale with six different grades reflects the detailed student evaluations for **Rated Resources**. Each rated resource receives a rating as follows:

A+	Excellent for boards review.
A A–	Very good for boards review; choose among the group.
B+ B	Good, but use only after exhausting better resources.
B–	Fair, but there are many better resources in the discipline; or low-yield subject material.

The Rating is meant to reflect the overall usefulness of the resource in helping medical students prepare for the USMLE Step 1. This is based on a number of factors, including:

- The cost
- The readability of the text
- The appropriateness and accuracy of the material
- The quality and number of sample questions
- The quality of written answers to sample questions
- The quality and appropriateness of the images and illustrations
- The quality of the user interface and learning experience, for web and mobile apps
- The length of the text (longer is not necessarily better)
- The quality and number of other resources available in the same discipline
- The importance of the discipline for the USMLE Step 1

Please note that ratings do not reflect the quality of the resources for purposes other than reviewing for the USMLE Step 1. Many books with

(c) ketabton.com: The Digital Library

lower ratings are well written and informative but are not ideal for boards preparation. We have not listed or commented on general textbooks available in the basic sciences.

Evaluations are based on the cumulative results of formal and informal surveys of thousands of medical students at many medical schools across the country. The summary comments and overall ratings represent a consensus opinion, but there may have been a broad range of opinion or limited student feedback on any particular resource.

Please note that the data listed are subject to change in that:

- Publisher and app store prices change frequently.
- Retail and online bookstores may set their own prices.
- New editions and app versions come out frequently, and the quality of updating varies.
- The same book may be reissued through another publisher.

We actively encourage medical students and faculty to submit their opinions and ratings of these basic science review materials so that we may update our database. In addition, we ask that publishers and authors submit for evaluation review copies of basic science review books, including new editions and books not included in our database. We also solicit reviews of new books, mobile apps, websites, flash cards, and commercial review courses.

Disclaimer/Conflict of Interest Statement

None of the ratings reflects the opinion or influence of the publisher. All errors and omissions will gladly be corrected if brought to the attention of the authors through our blog at www.firstaidteam.com. Please note that USMLE-Rx and the entire *First Aid for the USMLE* series are publications by certain authors of *First Aid for the USMLE Step 1*; the following ratings are based solely on recommendations from the student authors of *First Aid for the USMLE Step 1* as well as data from the student survey and feedback forms.

(c) ketabton.com: The Digital Library

▸ QUESTION BANKS AND BOOKS

A+

UWorld Qbank

$249–$749 Test/2400 q

UWORLD

www.uworld.com

Questions demand multistep reasoning and are often more difficult than those on the actual Step 1 exam. Offers detailed explanations with figures and tables. Features a number of test customization and analysis options. Users can see cumulative results both over time and compared to other test takers. In addition to a desktop version, it can be accessed through iOS or Android mobile apps.

A

NBME Practice Exams

$60 Test/200 q

NATIONAL BOARD OF MEDICAL EXAMINERS

www.nbme.org/students/sas/Comprehensive.html

The official practice exams published by the NBME are comprised of retired Step 1 questions. NBME research found that they show a "moderate correlation" with actual Step 1 performance. The exams will show you which questions you answered incorrectly, but they will not show any explanations. You will also not be able to review correctly answered questions. Students generally use these as rough gauges of their score progression over their study time. Note that you can sign up to for an in-person practice exam for an additional $75 to be taken at Prometric, for students who want to practice the logistics of exam day.

A–

AMBOSS

$9–$365 Test/3500 q

AMBOSS

www.amboss.com

Integrated question bank for Step 1 and Step 2 CK exams with an additional interactive online library of medical resources. Contains numerous illustrations within the clinical vignettes. Allows for the selection of questions by difficulty level. Includes personalized study plan. Free trial available, accessible through iOS or Android mobile apps.

A–

USMLE-Rx Qmax

$89–$339 Test/2300 q

USMLE-RX

www.usmle-rx.com

Offers Step 1–style questions accompanied by thorough explanations. Omits obscure material and distills high yield information. Each explanation includes references from *First Aid*. However, the proportion of questions covering a given subject area does not always reflect the actual exam's relative emphasis. Question stems occasionally rely on "buzzwords." Most useful to help memorize *First Aid* facts. Provides detailed performance analyses. Free trial available, accessible through iOS or Android mobile apps.

(c) ketabton.com: The Digital Library

B+

Kaplan Qbank

$99–$349 Test/2100 q

KAPLAN

www.kaptest.com

Covers most content found on Step 1, but sometimes emphasizes recall of low-yield details rather than integrative problem-solving skills. Test content and performance feedback can be organized by both organ system and discipline. Includes detailed explanations of all answer choices. Users can see cumulative results both over time and compared to other test takers. Accessible through iOS or Android mobile apps.

B

BoardVitals

$59–$179 Test/1750 q

www.boardvitals.com

Comprehensive question bank modeled closely after the format of the Step 1. Covers all subject areas and includes explanations for each answer choice. Users can create custom exams and compare their performance to national averages. Contains fewer image-based questions compared to similar platforms.

B

Kaplan USMLE Step 1 Qbook

$50 Test/850 q

KAPLAN

Kaplan, 2017, 468 pages, ISBN 9781506223544

Consists of over 850 exam-like questions organized by the traditional basic science disciplines. Similar to the Kaplan Qbank, and offers USMLE-style questions with clear, detailed explanations; however, lacks classic images typically seen on the exam. Also includes access to a sample online question bank and a guide on test-taking strategies.

B

Pastest

$79–$249 Test/2100 q

www.pastest.com

Questions appear to be simpler than board-style questions, with many first- and second-order questions. Explanations are accompanied by references to *First Aid* and short video clips to reinforce information. Accessible through iOS or Android mobile apps.

B

TrueLearn Review

$159–$399 Test/2200 q

www.truelearn.com

Includes over 2200 USMLE-style practice questions with topics mapped to the NBME blueprint. Uses national benchmarking to show students where they stand in comparison to peers.

(c) ketabton.com: The Digital Library

▶ WEB AND MOBILE APPS

A	***Anki*** **www.ankisrs.net**	***Free***	Flash cards
	Flash card–making resource designed for retention of facts through spaced repetition. Free access via desktop and smartphone for Windows, Mac, and Android. The iOS app must be purchased for $25. Available in different languages.		
A	***Boards and Beyond*** **www.boardsbeyond.com**	***$19–$249***	Review
	Includes over 400 videos averaging ~26 minutes each, covering the breadth of Step 1 material. Membership includes access to the companion books as PDFs. A collection of videos is offered as free samples on the website. Also includes over 1300 practice questions.		
A	***Physeo*** **www.physeo.com**	***$30–$150***	Review
	Online review containing 32 hours of review videos covering physiology. Accessible via website or mobile app. Includes a supplemental full-color PDF textbook. Videos are concise and focus on high-yield material, and board-style practice questions are included after each topic to help solidify understanding. Similar structure to Pathoma, but with physiology focus.		
A	***SketchyMedical*** **www.sketchymedical.com**	***$99–$369***	Review
	Video library of narrated lectures with thorough explanations that present microbiology, pharmacology, and pathology in a memorable style. Access to the entire gram-positive cocci section is free at signup. Additional content can be purchased on a subscription basis.		
A⁻	***Cram Fighter*** **www.cramfighter.com**	***$29–$159***	Study plan
	Helps organize a study schedule. Highly flexible with customizable settings. Supports more than 650 of the most popular books, video lectures, question banks, and flash cards. Mobile apps available for iOS and Android.		
A⁻	***First Aid Step 1 Express*** **www.usmle-rx.com**	***$69–$299***	Review/Test
	More than 80 hours of high-yield videos explaining material from *First Aid for the USMLE Step 1*. Videos include more than 600 extra images and multimedia clips. Step-by-step analysis of USMLE-style questions with each video. Subscription includes a color workbook with over 200 pages.		
B⁺	***First Aid Step 1 Flash Facts*** **www.usmle-rx.com**	***$29–$149***	Flash cards
	Access to 12,000+ flash cards with intelligent spaced repetition integrated with *First Aid for the USMLE Step 1*, of which 3500+ are case based. Updated each year to reflect the newest edition of the book; students can access the past 3 editions' worth of flash cards. Searchable by organ system, discipline, and topic.		

(c) ketabton.com: The Digital Library

B+	***Medbullets*** **www.medbullets.com** Free online learning and collaboration community for students preparing for their exams. Supplements medical school coursework and Step 1 studying with simplified, to-the-point online search platform that is best used as a reference. Recently added premium content for $80-$250 includes an online question bank and adaptive learning system.	***Free***	Review/ Test/1000 q
B+	***Medical School Pathology*** **www.medicalschoolpathology.com** Offers lectures and slides based on the Robbins *Pathology* textbook. Lectures can be downloaded.	***Free***	Review
B+	***OnlineMedEd*** **www.onlinemeded.org** A video lecture series covering primarily clinical science material, with recent addition of biochemistry, cell biology, and immunology topics. Video access is free with registration. A subscription of $10-$70/month gains access to ad-free videos, lecture notes, flash cards, question bank, and downloadable audio lectures.	***Free***	Review
B+	***Osmosis*** **www.osmosis.org** Web platform that includes exam study scheduling tool, 27,000+ variable quality multiple choice questions, flash cards with spaced repetition, and 3000+ curated concept cards with videos, memory anchors, and reference articles. Includes a curriculum analysis and search engine, collaboration features for study groups, and a mobile app with quizzes and videos.	***$179–$279***	Test
B+	***USMLE Step 1 Mastery*** **builtbyhlt.com/medical/usmle-step-1-mastery** Question bank accessible through website or via free mobile app. Covers all USMLE topics and includes vignettes, images, and mnemonics. Question formatting is generally less representative of actual USMLE questions compared with other widely used question banks. Mobile app contains supplemental flash cards for integrated learning.	***$2–$10***	Test/1400 q
B+	***WebPath: The Internet Pathology Laboratory*** **webpath.med.utah.edu** Features more than 2700 gross and microscopic images, clinical vignette questions, and case studies. Includes nine general pathology exams and 11 system-based pathology exams with approximately 1300 questions. Also features 170 questions associated with images. Questions are useful for reviewing boards content but are typically untimed, easier, and shorter. No multimedia practice questions. Not regularly updated with regard to high-yield Step 1 material.	***Free***	Review/ Test/1300 q
B	***Blue Histology*** **www.lab.anhb.uwa.edu.au/mb140** Provides access to 400+ histologic images with thorough explanations. Images searchable by topic, stain, keyword. Website also contains multiple choice practice questions.	***Free***	Review/Test

(c) ketabton.com: The Digital Library

B

Digital Anatomist Project: Interactive Atlases

Free Review

UNIVERSITY OF WASHINGTON

da.si.washington.edu/da.html

Contains an interactive neuroanatomy course along with a three-dimensional atlas of the brain, thorax, and knee. Atlases have computer-generated images and cadaver sections. Each atlas also has a quiz in which users identify structures in the slide images. However, questions do not focus on high-yield anatomy for Step 1.

B

Dr. Najeeb Lectures

$99 Review

www.drnajeeblectures.com

Hundreds of hours of video lectures covering basic medical sciences and clinical medicine with thousands of hand-drawn illustrations and mnemonics. Website provides mobile video support on smartphones and tablets. Free lectures accessible at www.drnajeeblectures.com/free-medical-videos.html.

B

Firecracker

$39–$660 Review/ Test/2800 q

FIRECRACKER INC.

firecracker.lww.com

Learning platform divided into modules. The Step 1 module is divided into organ systems and includes review of preclinical lecture material, periodic quizzes on flagged reviewed material, and USMLE-style questions in interface simulating the actual exam. Contains page references to *First Aid for the USMLE Step 1* and high-yield diagrams from various textbooks. Users can grade how well they remember the quiz answers (1–5), which allows the program to customize future quizzes. Features detailed performance analysis and a calendar for personalized study plans. Accessible on all smartphones and tablets. Comprehensive; best if started early in preclinical years.

B

KISSPrep

$99–$135 Review

www.kissprep.com

Online lecture videos focused on select subjects from the Step 1 exam. Focuses on harder-to-learn content and teaches it in a simple, easy-to-understand manner. Quizzes and other interactive tools are available to help with knowledge retention. Not all Step 1 content is covered on this platform.

B

Lecturio

$50–$300 Review/ Test/2150 q

www.lecturio.com

Online platform for comprehensive exam preparation, including over 250 hours of lectures, a flash card deck, quizzes, and a question bank. Organized by subject matter and allows users to customize their learning experience. Some content may be beyond the scope of the exam and better suited for medical school coursework. Lectures and quizzes may be accessed for free. iOS and Android apps are available.

B

Memorang

$19–$239 Flash cards

MEMORANG INC.

www.memorangapp.com

Platform utilizing spaced repetition, available both in website and app form. Utilizes custom and/or premade flash card "study sets" derived from 15,000 flash cards that focus on specific subject areas and are then tested via various games and quizzing methods. Free 7-day trial, or monthly/annual membership.

(c) ketabton.com: The Digital Library

B — ***Picmonic*** — ***$25–$480*** — Review

www.picmonic.com

Helpful resource for visual learners. Unique images and stories with daily quizzes and spaced repetition. Contains 1400 images and includes study guides, webinars, and infographics that help cover 15,000+ facts of Step 1 material. Offered via both web and mobile platforms.

B– — ***Radiopaedia.org*** — ***Free*** — Cases/Test

www.radiopaedia.org

A user-friendly website with thousands of well-organized radiology cases and articles. Encyclopedia entries contain high-yield bullet points of anatomy and pathology. Images contain detailed descriptions but no arrows to demarcate findings. Quiz mode allows students to make a diagnosis based on radiographic findings. Content may be too broad for boards review but is a good complement to classes and clerkships.

B– — ***The Pathology Guy*** — ***Free*** — Review

FRIEDLANDER

www.pathguy.com

Contains extensive but poorly organized information on a variety of fundamental concepts in pathology. A high-yield summary intended for USMLE review can be found at www.pathguy.com/meltdown.txt, but the information given is limited by a lack of images and frequent digressions.

(c) ketabton.com: The Digital Library

▸ COMPREHENSIVE

A — ***First Aid for the Basic Sciences: General Principles*** — ***$55*** — Review

LE

McGraw-Hill, 2019, 816 pages, ISBN 9781260143676

Comprehensive review of the basic sciences covered in year 1 of medical school. Similar to the first part of *First Aid*, organized by discipline, and includes hundreds of color images and tables. Best if started with first-year coursework and then used as a reference during boards preparation.

A — ***First Aid Cases for the USMLE Step 1*** — ***$50*** — Cases

LE

McGraw-Hill, 2018, 496 pages, ISBN 9781260143133

A recently updated series of hundreds of high-yield cases organized by organ system. Each case features a clinical vignette with relevant images, followed by questions and short, high-yield explanations. Offers coverage of many frequently tested concepts, and integrates subject matter in the discussion of the vignette. Helpful in reviewing material outlined in *First Aid for the USMLE Step 1*.

A− — ***First Aid for the Basic Sciences: Organ Systems*** — ***$72*** — Review

LE

McGraw-Hill, 2017, 912 pages, ISBN 9781259587030

A comprehensive review of the basic sciences covered in year 2 of medical school. Similar to the second part of *First Aid*, organized by organ system, and includes hundreds of color images and tables. Each organ system contains discussion of embryology and anatomy, physiology, pathology, and pharmacology. Best if started with second-year coursework and then used as a reference during boards preparation.

A− — ***Crush Step 1: The Ultimate USMLE Step 1 Review*** — ***$45*** — Review

O'CONNELL

Elsevier, 2017, 704 pages, 9780323481632

Detailed, text-heavy review book with practice questions included. Coverage of many high-yield topics but includes some outdated information. Best if used with coursework, but also recommended as a supplemental reference for boards review. Limited student feedback.

A− — ***Cracking the USMLE Step 1*** — ***$45*** — Review

PRINCETON REVIEW

Princeton Review, 2013, 832 pages, ISBN 9780307945068

Comprehensive review book with hundreds of illustrations, charts, and diagrams along with 2 full-length practice tests with detailed answer explanations available online. Limited student feedback.

B+ — ***USMLE Step 1 Secrets in Color*** — ***$43*** — Review

BROWN

Elsevier, 2016, 800 pages, ISBN 9780323396790

Clarifies difficult concepts in a concise, readable manner. Uses a case-based format and integrates information well. High-quality clinical images. Complements other boards study resources, with a focus on understanding preclinical fundamentals rather than on rote memorization. Slightly lengthy for last-minute studying.

(c) ketabton.com: The Digital Library

B+ | ***Step-Up to USMLE Step 1 2015*** | ***$50*** | Review

JENKINS

Lippincott Williams & Wilkins, 2014, 528 pages, ISBN 9781469894690

An organ system–based review text with clinical vignettes that is useful for integrating the basic sciences covered on Step 1. Composed primarily of outlines, charts, tables, and diagrams. Limited scope of material covered. Includes access to a sample online question bank.

B+ | ***USMLE Step 1 Lecture Notes 2018*** | ***$330*** | Review

KAPLAN

Kaplan Medical, 2018, ~2700 pages, ISBN 9781506221229

Extremely comprehensive review of Step 1 topics through videos and lecture notes. Split into individual sections covering pathology, pharmacology, physiology, biochemistry and medical genetics, immunology and microbiology. Generally best used to fill gaps in understanding and to review unfamiliar topics that one has not come across, and therefore the notes are commonly used by foreign medical graduates. Some sections are quite detailed and go beyond the scope of the Step 1 exam.

B+ | ***USMLE Images for the Boards: A Comprehensive Image-Based Review*** | ***$42*** | Review

TULLY

Elsevier, 2012, 296 pages, ISBN 9781455709038

Contains more than 300 color images of content likely to be tested on the Step 1. Contains a wide variety of images including ECGs and radiographic studies. Some images may be low yield for boards studying, but still excellent as a supplement to preclinical courses.

B | ***USMLE Step 1 Made Ridiculously Simple*** | ***$30*** | Review/Test 1000 q

CARL

MedMaster, 2017, 416 pages, ISBN 9781935660224

Concise, succinct text. Online access to more than 1000 practice questions. Uses a table and chart format organized by subject, but some charts are poorly labeled. Consider as an adjunct to more comprehensive sources.

B | ***medEssentials for the USMLE Step 1*** | ***$55*** | Review

MANLEY

Kaplan, 2012, 588 pages, ISBN 9781609780265

A comprehensive review divided into general principles and organ systems, organized using high-yield tables and figures. Helpful for visual learners, but can be overly detailed and time consuming. Includes color images in the back along with a monthly subscription to online interactive exercises, although these are of limited value for Step 1 preparation. Comes with a free mobile version.

(c) ketabton.com: The Digital Library

▸ ANATOMY, EMBRYOLOGY, AND NEUROSCIENCE

A– ***High-Yield Gross Anatomy*** — ***$43*** — Review
DUDEK
Lippincott Williams & Wilkins, 2014, 320 pages, ISBN 9781451190236

A good review of gross anatomy with some clinical correlations. Contains color clinical photos and well-labeled, high-yield radiographic images, but often goes into excessive detail that is beyond the scope of the boards.

A– ***Clinical Anatomy Made Ridiculously Simple*** — ***$30*** — Review
GOLDBERG
MedMaster, 2016, 175 pages, ISBN 9780940780972

An easy-to-read text offering simple diagrams along with numerous mnemonics, helpful charts, and amusing associations. The humorous style has variable appeal to students, so browse the content before purchasing. Offers good coverage of selected topics. Includes a CD-ROM atlas of normal radiographic anatomy. Best if used during coursework. Includes more detail than typically tested on Step 1.

B+ ***High-Yield Embryology*** — ***$56*** — Review
DUDEK
Lippincott Williams & Wilkins, 2013, 176 pages, ISBN 9781451176100

A concise review of a relatively less-tested subject. Offers excellent organization with clinical correlations. Includes a high-yield list of embryologic tissue origins and USMLE-style case studies after each chapter. May not be suitable for dedicated Step 1 studying.

B+ ***High-Yield Neuroanatomy*** — ***$40*** — Review/Test/50 q
FIX
Lippincott Williams & Wilkins, 2015, 208 pages, ISBN 9781451193435

An easy-to-read, straightforward format with excellent diagrams and illustrations. Features a useful atlas of brain and spinal cord images, a glossary of important terms, and an appendix of neurologic lesions. Overall, a great resource and quick read, but more detailed than what is required for Step 1.

B+ ***Anatomy—An Essential Textbook*** — ***$48*** — Text/Test/400 q
GILROY
Thieme, 2017, 528 pages, ISBN 9781626234390

A thorough, visually appealing approach to learning anatomy. Contains over 650 colorful, helpful illustrations. Presents material in bullet-point format and tables. Includes over 160 clinical correlates and 400 USMLE-style questions, with the opportunity to complete practice questions online.

B+ ***Netter's Anatomy Flash Cards*** — ***$40*** — Flash cards
HANSEN
Saunders, 2018, 688 flash cards, ISBN 9780323530507

Netter's illustrations in a question/answer column format that allows for self-testing. Each card includes commentary on the structures with a clinical correlation. More effective as a supplement to coursework, and much too detailed for boards preparation. Lack of embryology correlates limits Step 1 usefulness. Includes online access with additional bonus cards and more than 400 multiple choice questions. Note: an iOS app has a similar cost and additional functionality.

(c) ketabton.com: The Digital Library

B+ | ***Crash Course: Anatomy*** | ***$45*** | Review

STENHOUSE

Elsevier, 2015, 288 pages, ISBN 9780723438540

Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1 review questions. Contains an up-to-date self-assessment section. Provides a solid review of anatomy for Step 1. Best if started early.

B | ***BRS Embryology*** | ***$56*** | Review/Test/220 q

DUDEK

Lippincott Williams & Wilkins, 2014, 336 pages, ISBN 9781451190380

An outline-based review of embryology that is typical of the BRS series. Offers a good review and includes much more detail than is required for Step 1. A discussion of congenital malformations is included at the end of each chapter, along with over 220 USMLE-style questions with answers and explanations. The comprehensive exam at the end of the book is high yield. Includes access to a searchable online text on the free companion website, which also features interactive quizzing.

B | ***Anatomy Flash Cards: Anatomy on the Go*** | ***$60*** | Flash cards

GILROY

Thieme, 2013, 752 flash cards, ISBN 9781604069105

Flash card deck containing high-quality illustrations and a question/answer format that allows for self-testing. Occasional radiographic image. Best if used with coursework; too long for efficient boards preparation.

B | ***Clinical Neuroanatomy Made Ridiculously Simple*** | ***$26*** | Review/Test/Few q

GOLDBERG

MedMaster, 2014, 90 pages + CD-ROM, ISBN 9781935660194

An easy-to-read, memorable, and simplified format with clever diagrams. Offers a quick, high-yield review of clinical neuroanatomy, but not a comprehensive resource for boards review. Places appropriate emphasis on clinically relevant pathways, cranial nerves, and neurologic diseases. Includes a CD-ROM with CT and MR images, a tutorial on neurologic localization, and interactive quizzes covering classic neurology cases.

B | ***Netter's Anatomy Coloring Book*** | ***$20*** | Review

HANSEN

Elsevier, 2018, 392 pages, ISBN 9780323545037

An easy-to-understand, detailed, interactive book that is an excellent companion to traditional textbooks during preclinical anatomy coursework. Provides multiple views and magnifications of anatomic structures as well as dissection layers. The coloring aspect of the book can be highly beneficial for visual learners. Contains few clinical correlations, which limits its usefulness during dedicated studying for Step 1.

B | ***Case Files: Anatomy*** | ***$35*** | Cases

TOY

McGraw-Hill, 2014, 416 pages, ISBN 9780071794862

Review text that includes 58 well-chosen cases with discussion, comprehension questions, and take-home pearls. Tables are helpful, but schematics are black and white and not representative of Step 1. A reasonable book to work through for those who benefit from problem-based learning.

(c) ketabton.com: The Digital Library

B–

Case Files: Neuroscience

$35 Cases

TOY

McGraw-Hill, 2014, 432 pages, ISBN 9780071790253

Similar to other *Case Files* books, it includes 49 clinical cases with lengthy discussion and 3–5 multiple choice questions at the end of each case. Cases are helpful, but the discussion is too lengthy. Questions are not the most representative of those seen on boards. Limited student feedback.

▶ BEHAVIORAL SCIENCE

A–

BRS Behavioral Science

$52 Review/ Test/700 q

FADEM

Lippincott Williams & Wilkins, 2016, 384 pages, ISBN 9781496310477

An easy-to-read outline-format review of behavioral science. Offers detailed coverage of mostly high-yield topics, but at a level of depth that often exceeds what is tested on Step 1. Better used prior to dedicated study period. Incorporates tables and charts as well as a short but complete statistics chapter. Features over 700 review questions, including a 100-question comprehensive exam. References DSM-V criteria.

B+

High-Yield Biostatistics, Epidemiology, and Public Health

$43 Review

GLASER

Lippincott Williams & Wilkins, 2013, 168 pages, ISBN 9781451130171

A well-written, easy-to-read text that offers extensive coverage of epidemiology and biostatistics. Includes helpful review questions and tables, but somewhat lengthy given the low-yield nature of this subject on Step 1.

(c) ketabton.com: The Digital Library

BIOCHEMISTRY

A– | ***Pixorize*** | ***$100–$130*** | Review

www.pixorize.com

Visual mnemonic system focusing primarily on biochemistry. Step-by-step videos and interactive images aid studying and review. Compare to Sketchy and Picmonic.

B+ | ***Medical Biochemistry—An Illustrated Review*** | ***$40*** | Review/Test/400 q

PANINI

Thieme, 2013, 441 pages, ISBN 9781604063165

A comprehensive medical biochemistry study guide with an emphasis on images. Very detailed and may be better as a supplement to preclinical courses than as a review resource for the Step 1. Images and diagrams are helpful for solidifying knowledge. Online access available for additional content, including 400 USMLE-style practice questions.

B | ***Lange Flash Cards Biochemistry and Genetics*** | ***$40*** | Flash cards

BARON

McGraw-Hill, 2017, 196 flash cards, ISBN 9781259837210

Flash card deck featuring clinical vignettes on one side and concise discussions on the other. Each section contains 2–3 cards on biochemistry principles. High level of detail may make this less ideal for dedicated boards studying. Note that no carrying case for the cards is included.

B | ***Lippincott Illustrated Reviews: Biochemistry*** | ***$78*** | Review/Test/200 q

FERRIER

Lippincott Williams & Wilkins, 2017, 560 pages, ISBN 9781496344496

An integrative and comprehensive review of biochemistry that includes good clinical correlations and effective color diagrams. Extremely detailed and requires significant time commitment, so it should be started with first-year coursework. High-yield summaries at the end of each chapter. Comes with access to the companion website, which includes over 200 USMLE-style questions.

B | ***BRS Biochemistry, Molecular Biology, and Genetics*** | ***$54*** | Review/Test

LIEBERMAN

Lippincott Williams & Wilkins, 2013, 432 pages, ISBN 9781451175363

A highly detailed review featuring many figures and clinical correlations highlighted in colored boxes. The biochemistry portion includes much more detail than required for Step 1, but may be useful for students without a strong biochemistry background or as a reference text. The molecular biology section is more focused and high yield. Also offers a chapter on laboratory techniques and a comprehensive, 120-question exam. Questions are clinically oriented.

B | ***Case Files: Biochemistry*** | ***$35*** | Cases

TOY

McGraw-Hill, 2014, 480 pages, ISBN 9780071794886

Includes 51 clinical cases with comprehensive discussion and summary box, albeit too much depth and not enough breadth for boards preparation. Some cases will almost certainly *not* be tested. Questions at the end of each case are not representative of those on Step 1.

(c) ketabton.com: The Digital Library

B

PreTest Biochemistry and Genetics

$38 Test/500 q

WILSON

McGraw-Hill, 2017, 592 pages, ISBN 9780071791441

500 questions with detailed, well-referenced explanations. Features a high-yield introduction and appendix, but may be overly detailed in some cases. A solid supplement to preclinical courses and board studying.

▸ CELL BIOLOGY AND HISTOLOGY

B+

BRS Cell Biology and Histology

$54 Review/ Test/320 q

GARTNER

Lippincott Williams & Wilkins, 2018, 448 pages, ISBN 9781496396358

Covers concepts in cell biology and histology in an outline format. Can be used alone for cell biology study, but may have fewer histology images than some other resources. Includes more detail than is required for Step 1, and information is less high yield than that of other books in the BRS series. Interactive quizzes on the free companion website provide additional practice.

B+

Crash Course: Cell Biology and Genetics

$47 Review/Print + online

STUBBS

Elsevier, 2015, 216 pages, ISBN 9780723438762

Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1 review questions. Includes online access. High level of detail makes this resource best suited for coursework.

B

Wheater's Functional Histology

$83 Text

YOUNG

Elsevier, 2013, 464 pages, ISBN 9780702047473

A color atlas with more than 900 high-quality illustrations of normal histology with image captions and accompanying text. Far too detailed to use for boards studying given the low-yield nature of the material, but useful as a coursework text or boards reference. Provides online access to the entire atlas and USMLE-style self-assessment questions.

(c) ketabton.com: The Digital Library

MICROBIOLOGY AND IMMUNOLOGY

A– ***Basic Immunology*** — ***$70*** — Review

ABBAS

Elsevier, 2019, 336 pages, ISBN 9780323549431

A useful text that offers clear explanations of complex topics in immunology. Best if used in conjunction with coursework and later skimmed for quick Step 1 review. Includes colorful diagrams, images, tables, and a glossary for further study. Features online access.

A– ***Clinical Microbiology Made Ridiculously Simple*** — ***$38*** — Review

GLADWIN

MedMaster, 2019, 418 pages, ISBN 9781935660330

An excellent, easy-to-read, detailed review of microbiology that includes clever and memorable mnemonics. The sections on bacterial disease are most high yield, less emphasis placed on pharmacology. Recommended to read during coursework and review the concise charts at the end of each chapter during boards review. All images are cartoons; no microscopy images that appear on boards. Requires a supplemental source for immunology.

A– ***Medical Microbiology and Immunology Flash Cards*** — ***$40*** — Flash cards

ROSENTHAL

Elsevier, 2016, 192 flash cards, ISBN 9780323462242

Flash cards covering the microorganisms most commonly tested on Step 1. Each card features color microscopic images and clinical presentations on one side and relevant bug information in conjunction with a short case on the other side. Also includes Student Consult online access for extra features. Overemphasizes "trigger words" related to each bug. Not a comprehensive resource.

B+ ***Lippincott Illustrated Reviews: Immunology*** — ***$75*** — Reference/Test/Few q

DOAN

Lippincott Williams & Wilkins, 2012, 384 pages, ISBN 9781451109375

A clearly written, highly detailed review of basic concepts in immunology. Features many useful tables and review questions at the end of each chapter. More than 300 color annotated illustrations. Offers abbreviated coverage of immunodeficiencies and autoimmune disorders. Best if started with initial coursework and used as a reference during Step 1 study.

B+ ***Microcards: Microbiology Flash Cards*** — ***$53*** — Flash cards

HARPAVAT

Lippincott Williams & Wilkins, 2015, 312 flash cards, ISBN 9781451192353

A well-organized and complete resource for students who like to use flash cards for review. Cards feature the clinical presentation, pathobiology, diagnosis, treatment, and high-yield facts for a particular organism. Some cards also include excellent flow charts organizing important classes of bacteria or viruses. Overall, a good review resource, but at times it is overly detailed, requiring a significant time commitment. Also useful as an aid with coursework. Includes access to online USMLE-style questions with answers.

(c) ketabton.com: The Digital Library

B+ ***Review of Medical Microbiology and Immunology*** ***$63*** Review/ Test/654 q

LEVINSON

McGraw-Hill, 2018, 832 pages, ISBN 9781259644498

A clear, comprehensive text with outstanding diagrams and tables. Includes an excellent immunology section. Contains a chapter summarizing details on medically important organisms. Can be used as reference for reviewing immunology concepts. Can be detailed and dense at points, so best if started early with coursework. Includes practice questions of mixed quality and does not provide detailed explanation of answers. Compare with *Lippincott Illustrated Reviews: Microbiology*.

B+ ***How the Immune System Works*** ***$50*** Review

SOMPAYRAC

Wiley-Blackwell, 2019, 168 pages, ISBN 9781119542124

A short overview of high-yield immunology designed for those with no prior immunology knowledge. Analogies and images create a "storybook" feel to spruce up a relatively dry subject. The 15 chapters offer a general overview with good supporting details.

B ***Case Studies in Immunology: Clinical Companion*** ***$62*** Cases

GEHA

W. W. Norton & Company, 2016, 384 pages, ISBN 9780815345121

A text that was originally designed as a clinical companion to *Janeway's Immunobiology*. Provides a great synopsis of the major disorders of immunity in a clinical vignette format. Integrates basic and clinical sciences. Features excellent images and illustrations from Janeway, as well as questions and discussions.

B ***Pretest: Microbiology*** ***$38*** Test/500 q

KETTERING

McGraw-Hill, 2013, 480 pages, ISBN 9780071791045

Includes a short section on high-yield facts followed by 500 questions in a clinical vignette format. Questions are more difficult than encountered on the boards and some topics discussed are not likely to be tested. A good book to work through with coursework but too low yield for review purposes.

B ***Case Files: Microbiology*** ***$36*** Cases

TOY

McGraw-Hill, 2014, 416 pages, ISBN 9780071820233

Provides 54 clinical microbiology cases followed by a clinical correlation, a discussion with boldfaced buzzwords, and questions. Cases are well chosen, but the text lacks the high-yield charts and tables found in other books in the Case Files series. Images are sparse, black and white, and of poor quality.

B ***Lange Microbiology and Infectious Diseases Flash Cards, 3e*** ***$46*** Flash cards

Somers, 2017

Clinical vignettes presented on one side of the card as a mini-case study of the disease and the flip side presents the etiology and epidemiology, pathogenesis, clinical manifestations, laboratory diagnosis, and treatment and prevention of the disorder. Good for reviewing clinical aspects of many infectious diseases including those caused by bacteria, viruses, and fungi.

(c) ketabton.com: The Digital Library

B– ***Lippincott Illustrated Reviews: Microbiology*** — ***$73*** — Review/Test/Few q

CORNELISSEN

Lippincott Williams & Wilkins, 2019, 448 pages, ISBN 9781496395856

A comprehensive, highly illustrated review of microbiology that is similar in style to other titles in the Illustrated Reviews series. Has more than 400 color illustrations and color-coded summaries to help visual learners. Contains several hundred USMLE-style review questions to help with exam preparation. Compare with Levinson's *Review of Medical Microbiology and Immunology*.

PATHOLOGY

A+ ***Pathoma: Fundamentals of Pathology*** — ***$85–$120*** — Review/Lecture

SATTAR

Pathoma, 2019, 218 pages, ISBN 9780983224631

Integrated approach to pathology review, combining a focused textbook with 35+ hours of online lectures. Book contains more than 350 color images. Videos combine "chalk talk" and slide formats to explain pathogenesis in an easy-to-understand manner. Online subscription needed for full access.

A– ***Rapid Review: Pathology*** — ***$65*** — Review/Test/500 q

GOLJAN

Elsevier, 2018, 864 pages, ISBN 9780323476683

A comprehensive source for key concepts in pathology, presented in a bulleted outline format with many high-yield tables and color figures. Features detailed explanations of disease mechanisms. Integrates concepts across disciplines with a strong clinical orientation. Lengthy, so best if started early with coursework. Includes access to online question bank with more than 500 questions. Covers material for both Step 1 and Step 2 exams.

A– ***Robbins and Cotran Review of Pathology*** — ***$55*** — Test/1100 q

KLATT

Elsevier, 2014, 504 pages, ISBN 9781455751556

A question book that follows the main Robbins textbooks. Questions are more detailed, difficult, and arcane than those on the actual Step 1 exam, but the text offers a great review of pathology integrated with more than 1100 images. Thorough answer explanations reinforce key points. Requires significant time commitment, so best if started with coursework. 2014 edition table of contents closely follows the organization of *Robbins and Cotran Pathologic Basis of Disease*, 8th edition.

A– ***Crash Course: Pathology*** — ***$40*** — Review

XIU

Elsevier, 2019, 438 pages, ISBN 9780702073540

Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1 review questions. Includes online access. Best if started during coursework.

(c) ketabton.com: The Digital Library

B

High-Yield Histopathology

$36 Review

DUDEK

Lippincott Williams & Wilkins, 2017, 320 pages, ISBN 9781496353344

Reviews the relationship of basic histology to the pathology, physiology, and pharmacology of clinical conditions that are tested on Step 1. Includes case studies, numerous light and electron micrographs, and pathology photographs. Given its considerable length, should be started with coursework or used as a reference to better identify images.

B

Pathophysiology of Disease: Introduction to Clinical Medicine

$90 Text

HAMMER

McGraw-Hill, 2018, 832 pages, ISBN 9781260026504

An interdisciplinary text useful for understanding the pathophysiology of clinical symptoms. Effectively integrates the basic sciences with mechanisms of disease. Features great graphs, diagrams, and tables. In view of its length, most useful if started during coursework. Includes 120 case studies, checkpoint questions that appear in every chapter, and a few non–boards-style questions. The text's clinical emphasis nicely complements *BRS Pathology*.

B

Haematology at a Glance

$49 Review

MEHTA

Blackwell Science, 2014, 136 pages, ISBN 9781119969228

A resource that covers common hematologic issues. Includes color illustrations. Presented in a logical sequence that is easy to read. Good for use with coursework.

B

Pocket Companion to Robbins and Cotran Pathologic Basis of Disease

$40 Review

MITCHELL

Elsevier, 2016, 896 pages, ISBN 9781455754168

A condensed version of *Robbins and Cotran Pathologic Basis of Disease* that is good for reviewing keywords associated with most important diseases. Presented in a highly condensed format, but the text is complete and easy to understand. Contains no photographs or illustrations but does include tables. Useful as a quick reference.

B

BRS Pathology

$54 Review/ Test/450 q

SCHNEIDER

Lippincott Williams & Wilkins, 2013, 480 pages, ISBN 9781451115871

An excellent, concise review with appropriate content emphasis. Chapters are organized by organ system and feature an outline format with boldfacing of key facts. Includes good questions with explanations at the end of each chapter plus a comprehensive exam at the end of the book. Offers well-organized tables and diagrams as well as photographs representative of classic pathology. Contains a chapter on lab testing and "key associations" with each disease. Contains excellent color images and access to an online test and interactive question bank. Most effective if started early in conjunction with coursework, as it does not discuss detailed mechanisms of disease pathology.

(c) ketabton.com: The Digital Library

PHARMACOLOGY

B+ | ***Crash Course: Pharmacology*** | ***$40*** | Review

BATTISTA

Elsevier, 2019, 336 pages, ISBN 9780702073441

Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1–style review questions with a self-assessment section. Includes online access. Gives a solid, easy-to-follow overview of pharmacology.

B+ | ***Master the Boards USMLE Step 1 Pharmacology Flashcards*** | ***$55*** | Flash cards

FISCHER

Kaplan, 2015, 200 flash cards, ISBN 9781618657947

Easy-to-read flash cards with drug and questions on one side and discussion on the other. Useful for a quick pharmacology review. Some drugs/material may be beyond the scope of the Step 1, or more appropriate at the Step 2 level.

B+ | ***BRS Pharmacology*** | ***$55*** | Review/ Test/200 q

ROSENFELD

Lippincott Williams & Wilkins, 2019, 384 pages, ISBN 9781975105495

Features two-color tables and figures that summarize essential information for quick recall. A list of drugs organized by drug family is included in each chapter. Too detailed for boards review; best used as a reference. Also offers end-of-chapter review tests with Step 1–style questions and a comprehensive exam with explanations of answers. An additional question bank is available online.

B | ***Lange Pharmacology Flash Cards*** | ***$39*** | Flash cards

BARON

McGraw-Hill, 2017, 266 flash cards, ISBN 9781259837241

A total of 230 pocket-sized flash cards of relevant drugs formatted with clinical vignettes on one side and relevant information on the other side (eg, mode of action, adverse effects, clinical uses). Particularly high-yield information is highlighted in bold. Mainly useful as a supplement for pharmacology knowledge, rather than as a primary resource. Printed on less durable material.

B | ***Pharmacology Flash Cards*** | ***$45*** | Flash cards

BRENNER

Elsevier, 2017, 230 flash cards, ISBN 9780323355643

Flash cards for more than 200 of the most commonly tested drugs. Cards include the name of the drug (both generic and brand) on the front and basic drug information on the back, with occasional cards covering high-yield pharmacology pathways. Divided and color coded by class, and comes with a compact carrying case. Lacks figures and clinical vignettes.

(c) ketabton.com: The Digital Library

B

Katzung & Trevor's Pharmacology: Examination and Board Review

$54 Review/Test/800 q

TREVOR

McGraw-Hill, 2018, 592 pages, ISBN 9781259641022

A well-organized text with concise explanations. Features good charts and tables; the crammable list in Appendix I is especially high yield for Step 1 review. Also good for reviewing drug interactions and toxicities. Offers two 100-question practice exams. Text includes many low-yield/obscure drugs. Compare with *Lippincott Illustrated Reviews: Pharmacology*, both of which are better suited to complementing coursework than last-minute studying for boards.

B

Lippincott Illustrated Reviews: Pharmacology

$75 Review/Test/380 q

WHALEN

Lippincott Williams & Wilkins, 2018, 576 pages, ISBN 9781496384133

A resource presented in outline format with practice questions, many excellent illustrations, and comparison tables. Effectively integrates pharmacology and pathophysiology. Best started alongside coursework, as it is highly detailed and requires significant time commitment. Focuses on basic principles.

▸ PHYSIOLOGY

A–

BRS Physiology

$54 Review/Test/350 q

COSTANZO

Lippincott Williams & Wilkins, 2018, 304 pages, ISBN 9781496367617

A clear, concise review of physiology that is both comprehensive and efficient, making for fast, easy reading. Includes excellent high-yield charts and tables, but lacks some figures from Costanzo's *Physiology*. Features high-quality practice questions with explanations in each chapter along with a clinically oriented final exam. An excellent reference during times of focused Step 1 studying, but best if started early in combination with coursework. Respiratory and acid-base sections are comparatively weak.

A–

Pathophysiology of Heart Disease

$57 Review

LILLY

Lippincott Williams & Williams, 2015, 480 pages, ISBN 9781451192759

Great resource that outlines an in-depth explanation of both cardiac physiology and pathology. Best used as a supplement when learning the material for the first time, as it helps build a strong foundation. Because the book itself is rather dense, it is not recommend as a primary resource during focused boards studying period.

A–

PreTest Physiology

$38 Test/500 q

METTING

McGraw-Hill, 2013, 528 pages, ISBN 9780071791427

Contains questions with detailed, well-written explanations. One of the best of the PreTest series. Best for use by the motivated student after extensive review of other sources. Includes a high-yield facts section with useful diagrams and tables.

(c) ketabton.com: The Digital Library

A– | ***Color Atlas of Physiology*** | ***$50*** | Review

SILBERNAGL

Thieme, 2015, 472 pages, ISBN 9783135450070

Contains more than 180 high-quality illustrations of disturbed physiologic processes that lead to dysfunction. An alternative to standard texts, but not high yield for boards review.

B+ | ***BRS Physiology Cases and Problems*** | ***$58*** | Cases

COSTANZO

Lippincott Williams & Wilkins, 2012, 368 pages, ISBN 9781451120615

Presents 62 classic cases in vignette format with several questions per case. Includes exceptionally detailed explanation of answers along with supplemental diagrams. For students interested in an in-depth discussion of physiology concepts.

B+ | ***Physiology*** | ***$60*** | Text

COSTANZO

Saunders, 2017, 528 pages, ISBN 9780323478816

A comprehensive, clearly written text that covers concepts outlined in *BRS Physiology* in greater detail. Offers excellent color diagrams and charts. Each systems-based chapter features a detailed summary of objectives and a Step 1–relevant clinical case. Includes access to online interactive extras. Requires time commitment, but helps develop a strong foundation in physiology concepts. Best if started alongside coursework. Practice questions at end of each chapter.

B+ | ***Vander's Renal Physiology*** | ***$49*** | Text

EATON

McGraw-Hill, 2018, 224 pages, ISBN 9781260019377

Well-written text on renal physiology, with helpful but sparse diagrams and practice questions at the end of each chapter. Too detailed for Step 1 review, however. Best if used with organ-based coursework to understand the principles of renal physiology.

B+ | ***Acid-Base, Fluids, and Electrolytes Made Ridiculously Simple*** | ***$24*** | Review

PRESTON

MedMaster, 2017, 166 pages, ISBN 9781935660293

A resource that covers major acid-base and renal physiology concepts. Provides information beyond the scope of Step 1, but remains a useful companion for studying kidney function, electrolyte disturbances, and fluid management. Includes scattered diagrams and questions at the end of each chapter. Consider using after exhausting more high-yield physiology review resources.

B+ | ***Pulmonary Pathophysiology: The Essentials*** | ***$57*** | Review/ Test/75 q

WEST

Lippincott Williams & Wilkins, 2017, 264 pages, ISBN 9781496339447

A volume offering comprehensive coverage of respiratory physiology. Clearly organized with useful charts and diagrams. Review questions at the end of each chapter provide answers but no explanations. Best used as a course supplement during the second year, less ideal for use immediately prior to Step 1.

(c) ketabton.com: The Digital Library

B | ***Rapid Review: Physiology*** | ***$39*** | Test/350 q

BROWN

Elsevier, 2011, 384 pages, ISBN 9780323072601

Offers a good review of physiology in a format typical of the Rapid Review series, albeit with more images. Includes online access to 350 questions with concise explanations, along with other extras. Compare with Robbins *Physiology*.

B | ***Endocrine Physiology*** | ***$59*** | Review

MOLINA

McGraw-Hill, 2018, 320 pages, ISBN 9781260019353

Questions at the end of each chapter are helpful solidify knowledge, but some are not representative of Step 1 questions. Provides more detailed explanations of endocrine physiology than Costanzo review offers, but much too lengthy for Step 1 review. May be useful as a coursework adjunct.

B– | ***Netter's Physiology Flash Cards*** | ***$40*** | Flash cards

MULRONEY

Saunders, 2015, 450 flash cards, ISBN 9780323359542

Flash cards contain a high-quality illustration on one side with question and commentary on the other. Good for self-testing, but too fragmented for learning purposes and not comprehensive enough for boards.

Get more e-books from www.ketabton.com
Ketabton.com: The Digital Library